

A final word on FCNC-Baryogenesis from two Higgs doublets

Wei-Shu Hou¹, Tanmoy Modak^{2*} and Tilman Plehn²

¹ Department of Physics, National Taiwan University, Taipei, Taiwan

² Institut für Theoretische Physik, Universität Heidelberg, Germany

* modak@thphys.uni-heidelberg.de

Abstract

Electroweak baryogenesis in a two-Higgs doublet model is a well-motivated and testable scenario for physics beyond the Standard Model. An attractive way of providing CP violation is through flavor-changing Higgs couplings, where the top-charm coupling is hardly constrained. This minimal scenario can be tested by searching for heavy charged and neutral Higgs bosons at the LHC. While the charged Higgs signature requires a dedicated analysis, the neutral Higgs signature will be covered by a general search for same-sign top pairs. Together, they provide a conclusive test of this kind of baryogenesis.



Copyright W. Hou *et al.*

This work is licensed under the Creative Commons

[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Published by the SciPost Foundation.

Received 16-12-2020

Accepted 16-06-2021

Published 21-06-2021

doi:[10.21468/SciPostPhys.10.6.150](https://doi.org/10.21468/SciPostPhys.10.6.150)



Check for updates

Contents

1	Introduction	1
2	Model and parameter space	3
3	Charged Higgs production	5
4	Neutral Higgs production	7
5	Outlook	10
	References	11

1 Introduction

The Higgs discovery [1,2] and subsequent measurements of the Higgs Lagrangian at Run 2 [3–11] indicate that the Standard Model is the correct effective theory around the electroweak

scale. While there exists no experimental evidence for physics beyond the Standard Model so far, extended Higgs sectors are motivated by theoretical considerations, like mass generation of up-type and down-type fermions, neutrino mass generation, electroweak baryogenesis, or dark matter. In particular, two-Higgs doublet models (2HDMs) [12–14] are an integral part of well-defined models for physics beyond the Standard Model, including MSSM [15], composite Higgs models [16], little Higgs models [17, 18], or GUTs [19–22].

If we use baryogenesis [23] as a guiding principle to new physics searches at the LHC, a 2HDM is an attractive and minimal choice. It can provide both, new scalar degrees of freedom [24, 25] and CP -violation. In the general [26, 27] or type-III [28] 2HDM, the new particles can be close in mass to the SM-Higgs [29, 30]. Sufficiently large CP -violation is non-trivial to achieve, in our model we rely on the Yukawa sector. If both doublets couple to up-type and down-type quarks, they define two separate Yukawa matrices. After diagonalizing the quark mass matrices we find the real, diagonal couplings $\lambda_{ii} = \sqrt{2}m_i/v$ and the complex, non-diagonal couplings ρ_{ij} . While flavor-changing neutral couplings are generally well-constrained, it is possible to have electroweak baryogenesis (EWBG) driven by a single, order-one, complex coupling ρ_{tc} [31],

$$\text{Im } \rho_{tc} \gtrsim 0.5 \quad \text{and} \quad |\cos \gamma| \gtrsim 0.1, \quad (1)$$

where γ is the mixing angle between the two CP -even Higgs states.

It has been shown [32–36] that the coupling ρ_{tc} can be discovered in the LHC process

$$cg \rightarrow tA/H \rightarrow t(t\bar{c}), \quad (2)$$

where this process retains a very mild dependence on $\cos \gamma$ and is especially useful for small values of $\cos \gamma$. To attribute this signal to EWBG requires information on the mixing angle $\cos \gamma$, for instance through b -associated charged Higgs production [37]

$$cg \rightarrow bH^+ \rightarrow b(W^+h). \quad (3)$$

Here the production process is induced by ρ_{tc} [38, 39], while the decay amplitude is proportional to the mixing angle $\cos \gamma$. Even in the absence of complex phase information, such a search can test the required particle content and parameter space for the ρ_{tc} -EWBG scenario. Finally, the exotic top decay

$$t \rightarrow ch \quad (4)$$

is induced by the coupling ρ_{tc} combined with non-vanishing $\cos \gamma$ [28] and is searched for by CMS [40] and ATLAS [41].

In this paper we show how the two LHC searches for charged and neutral heavy Higgs bosons can conclusively probe the parameter region required for ρ_{tc} -EWBG in the general 2HDM (g2HDM). The paper is organized as follows: in Sec. 2 we discuss the model and its preferred parameter space, and then compare it to the reach of the charged Higgs channel in Sec. 3. Section 4 is dedicated to same-sign top production from neutral Higgs production and its complementarity to the charged Higgs signature. We summarize our results in Sec. 5.

2 Model and parameter space

The general CP -conserving two Higgs doublet potential can be written as [42, 43]

$$V(\Phi, \Phi') = \mu_{11}^2 |\Phi|^2 + \mu_{22}^2 |\Phi'|^2 - (\mu_{12}^2 \Phi^\dagger \Phi' + \text{h.c.}) + \frac{\eta_1}{2} |\Phi|^4 + \frac{\eta_2}{2} |\Phi'|^4 + \eta_3 |\Phi|^2 |\Phi'|^2 + \eta_4 |\Phi^\dagger \Phi'|^2 + \left[\frac{\eta_5}{2} (\Phi^\dagger \Phi')^2 + (\eta_6 |\Phi|^2 + \eta_7 |\Phi'|^2) \Phi^\dagger \Phi' + \text{h.c.} \right]. \quad (5)$$

In the Higgs basis, the VEV $v = 246$ GeV is generated by the doublet Φ , while Φ' does not develop a VEV, hence $\mu_{22}^2 > 0$. The minimization conditions in the two field directions lead to $\mu_{11}^2 = -\eta_1 v^2/2$ and $\mu_{12}^2 = \eta_6 v^2/2$. The mixing angle γ diagonalizes the CP -even mass matrix to define the mass eigenstates h and H [42, 43],

$$c_\gamma^2 = \cos^2 \gamma = \frac{\eta_1 v^2 - m_h^2}{m_H^2 - m_h^2} \quad \text{and} \quad s_{2\gamma} = \sin(2\gamma) = \frac{2\eta_6 v^2}{m_H^2 - m_h^2}, \quad (6)$$

where $c_\gamma \rightarrow 0$ in the alignment limit ($\eta_6 = 0$ and $\eta_1 = m_h^2/v^2 \sim 1/4$). To satisfy the first Sakharov condition [23], a new scalar degree of freedom close in mass with the SM-Higgs [44–46] can trigger a strong first-order phase transition. Following Eq.(6) this is guaranteed by finite c_γ and perturbatively stable $\eta_i = \mathcal{O}(1)$, for instance $\eta_6 = \mathcal{O}(1)$ and $\eta_1 = \mathcal{O}(1) > m_h^2/v^2$ [43].

Next, baryogenesis requires a complex phase in the Higgs or Yukawa sectors [23]. Many analyses have studied a complex Higgs potential, which tends to be strongly constrained by EDM measurements [47–50]. We look at the alternative option of CP -violation arising from the Yukawa sector [31, 42, 51]

$$\mathcal{L} \supset -\frac{1}{\sqrt{2}} \sum_{F=U,D,L} \bar{F}_i \left[(-\lambda_{ij}^F s_\gamma + \rho_{ij}^F c_\gamma) h + (\lambda_{ij}^F c_\gamma + \rho_{ij}^F s_\gamma) H - i \operatorname{sgn}(Q_F) \rho_{ij}^F A \right] P_R F_j - \bar{U}_i [(V \rho^D)_{ij} P_R - (\rho^{U\dagger} V)_{ij} P_L] D_j H^+ - \bar{\nu}_i \rho_{ij}^L P_R L_j H^+ + \text{h.c.}, \quad (7)$$

where $i, j = 1, 2, 3$ are generation indices, $P_{L,R} \equiv (1 \mp \gamma_5)/2$, and V is the CKM matrix. In flavor space, the fermion fields F are defined as $U = (u, c, t)$, $D = (d, s, b)$, $L = (e, \mu, \tau)$ and $\nu = (\nu_e, \nu_\mu, \nu_\tau)$. While the mass matrices are diagonalized as in the Standard Model, one cannot rotate away CP -violating phases of the second set of ρ^F matrices in the general 2HDM even in the Higgs basis. That is, the two coupling matrices are

$$\lambda_{ij}^F = \sqrt{2} \frac{m_i^F}{v} \delta_{ij} \in \mathbb{R} \quad \text{and} \quad \rho_{ij}^F \in \mathbb{C}. \quad (8)$$

The complex coupling matrices ρ^F are, strictly speaking, not related to the fermion masses. On the other hand, given experimental constraints and a possible order-of-magnitude correspondence in the values of ρ^F and λ^F lead us to consider ρ_{ij}^U or ρ_{tt}^U .

In principle, a complex ρ_{tt} can robustly drive EWBG [31], which motivates search for channels like $gg \rightarrow H \rightarrow t\bar{t}$ or $gg \rightarrow Ht\bar{t} \rightarrow 4t$ [52–54]. In this paper we focus instead on complex off-diagonal entries ρ_{tj} , specifically ρ_{tc} . With a large phase, this FCNC coupling can also account for the observed baryon asymmetry [31]. One of its merits is that ρ_{tc} does not generate an electron EDM through the Barr-Zee [55] two-loop mechanism, and can therefore more easily [56] evade the ACME bound [57] $d_e < 1.1 \times 10^{-29}$ e cm. Moreover, if we assume ρ_{ct} to be small, the constraint on ρ_{tc} from the charm chromo-EDM also vanishes [58]. We therefore define our specific baryogenesis scenario as [31]

$$|\rho_{tc}| \gtrsim 0.5 \quad \text{and} \quad |c_\gamma| \gtrsim 0.1, \quad (9)$$




Figure 1: Parameter space allowed by perturbativity, positivity, unitarity, and electroweak precision measurements in the m_A - m_H and m_A - m_{H^\pm} planes. The purple and orange scanned points corresponds to $c_\gamma = 0.1$ and 0.3 respectively.

with a sufficiently large complex phase. A strong first-order phase transition is then possible for [59–68]

$$m_{A,H,H^\pm} \sim 300 \dots 600 \text{ GeV}. \tag{10}$$

This mass range is allowed by perturbativity, positivity, unitarity, and electroweak precision data. We rely on 2HDMC [69] to provide the results of Fig. 1 for $c_\gamma = 0.1$ and 0.3 . The 2HDMC parameters in the Higgs basis are $\eta_{1,\dots,7}$ and m_{H^\pm} . To save computing time we actually scan $\mu_{22} \in [0, 1000]$ GeV, $m_A \in [300, 600]$ GeV, $m_H \in [300, 600]$ GeV, $m_{H^\pm} \in [300, 600]$ GeV, $\eta_2 \in [0, 6]$, and $\eta_7 \in [-6, 6]$, and express them in terms of the η_i . To match the 2HDMC conventions we define $\gamma \in [-\pi/2, \pi/2]$. We refer readers to Refs. [39, 51, 70–72] for further details on the parameter scan.

As the first constraint on c_γ and the set of ρ_{ij} we consider measurements of the SM-like Higgs. Higgs coupling measurements constrain the Higgs mixing angle to $c_\gamma \leq 0.3$ and 95%CL. Our choice of ρ_{tc} as the source of CP -violation is motivated by its much weaker constraints, because it hardly affects SM-like Higgs production and decay. The relevant constraints on ρ_{tc} are indirect. For flavor observables, ρ_{tc} enters through loops with charm quarks and a charged Higgs into $B_s - \bar{B}_s$ mixing and $\mathcal{B}(B \rightarrow X_s \gamma)$. Reinterpreting the limit from Ref. [73] we find

$$|\rho_{tc}| \lesssim 1, \quad \text{for } m_{H^\pm} = 300 \text{ GeV}, \tag{11}$$

and its counterpart for $m_{H^\pm} = 500$ GeV is illustrated in Fig. 2, alongside with the EWBG-region. The limit is relatively weak in our general model, in contrast to the type-II 2HDM, and for larger m_{H^\pm} it rapidly becomes irrelevant. Finally, finite c_γ in combination with ρ_{tc} [26] leads to anomalous top decays $t \rightarrow ch$ [28], forbidden at tree level in the SM. The current Run 2 limits at 95%CL are

$$\mathcal{B}(t \rightarrow ch) \approx \frac{c_\gamma^2 |\rho_{tc}|^2}{7.66 + c_\gamma^2 |\rho_{tc}|^2} < \begin{cases} 1.1 \times 10^{-3} & \text{ATLAS [41]} \\ 4.7 \times 10^{-3} & \text{CMS [40]}. \end{cases} \tag{12}$$

They get weaker for smaller c_γ and vanish in the alignment limit. We illustrate the stronger ATLAS [41] constraint also in Fig. 2, along with the projected HL-LHC 95%CL upper limit $\mathcal{B}(t \rightarrow ch) < 1.0 \times 10^{-4}$ [74]. While an observation of this anomalous decay could point to a large value of $|\rho_{tc}|$, if would not provide a link to baryogenesis. A natural step towards solving the baryogenesis puzzle would be to search for new scalar degrees of freedom related to this flavor-changing coupling.




Figure 2: Indirect constraints from $B_s - \bar{B}_s$ mixing (blue), $\mathcal{B}(t \rightarrow ch)$ (purple) in the $\rho_{tc}-c_\gamma$ plane for two H^+ mass values, together with the baryogenesis region (orange).

While we will focus on ρ_{tc} throughout this paper, we point out that ρ_{tu} can be tested using a very similar strategy. For the LHC processes discussed in the coming sections, there is always a corresponding process with an up-quark replacing the charm-quark. One difference between the two FCNC scenarios is that ρ_{tu} can induce observable effects in $\mathcal{B}(B \rightarrow \mu\nu)$ [75], within the reach of Belle-II [76]. The combination of ρ_{tc} and ρ_{tu} is subject to very strong constraints from $D-\bar{D}$ mixing [73], and we will assume only one of the two, but not both at the same time.

3 Charged Higgs production

In the EWBG parameter region of Eq.(9), the partonic process at LHC

$$cg \rightarrow bH^+ \rightarrow b(W_\ell^+ h) \rightarrow b W_\ell^+ W_\ell^+ W_\ell^-, \tag{13}$$

probes ρ_{tc} in H^+ -production and c_γ in the decay $H^+ \rightarrow W^+ h$. The production benefits from the relatively large charm density in the proton, as well as the combination [39] with the CKM matrix element V_{tb} following Eq.(7). The leading-order Feynman diagrams are presented in Fig. 3. While we will require a tagged b -jet, the b -inclusive production process could also be defined as $c\bar{b} \rightarrow H^+$ [77,78]. For a clean analysis, we assume that all three W -bosons decay to either electrons or muons. The same process is induced by ρ_{ct} , but this coupling is constrained to be much smaller [79] by flavor constraints.

Table 1: Charged Higgs properties for the two benchmark points with $\rho_{tc} = 0.35$ and $c_\gamma = 0.25$. The quoted LHC cross sections include the decay $H^+ \rightarrow Wh$ in the fully leptonic mode, as shown in Eq.(13), as well as selection and background rejection cuts.

$m_{H^+}[\text{GeV}]$	$\Gamma_{H^+}[\text{GeV}]$	$\mathcal{B}(H^+ \rightarrow c\bar{b})$	$\mathcal{B}(H^+ \rightarrow W^+ h)$	$\sigma(cg \rightarrow bH^+)[\text{fb}]$
350	2.2	0.85	0.15	0.126
500	3.9	0.66	0.34	0.113




Figure 3: Leading-order Feynman diagrams for the ρ_{tc} -induced $cg \rightarrow bH^+$ and $cg \rightarrow tA/tH$ processes.

The H^+W^-h coupling, modulated by c_γ , arises from [13, 14]

$$\mathcal{L} \supset -\frac{g_2}{2}c_\gamma(h\partial^\mu H^+ - H^+\partial^\mu h)W_\mu^- + \text{h.c.}, \tag{14}$$

where g_2 is the $SU(2)$ gauge coupling. To estimate the reach of our charged Higgs signal, we choose two allowed benchmark points,

$$\rho_{tc} = 0.35, \quad c_\gamma = 0.25, \quad m_{H^+} = 350, 500 \text{ GeV}, \tag{15}$$

as given in Tab. 1. For the branching ratios, we ignore the loop-induced decays $H^+ \rightarrow W^+\gamma$ and $H^+ \rightarrow W^+Z$. We generate signal and background events for $\sqrt{s} = 14$ TeV at leading order with MadGraph5_aMC@NLO [80]. The effective model is implemented in the FeynRules [81] framework, and for parton densities we use NN23LO1 [82]. The events are showered and hadronized with PYTHIA6.4 [83] and then handed to Delphes 3.4.2 [84] for a fast detector simulation with the default ATLAS card. Jets are reconstructed with an $R = 0.6$ anti- k_T algorithm [85] in FastJet [86]. For b -tagging as well as c -jet and light-jet rejections, we also rely on the default ATLAS card. To allow for extra jets we apply MLM matching [87, 88] with the default MadGraph5_aMC run card. The signal is generated with up to two additional jets, do account for higher-order effects in the event kinematics.

The dominant SM-backgrounds are $t\bar{t}W$ and $t\bar{t}Z$ production, followed by WZ + jets, $4t$, $t\bar{t}h$, tZj , tWZ , and ZZ + jets. Furthermore, we find the backgrounds $3t$, $3t+W$, and $3W$ to be negligible, so we ignore them in our analysis. However, given a mis-identification probability for a jet as a lepton around 10^{-4} [89, 90], $t\bar{t}$ production will lead to non-trivial background contributions. For all backgrounds, we use the same simulation chain as for the signal, with up to one additional jet for $t\bar{t}W$, $t\bar{t}Z$, WZ + jets, ZZ + jets, tZ + jets, $t\bar{t}$ + jets, and no QCD jets for the high-multiplicity backgrounds $4t$, tWZ and $t\bar{t}h$. To approximately account for QCD corrections in addition to the jet emission, we attach NLO K -factors to the dominant $t\bar{t}V$ backgrounds, namely 1.35 (W^-), 1.27 (W^+) [91], and 1.56 (Z) [92]. We also correct the WZ + jets and $t\bar{t}$ + jets background normalizations to NNLO by factors 2.07 [93] and 1.84 [94] respectively. Furthermore, we adjust the $4t$, $t\bar{t}h$, and $\bar{t}Z$ + jets rates to NLO through the K -factors 2.04 [80], 1.27 [95] and 1.44 [80]. The cross sections for the signal and tWZ are kept at LO for simplicity. Here, we simply assume the QCD correction factors for the W^+Z + jets and tZ + jets processes to be the same as their respective charge-conjugate processes.

To suppress the backgrounds, we adopt a simple set of requirements. We start with events containing at least three charged leptons and at least one tagged b -jet passing

$$\begin{aligned} p_{T,\ell} &> 20 \text{ GeV}, & |\eta_\ell| &< 2.5, \\ p_{T,b} &> 20 \text{ GeV}, & |\eta_b| &< 2.5, \\ \Delta R_{ij} &> 0.4, & - & (i, j = \ell, b), \\ \cancel{E}_T &> 35 \text{ GeV}, & m_{\ell^+\ell^-} &\notin [76, 110] \text{ GeV}, \quad (\ell = e, \mu). \end{aligned} \tag{16}$$

Table 2: Background cross sections for the charged Higgs process after cuts.

	$t\bar{t}W$	$t\bar{t}\bar{Z}$	$WZ+\text{jets}$	$4t$	$t\bar{t}h$	$tZ+\text{jets}$	tWZ	$ZZ+\text{jets}$	$t\bar{t}+\text{jets}$	sum bkg
merged jets	1	1	1	0	0	1	0	1	1	
K -factor	NLO	NLO	NNLO	NLO	NLO	NLO	LO	LO	NNLO	
σ_{bkg} [fb]	0.685	0.279	0.101	0.074	0.026	0.017	0.02	0.001	0.304	1.504

The same-flavor opposite-sign dilepton veto reduces the dominant $t\bar{t}\bar{Z}$ background. In case more than one such $\ell^+\ell^-$ pair exists, we select the combination closest to the Z -mass for rejection. The remaining signal rate is given in Tab. 1, while the background rates are summarized in Tab. 2.

For discovery reach and exclusion limits, we compute the significance using the likelihood for a simple counting experiment [96]. If we observe n events with n_{pred} predicted, the agreement between observation and prediction is given by

$$Z(n|n_{\text{pred}}) = \sqrt{-2 \ln \frac{L(n|n_{\text{pred}})}{L(n|n)}}, \quad \text{with} \quad L(n|\bar{n}) = \frac{e^{-\bar{n}} \bar{n}^n}{n!}. \quad (17)$$

For discovery, we compare the observed signal plus background with the background prediction and require $Z(s+b|b) > 5$. For exclusion, we assume a background-consistent measurement after predicting a signal on top of the background, such that $Z(b|s+b) > 2$. For instance, assuming an HL-LHC data set with 3000 fb^{-1} and the signal and background cross sections in Tabs. 1 and 2, we find a significance of $\sim 5.6\sigma$ for $m_{H^+} = 350 \text{ GeV}$ and $\sim 5\sigma$ for $m_{H^+} = 500 \text{ GeV}$.

We illustrate in Fig. 4 the Run 3 and HL-LHC reach for the charged Higgs signature in the $|\rho_{tc}|$ - c_γ plane. We see from the left panel that Run 3 can exclude $|\rho_{tc}| > 0.3$ and $|c_\gamma| = 0.27$ for $m_{H^\pm} = 350 \text{ GeV}$, while the HL-LHC will be sensitive to $|\rho_{tc}| > 0.2$ and $|c_\gamma| = 0.14$. For larger Higgs masses, the expected limits become only slightly weaker. The b -associated charged Higgs channel covers the $|\rho_{tc}|$ range preferred by EWBG, but there remains a slice of EWBG parameter space with $|c_\gamma| \lesssim 0.14$. This follows as an effect of decreasing $\mathcal{B}(H^\pm \rightarrow W^\pm h)$ with smaller c_γ . Unfortunately, this hole is unlikely to be filled by other charged Higgs decays, because for instance the standard signature $H^+ \rightarrow t\bar{b}$ requires large production rates. Here, utilizing the expression from Ref. [73], the limit from B_s in the left panel of Fig. 4 is plotted for $m_{H^+} = 350 \text{ GeV}$ to conform with the benchmark charged Higgs mass for $pp \rightarrow bH^+ \rightarrow bW^+h$ signature.

4 Neutral Higgs production

To cover the parameter region $|c_\gamma| < 0.14$, left open by the charged Higgs signature, we turn to the neutral Higgs channel,

$$cg \rightarrow tH/tA \rightarrow t(t\bar{c}), \quad (18)$$

also given in Fig. 3, where production and decay are both mediated by ρ_{tc} . A very slight c_γ -dependence of the $cg \rightarrow tH/tA \rightarrow t\bar{t}\bar{c}$ process arises from the heavy Higgs branching ratios. Non-resonant and t -channel diagrams with H/A exchange leading to $cc \rightarrow tt$ scattering as well as $gg \rightarrow t\bar{t}\bar{c}\bar{c}$, though small, are included in our signal analysis.




Figure 4: Projected 300 fb^{-1} exclusion (dot-dashed) and HL-LHC discovery (solid) and exclusion (dashed) contours for the charged Higgs signature $pp \rightarrow bH^+ \rightarrow bW^+h$, along with EWBG-favored region and the indirect constraints from Fig. 2.

For small c_γ , the neutral Higgs production process currently leads to the most stringent limit on ρ_{tc} [33, 97], because it affects the SM control region of the Run 2 $t\bar{t}\bar{\ell}\bar{\ell}$ ($4t$) analysis by CMS [98]. Based on the number of b -jets and leptons, CMS divides its analysis into several signal and two control regions. The most stringent constraint on ρ_{tc} arises from the $t\bar{t}W$ control region (CRW) [32, 33]. The CMS baseline selection includes two same-sign leptons with

$$p_{T,\ell} > 25, 20 \text{ GeV} \quad \text{and} \quad |\eta_e| < 2.5, \quad |\eta_\mu| < 2.4, \quad (19)$$

where the charge-misidentified Drell-Yan background is reduced by vetoing same-sign electron pairs with $m_{ee} < 12 \text{ GeV}$. The CRW then requires two to five jets, two of them b -tagged. All jets have to fulfill $|\eta_j| < 2.4$, and events are selected if they fulfill any one of

$$\begin{aligned} \text{(i)} \quad & p_{T,b_1} > 40 \text{ GeV}, & p_{T,b_2} > 40 \text{ GeV}, & - \\ \text{(ii)} \quad & p_{T,b_1} > 20 \text{ GeV}, & p_{T,b_2} = 20 \dots 40 \text{ GeV}, & p_{T,j_3} > 40 \text{ GeV}, \\ \text{(iii)} \quad & p_{T,b_{1,2}} = 20 \dots 40 \text{ GeV}, & - & p_{T,j_{3,4}} > 40 \text{ GeV}. \end{aligned} \quad (20)$$

Finally, the analysis requires [98]

$$H_T = \sum_{\text{jets}} p_{T,j} > 300 \text{ GeV} \quad \text{and} \quad \cancel{E}_T > 50 \text{ GeV}. \quad (21)$$

With this selection, CMS observes 338 events with 335 ± 18 events expected from SM-backgrounds plus $4t$ signal. To estimate the CRW limits on ρ_{tc} , we generate both neutral Higgs processes with the decay $H/A \rightarrow t\bar{c}$, followed by lepton-hadron combinations of the top decays at $\sqrt{s} = 13 \text{ TeV}$. We use the same setup as for the charged Higgs simulations, except that we use the default CMS detector card in Delphes 3.4.2. Remaining uncertainties on our simulation affect the c -initiated processes $cg \rightarrow bH^+$ and $cg \rightarrow tA/tH$, such as from parton densities and scale dependence [78, 99–101]. We expect them to be small, and do not include them, just as we do not account for non-prompt and fake backgrounds.

There exist a similar ATLAS search [102], but it is less constraining [103]. This is primarily due to the definition of signal regions and selection criteria. Furthermore, searches for squark

Table 3: Background cross sections for the dedicated same-sign top search after selection cuts at $\sqrt{s} = 14$ TeV.

background	σ [fb]	background	σ [fb]	background	σ [fb]
$t\bar{t}W$	1.31	$t\bar{t}Z$	1.97	$tZ + \text{jets}$	0.007
$4t$	0.092	$3t + W$	0.001	$3t + j$	0.0004
$t\bar{t}h$	0.058	charge-flip	0.024	non-prompt	$1.5 \times t\bar{t}W$

pair production in R -parity violating supersymmetry [104] and exotics searches for same-sign dileptons and b -jets [105] involve similar final states, but their selection cuts are too model-specific to be applied to our signature.

To judge the impact of the existing CMS CRW limits from $4t$ search, we focus on the border of the EWBG-region with $c_\gamma = 0.1$ and $|\rho_{tc}| = 0.5$. We stick to our two charged Higgs masses, assume $m_A \approx m_{H^\pm} = 350, 500$ GeV for the pseudoscalar, and decouple the heavy scalar H . In this scenario, the same-sign top contribution to the CRW arises from $cg \rightarrow tA \rightarrow t\bar{t}\bar{c}$. We demand that the combination of SM-backgrounds and heavy neutral Higgs production agree with observed within 2σ and give the excluded regions in Fig. 5. To scan the parameter space we use a simplified scaling $|\rho_{tc}|^2 \mathcal{B}(A \rightarrow t\bar{c})$, such that $\Gamma_A = 3.05(6.08)$ GeV for $m_A = 350(500)$ GeV. The exclusion covers most of the EWBG-region except for small values of $|\rho_{tc}|$.

A dedicated same-sign top search, such as the $pp \rightarrow tA + X \rightarrow t\bar{t}\bar{c} + X$ study of Ref. [103], can probe the nominal parameter space of ρ_{tc} -EWBG. This process can be searched for in events containing same-sign dileptons ($ee, \mu\mu, e\mu$), at least three jets with at least two b -tag, and some \cancel{E}_T . The dominant backgrounds are $t\bar{t}Z, t\bar{t}W, 4t$, while $t\bar{t}h$, with $tZ + \text{jets}, 3t + W$ and $3t + j$ give subdominant contributions, and the non-prompt background can be 1.5 times the rate of $t\bar{t}W$. In addition, if a lepton charge gets misidentified, the $t\bar{t} + \text{jets}$ and $Z/\gamma^* + \text{jets}$ processes will also contribute. For further details of the QCD correction factors for different backgrounds, we refer to Ref. [103]. To reduce backgrounds, we applied an event selection different from the CRW of Ref. [98]: the leading and sub-leading same-sign leptons should have $p_T > 25(20)$ GeV and $|\eta| < 2.5$. All three jets are required to have $p_T > 20$ GeV and $|\eta| < 2.5$. All jets and leptons are separated by $\Delta R_{ij} > 0.4$. The all event should have $\cancel{E}_T > 35$ GeV and $H_T > 300$ GeV, where the latter includes the two leading same-sign leptons. The background cross sections after selection cuts are summarized in Tab. 3.

For the reference values $|\rho_{tc}| = 0.5$ and $c_\gamma = 0.1$, we generate the same-sign top cross sections for $m_A = 350$ and 500 GeV. Based on the background rates of Tab. 3 and Eq.(17), rescaling the signal cross section by $|\rho_{tc}|^2 \mathcal{B}(A \rightarrow t\bar{c})$, we find the exclusion (green dashed) and discovery (green solid) contours in the $|c_\gamma| - |\rho_{tc}|$ plane as given in Fig. 5.

A loop hole in the neutral Higgs analysis appears though the destructive interference of $cg \rightarrow tH \rightarrow t\bar{t}\bar{c}$ and $cg \rightarrow tA \rightarrow t\bar{t}\bar{c}$. If the widths and masses of the two heavy neutral Higgses become degenerate, the two production processes completely cancel [32, 33] and the same-sign top signature vanishes. Our limits derived from A -production would be similar for H -production with $m_A \gg m_H$. We now illustrate limits for $m_A \sim m_H$ with a case where the three heavy Higgs masses are of similar size, specifically $m_{H^\pm} = 350(500)$ GeV, $m_A = 343(524)$ GeV, and $m_H = 355(501)$ GeV. The self-couplings are $\eta_1 = 0.276(0.297)$, $\eta_2 = 1.335(2.762)$, $\eta_3 = 1.66(1.21)$, $\eta_4 = -0.04(0.398)$, $\eta_5 = 0.121(-0.428)$, $\eta_6 = -0.181(-0.386)$, $\eta_7 = 0.605(-0.095)$, and $\mu_{22}^2/v^2 = 1.189(3.516)$, in agreement with perturbativity, positivity, unitarity, and electroweak precision data [69]. The relevant decays are $A \rightarrow t\bar{c}, Zh$ and $H \rightarrow t\bar{c}, hh, ZZ, WW$, with mild contributions from the $\lambda_f c_\gamma$ -dependent fermionic decays to $b\bar{b}$ and $t\bar{t}$. For $\rho_{tc} = 0.5$ and $c_\gamma = 0.1$, the total widths are $\Gamma_A = 3.28(7.37)$ GeV and




Figure 5: Exclusion regions from neutral Higgs production contributing to the CMS CRW [98] (gray shades), as well as HL-LHC expectations from a dedicated same-sign top search [103] (green). We also show the EWBG region and the indirect constraints from Fig. 2 and the HL-LHC charged Higgs reach from Fig. 4.

$\Gamma_H = 2.91(6.56)$ GeV, and the combined contributions to the CRW rates are 0.467 fb and 0.261 fb, corresponding to 64 and 35.8 events. Demanding that the combination of events expected in the SM and from the neutral Higgs channels agree within 2σ of the observed number, we find that $|\rho_{tc}| = 0.5$ is already excluded for $m_{H^\pm} = 350$ GeV and $c_\gamma = 0.1$, and barely allowed for $m_{H^\pm} = 500$ GeV. We see that, due to the choice of parameters, the cancellation between $c_g \rightarrow tH \rightarrow tt\bar{c}$ and $c_g \rightarrow tA \rightarrow tt\bar{c}$ is not exact, and the CRW limit is stronger than the H (or A) decoupled case.

As mentioned in the introduction, we ignore all ρ_{ij} couplings except for ρ_{tc} , so before closing we should discuss the impact of this assumption. It may well be that the $\rho^{U,D,L}$ matrices share the flavor-ordering of the Yukawa couplings, $\rho_{tt} \sim \lambda_t$, $\rho_{bb} \sim \lambda_b$ and $\rho_{\tau\tau} \sim \lambda_\tau$. Current data still allows $\rho_{tt} \lesssim 0.5$ [39] and $\rho_{bb} \sim 0.1$ [71, 72] for sub-TeV scalars, and both parameters can account for the observed baryon asymmetry. The extra top Yukawa coupling ρ_{tt} can be searched for in signatures such as $gg \rightarrow A/H \rightarrow t\bar{t}$ [106, 107] $gg \rightarrow A/Ht\bar{t} \rightarrow t\bar{t}t\bar{t}$ [98] and $gb \rightarrow \bar{t}H^+ \rightarrow \bar{t}t\bar{b}$ [77, 78, 108, 109], while rare decays $\mathcal{B}(B \rightarrow X_s\gamma)$ and $B_{d,s}$ mixing provide indirect probes [79]. In general, a large value for ρ_{tt} dilutes the decays $A/H \rightarrow t\bar{c}$ and $H^+ \rightarrow W^+h$ through $A/H \rightarrow t\bar{t}$ and $H^+ \rightarrow t\bar{b}$. However, the combination with ρ_{tc} opens additional discovery modes such as $cg \rightarrow tA/tH \rightarrow tt\bar{t}$ [32] and $cg \rightarrow bH^+ \rightarrow bt\bar{b}$ [39]. There also exist several direct and indirect constraints on ρ_{bb} [71, 72]. Finally, a large allowed value of ρ_{tu} [103] combined with non-vanishing ρ_{tc} will be constrained by D -meson mixing [73, 79]. Similarly, constraints on ρ_{tt} , ρ_{bb} , $\rho_{\tau\tau}$ from flavor physics and low energy observables as discussed in Refs. [38, 72, 73, 79], and their detailed impact on the ρ_{tc} -EWBG would be an interesting future direction.

5 Outlook

Electroweak baryogenesis is an attractive target for experimental analysis, because it can be tested by a variety of measurements. Specific models typically combine new bosonic degrees of freedom with extra CP -violation. In our case, the new degrees of freedom are provided by

a general or type-III 2HDM. If the Higgs self-couplings are sufficiently large, the heavy Higgs states can be relatively heavy, so we use $m_{H^\pm} = 350$ and 500 GeV as benchmark scenarios. The complex phase is given by an FCNC top–charm coupling with $|\rho_{tc}| \gtrsim 0.5$, combined with a CP -even Higgs mixing angle $c_\gamma \gtrsim 0.1$. At the LHC, ρ_{tc} has the advantage that we can test it in processes mediated by this large top Yukawa, but with a charm quark in the initial state, while it easily evades EDM constraints.

In the allowed 2HDM parameter space, the charged Higgs has to be relatively light, which means we can search for it via $cg \rightarrow bH^+$ with a subsequent $H^+ \rightarrow W^+h$ decay. Our proposed analysis is relatively straightforward and probes most of the EWBG parameter space at the HL-LHC, with the exception of small values of $c_\gamma \sim 0.1 \dots 0.12$, when $H^+ \rightarrow W^+h$ decay becomes too suppressed by CP -even Higgs boson mixing.

A complementary channel that can survive small CP -even Higgs boson mixing is heavy neutral Higgs production, $cg \rightarrow tA/tH$, together with $A/H \rightarrow t\bar{c}$ decay. In this case, production and decay are both mediated by ρ_{tc} without being suppressed by small c_γ , providing strong limits on ρ_{tc} even for small c_γ values. The search channel at the LHC is same-sign top pairs, allowing us to extract limits already from Run 2. At the HL-LHC, the decay $t \rightarrow ch$, charged heavy Higgs searches, and neutral heavy Higgs searches guarantee a comprehensive coverage of the ρ_{tc} -EWBG parameter space in the general 2HDM, leaving us with the challenge of observing the CP -violating phase in a dedicated analysis.

Acknowledgments

First, we would like to thank Kai-Feng (Jack) Chen for discussions and for clarifications on Eq.(20). We are also grateful to Eibun Senaha and Margarete Mühlleitner for very helpful discussions and comments. WSH is supported by MOST 109-2112-M-002-015-MY3 of Taiwan and NTU 109L104019. TM is supported by a Postdoctoral Research Fellowship from Alexander von Humboldt Foundation. The research of TP is supported by the Deutsche Forschungsgemeinschaft (German Research Foundation) under grant 396021762 – TRR 257 *Particle Physics Phenomenology after the Higgs Discovery*.

References

- [1] G. Aad et al., *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, Phys. Lett. B **716**, 1 (2012), doi:[10.1016/j.physletb.2012.08.020](https://doi.org/10.1016/j.physletb.2012.08.020).
- [2] S. Chatrchyan et al., *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, Phys. Lett. B **716**, 30 (2012), doi:[10.1016/j.physletb.2012.08.021](https://doi.org/10.1016/j.physletb.2012.08.021).
- [3] G. Aad et al., *Combined measurements of Higgs boson production and decay using up to 80 fb⁻¹ of proton-proton collision data at $\sqrt{s} = 13$ TeV collected with the ATLAS experiment*, Phys. Rev. D **101**, 012002 (2020), doi:[10.1103/PhysRevD.101.012002](https://doi.org/10.1103/PhysRevD.101.012002).
- [4] A. M. Sirunyan et al., *Combined measurements of Higgs boson couplings in proton-proton collisions at $\sqrt{s} = 13$ TeV*, Eur. Phys. J. C **79**, 421 (2019), doi:[10.1140/epjc/s10052-019-6909-y](https://doi.org/10.1140/epjc/s10052-019-6909-y).

- [5] A. Biekötter, T. Corbett and T. Plehn, *The Gauge-Higgs legacy of the LHC run II*, SciPost Phys. **6**, 064 (2019), doi:[10.21468/SciPostPhys.6.6.064](https://doi.org/10.21468/SciPostPhys.6.6.064).
- [6] J. Ellis, C. W. Murphy, V. Sanz and T. You, *Updated global SMEFT fit to Higgs, diboson and electroweak data*, J. High Energ. Phys. **06**, 146 (2018), doi:[10.1007/JHEP06\(2018\)146](https://doi.org/10.1007/JHEP06(2018)146).
- [7] E. da Silva Almeida, A. Alves, N. Rosa-Agostinho, O. J. P. Éboli and M. C. Gonzalez-Garcia, *Electroweak sector under scrutiny: A combined analysis of LHC and electroweak precision data*, Phys. Rev. D **99**, 033001 (2019), doi:[10.1103/PhysRevD.99.033001](https://doi.org/10.1103/PhysRevD.99.033001).
- [8] S. Kraml, T. Quang Loc, D. Thi Nhung and L. Duc Ninh, *Constraining new physics from Higgs measurements with Lilith: update to LHC Run 2 results*, SciPost Phys. **7**, 052 (2019), doi:[10.21468/SciPostPhys.7.4.052](https://doi.org/10.21468/SciPostPhys.7.4.052).
- [9] S. van Beek, E. Roberto Nocera, J. Rojo and E. Slade, *Constraining the SMEFT with Bayesian reweighting*, SciPost Phys. **7**, 070 (2019), doi:[10.21468/SciPostPhys.7.5.070](https://doi.org/10.21468/SciPostPhys.7.5.070).
- [10] S. Dawson, S. Homiller and S. D. Lane, *Putting standard model EFT fits to work*, Phys. Rev. D **102**, 055012 (2020), doi:[10.1103/PhysRevD.102.055012](https://doi.org/10.1103/PhysRevD.102.055012).
- [11] J. Ellis, M. Madigan, K. Mimasu, V. Sanz and T. You, *Top, Higgs, diboson and electroweak fit to the Standard Model effective field theory*, J. High Energ. Phys. **04**, 279 (2021), doi:[10.1007/JHEP04\(2021\)279](https://doi.org/10.1007/JHEP04(2021)279).
- [12] T. D. Lee, *A theory of spontaneous T violation*, Phys. Rev. D **8**, 1226 (1973), doi:[10.1103/PhysRevD.8.1226](https://doi.org/10.1103/PhysRevD.8.1226).
- [13] A. Djouadi, *The anatomy of electroweak symmetry breaking*, Phys. Rep. **457**, 1 (2008), doi:[10.1016/j.physrep.2007.10.004](https://doi.org/10.1016/j.physrep.2007.10.004).
- [14] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher and J. P. Silva, *Theory and phenomenology of two-Higgs-doublet models*, Phys. Rep. **516**, 1 (2012), doi:[10.1016/j.physrep.2012.02.002](https://doi.org/10.1016/j.physrep.2012.02.002).
- [15] A. Djouadi, *The anatomy of electroweak symmetry breaking Tome II: The Higgs bosons in the Minimal Supersymmetric Model*, Phys. Rep. **459**, 1 (2008), doi:[10.1016/j.physrep.2007.10.005](https://doi.org/10.1016/j.physrep.2007.10.005).
- [16] C. T. Hill and E. H. Simmons, *Strong dynamics and electroweak symmetry breaking*, Phys. Rep. **381**, 235 (2003), doi:[10.1016/S0370-1573\(03\)00140-6](https://doi.org/10.1016/S0370-1573(03)00140-6).
- [17] M. Schmaltz and D. Tucker-Smith, *Little Higgs theories*, Annu. Rev. Nucl. Part. Sci. **55**, 229 (2005), doi:[10.1146/annurev.nucl.55.090704.151502](https://doi.org/10.1146/annurev.nucl.55.090704.151502).
- [18] M. Perelstein, *Little Higgs models and their phenomenology*, Prog. Part. Nucl. Phys. **58**, 247 (2007), doi:[10.1016/j.ppnp.2006.04.001](https://doi.org/10.1016/j.ppnp.2006.04.001).
- [19] J. C. Pati and A. Salam, *Lepton number as the fourth "color"*, Phys. Rev. D **10**, 275 (1974), doi:[10.1103/PhysRevD.10.275](https://doi.org/10.1103/PhysRevD.10.275).
- [20] F. Gürsey, P. Ramond and P. Sikivie, *A universal gauge theory model based on E_6* , Phys. Lett. B **60**, 177 (1976), doi:[10.1016/0370-2693\(76\)90417-2](https://doi.org/10.1016/0370-2693(76)90417-2).
- [21] Y. Achiman and B. Stech, *Quark-lepton symmetry and mass scales in an E_6 unified gauge model*, Phys. Lett. B **77**, 389 (1978), doi:[10.1016/0370-2693\(78\)90584-1](https://doi.org/10.1016/0370-2693(78)90584-1).

- [22] R. Barbieri, D. V. Nanopoulos and A. Masiero, *Hierarchical fermion masses in E_6* , Phys. Lett. B **104**, 194 (1981), doi:[10.1016/0370-2693\(81\)90589-X](https://doi.org/10.1016/0370-2693(81)90589-X).
- [23] A. D. Sakharov, *Violation of CP Invariance, C asymmetry, and baryon asymmetry of the universe*, Sov. Phys. Usp. **34**, 392 (1991), doi:[10.1070/PU1991v034n05ABEH002497](https://doi.org/10.1070/PU1991v034n05ABEH002497).
- [24] A. I. Bochkarev, S. V. Kuzmin and M. E. Shaposhnikov, *Electroweak baryogenesis and the Higgs boson mass problem*, Phys. Lett. B **244**, 275 (1990), doi:[10.1016/0370-2693\(90\)90069-I](https://doi.org/10.1016/0370-2693(90)90069-I).
- [25] N. Turok and J. Zadrozny, *Electroweak baryogenesis in the two-doublet model*, Nucl. Phys. B **358**, 471 (1991), doi:[10.1016/0550-3213\(91\)90356-3](https://doi.org/10.1016/0550-3213(91)90356-3).
- [26] K.-F. Chen, W.-S. Hou, C. Kao and M. Kohda, *When the Higgs meets the top: Search for $t \rightarrow ch^0$ at the LHC*, Phys. Lett. B **725**, 378 (2013), doi:[10.1016/j.physletb.2013.07.060](https://doi.org/10.1016/j.physletb.2013.07.060).
- [27] P. Chang, K.-F. Chen and W.-S. Hou, *Flavor physics and CP violation*, Prog. Part. Nucl. Phys. **97**, 261 (2017), doi:[10.1016/j.pnpnp.2017.07.001](https://doi.org/10.1016/j.pnpnp.2017.07.001).
- [28] W.-S. Hou, *Tree level $t \rightarrow ch^0$ or $h^0 \rightarrow t\bar{c}$ or decays*, Phys. Lett. B **296**, 179 (1992), doi:[10.1016/0370-2693\(92\)90823-M](https://doi.org/10.1016/0370-2693(92)90823-M).
- [29] D. López-Val, T. Plehn and M. Rauch, *Measuring extended Higgs sectors as a consistent free couplings model*, J. High Energ. Phys. **10**, 134 (2013), doi:[10.1007/JHEP10\(2013\)134](https://doi.org/10.1007/JHEP10(2013)134).
- [30] J. Haller, A. Hoecker, R. Kogler, K. Mönig, T. Peiffer and J. Stelzer, *Update of the global electroweak fit and constraints on two-Higgs-doublet models*, Eur. Phys. J. C **78**, 675 (2018), doi:[10.1140/epjc/s10052-018-6131-3](https://doi.org/10.1140/epjc/s10052-018-6131-3).
- [31] K. Fuyuto, W.-S. Hou and E. Senaha, *Electroweak baryogenesis driven by extra top Yukawa couplings*, Phys. Lett. B **776**, 402 (2018), doi:[10.1016/j.physletb.2017.11.073](https://doi.org/10.1016/j.physletb.2017.11.073).
- [32] M. Kohda, T. Modak and W.-S. Hou, *Searching for new scalar bosons via triple-top signature in $cg \rightarrow tS^0 \rightarrow tt\bar{t}$* , Phys. Lett. B **776**, 379 (2018), doi:[10.1016/j.physletb.2017.11.056](https://doi.org/10.1016/j.physletb.2017.11.056).
- [33] W.-S. Hou, M. Kohda and T. Modak, *Constraining a lighter exotic scalar via same-sign top*, Phys. Lett. B **786**, 212 (2018), doi:[10.1016/j.physletb.2018.09.046](https://doi.org/10.1016/j.physletb.2018.09.046).
- [34] W.-S. Hou, G.-L. Lin, C.-Y. Ma and C.-P. Yuan, *Probing flavor changing neutral Higgs couplings at LHC*, Phys. Lett. B **409**, 344 (1997), doi:[10.1016/S0370-2693\(97\)00921-0](https://doi.org/10.1016/S0370-2693(97)00921-0).
- [35] W. Altmannshofer, J. Eby, S. Gori, M. Lotito, M. Martone and D. Tuckler, *Collider signatures of flavorful Higgs bosons*, Phys. Rev. D **94**, 115032 (2016), doi:[10.1103/PhysRevD.94.115032](https://doi.org/10.1103/PhysRevD.94.115032).
- [36] W. Altmannshofer, B. Maddock and D. Tuckler, *Rare top decays as probes of flavorful Higgs bosons*, Phys. Rev. D **100**, 015003 (2019), doi:[10.1103/PhysRevD.100.015003](https://doi.org/10.1103/PhysRevD.100.015003).
- [37] S. Gori, C. Grojean, A. Juste and A. Paul, *Heavy Higgs searches: flavour matters*, J. High Energ. Phys. **01**, 108 (2018), doi:[10.1007/JHEP01\(2018\)108](https://doi.org/10.1007/JHEP01(2018)108).
- [38] S. Iguro and K. Tobe, *$R(D^{(*)})$ in a general two Higgs doublet model*, Nucl. Phys. B **925**, 560 (2017), doi:[10.1016/j.nuclphysb.2017.10.014](https://doi.org/10.1016/j.nuclphysb.2017.10.014).

- [39] D. K. Ghosh, W.-S. Hou and T. Modak, *Sub-TeV H^+ boson production as probe of extra top Yukawa couplings*, Phys. Rev. Lett. **125**, 221801 (2020), doi:[10.1103/PhysRevLett.125.221801](https://doi.org/10.1103/PhysRevLett.125.221801).
- [40] A. M. Sirunyan et al., *Search for the flavor-changing neutral current interactions of the top quark and the Higgs boson which decays into a pair of b quarks at $\sqrt{s} = 13$ TeV*, J. High Energ. Phys. **06**, 102 (2018), doi:[10.1007/JHEP06\(2018\)102](https://doi.org/10.1007/JHEP06(2018)102).
- [41] M. Aaboud et al., *Search for top-quark decays $t \rightarrow Hq$ with $36fb^{-1}$ of pp collision data at $\sqrt{s} = 13$ TeV with the ATLAS detector*, J. High Energ. Phys. **05**, 123 (2019), doi:[10.1007/JHEP05\(2019\)123](https://doi.org/10.1007/JHEP05(2019)123).
- [42] S. Davidson and H. E. Haber, *Basis-independent methods for the two-Higgs-doublet model*, Phys. Rev. D **72**, 035004 (2005), doi:[10.1103/PhysRevD.72.035004](https://doi.org/10.1103/PhysRevD.72.035004), [Erratum: Phys. Rev. D **72**, 099902 (2005), doi:[10.1103/PhysRevD.72.099902](https://doi.org/10.1103/PhysRevD.72.099902)].
- [43] W.-S. Hou and M. Kikuchi, *Approximate alignment in two-Higgs-doublet model with extra Yukawa couplings*, Europhys. Lett. **123**, 11001 (2018), doi:[10.1209/0295-5075/123/11001](https://doi.org/10.1209/0295-5075/123/11001).
- [44] P. Basler and M. Mühlleitner, *BSMPT (Beyond the Standard Model Phase Transitions): A tool for the electroweak phase transition in extended Higgs sectors*, Comput. Phys. Commun. **237**, 62 (2019), doi:[10.1016/j.cpc.2018.11.006](https://doi.org/10.1016/j.cpc.2018.11.006).
- [45] P. Basler, M. Mühlleitner and J. Müller, *Electroweak phase transition in non-minimal Higgs sectors*, J. High Energ. Phys. **05**, 016 (2020), doi:[10.1007/JHEP05\(2020\)016](https://doi.org/10.1007/JHEP05(2020)016).
- [46] P. Basler, M. Mühlleitner and J. Müller, *BSMPT v2 A tool for the electroweak phase transition and the Baryon asymmetry of the universe in extended Higgs sectors*, (2020), [arXiv:2007.01725](https://arxiv.org/abs/2007.01725).
- [47] J. M. Cline, K. Kainulainen and A. P. Vischer, *Dynamics of two-Higgs-doublet CP violation and baryogenesis at the electroweak phase transition*, Phys. Rev. D **54**, 2451 (1996), doi:[10.1103/PhysRevD.54.2451](https://doi.org/10.1103/PhysRevD.54.2451).
- [48] K. Funakubo, A. Kakuto, S. Otsuki, K. Takenaga and F. Toyoda, *CP-violating profile of the electroweak bubble wall*, Progr. Theor. Phys. **94**, 845 (1995), doi:[10.1143/PTP.94.845](https://doi.org/10.1143/PTP.94.845).
- [49] G. C. Dorsch, S. J. Huber, T. Konstandin and J. M. No, *A second Higgs doublet in the early universe: baryogenesis and gravitational waves*, J. Cosmol. Astropart. Phys. **052** (2017), doi:[10.1088/1475-7516/2017/05/052](https://doi.org/10.1088/1475-7516/2017/05/052).
- [50] P. Basler, M. Mühlleitner and J. Wittbrodt, *The CP-violating 2HDM in light of a strong first order electroweak phase transition and implications for Higgs pair production*, J. High Energ. Phys. **03**, 061 (2018), doi:[10.1007/JHEP03\(2018\)061](https://doi.org/10.1007/JHEP03(2018)061).
- [51] W.-S. Hou and T. Modak, *Production of tZH and $t\bar{Z}h$ in 2HDM: Prospects for discovery at the LHC*, Phys. Rev. D **101**, 035007 (2020), doi:[10.1103/PhysRevD.101.035007](https://doi.org/10.1103/PhysRevD.101.035007).
- [52] N. Craig, F. D'Eramo, P. Draper, S. Thomas and H. Zhang, *The hunt for the rest of the Higgs bosons*, J. High Energ. Phys. **06**, 137 (2015), doi:[10.1007/JHEP06\(2015\)137](https://doi.org/10.1007/JHEP06(2015)137).
- [53] S. Kanemura, H. Yokoya and Y.-J. Zheng, *Searches for additional Higgs bosons in multi-top-quarks events at the LHC and the International Linear Collider*, Nucl. Phys. B **898**, 286 (2015), doi:[10.1016/j.nuclphysb.2015.07.005](https://doi.org/10.1016/j.nuclphysb.2015.07.005).

- [54] S. Gori, I.-W. Kim, N. R. Shah and K. M. Zurek, *Closing the wedge: Search strategies for extended Higgs sectors with heavy flavor final states*, Phys. Rev. D **93**, 075038 (2016), doi:[10.1103/PhysRevD.93.075038](https://doi.org/10.1103/PhysRevD.93.075038).
- [55] S. M. Barr and A. Zee, *Electric dipole moment of the electron and of the neutron*, Phys. Rev. Lett. **65**, 21 (1990), doi:[10.1103/PhysRevLett.65.21](https://doi.org/10.1103/PhysRevLett.65.21), [Erratum: Phys. Rev. Lett. **65**, 2920 (1990), doi:[10.1103/PhysRevLett.65.2920](https://doi.org/10.1103/PhysRevLett.65.2920)].
- [56] K. Fuyuto, W.-S. Hou and E. Senaha, *Cancellation mechanism for the electron electric dipole moment connected with the baryon asymmetry of the Universe*, Phys. Rev. D **101**, 011901 (2020), doi:[10.1103/PhysRevD.101.011901](https://doi.org/10.1103/PhysRevD.101.011901).
- [57] V. Andreev et al., *Improved limit on the electric dipole moment of the electron*, Nature **562**, 355 (2018), doi:[10.1038/s41586-018-0599-8](https://doi.org/10.1038/s41586-018-0599-8).
- [58] F. Sala, *A bound on the charm chromo-EDM and its implications*, J. High Energ. Phys. **03**, 061 (2014), doi:[10.1007/JHEP03\(2014\)061](https://doi.org/10.1007/JHEP03(2014)061).
- [59] K. Funakubo, A. Kakuto and K. Takenaga, *The Effective potential of electroweak theory with two massless Higgs doublets at finite temperature*, Prog. Theor. Phys. **91**, 341 (1994), doi:[10.1143/PTP91.341](https://doi.org/10.1143/PTP91.341).
- [60] J. M. Cline and P.-A. Lemieux, *Electroweak phase transition in two Higgs doublet models*, Phys. Rev. D **55**, 3873 (1997), doi:[10.1103/PhysRevD.55.3873](https://doi.org/10.1103/PhysRevD.55.3873).
- [61] S. Kanemura, Y. Okada and E. Senaha, *Electroweak baryogenesis and quantum corrections to the triple Higgs boson coupling*, Phys. Lett. B **606**, 361 (2005), doi:[10.1016/j.physletb.2004.12.004](https://doi.org/10.1016/j.physletb.2004.12.004).
- [62] L. Fromme, S. J. Huber and M. Seniuch, *Baryogenesis in the two-Higgs doublet model*, J. High Energ. Phys. **11**, 038 (2006), doi:[10.1088/1126-6708/2006/11/038](https://doi.org/10.1088/1126-6708/2006/11/038).
- [63] D. Borah and J. M. Cline, *Inert doublet dark matter with strong electroweak phase transition*, Phys. Rev. D **86**, 055001 (2012), doi:[10.1103/PhysRevD.86.055001](https://doi.org/10.1103/PhysRevD.86.055001).
- [64] J. M. Cline and K. Kainulainen, *Improved electroweak phase transition with subdominant inert doublet dark matter*, Phys. Rev. D **87**, 071701 (2013), doi:[10.1103/PhysRevD.87.071701](https://doi.org/10.1103/PhysRevD.87.071701).
- [65] G. C. Dorsch, S. J. Huber and J. M. No, *A strong electroweak phase transition in the 2HDM after LHC8*, J. High Energ. Phys. **10**, 029 (2013), doi:[10.1007/JHEP10\(2013\)029](https://doi.org/10.1007/JHEP10(2013)029).
- [66] N. Blinov, S. Profumo and T. Stefaniak, *The electroweak phase transition in the Inert Doublet Model*, J. Cosmol. Astropart. Phys. **028** (2015), doi:[10.1088/1475-7516/2015/07/028](https://doi.org/10.1088/1475-7516/2015/07/028).
- [67] K. Fuyuto and E. Senaha, *Sphaleron and critical bubble in the scale invariant two Higgs doublet model*, Phys. Lett. B **747**, 152 (2015), doi:[10.1016/j.physletb.2015.05.061](https://doi.org/10.1016/j.physletb.2015.05.061).
- [68] P. Basler, M. Krause, M. Mühlleitner, J. Wittbrodt and A. Wlotzka, *Strong first order electroweak phase transition in the CP-conserving 2HDM revisited*, J. High Energ. Phys. **02**, 121 (2017), doi:[10.1007/JHEP02\(2017\)121](https://doi.org/10.1007/JHEP02(2017)121).
- [69] D. Eriksson, J. Rathsman and O. Stål, *2HDMC: Two-Higgs-Doublet model calculator*, Comput. Phys. Commun. **181**, 189 (2010), doi:[10.1016/j.cpc.2009.09.011](https://doi.org/10.1016/j.cpc.2009.09.011).

- [70] W.-S. Hou, M. Kohda and T. Modak, *Top-assisted di-Higgs boson production motivated by baryogenesis*, Phys. Rev. D **99**, 055046 (2019), doi:[10.1103/PhysRevD.99.055046](https://doi.org/10.1103/PhysRevD.99.055046).
- [71] T. Modak, *Probing an additional bottom Yukawa coupling via $bg \rightarrow bA \rightarrow bZH$ signature*, Phys. Rev. D **100**, 035018 (2019), doi:[10.1103/PhysRevD.100.035018](https://doi.org/10.1103/PhysRevD.100.035018).
- [72] T. Modak and E. Senaha, *Probing electroweak baryogenesis induced by extra bottom Yukawa coupling via EDMs and collider signatures*, J. High Energ. Phys. **11**, 025 (2020), doi:[10.1007/JHEP11\(2020\)025](https://doi.org/10.1007/JHEP11(2020)025).
- [73] A. Crivellin, C. Greub and A. Kokulu, *Flavor-phenomenology of two-Higgs-doublet models with generic Yukawa structure*, Phys. Rev. D **87**, 094031 (2013), doi:[10.1103/PhysRevD.87.094031](https://doi.org/10.1103/PhysRevD.87.094031).
- [74] ATLAS Collaboration, *Expected sensitivity of ATLAS to FCNC top quark decays $t \rightarrow Zu$ and $t \rightarrow Hq$ at the High Luminosity LHC*, Tech. Rep. ATL-PHYS-PUB-2016-019, CERN, Geneva (2016).
- [75] W.-S. Hou, M. Kohda, T. Modak and G.-G. Wong, *Enhanced $B \rightarrow \mu \bar{\nu}$ decay at tree level as probe of extra Yukawa couplings*, Phys. Lett. B **800**, 135105 (2020), doi:[10.1016/j.physletb.2019.135105](https://doi.org/10.1016/j.physletb.2019.135105).
- [76] W. Altmannshofer et al., *The Belle II Physics Book*, Prog. Theor. Exp. Phys. 123C01 (2019), doi:[10.1093/ptep/ptz106](https://doi.org/10.1093/ptep/ptz106).
- [77] T. Plehn, *Charged Higgs boson production in bottom-gluon fusion*, Phys. Rev. D **67**, 014018 (2003), doi:[10.1103/PhysRevD.67.014018](https://doi.org/10.1103/PhysRevD.67.014018).
- [78] E. Boos and T. Plehn, *Higgs-boson production induced by bottom quarks*, Phys. Rev. D **69**, 094005 (2004), doi:[10.1103/PhysRevD.69.094005](https://doi.org/10.1103/PhysRevD.69.094005).
- [79] B. Altunkaynak, W.-S. Hou, C. Kao, M. Kohda and B. McCoy, *Flavor changing heavy Higgs interactions at the LHC*, Phys. Lett. B **751**, 135 (2015), doi:[10.1016/j.physletb.2015.10.024](https://doi.org/10.1016/j.physletb.2015.10.024).
- [80] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, J. High Energ. Phys. **07**, 079 (2014), doi:[10.1007/JHEP07\(2014\)079](https://doi.org/10.1007/JHEP07(2014)079).
- [81] A. Alloul, N. D. Christensen, C. Degrande, C. Duhr and B. Fuks, *FeynRules 2.0 — A complete toolbox for tree-level phenomenology*, Comput. Phys. Commun. **185**, 2250 (2014), doi:[10.1016/j.cpc.2014.04.012](https://doi.org/10.1016/j.cpc.2014.04.012).
- [82] R. D. Ball, V. Bertone, S. Carrazza, L. Del Debbio, S. Forte, A. Guffanti, N. P. Hartland and J. Rojo, *Parton distributions with QED corrections*, Nucl. Phys. B **877**, 290 (2013), doi:[10.1016/j.nuclphysb.2013.10.010](https://doi.org/10.1016/j.nuclphysb.2013.10.010).
- [83] T. Sjöstrand, S. Mrenna and P. Skands, *PYTHIA 6.4 physics and manual*, J. High Energ. Phys. **05**, 026 (2006), doi:[10.1088/1126-6708/2006/05/026](https://doi.org/10.1088/1126-6708/2006/05/026).
- [84] J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaître, A. Mertens and M. Selvaggi, *DELPHES 3: a modular framework for fast simulation of a generic collider experiment*, J. High Energ. Phys. **02**, 057 (2014), doi:[10.1007/JHEP02\(2014\)057](https://doi.org/10.1007/JHEP02(2014)057).
- [85] M. Cacciari, G. P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, J. High Energ. Phys. **04**, 063 (2008), doi:[10.1088/1126-6708/2008/04/063](https://doi.org/10.1088/1126-6708/2008/04/063).

- [86] M. Cacciari, G. P. Salam and G. Soyez, *FastJet user manual*, Eur. Phys. J. C **72**, 1896 (2012), doi:[10.1140/epjc/s10052-012-1896-2](https://doi.org/10.1140/epjc/s10052-012-1896-2).
- [87] M. L. Mangano, M. Moretti, F. Piccinini and M. Treccani, *Matching matrix elements and shower evolution for top-pair production in hadronic collisions*, J. High Energ. Phys. **01**, 013 (2007), doi:[10.1088/1126-6708/2007/01/013](https://doi.org/10.1088/1126-6708/2007/01/013).
- [88] J. Alwall et al., *Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions*, Eur. Phys. J. C **53**, 473 (2007), doi:[10.1140/epjc/s10052-007-0490-5](https://doi.org/10.1140/epjc/s10052-007-0490-5).
- [89] G. Aad et al., *Search for supersymmetry at $\sqrt{s} = 13$ TeV in final states with jets and two same-sign leptons or three leptons with the ATLAS detector*, Eur. Phys. J. C **76**, 259 (2016), doi:[10.1140/epjc/s10052-016-4095-8](https://doi.org/10.1140/epjc/s10052-016-4095-8).
- [90] E. Alvarez, D. A. Faroughy, J. F. Kamenik, R. Morales and A. Szykman, *Four tops for LHC*, Nucl. Phys. B **915**, 19 (2017), doi:[10.1016/j.nuclphysb.2016.11.024](https://doi.org/10.1016/j.nuclphysb.2016.11.024).
- [91] J. M. Campbell and R. Keith Ellis, *$t\bar{t}W^\pm$ production and decay at NLO*, J. High Energ. Phys. **07**, 052 (2012), doi:[10.1007/JHEP07\(2012\)052](https://doi.org/10.1007/JHEP07(2012)052).
- [92] J. Campbell, R. Keith Ellis and R. Röntsch, *Single top production in association with a Z boson at the LHC*, Phys. Rev. D **87**, 114006 (2013), doi:[10.1103/PhysRevD.87.114006](https://doi.org/10.1103/PhysRevD.87.114006).
- [93] M. Grazzini, S. Kallweit, D. Rathlev and M. Wiesemann, *$W^\pm Z$ production at hadron colliders in NNLO QCD*, Phys. Lett. B **761**, 179 (2016), doi:[10.1016/j.physletb.2016.08.017](https://doi.org/10.1016/j.physletb.2016.08.017).
- [94] ATLAS Collaboration, *twiki: ATLAS-CMS recommended $t\bar{t}$ cross section predictions*.
- [95] ATLAS Collaboration, *twiki: SM Higgs production cross sections at $\sqrt{s} = 14$ TeV*.
- [96] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, Eur. Phys. J. C **71**, 1554 (2011), doi:[10.1140/epjc/s10052-011-1554-0](https://doi.org/10.1140/epjc/s10052-011-1554-0).
- [97] W.-S. Hou, M. Kohda and T. Modak, *Implications of four-top and top-pair studies on triple-top production*, Phys. Lett. B **798**, 134953 (2019), doi:[10.1016/j.physletb.2019.134953](https://doi.org/10.1016/j.physletb.2019.134953).
- [98] A. M. Sirunyan et al., *Search for production of four top quarks in final states with same-sign or multiple leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV*, Eur. Phys. J. C **80**, 75 (2020), doi:[10.1140/epjc/s10052-019-7593-7](https://doi.org/10.1140/epjc/s10052-019-7593-7).
- [99] M. Buza, Y. Matiounine, J. Smith and W. L. van Neerven, *Charm electroproduction viewed in the variable-flavour number scheme versus fixed-order perturbation theory*, Eur. Phys. J. C **1**, 301 (1998), doi:[10.1007/BF01245820](https://doi.org/10.1007/BF01245820).
- [100] F. Maltoni, G. Ridolfi and M. Ubiali, *Erratum: b-initiated processes at the LHC: a reappraisal*, J. High Energ. Phys. **04**, 095 (2013), doi:[10.1007/JHEP04\(2013\)095](https://doi.org/10.1007/JHEP04(2013)095).
- [101] J. Butterworth et al., *PDF4LHC recommendations for LHC Run II*, J. Phys. G: Nucl. Part. Phys. **43**, 023001 (2016), doi:[10.1088/0954-3899/43/2/023001](https://doi.org/10.1088/0954-3899/43/2/023001).
- [102] G. Aad et al., *Evidence for $t\bar{t}t\bar{t}$ production in the multilepton final state in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, Eur. Phys. J. C **80**, 1085 (2020), doi:[10.1140/epjc/s10052-020-08509-3](https://doi.org/10.1140/epjc/s10052-020-08509-3).

- [103] W.-S. Hou, T.-H. Hsu and T. Modak, *Constraining the $t \rightarrow u$ flavor changing neutral Higgs coupling at the LHC*, Phys. Rev. D **102**, 055006 (2020), doi:[10.1103/PhysRevD.102.055006](https://doi.org/10.1103/PhysRevD.102.055006).
- [104] G. Aad et al., *Search for squarks and gluinos in final states with same-sign leptons and jets using 139 fb^{-1} of data collected with the ATLAS detector*, J. High Energ. Phys. **06**, 046 (2020), doi:[10.1007/JHEP06\(2020\)046](https://doi.org/10.1007/JHEP06(2020)046).
- [105] M. Aaboud et al., *Search for new phenomena in events with same-charge leptons and b-jets in pp collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector*, J. High Energ. Phys. **12**, 039 (2018), doi:[10.1007/JHEP12\(2018\)039](https://doi.org/10.1007/JHEP12(2018)039).
- [106] M. Aaboud et al., *Search for heavy Higgs bosons A/H decaying to a top quark pair in pp collisions at $\sqrt{s} = 8 \text{ TeV}$ with the ATLAS detector*, Phys. Rev. Lett. **119**, 191803 (2017), doi:[10.1103/PhysRevLett.119.191803](https://doi.org/10.1103/PhysRevLett.119.191803).
- [107] A. M. Sirunyan et al., *Search for heavy Higgs bosons decaying to a top quark pair in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$* , J. High Energ. Phys. **04**, 171 (2020), doi:[10.1007/JHEP04\(2020\)171](https://doi.org/10.1007/JHEP04(2020)171).
- [108] Atlas Collaboration, *Search for charged Higgs bosons decaying into a top-quark and a bottom-quark at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector*, Tech. Rep. ATLAS-CONF-2020-039, CERN, Geneva (2020).
- [109] A. M. Sirunyan et al., *Search for charged Higgs bosons decaying into a top and a bottom quark in the all-jet final state of pp collisions at $\sqrt{s} = 13 \text{ TeV}$* , J. High Energ. Phys. **07**, 126 (2020), doi:[10.1007/JHEP07\(2020\)126](https://doi.org/10.1007/JHEP07(2020)126).