

Strings and membranes from \mathcal{A} -theory five brane

Machiko Hatsuda^{1,2*}, Ondřej Hulík³, William D. Linch⁴,
Warren D. Siegel^{5,6}, Di Wang⁶ and Yu-Ping Wang⁶

¹ Department of Radiological Technology, Faculty of Health Science,
Juntendo University, Yushima, Bunkyo-ku, Tokyo 113-0034, Japan

² KEK Theory Center, High Energy Accelerator Research Organization,
Tsukuba, Ibaraki 305-0801, Japan

³ Institute for Mathematics Ruprecht-Karls-Universität Heidelberg,
69120 Heidelberg, Germany

⁴ Thomas Jefferson High School for Science and Technology, Alexandria, VA 22312, USA
⁵ C. N. Yang Institute for Theoretical Physics. Stony Brook, NY 11794, USA

⁶ Department of Physics, SUNY Stony Brook University, Stony Brook, NY 11794, USA

★ mhatsuda@juntendo.ac.jp

Abstract

The \mathcal{A} -theory takes U-duality symmetry as a guiding principle, with the $SL(5)$ U-duality symmetry being described as the world-volume theory of a 5-brane. Furthermore, by unifying the 6-dimensional world-volume Lorentz symmetry with the $SL(5)$ spacetime symmetry, it extends to $SL(6)$ U-duality symmetry. The $SL(5)$ spacetime vielbein fields and the 5-brane world-volume vielbein fields are mixed under the $SL(6)$ U-duality transformation. We demonstrate that consistent sectionings of the $SL(6)$ $\mathcal{A}5$ -brane world-volume Lagrangian yield Lagrangians of the \mathcal{T} -string with $O(D,D)$ T-duality symmetry, the conventional string, the $\mathcal{M}5$ -brane with $GL(4)$ duality symmetry, and the non-perturbative $M2$ -brane in supergravity theory. The $GL(4)$ covariant Lagrangian of the $\mathcal{M}5$ -brane derived in this manner is a new, perturbatively quantizable theory.



Copyright M. Hatsuda *et al.*

This work is licensed under the Creative Commons

[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Published by the SciPost Foundation.

Received 2025-01-27

Accepted 2025-06-17

Published 2025-07-07

doi:[10.21468/SciPostPhys.19.1.009](https://doi.org/10.21468/SciPostPhys.19.1.009)



Check for
updates

Contents

1	Introduction	2
1.1	An overview of \mathcal{A} -theory formalism	2
1.2	Summary	4
1.3	Dimensional reduction procedure	5
2	Theories with manifest duality symmetries and sectionings	8
2.1	Diamond diagrams	8
2.2	Representations	8
2.3	Constraints and sectionings	11

3	$\mathcal{A}5$-brane Lagrangians	16
3.1	$\mathcal{A}5$ -brane Lagrangian with $SL(5)$ U-duality symmetry	16
3.2	World-volume covariant $\mathcal{A}5$ -brane Lagrangian with $SL(6)$ duality symmetry	17
4	Lagrangian of D-dimensional string from $O(D,D)$ \mathcal{T}-string	18
4.1	$O(D,D)$ \mathcal{T} -string	18
4.2	String from $O(D,D)$ \mathcal{T} -string	20
5	Lagrangians of string via \mathcal{T}-string from $\mathcal{A}5$-brane	22
5.1	\mathcal{T} -string from $\mathcal{A}5$ -brane	22
5.2	String from \mathcal{T} -string	24
6	Lagrangians of M2-brane via $\mathcal{M}5$-brane from $\mathcal{A}5$-brane	25
6.1	$\mathcal{M}5$ -brane from $\mathcal{A}5$ -brane	25
6.2	Non-perturbative M2-brane from $\mathcal{M}5$ -brane	29
7	Discussion	31
	References	32

1 Introduction

1.1 An overview of \mathcal{A} -theory formalism

Superstring theory is regarded as a strong candidate for a unified theory that encompasses all four fundamental interactions. However, rather than a single theory, it currently exists in the form of six distinct theories: five superstring theories and M-theory. These six theories are intricately connected via S-duality and T-duality, forming a hexagonal network of dualities. Each of these six theories is defined on its own characteristic brane, and together they provide complementary descriptions of the broader structure of superstring theory. A key open question is what kind of theoretical framework can provide a unified formulation of these six theories.

S-duality and T-duality are unified as U-duality, whose structure reflects the group-theoretic inclusiveness of these dualities. The symmetry group of a theory with manifest S-duality is given by $GL(D+1)$ which extends the spacetime diffeomorphism group $GL(D)$ by incorporating the S-duality group $SL(2)$ as a subgroup. $GL(D+1)$ and the T-duality symmetry $O(D,D)$ are embedded within the exceptional group E_{D+1} which serves as the symmetry group of U-duality. U-duality is therefore expected to relate all six superstring theories in a coherent manner. A theory that explicitly manifests this U-duality symmetry is referred to as “ \mathcal{A} -theory”, and major objective is to construct a perturbative formulation of this theory that facilitates a systematic analysis of its quantum aspects (see [1] for review).

In 1995, Witten proposed M-theory as a strong coupling limit of the type IIA superstring theory via S-duality [2]. The low-energy effective theory of M-theory is 11-dimensional supergravity, whose diffeomorphism symmetry is $GL(11)$, combining the 10-dimensional diffeomorphism and the $SL(2)$ S-duality symmetry. We refer to a world-volume theories that exhibit manifest $GL(D+1)$ symmetry as \mathcal{M} -theory. In 1996, Vafa proposed F-theory as a framework related to the type IIB superstring theory via S-duality [3]. For further discussion, see, for example [4].

In 1993, Siegel proposed a string theory guided by the T-duality symmetry $O(D,D)$ [5–7]. This framework employs the idea of doubling spacetime coordinates, where the 2D-dimensional spacetime coordinates are treated as vectors in the $O(D,D)$ representation [8]. The geometry generated by the $O(D,D)$ covariant current algebra is the stringy gravity theory with B -field, where gauge fields are parameters of the coset $O(D,D)$ over the doubled Lorentz group. We refer to this as \mathcal{T} -theory and the T-duality covariant string as \mathcal{T} -string. Later Hull, Zwiebach, and Hohm proposed a theory of $O(D,D)$ covariant background fields, known as Double Field Theory (DFT) [9, 10]. This theory serves as the low-energy effective gravitational description of \mathcal{T} -string theory. In order to consistently reduce the doubled 2D-dimensional spacetime to the physical D -dimensional spacetime, the section condition was employed, which corresponds to the zero-mode of the Virasoro constraint $S=0$. For detailed reviews, see [11, 12]. The \mathcal{T} -theory was extended to incorporate $\mathcal{N} = 2$ supersymmetry including Ramond–Ramond gauge fields based on the doubled non-degenerate super-Poincaré group [13–17]. All bosonic component fields represent supersymmetrized $O(D,D)$ while fermions represent only $O(D-1,1)^2$. Incorporating S-duality requires extending the symmetry to the exceptional U-duality group. Consequently, the exceptional group E_{D+1} is expected to be related to a subgroup of the doubled non-degenerate super-Poincaré group.

Siegel, Linch and Polacek subsequently proposed brane theories guided by the U-duality symmetry (E_{D+1}) [13, 18, 19]. Since the representation of exceptional groups varies with dimension, the theory is labeled by the spacetime dimension D when reduced to string theory. This is called $D = D$ \mathcal{A} -theory. According to the classification of Lie algebras, removing a specific node in the Dynkin diagram of E_{D+1} reduces it to the $GL(D+1)$ Dynkin diagram. This corresponds to reducing the spacetime dimensions of \mathcal{A} -theory using the Virasoro constraint, recovering the aforementioned \mathcal{M} -theory. Conversely, removing another node in the E_{D+1} Dynkin diagram reduces it to the $O(D,D)$ Dynkin diagram. This corresponds to reducing the world-volume dimensions of \mathcal{A} -theory using the Gauß law constraint $\mathcal{U}=0$, recovering the aforementioned \mathcal{T} -theory. As a consequence of these dual reductions, \mathcal{A} -theory is consistently described by branes covariant under the exceptional group. Both the spacetime coordinates and world-volume coordinates of the branes are representations of the exceptional group, ensuring that the brane current algebra is covariant under the exceptional group. Moreover, a perturbative Lagrangian describing these branes on their world-volume has been constructed. In the previous papers [18–24, 24–31] we refer these models as “F-theory”, but we have renamed our formulation from “F-theory” to \mathcal{A} -theory in our most recent work [1], in order to present it as a general framework that includes all spacetime dimensions and accommodates all duality symmetries.

Exceptional Field Theory (EFT) applies DFT concepts to exceptional groups [32–44]. The symmetry of exceptional groups was initially discovered as a partial dimensional symmetry of background fields in 11-dimensional supergravity [45, 46]. The generalized diffeomorphism in EFT is characterized by the “Y-tensor”, which reflects the structure of the exceptional group. This Y-tensor, $Y^{MN}{}_{PQ}$, is related to the group invariant metric in \mathcal{A} -theory, denoted by η^{MNm} , through the relation $Y^{MN}{}_{PQ} = \eta^{MNm} \eta_{PQm}$ where M, N, \dots are the spacetime indices and m is the world-volume index. These indices correspond to different representations of the exceptional group. The origin of the Y-tensor lies in the Schwinger term of the current algebra, $\{\triangleright_M(\sigma), \triangleright_N(\sigma')\} = \eta_{MNm} \partial^m \delta(\sigma - \sigma')$, where the world-volume derivative ∂^m is defined through the commutator with the Virasoro constraint, $S^m = \frac{1}{2} \eta^{MNm} \triangleright_M \triangleright_N$. The section condition given by the Y-tensor, $Y^{MN}{}_{PQ} \partial_M \partial_N = 0$, is related to the zero-mode of the Virasoro constraint in \mathcal{A} -theory. Specifically, the zero-mode component of the constraint S^m takes the form $S^m|_{0\text{-modes}} = \eta^{MNm} \partial_M \partial_N = 0$, establishing a direct connection between the section condition in EFT and the Virasoro structure of \mathcal{A} -theory. Active research continues on expressing exceptional groups through brane current algebra [47–54].

This paper focuses on the 5-brane that describes D=3 \mathcal{A} -theory with $SL(5)$ U-duality symmetry. In the case of D=3, the theory provides a nontrivial yet tractable example that includes various types of branes and permits explicit computations. The Virasoro algebra, when extended to incorporate brane degrees of freedom, involves Gauß law-type constraints that are intrinsic to the brane. These constraints facilitate the dimensional reduction of the brane world-volume. Such an extended Virasoro algebra serves as a useful prototype for generalization to higher dimensions. Furthermore, the construction of the Lagrangian in this framework constitutes a significant milestone toward extending the formulation to higher-dimensional cases. We clarify how the usual string, and \mathcal{T} -string and membrane of 11-dimensional supergravity (M2) emerge. Specifically, we derive the Lagrangians for the conventional non-perturbative M2-brane, the $O(D,D)$ -covariant \mathcal{T} -string, and the conventional string from the \mathcal{A} -theory 5-brane ($\mathcal{A}5$) Lagrangian.

1.2 Summary

In this paper, we focus on the $SL(5)$ U-duality symmetry and clarify the relation between the \mathcal{A} -theory five-brane ($\mathcal{A}5$ -brane) and conventional branes. The $\mathcal{A}5$ -brane is a 5-brane that exhibits manifest $SL(5)$ U-duality symmetry, and it is described by a perturbative Lagrangian described using the $SL(5)$ rank-two antisymmetric tensor coordinate [1, 18, 30]. The $\mathcal{A}5$ -brane theory is reduced, through the duality reduction procedure, to the \mathcal{T} -string with $O(3,3)$ T-duality symmetry, the $\mathcal{M}5$ -brane with $GL(4)$ duality symmetry, and the String with $GL(3)$ symmetry. The interrelation among these theories forms a diamond-shaped diagram. In this paper, we present the sectioning procedure along this diamond contour. Additionally, we detail reduction procedures from the \mathcal{T} -string Lagrangian to the conventional string Lagrangian, and from the $\mathcal{M}5$ -brane Lagrangian to the conventional non-perturbative M2-brane Lagrangian [55]. We further generalize the dimensional reduction of S-duality, as applied to 11-dimensional supergravity and its relation to type IIA supergravity [2], to the case of T-duality, as discussed in subsection 1.3. For S-duality, $\lambda_{\text{string}} \leftrightarrow 1/\lambda_{\text{string}}$, the dimensional reduction is realized by taking the limit $\lambda_{\text{string}} \ll 1$ in the metric. For T-duality, characterized by $R/\sqrt{\alpha'} \leftrightarrow \sqrt{\alpha'}/R$ with string length $l_{\text{string}} = \sqrt{\alpha'}$, the dimensional reduction is achieved by taking the large-radius limit $R \gg \sqrt{\alpha'}$, such that the $O(D,D)$ spacetime for the \mathcal{T} -string reduces to the conventional D-dimensional spacetime. Finally, we propose a perturbative $\mathcal{M}5$ -brane Lagrangian in a supergravity background derived from the $\mathcal{A}5$ -brane, as given in equation (109).

In section 2 the relationships among the $\mathcal{A}5$ -brane, $\mathcal{M}5$ -brane, \mathcal{T} -string, and String theories are elucidated through diamond diagrams based on duality symmetries. The diamond diagram represents a contour in the U-duality plane, which is parametrized by two quantities: the string coupling and a scale defined by the string length. The branes are described in terms of field strengths, where the spacetime coordinates possess the gauge symmetry generated by the Gauß law constraint. These field strengths and their associated gauge parameters, along with the world-volume and spacetime coordinates, transform as representations under the relevant duality symmetries. The world-volume diffeomorphism is generated by the Virasoro constraints $\mathcal{S} = 0$. Dimensions to reduce are determined by solving the Virasoro constraints $\mathcal{S} = 0$ for spacetime and by solving the Gauß law constraints $\mathcal{U} = 0$ for world-volume.

In section 3 both the $SL(5)$ and the $SL(6)$ covariant Lagrangians of the $\mathcal{A}5$ -brane are given, where the $SL(6)$ manifests the 5-brane world-volume Lorentz symmetry. Duff and Lu [56, 57] showed that the membrane theory exhibits the $SL(5)$ duality symmetry by the Gaillard-Zumino approach. In general, the symmetry of Lagrangian formulation is larger than that of the corresponding Hamiltonian formulation. In \mathcal{A} -theory, the U-duality symmetry in the Hamiltonian formulation, G-symmetry, is enhanced to a novel duality symmetry in the Lagrangian formulation, A-symmetry. This symmetry enhancement in higher-dimensional cases ($D < 6$) is

summarized on page 6 of [22] and page 14 of [1]. It was shown that the brane world-volume metric is also transformed conformally under the $SL(5)$ duality transformation as well as the spacetime background fields in [56, 57]. This mixing between spacetime and world-volume is a manifestation of the extended $SL(6)$ duality symmetry transformation. The $SL(6)$ vielbein includes both the $SL(5)$ spacetime vielbein and the 6-dimensional world-volume vielbein, so the spacetime and world-volume are mixed under the new duality symmetry $SL(6)$. The $SL(6)$ formulation is useful to reduce to other branes: Since the string world-sheet directions and the spacetime directions are direct sum, the $SL(6)$ vielbein is in a block diagonal form as shown in subsection 5.1. On the other hand, the brane world-volume directions share the spacetime directions unlike the string as shown in subsection 6.1.

In section 4, we begin with the $O(D,D)$ string Hamiltonian and apply the double zweibein method [58, 59] to derive the \mathcal{T} -string Lagrangian. We then present the reduction procedure from the $O(D,D)$ \mathcal{T} -string Lagrangian to the conventional string Lagrangian in D dimensions, following an approach analogous to that in subsection 1.3. It is shown that the Wess–Zumino term can be obtained by adding a total derivative term.

In section 5, we start from the $\mathcal{A}5$ -brane Lagrangian and present the reduction procedure leading to the \mathcal{T} -string Lagrangian. The $O(D,D)$ background gauge field is reformulated using $SL(4)$ tensor indices in such a way that it couples naturally to the $SL(4)$ tensor coordinates of the \mathcal{T} -string. Subsequently, by applying the procedure described in section 4, we derive the conventional string Lagrangian.

In section 6 we begin with the $SL(6)$ covariant $\mathcal{A}5$ -brane Lagrangian which leads to a new perturbative $\mathcal{M}5$ -brane Lagrangian. We further reduce it to the conventional $\mathcal{M}2$ -brane Lagrangian. The “perturbative” $\mathcal{M}5$ -brane Lagrangian is formulated as a bilinear expression in terms of currents, while the “non-perturbative” $\mathcal{M}2$ -brane Lagrangian comprises the sum of the Nambu–Goto and Wess–Zumino terms. The dimensional reduction from the $\mathcal{M}5$ -brane to the $\mathcal{M}2$ -brane is implemented via the “non-perturbative projection” $\partial^m = \epsilon^{ij} \partial_j x^m \partial_i$, in (114) and the gauge fixing of the world-volume metric in (120). The Nambu–Goto Lagrangian is obtained by the gauge choice of the world-volume vielbein, while the Wess–Zumino term is obtained by adding the total derivative term.

1.3 Dimensional reduction procedure

In [2], it was pointed out that under an S-duality transformation between the 10-dimensional type IIA theory and the 11-dimensional supergravity theory, the structure of the supersymmetry algebra remains invariant, although the interpretation of the central charge changes. The global superalgebra, involving supercharges Q and Q' of the opposite chirality, 10-dimensional momenta P , and a central charge W , is given by $\{Q, Q'\} \sim P \sim \{Q', Q'\}$, $\{Q_\alpha, Q'_\beta\} \sim \delta_{\alpha\beta} W$. The central charge W is interpreted as the Ramond–Ramond (RR) D0-brane charge in 10 dimensions, and as the momentum in the 11th dimension in 11-dimensional supergravity. The 11-dimensional spacetime reduces into the 10-dimensional spacetime in the weak coupling limit $e^{2\phi} \ll 1$,

$$ds_{11}^2 = g_{mn}^{[10]} dx^m dx^n + e^{2\phi} (dy - A_m dx^m)^2 \xrightarrow[\text{reduction}]{\text{dimensional}} ds_{10}^2 = g_{mn}^{[10]} dx^m dx^n, \quad (1)$$

with the 11-th dimensional coordinate y and the string coupling $\lambda_{\text{string}} = e^{3\phi/2}$. The 11-dimensional momentum W is maintained as the D0-brane charge in the 10-dimensional IIA theory after the dimensional reduction.

This framework is generalized to incorporate T-duality. We compare the superalgebra of the $2D$ -dimensional \mathcal{T} -string theory, which exhibits manifest T-duality, with the type II superalgebra of conventional string theory in D dimensions. The global type II superalgebra involves two supercharges, Q and Q' , and the D -dimensional momentum P , and is expressed

as $\{Q, Q\} \sim (P + \tilde{P})$, $\{Q', Q'\} \sim (P - \tilde{P})$, where \tilde{P} is a central charge. In D dimensions, \tilde{P} is interpreted as an NS–NS charge, while in $2D$ dimensions, it corresponds to the additional momenta associated with the extended spacetime. By restoring the α' dependence in the $O(D, D)$ momentum–winding vector, $(p_m, \frac{1}{\alpha'} \partial_\sigma x^m) \rightarrow (p_m, \frac{1}{\alpha'} \tilde{p}^m)$, the canonical conjugate coordinates become $(x^m, \alpha' y_m)$. At small compactification scales ($R \ll \sqrt{\alpha'}$), the winding modes become light and are readily excited, whereas at large scales ($R \gg \sqrt{\alpha'}$), they become heavy and only the momentum modes remain dynamically relevant. Manifest T-duality is broken by choosing a specific background such that the $2D$ -dimensional spacetime effectively reduces to the D -dimensional one in the limit $R \gg \sqrt{\alpha'}$:

$$ds_{2D}^2 = g_{mn} dx^m dx^n + \alpha'^2 (dy_m - dx^l B_{lm}) g^{mn} (dy_n - dx^k B_{kn}) \xrightarrow[\text{reduction}]{\text{dimensional}} ds_D^2 = g_{mn} dx^m dx^n. \quad (2)$$

Here, y_m denotes the additional D -dimensional coordinates, and B_{mn} is the NS–NS gauge field. The additional momenta in the extended dimensions are preserved as NS–NS charges after the dimensional reduction.

This dimensional reduction procedure corresponds to the gauge fixing of the dimensional reduction constraint which is the first class constraint in Hamiltonian formulation. The dimensional reduction constraint and the gauge fixing to reduce the conventional string are discussed in [60] for a flat space case. The dimensional reduction constraint is the y component of the symmetry generator, $\tilde{\mathfrak{P}}_y = 0$. The gauge fixing condition $\partial_\sigma y = 0$ reduces the set of conventional string operators, the physical momentum $P_x \neq 0$ and left/right covariant derivatives $P_x \pm \partial_\sigma x$. In Lagrangian formulation the momentum is replaced by $P_x = \partial L / \partial \dot{X}$. It is generalized to the brane case, and then the dimensional reduction constraint turns out to be the Virasoro constraint in which one of the momenta is replaced by the 0-mode [19]. For zero-mode momenta p_x , p_y and momenta including all modes P_x , P_y , the dimensional reduction constraint $\tilde{\mathfrak{P}}_y = 0$ is expressed by the Virasoro operator as $p_y \cdot P_x + p_x \cdot P_y = 0 \rightarrow P_y = 0$.

Although the equation of motion derived from the doubled Lagrangian, when combined with the self-duality condition, coincides with that obtained from the original Lagrangian, the self-duality condition causes the doubled Lagrangian to vanish [61]. In particular, the self-duality condition $\partial_\mu x = \epsilon_{\mu\nu} \partial^\nu y$ reduces the Lagrangian of the $O(D, D)$ \mathcal{T} -string in flat space to zero, as follows

$$L = \frac{1}{2} (\dot{x}^2 - x'^2 + \dot{y}^2 - y'^2) \xrightarrow{\text{selfduality}} 0.$$

It is also mentioned that the naive section $y = 0$ the \mathcal{T} -string Lagrangian in curved background does not reduce to the expected string Lagrangian in curved background as

$$L = \frac{1}{2} (\partial_+ x^m \partial_+ y_m) \begin{pmatrix} g_{mn} - B_{ml} g^{lk} B_{kn} & -B_{ml} g^{ln} \\ g^{ml} B_{ln} & g^{mn} \end{pmatrix} \begin{pmatrix} \partial_- x^n \\ \partial_- y_n \end{pmatrix} \xrightarrow{y=0} \frac{1}{2} \partial_+ x^m (g_{mn} - B_{ml} g^{lk} B_{kn}) \partial_- x^n.$$

The following points are also noteworthy. Integrating out the $(dy + \dots)^2$ term is possible in the case of a constant background. However, for a general non-constant background, the path integral over dy in $\exp[-\int (dy + \dots) g(dy + \dots)]$ yields a Jacobian factor \sqrt{g} in the path integral measure, which in turn generates an additional term in the effective action. Using the equation of motion is again valid in the constant background case, but it does not reproduce the conventional string Lagrangian when the background fields $g(x, y)$ and $B(x, y)$ are non-constant. Furthermore, imposing both conditions $\partial_+ y - \partial_+ x B = 0$ and $\partial_- y - \partial_- x B = 0$ is inconsistent, since the integrability condition is violated in curved backgrounds where $[\partial_+, \partial_-]y \neq 0$. Several studies have been devoted to refining the reduction to the conventional string Lagrangian, resulting in a variety of interesting approaches [43, 50, 61–64].

Instead we propose the reduction procedure from \mathcal{T} -string Lagrangian to the conventional string Lagrangian: (1) adding the total derivative term $-\partial_\mu(\epsilon^{\mu\nu}x^m\partial_\nu y_m)$ to derive the Wess-Zumino term, then (2) the dimensional reduction (2) as

$$L = \frac{1}{2}(\partial_+ x^m \partial_+ y_m) \begin{pmatrix} g_{mn} - B_{ml}g^{lk}B_{kn} & -B_{ml}g^{ln} \\ g^{ml}B_{ln} & g^{mn} \end{pmatrix} \begin{pmatrix} \partial_- x^n \\ \partial_- y_n \end{pmatrix} - \partial_\mu(\epsilon^{\mu\nu}x^m\partial_\nu y_m) \quad (3)$$

$$\xrightarrow[\text{reduction (1.2)}]{\text{dimensional}} \partial_+ x^m(g_{mn} + B_{mn})\partial_- x^n.$$

This is the expected string Lagrangian up to the normalization factor two which can be absorbed by the Lagrange multiplier. The section conditions of spacetime fields $\Phi(x, y)$ are consistent with the Lagrangian where the section $y = 0$ can be chosen as $\Phi(x)$.

This procedure is similar to the usual dimensional reduction where the reduction is done in the local flat Lorentz coordinate. i.e. Suppose that we have a line element $dx^A \equiv dx^M E_M^A$. We decompose the doubled coordinate dx^A into dx^a and dy_a in the local Lorenz frame, and then discard $(dy_a)^2$. Since the metric ($\hat{\eta}$ -tensor) in local flat spacetime is already diagonal, in practice we can just apply this reduction by deleting certain blocks of $\hat{\eta}$ -tensor similar to (2).

The main purpose of this paper is to carry out the above reduction procedure in several specific cases. In general, however, the procedure can be schematically summarized as follows.

1. We start with the current algebra defined on an extended space of coordinates, where both momentum and winding modes have their corresponding conjugate coordinates. The Hamiltonian is written as a sum of self-dual and anti-self-dual constraints: $H = g\mathcal{H} + \tilde{g}\tilde{\mathcal{H}} + s_m\mathcal{S}^m + \tilde{s}_m\tilde{\mathcal{S}}^m + Y^m\mathcal{U}_m$, where $\mathcal{H}, \tilde{\mathcal{H}}$ are the τ component of the Virasoro constraints and its dual counterpart, $\mathcal{S}^m, \tilde{\mathcal{S}}^m$ are the σ^m components of the Virasoro constraints and its dual counterpart. \mathcal{U}_m is the Gauß law constraint specific to branes.
2. The Lagrangian is obtained via a Legendre transformation of the Hamiltonian H . This has been performed in previous works for various theories [1]. Schematically, the Lagrangian takes the form $L = \Phi J_{\text{SD}} \cdot \hat{\eta} \cdot J_{\text{SD}} + \Lambda \cdot J_{\text{SD}} \cdot \hat{\eta} \cdot J_{\text{SD}} + \dots$, where $J_{\text{SD}/\text{SD}}^A$ are the selfdual and anti-selfdual currents. They are coupled with vielbein, and thus they have flat indices. Φ and Λ are Lagrange multipliers which are functions of $g, s^m, \tilde{g}, \tilde{s}^m$. One can gauge fix $\Lambda = 0$ by the suitable choice of original parameters.
3. Separate coordinates and currents into the physical part and the auxiliary part as $X^M \rightarrow x^m, y^\mu$, and $J^A \rightarrow J^a, J^\alpha$ where x^m represents the physical coordinates for the target string or brane theory, and y^μ denotes auxiliary coordinates. Dimensional reduction is then performed according to equation (2).
4. The reduced Lagrangian $L' = J^a_{\text{SD}} \hat{\eta}_{ab} J^b_{\text{SD}}$ can be shown to reproduce the desired string or brane action, up to the absence of the Wess–Zumino (WZ) term. We find that adding a total derivative term to the Lagrangian restores the WZ term

$$L + \text{Total derivative} = J^a_{\text{SD}} \hat{\eta}_{ab} J^b_{\text{SD}} + \tilde{J}^\alpha_{\text{SD}} \hat{\eta}_{\alpha\beta} \tilde{J}^\beta_{\text{SD}} + L_{\text{WZ}}.$$

Here, the current \tilde{J}^α is modified by the addition of the total derivative term and is subsequently eliminated through the dimensional reduction (2). This procedure yields the correct WZ term.




Figure 1: G -symmetries of $D = D$ theories and spacetime dimensions.

2 Theories with manifest duality symmetries and sectionings

2.1 Diamond diagrams

The duality web of the G -symmetry in \mathcal{A} -theory is represented by the diamond diagram shown in Fig. 1, as studied in [19]. The G -symmetry, associated with the coset group G/H , plays the role of a duality symmetry. The coset parameter serves as the gauge field of the duality-covariant geometry, incorporating the spacetime vielbein as well as the NS–NS and R–R gauge fields of superstring theory.

The relationships among these duality groups are illustrated using Dynkin diagrams, as discussed in [1]. Removing a single node from the Dynkin diagram of $E_{D+1(D+1)}$ reduces it to that of either $GL(D+1)$ or $O(D,D)$, depending on which node is removed. Further removing one more node from the Dynkin diagram of $GL(D+1)$ or $O(D,D)$ leads to that of $GL(D)$.

In this paper we focus on $D=3$ case where the G -symmetry is $SL(5)$ and the diamond diagram becomes Fig. 2. This $SL(5)$ duality symmetry is enlarged to $SL(6)$ for the $(5+1)$ -dimensional world-volume covariance in Lagrangian [1]. We named this enlarged symmetry “ A -symmetry”. This \mathcal{A} -theory unifies the spacetime and the world-volume, in a sense that the coset parameter of $A/L=SL(6)/GL(4)$ includes not only the spacetime vielbein field but also the world-volume vielbein field.

In this paper, we focus on the $D=3$ case, where the G -symmetry is $SL(5)$ and the diamond diagram corresponds to Fig. 2. This $SL(5)$ duality symmetry is further enhanced to $SL(6)$ in order to accommodate the $(5+1)$ -dimensional world-volume covariance in the Lagrangian formulation [1]. We refer to this enlarged symmetry as the “ A -symmetry”. The resulting \mathcal{A} -theory unifies the spacetime and world-volume structures, in the sense that the coset parameter of $A/L=SL(6)/GL(4)$ includes not only the spacetime vielbein field but also the world-volume vielbein field. We note that $H=SO(D)$ is used instead of $SO(D-1,1)$ for simplicity. Consequently, a Wick rotation is required to properly account for the time component in this section and elsewhere.

2.2 Representations

In duality covariant theories, spacetime and world-volume coordinates transform as representations of the duality symmetry (A -symmetry or G -symmetry), which determines the world-volume dimension. The Gauß law constraint generates gauge symmetry of the duality covariant spacetime coordinate, making the brane current correspond to a field strength.




Figure 2: A- and G-symmetries of D = 3 theories and spacetime dimensions.

The D=3 \mathcal{M} , \mathcal{T} , \mathcal{S} -theories are obtained from the D=3 \mathcal{A} -theory [18, 30]. We list representations of duality groups in Tab. 1; the world-volume derivative ∂^m , the gauge parameter λ^m , the spacetime coordinate X^M , and the field strength (the current) $F_M = \eta_{MNm} \partial^m X^N$ ($J_\mu^M = \partial_\mu X^M$, $\mu = (\tau, \sigma)$). η_{MNm} is the G -symmetry invariant tensor which enters the current algebra, where the $SL(5)$ invariant metric is $\eta_{MNI} = \eta_{m_1 m_2 n_1 n_2 l} = \epsilon_{m_1 m_2 n_1 n_2 l}$. The world-volume dimension of \mathcal{M} -theory is still $1 \oplus 5$ where four dimensions are embedded in the 4 spacetime x^m and one dimension is embedded in the internal space, so we denote as $1 \oplus 4(5)$.

The field strengths and currents together with the gauge transformations are given concretely as follows.

1. $\mathcal{A}5$ -brane field strengths

(a) World-volume covariant $\mathcal{A}5$ -brane field strength

The $SL(6)$ A -symmetry covariant \mathcal{A} -theory is described by a 5-brane with the manifest $SL(6)$ new duality symmetry which manifests 6-dimensional world-volume Lorentz symmetry, namely world-volume covariant $\mathcal{A}5$ -brane.

$$F^{\hat{m}\hat{n}\hat{p}} = \frac{1}{2} \partial^{[\hat{m}} X^{\hat{n}\hat{p}]}, \quad \delta_\lambda X^{\hat{m}\hat{n}} = \partial^{[\hat{m}} \lambda^{\hat{n}]}, \quad \hat{m} = 0, 1, \dots, 5. \quad (4)$$

(b) $\mathcal{A}5$ -brane field strength

The $SL(5)$ G -symmetry covariant \mathcal{A} -theory is described by a 5-brane with manifest $SL(5)$ U-duality symmetry, namely $\mathcal{A}5$ -brane.

$$\left\{ \begin{array}{l} F_\tau^{mn} = \dot{X}^{mn} - \partial^{[m} Y^{n]}, \\ F_{\sigma; m_1 m_2} = \frac{1}{2} \epsilon_{m_1 \dots m_5} \partial^{m_3} X^{m_4 m_5}, \\ m = 1, \dots, 5, \end{array} \right\} \quad \left\{ \begin{array}{l} \delta_\lambda X^{mn} = \partial^{[m} \lambda^{n]}, \\ \delta_\lambda Y^m = \dot{\lambda}^m - \partial^m \lambda^0. \end{array} \right. \quad (5)$$

2. $\mathcal{M}5$ -brane field strength

The $GL(4)$ \mathcal{M} -theory is described by a 5-brane with the manifest $GL(4)$ duality symmetry, namely $\mathcal{M}5$ -brane. We focus only on 4-dimensional subspace of the 5-dimensional

Table 1: Representations of the duality groups of theories.

Theories Groups	World-volumes ∂	Gauges λ	Spacetimes X	Field strengths F
\mathcal{A} -theory SL(6)	6 $\partial^{\hat{m}}, \hat{m}=0, \dots, 5$	6 $\lambda^{\hat{m}}$	15 $X^{\hat{m}\hat{n}}$	20 $F^{\hat{m}\hat{n}\hat{p}}$
\mathcal{A} -theory SL(5)	$1 \oplus 5$ $\partial^0, \partial^m, m=1, \dots, 5$	$1 \oplus 5$ λ^0, λ^m	$10 \oplus 5$ X^{mn}, Y^m	$10 \oplus 10'$ $F_{\tau}{}^{mn}, F_{\sigma mn}$
\mathcal{M} -theory GL(4)	$1 \oplus 4(5)$ $\partial^0, \partial^{\underline{m}}, \underline{m}=1, \dots, 4$	$1 \oplus 4$ $\lambda^0, \lambda^{\underline{m}}$	$4 \oplus 1$ $x^{\underline{m}}, Y$	$4 \oplus 6$ $F_{\tau}{}^{\underline{m}}, F_{\sigma \underline{m}\underline{n}}$
\mathcal{T} -theory O(3, 3)	$1 \oplus 1$ $\partial^0, \partial^{\sigma}, \bar{m}=1, 2, 3$	0	$3 \oplus 3'$ $x^{\bar{m}}, y^{\bar{m}\bar{n}}$	$3 \oplus 3 \oplus 3' \oplus 3'$ $J_{\tau}{}^{\bar{m}}, J_{\sigma \bar{m}\bar{n}}; J_{\tau}{}^{\bar{m}}, J_{\sigma \bar{m}},$
S-theory GL(3)	$1 \oplus 1$ $\partial^0, \partial^{\sigma}, \bar{m}=1, 2, 3$	0	3 $x^{\bar{m}}$	$3 \oplus 3$ $J_{\tau}{}^{\bar{m}}, J_{\sigma \bar{m}\bar{n}}$

world-volume which is embedded in the 4-dimensional spacetime. This $\mathcal{M}5$ -brane extends over both the main space (i.e., the duality-covariant space) and the internal space. Four of its world-volume directions lie in the main space while the remaining directions lie in the internal space, specifically one world-volume direction in the Hamiltonian formalism, or two in the Lagrangian formalism. Considering the critical string action in the full spacetime structure is an interesting subject, although it lies beyond the scope of the present discussion. The relationship between the main space and the internal space is schematically illustrated in Figure 2 the “slug diagram” (see page 27 in [20] or page 14 in [1]). In the case of $D = 3$ the main space coordinate is represented by a bispinor $X^{\alpha\beta}$, and the world-volume coordinate by an antisymmetric bispinor $\sigma^{[\alpha\beta]}$, with $\alpha = 1, \dots, 4$. The internal space coordinate is given by a bispinor $Y^{[\alpha'\beta']}$, where $\alpha' = 1, \dots, 8$. The total number of supersymmetries is 32, which corresponds to the product of the dimensions of the spinor indices $32 = 4 \times 8$. It is noted that the assignment of the duality symmetric space in \mathcal{A} -theory differs from that in conventional formulations. In \mathcal{A} -theory, the duality-symmetric space is assigned to the main “spacetime” rather than the internal space, such that all tensor gauge fields are automatically incorporated into the coset parameter of E_{D+1}/H .

Physical currents are as follows.

$$\left\{ \begin{array}{l} F_{\tau}{}^{\underline{m}} = \dot{x}^{\underline{m}} + \partial^{\underline{m}} Y, \\ F_{\sigma; \underline{m}_1 \underline{m}_2} = -\epsilon_{\underline{m}_1 \dots \underline{m}_4} \partial^{\underline{m}_3} x^{\underline{m}_4}, \\ \underline{m} = 1, \dots, 4, \end{array} \right. \quad \left\{ \begin{array}{l} \delta_{\lambda} x^{\underline{m}} = \partial^{\underline{m}} \lambda, \\ \delta_{\lambda} Y = -\dot{\lambda}. \end{array} \right. \quad (6)$$

The following currents are auxiliary written by auxiliary coordinates y^{mn} , Y^m .

$$\left\{ \begin{array}{l} F_{\tau}^{mn} = \dot{y}^{mn} - \partial^{[m} Y^{n]}, \\ F_{\sigma; \underline{m}_1} = \frac{1}{2} \epsilon_{\underline{m}_1 \dots \underline{m}_4} \partial^{\underline{m}_2} y^{\underline{m}_3 \underline{m}_4}, \end{array} \right. \left\{ \begin{array}{l} \delta_{\lambda} y^{mn} = \partial^{[m} \lambda^{n]}, \\ \delta_{\lambda} Y^m = \dot{\lambda}^m, \\ \delta_{\lambda} \lambda^m = \partial^m \lambda. \end{array} \right. \quad (7)$$

These currents constitute the SL(5) A-symmetry together with (6), and they are used to lead the non-perturbative M2-brane Lagrangian.

3. \mathcal{T} -string currents

The O(3,3) \mathcal{T} -theory is described by a string with the manifest O(3,3) T-duality symmetry, namely \mathcal{T} -string.

$$\left\{ \begin{array}{l} J_{\tau}^{\underline{m}_1 \underline{m}_2} = \dot{X}^{\underline{m}_1 \underline{m}_2}, \\ J_{\sigma \underline{m}_1 \underline{m}_2} = \frac{1}{2} \epsilon_{\underline{m}_1 \dots \underline{m}_4} \partial_{\sigma} X^{\underline{m}_3 \underline{m}_4}. \end{array} \right. \quad (8)$$

It is convenient to represent in terms of $x^{\bar{m}}$ and $y_{\bar{m}} = \frac{1}{2} \epsilon_{\bar{m} \bar{n} \bar{l}} y^{\bar{n} \bar{l}}$.

$$\left\{ \begin{array}{l} J_{\tau}^{\bar{m}} = \dot{x}^{\bar{m}}, \\ J_{\sigma \bar{m}_1 \bar{m}_2} = -\epsilon_{\bar{m}_1 \bar{m}_2 \bar{m}_3} \partial_{\sigma} x^{\bar{m}_3}, \\ J_{\tau}^{\bar{m}_1 \bar{m}_2} = \dot{y}^{\bar{m}_1 \bar{m}_2}, \\ J_{\sigma \bar{m}_1} = \frac{1}{2} \epsilon_{\bar{m}_1 \bar{m}_2 \bar{m}_3} \partial_{\sigma} y^{\bar{m}_2 \bar{m}_3}, \\ \bar{m} = 1, 2, 3. \end{array} \right. \quad (9)$$

4. S -string currents

The GL(3) S -theory is described by a string with the manifest GL(3) spacetime diffeomorphism symmetry, namely a 3-dimensional string.

$$\left\{ \begin{array}{l} J_{\tau}^{\bar{m}} = \dot{x}^{\bar{m}}, \\ J_{\sigma \bar{m}_1 \bar{m}_2} = -\epsilon_{\bar{m}_1 \bar{m}_2 \bar{m}_3} \partial_{\sigma} x^{\bar{m}_3}. \end{array} \right. \quad (10)$$

Some minus signs come from the mere notation $\epsilon_{1234} = 1 = -\epsilon_{4123}$. It is denoted that these currents are flat currents, and in later sections flat current symbols $\overset{\circ}{F}$ or $\overset{\circ}{J}$ will be used to distinguish from curved background currents.

2.3 Constraints and sectionings

The theories in Hamiltonian formulation are constructed by the current algebra with manifest duality symmetries [18]. The Spacetime translation is generated by the covariant derivative $\triangleright_M(\sigma)$. The p -brane current algebra with G -symmetry covariance is given by

$$[\triangleright_M(\sigma), \triangleright_N(\sigma')] = 2if_{MN}{}^L \triangleright_L(\sigma) \delta(\sigma - \sigma') + 2i\eta_{MNm} \partial^m \delta^{(p)}(\sigma - \sigma'). \quad (11)$$




Figure 3: Diamond diagram of Sectionings of branes of $D = 3$ theories.

Branes are governed by the brane Virasoro constraints $\mathcal{S}^m = \frac{1}{2} \mathfrak{P}_M \eta^{MN} \mathfrak{P}_N = 0$ and $\mathcal{H} = \frac{1}{2} \mathfrak{P}_M \hat{\eta}^{MN} \mathfrak{P}_N = 0$ together with the Gauß law constraints $\mathcal{U}_m = 0$ which is required by the closure of the Virasoro algebra. $\hat{\eta}^{MN}$ is the H -invariant metric. Theories are related by sectionings; The Virasoro constraint $\mathcal{S}^m = 0$ gives the section conditions to reduce the space-time dimensions, and the Gauß law constraint $\mathcal{U}_m = 0$ is used to reduce the world-volume dimension as Fig. 3.

The spacetime covariant derivatives, constraints and section conditions are given [18, 30] in Fig. 3 concretely as follows.

1. $\mathcal{A}5$ -brane in 10-dimensional spacetime

The 10-dimensional spacetime is described by the rank-two anti-symmetric tensor covariant derivative $\mathfrak{P}_{m_1 m_2}$ as

$$\mathfrak{P}_{m_1 m_2} = P_{m_1 m_2} + \frac{1}{2} \epsilon_{m_1 \dots m_5} \partial^{m_3} X^{m_4 m_5}, \quad (12)$$

where P_{mn} is canonical conjugate of X^{mn} with $[P_{mn}(\sigma), X^{lk}(\sigma')] = \frac{1}{i} \delta_m^{[l} \delta_n^{k]} \delta^{(5)}(\sigma - \sigma')$ and $m = 1, \dots, 5$.

The $SL(5)$ covariant current algebra of $\mathcal{A}5$ -brane is given by

$$[\mathfrak{P}_{m_1 m_2}(\sigma), \mathfrak{P}_{m_3 m_4}(\sigma')] = 2i \epsilon_{m_1 \dots m_5} \partial^{m_5} \delta^{(5)}(\sigma - \sigma'). \quad (13)$$

The 5-dimensional world-volume diffeomorphism is generated by the Virasoro constraints $\mathcal{S}^m = 0$ while the world-volume time diffeomorphism is generated by $\mathcal{H} = 0$. The Gauß law constraint $\mathcal{U}_m = 0$ generates the gauge symmetry of the spacetime coordinate. These constraints are given by [1] as:

$$\begin{cases} \mathcal{S}^m = \frac{1}{16} \mathfrak{P}_{m_1 m_2} \epsilon^{m m_1 \dots m_4} \mathfrak{P}_{m_3 m_4} = 0, \\ \mathcal{H} = \frac{1}{16} \mathfrak{P}_{m_1 m_2} \delta^{m_1 [n_1} \delta^{n_2] m_2} \mathfrak{P}_{n_1 n_2} = 0, \\ \mathcal{U}_m = \partial^n \mathfrak{P}_{mn} = 0. \end{cases} \quad (14)$$

The $SL(5)$ covariant constraints $\mathcal{S}^m = 0$ and $\mathcal{U}_m = 0$ are background independent. These constraints are used as the dimensional reduction and section condition by replacing

Table 2: Constraints, dimensional reduction conditions and section conditions of $\mathcal{M}5$ -brane.

	Virasoro : \mathcal{S}^m	Gauß law : \mathcal{U}_m
dimensional reduction	$\epsilon^{mm_1 \dots m_4} P_{m_1 m_2}(\sigma) \frac{\partial}{\partial X_0^{m_3 m_4}}$	$P_{mn}(\sigma) \frac{\partial}{\partial \sigma_n}$
section condition	$\epsilon^{mm_1 \dots m_4} \frac{\partial}{\partial X_0^{m_1 m_2}} \frac{\partial}{\partial X_0^{m_3 m_4}}$	$\frac{\partial}{\partial \sigma_n} \frac{\partial}{\partial X_0^{mn}}$

the spacetime momentum $P_M(\sigma)$ with the derivative of the 0-mode of the spacetime coordinate X_0^M . These operators act on fields $\Phi(X)$ and $\Psi(X)$ as

$$\begin{aligned} \frac{\partial}{\partial X_0^M} \frac{\partial}{\partial X_0^N} \Phi(X_0) &= 0 = \frac{\partial}{\partial X_0^M} \Phi(X_0) \frac{\partial}{\partial X_0^N} \Psi(X_0), \\ \frac{\partial}{\partial \sigma_n} \frac{\partial}{\partial X^M} \Phi(\sigma, X(\sigma)) &= 0 = \frac{\partial}{\partial \sigma_n} \Phi(\sigma, X(\sigma)) \frac{\partial}{\partial X^M} \Psi(\sigma, X(\sigma)), \end{aligned} \quad (15)$$

where fields may be functions on σ as $\Phi(\sigma, X(\sigma))$ and $\Psi(\sigma, X(\sigma))$.

2. $\mathcal{M}5$ -brane in 4-dimensional spacetime

The dimensional reduction of the spacetime is obtained by solving the Virasoro constraint in Tab. 2 as

$$P_{\underline{mn}}(\sigma) = 0, \quad \underline{m} = 1, \dots, 4 \quad \Rightarrow \quad \epsilon^{mm_1 \dots m_4} \frac{\partial}{\partial X_0^{m_1 m_2}} P_{m_3 m_4}(\sigma) = 0. \quad (16)$$

This condition makes $X^{\underline{m}_1 \underline{m}_2} = y^{\underline{m}_1 \underline{m}_2}$ to be non-dynamical and reduced dimensionally. The remaining spacetime is 4 dimensions $P_{5\underline{m}} = p_{\underline{m}} \neq 0$.

The 4-dimensional spacetime is described by the covariant derivative $\triangleright_{\underline{m}}$. The 6-dimensional covariant derivative $\triangleright_{\underline{mn}}$ is maintained to construct $SL(5)$ current algebra

$$\left\{ \begin{array}{l} \triangleright_{\underline{m}} = p_{\underline{m}}, \\ \triangleright_{\underline{m}_1 \underline{m}_2} = -\epsilon_{\underline{m}_1 \dots \underline{m}_4} \partial^{\underline{m}_3} x^{\underline{m}_4}, \end{array} \right. \quad (17)$$

with $X^{5\underline{m}} = x^{\underline{m}}$ and $P_{5\underline{m}} = p_{\underline{m}}$ which is not confused with the 0-mode momentum.

The $SL(5)$ current algebra of $\mathcal{M}5$ -brane is

$$\left\{ \begin{array}{l} [\triangleright_{\underline{m}}(\sigma), \triangleright_{\underline{n}}(\sigma')] = 0, \\ [\triangleright_{\underline{m}_1}(\sigma), \triangleright_{\underline{m}_2 \underline{m}_3}(\sigma')] = 2i \epsilon_{\underline{m}_1 \dots \underline{m}_4} \partial^{\underline{m}_4} \delta^{(5)}(\sigma - \sigma'), \\ [\triangleright_{\underline{m}_1 \underline{m}_2}(\sigma), \triangleright_{\underline{m}_3 \underline{m}_4}(\sigma')] = 0, \end{array} \right. \quad (18)$$

where the last algebra forces to $\partial^5 = 0$.

Table 3: Constraints, dimensional reduction conditions and section conditions of $\mathcal{M}5$ -brane.

	Virasoro : $\mathcal{S}^{\underline{m}}$	Gauß law : $\mathcal{U}_{\underline{m}}$
dimensional reduction	$\frac{1}{2}p_{\underline{m}}(\sigma)(\partial^{[\underline{m}x^{\underline{n}]})$	$p_{\underline{m}}(\sigma)\partial^{\underline{m}}$
section condition	none	$\partial^{\underline{m}}\frac{\partial}{\partial x_0^{\underline{m}}}$

The Virasoro operators of $\mathcal{M}5$ -brane are

$$\left\{ \begin{array}{l} \mathcal{S}^{\underline{m}} = \frac{1}{2}\partial^{[\underline{m}x^{\underline{n}]}p_{\underline{n}}, \\ \mathcal{S}^5 = \frac{1}{4}\epsilon_{\underline{m}_1\cdots\underline{m}_4}(\partial^{\underline{m}_1}x^{\underline{m}_2})(\partial^{\underline{m}_3}x^{\underline{m}_4}), \\ \mathcal{H} = \frac{1}{4}p_{\underline{m}}\hat{\eta}^{\underline{mn}}p_{\underline{n}} + \frac{1}{16}(\partial^{[\underline{m}_1}x^{\underline{m}_2]})\hat{\eta}_{\underline{m}_1[\underline{n}_1}\hat{\eta}_{\underline{n}_2]\underline{m}_2}(\partial^{[\underline{n}_1}x^{\underline{n}_2]}), \\ \mathcal{U}_5 = \partial^{\underline{n}}p_{\underline{n}}. \end{array} \right. \quad (19)$$

These constraints lead to the following dimensional reductions and section conditions.

3. \mathcal{T} -string in 6-dimensional spacetime

The dimensional reduction condition of the world-volume is obtained by solving the Gauß law constraint in Tab. 2 as

$$\partial^{\underline{n}} = 0, \quad P_{\underline{m}}(\sigma) = 0 \quad \Rightarrow \quad \partial^{\underline{n}}P_{\underline{mn}}(\sigma) = 0. \quad (20)$$

These conditions make $\partial^5 = \frac{\partial}{\partial\sigma} \neq 0$ and $X^{\underline{m}}$ to be non-dynamical (constant). The remaining spacetime is 6 dimensional $P_{\underline{mn}} = p_{\underline{mn}} \neq 0$.

The 6-dimensional spacetime is described by the covariant derivative $\triangleright_{\underline{mn}}$. The 4-dimensional covariant derivative vanishes $\triangleright_{\underline{m}} = 0$

$$\triangleright_{\underline{m}_1\underline{m}_2} = p_{\underline{m}_1\underline{m}_2} + \frac{1}{2}\epsilon_{\underline{m}_1\cdots\underline{m}_4}\partial^5x^{\underline{m}_3\underline{m}_4}, \quad (21)$$

with $X^{\underline{mn}} = x^{\underline{mn}}$.

The $O(3,3)$ current algebra of \mathcal{T} -string is

$$[\triangleright_{\underline{m}_1\underline{m}_2}(\sigma), \triangleright_{\underline{m}_3\underline{m}_4}(\sigma')] = 2i\epsilon_{\underline{m}_1\cdots\underline{m}_4}\partial_\sigma\delta(\sigma - \sigma'),$$

with $\partial^5 = \partial_\sigma$.

The Virasoro operators of \mathcal{T} -string are

$$\left\{ \begin{array}{l} \mathcal{S}^5 = \frac{1}{16}\triangleright_{\underline{m}_1\underline{m}_2}\epsilon^{\underline{m}_1\cdots\underline{m}_4}\triangleright_{\underline{m}_3\underline{m}_4}, \\ \mathcal{H} = \frac{1}{16}\triangleright_{\underline{m}_1\underline{m}_2}\hat{\eta}^{\underline{m}_1[\underline{n}_1}\hat{\eta}^{\underline{n}_2]\underline{m}_2}\triangleright_{\underline{n}_1\underline{n}_2}, \end{array} \right. \quad (22)$$

with $\mathcal{S}^m = 0 = \mathcal{U}_m$.

The Virasoro constraint $\mathcal{S}^5 = 0$ leads to the following dimensional reduction and the section condition.

4. String in 3-dimensional spacetime

(a) From \mathcal{T} -string to String

The dimensional reduction of the spacetime is obtained by solving the Virasoro constraint in Tab. 4 as

$$P_{\bar{m}\bar{n}}(\sigma) = 0, \quad \bar{m} = 1, 2, 3 \quad \Rightarrow \quad P_{\underline{m}_1 \underline{m}_2}(\sigma) \epsilon^{\underline{m}_1 \dots \underline{m}_4} \frac{\partial}{\partial x_0^{\underline{m}_3 \underline{m}_4}} = 0. \quad (23)$$

This condition makes $X^{\bar{m}_1 \bar{m}_2} = y^{\bar{m}_1 \bar{m}_2}$ to be non-dynamical (constant). The remaining spacetime is 3 dimensions $P_{4\bar{m}} = p_{\bar{m}} \neq 0$.

(b) From $\mathcal{M}5$ -brane to String

The dimensional reduction condition of the world-volume is obtained by solving the Gauß law constraint in Tab. 3 as

$$\partial^{\bar{n}} = 0, \quad P_4(\sigma) = 0 \quad \Rightarrow \quad P_{\underline{m}} \partial^{\underline{m}} = 0. \quad (24)$$

In the 4-dimensional spacetime ∂^5 is considered to be 0. These conditions make $\partial^4 = \frac{\partial}{\partial \sigma} = \partial_\sigma \neq 0$ and X^{54} to be non-dynamical (constant). The remaining spacetime is 3 dimensions $P_{4\bar{m}} = p_{\bar{m}} \neq 0$.

The 3-dimensional spacetime is described by the covariant derivative $\triangleright_{4\bar{m}}$. The 3-dimensional covariant derivative vanishes $\triangleright_{\bar{m}_1 \bar{m}_2}$

$$\left\{ \begin{array}{l} \triangleright_{\bar{m}} = p_{\bar{m}}, \\ \triangleright_{\bar{m}_1 \bar{m}_2} = \epsilon_{\bar{m}_1 \bar{m}_2 \bar{m}} \partial_\sigma x^{\bar{m}}, \end{array} \right. \quad (25)$$

with $X^{4\bar{m}} = x^{\bar{m}}$ and $\sigma^5 = \sigma$ via \mathcal{T} -string and $X^{5\bar{m}} = x^{\bar{m}}$ and $\sigma^4 = \sigma$ via $\mathcal{M}5$ -brane.

The $\text{GL}(3)$ current algebra becomes

$$[\triangleright_{\bar{m}_1}(\sigma), \triangleright_{\bar{m}_2 \bar{m}_3}(\sigma')] = i \epsilon_{\bar{m}_1 \bar{m}_2 \bar{m}_3} \partial_\sigma \delta(\sigma - \sigma'). \quad (26)$$

Table 4: Constraints, dimensional reduction conditions and section conditions of \mathcal{T} -string.

	Virasoro : \mathcal{S}^m	Gauß law : \mathcal{U}_m
dimensional reduction	$p_{\underline{m}_1 \underline{m}_2}(\sigma) \epsilon^{\underline{m}_1 \dots \underline{m}_4} \frac{\partial}{\partial x_0^{\underline{m}_3 \underline{m}_4}}$	none
section condition	$\frac{\partial}{\partial x_0^{\underline{m}_1 \underline{m}_2}} \epsilon^{\underline{m}_1 \dots \underline{m}_4} \frac{\partial}{\partial x_0^{\underline{m}_3 \underline{m}_4}}$	none

This is equivalent to the $O(D,D)$ current algebra which is given by $\triangleright_M = (\triangleright_{\bar{m}}, \triangleright^{\bar{m}})$ with $\triangleright^{\bar{m}} = \frac{1}{2}\epsilon^{\bar{m}\bar{m}_1\bar{m}_2}\triangleright_{\bar{m}_1\bar{m}_2}$ and the $O(D,D)$ invariant metric $\eta_{MN} = \epsilon_{\bar{m}_1\bar{m}_2\bar{m}_3}$ as

$$[\triangleright_M(\sigma), \triangleright_N(\sigma')] = i\eta_{MN}\partial_\sigma\delta(\sigma - \sigma'). \quad (27)$$

The Virasoro operators become

$$\begin{cases} \mathcal{S} = p_{\bar{m}}\partial_\sigma x^{\bar{m}} = \frac{1}{2}\triangleright_M\eta^{MN}\triangleright_N, \\ \mathcal{H} = \frac{1}{2}p_{\bar{m}}\hat{\eta}^{\bar{m}\bar{n}}p_{\bar{n}} + \frac{1}{2}\partial_\sigma x^{\bar{m}}\hat{\eta}_{\bar{m}\bar{n}}\partial_\sigma x^{\bar{n}} = \frac{1}{2}\triangleright_M\hat{\eta}^{MN}\triangleright_N, \end{cases} \quad (28)$$

with the double Lorentz invariant metric $\hat{\eta}^{MN}$. There are no further conditions of the Virasoro and the Gauß law constraints; $\mathcal{S}^m = 0 = \mathcal{U}_m$.

3 $\mathcal{A}5$ -brane Lagrangians

3.1 $\mathcal{A}5$ -brane Lagrangian with $SL(5)$ U-duality symmetry

The $SL(5)$ U-duality symmetry is manifestly realized by the $\mathcal{A}5$ -brane. The spacetime background is described by the vielbein which is a $SL(5)/SO(5)$ coset element E_m^a satisfying

$$E_{m_1}^{a_1}E_{m_2}^{a_2}E_{m_3}^{a_3}E_{m_4}^{a_4}E_{m_5}^{a_5}\epsilon_{a_1a_2a_3a_4a_5} = \epsilon_{m_1m_2m_3m_4m_5}, \quad (29)$$

with $m, a = 1, \dots, 5$. The background metrics with tensor indices are

$$\begin{aligned} G_{mn} &= E_m^a\hat{\eta}_{ab}E_n^b, \\ G_{m_1m_2;n_1n_2} &= E_{m_1}^{a_1}E_{m_2}^{a_2}\hat{\eta}_{a_1[b_1}\hat{\eta}_{b_2]a_2}E_{n_1}^{b_1}E_{n_2}^{b_2}. \end{aligned} \quad (30)$$

The selfdual and anti-selfdual currents in a flat background $\overset{\circ}{F}_{SD/\overline{SD}}{}^{mn}$ and in a curved background $F_{SD/\overline{SD}}{}^{ab}$ in terms of (5) are given as

$$\begin{cases} \overset{\circ}{F}_{SD}{}^{m_1m_2} = F_\tau^{m_1m_2} - \frac{1}{2}\epsilon^{m_1\dots m_5}s_{m_3}F_{\sigma;m_4m_5} + g\hat{\eta}^{m_1n_1}\hat{\eta}^{m_2n_2}F_{\sigma;n_1n_2}, \\ \overset{\circ}{F}_{\overline{SD}}{}^{m_1m_2} = F_\tau^{m_1m_2} - \frac{1}{2}\epsilon^{m_1\dots m_5}s_{m_3}F_{\sigma;m_4m_5} - g\hat{\eta}^{m_1n_1}\hat{\eta}^{m_2n_2}F_{\sigma;n_1n_2}, \end{cases} \quad (31)$$

$$F_{SD/\overline{SD}}{}^{a_1a_2} = \overset{\circ}{F}_{SD/\overline{SD}}{}^{m_1m_2}E_{m_1}^{a_1}E_{m_2}^{a_2}, \quad (32)$$

where $\hat{\eta}^{mn}$ becomes G^{mn} in a curved background. g and s_m are 5-brane world-volume vielbein fields which are introduced as Lagrange multipliers of Virasoro constraints.

The Lagrangian of the $\mathcal{A}5$ -brane $L_{SL(5)}$ is given [1] as

$$\begin{aligned} I_{SL(5)} &= \int d\tau d^5\sigma L_{SL(5)}, \\ L_{SL(5)} &= \frac{1}{2}\phi F_{SD}{}^{ab}F_{\overline{SD}ab} + \frac{1}{2}\bar{\phi}(F_{\overline{SD}}{}^{ab})^2 + \frac{1}{2}\lambda_{ab}F_{SD}{}^{ac}F_{SD}{}^b{}_c - \frac{1}{4}\epsilon_{a_1\dots a_5}\lambda^{a_1}F_{SD}{}^{a_2a_3}F_{SD}{}^{a_4a_5} \\ &= \frac{\phi}{4}\overset{\circ}{F}_{SD}{}^{m_1m_2}G_{m_1m_2;m_3m_4}\overset{\circ}{F}_{SD}{}^{m_3m_4} + \frac{\bar{\phi}}{8}\overset{\circ}{F}_{SD}{}^{m_1m_2}G_{m_1m_2;m_3m_4}\overset{\circ}{F}_{SD}{}^{m_3m_4} \\ &\quad + \frac{1}{2}\overset{\circ}{\lambda}_{mn}F_{SD}{}^{ml_1}G_{l_1l_2}\overset{\circ}{F}_{SD}{}^{nl_2} + \frac{1}{8}\epsilon_{m_1\dots m_5}\overset{\circ}{\lambda}^{m_1}F_{SD}{}^{m_2m_3}\overset{\circ}{F}_{SD}{}^{m_4m_5}, \end{aligned} \quad (33)$$

with symmetric traceless tensors $\lambda_{\hat{m}\hat{n}}$'s.

3.2 World-volume covariant $\mathcal{A}5$ -brane Lagrangian with $SL(6)$ duality symmetry

The $G=SL(5)$ U-duality symmetry is enlarged to $A=SL(6)$ by cooperating with the 6-dimensional world-volume Lorentz covariance. The $SL(6)/SO(6)$ coset parameter includes not only the target space vielbein $SL(5)/SO(5)$ but also 6 components of the world-volume vielbein. The background vielbein $E_{\hat{m}}^{\hat{a}} \in SL(6)/SO(6)$ satisfies

$$E_{\hat{m}_1}^{\hat{a}_1} E_{\hat{m}_2}^{\hat{a}_2} E_{\hat{m}_3}^{\hat{a}_3} E_{\hat{m}_4}^{\hat{a}_4} E_{\hat{m}_5}^{\hat{a}_5} E_{\hat{m}_6}^{\hat{a}_6} \epsilon_{\hat{a}_1 \hat{a}_2 \hat{a}_3 \hat{a}_4 \hat{a}_5 \hat{a}_6} = \epsilon_{\hat{m}_1 \hat{m}_2 \hat{m}_3 \hat{m}_4 \hat{m}_5 \hat{m}_6}, \quad (34)$$

with $\hat{m}, \hat{a} = 0, 1, \dots, 5$.

This $SL(6)$ covariant vielbein (34) includes the 5-brane world-volume vielbein fields g and s_m as

$$E_{\hat{m}}^{\hat{a}} = \begin{pmatrix} E_0^{\hat{0}} & E_0^a \\ E_m^{\hat{0}} & E_m^a \end{pmatrix} = \begin{pmatrix} \frac{1}{g} & 0 \\ -\frac{s_m}{g} & E_m^a \end{pmatrix}, \quad (35)$$

with $\hat{m} = (0, m)$, $\hat{a} = (\hat{0}, a)$ and $m, a = 1 \dots, 5$. It is denoted that the E_m^a component of $SL(6)$ vielbein (34) is different from the $SL(5)$ vielbein E_m^a in (35) up to the determinant factor. The number of degrees of freedom of the $SL(6)$ vielbein is sum of the spacetime vielbein and the world-volume vielbein as

$$(6^2 - 1) - \frac{6 \times 5}{2} = \left((5^2 - 1) - \frac{5 \times 4}{2} \right) + 6. \quad (36)$$

This is generalized for a p -brane of \mathcal{A} -theory symmetry with A/L coset as

$$\dim \frac{A}{L} = \dim \frac{G}{H} + (p + 1). \quad (37)$$

The $SL(6)$ covariant field strengths are given by a simple form; the one in a flat background $\overset{\circ}{F}^{\hat{m}\hat{n}\hat{l}}$ (the same as (4)) and the one in a curved background $F^{\hat{a}\hat{b}\hat{c}}$ as

$$\overset{\circ}{F}^{\hat{m}\hat{n}\hat{l}} = \frac{1}{2} \partial^{[\hat{m}} X^{\hat{n}\hat{l}]}, \quad F^{\hat{a}\hat{b}\hat{c}} = \overset{\circ}{F}^{\hat{m}\hat{n}\hat{l}} E_{\hat{m}}^{\hat{a}} E_{\hat{n}}^{\hat{b}} E_{\hat{l}}^{\hat{c}}. \quad (38)$$

The selfdual and the anti-selfdual field strength (31) and (32) are written in terms of the $SL(6)$ current (38) with $\partial^{\hat{0}} = \partial_{\tau}$ and $\epsilon^{\hat{0}12345} = 1$ as

$$F_{SD/\overline{SD}}^{a_1 a_2} = g \left(F^{\hat{0} a_1 a_2} \pm \frac{1}{6} \epsilon^{\hat{0} a_1 a_2}{}_{a_3 a_4 a_5} F^{a_3 a_4 a_5} \right). \quad (39)$$

Then the $\mathcal{A}5$ -brane Lagrangian (33) is rewritten in terms of the $SL(6)$ covariant field strength (38). The world-volume covariant $\mathcal{A}5$ -brane Lagrangian $L_{SL(6)}$ is given [1] as

$$\begin{aligned} I_{SL(6)} &= \int d^6 \sigma L_{SL(6)}, \\ L_{SL(6)} &= -\frac{1}{12} \Phi F^{\hat{a}_1 \hat{a}_2 \hat{a}_3} F_{\hat{a}_1 \hat{a}_2 \hat{a}_3} + \frac{1}{2} \Lambda_{\hat{a}\hat{b}} F^{\hat{a}\hat{c}_1 \hat{c}_2} F^{\hat{b}}{}_{\hat{c}_1 \hat{c}_2} + \frac{1}{12} \epsilon_{\hat{a}_1 \dots \hat{a}_6} \tilde{\Lambda}_{\hat{b}}^{\hat{a}_1} F^{\hat{a}_2 \hat{a}_3 \hat{a}_4} F^{\hat{a}_5 \hat{a}_6 \hat{b}} \\ &= -\frac{1}{72} \Phi \overset{\circ}{F}^{\hat{m}_1 \hat{m}_2 \hat{m}_3} G_{\hat{m}_1 \hat{m}_2 \hat{m}_3; \hat{m}_4 \hat{m}_5 \hat{m}_6} \overset{\circ}{F}^{\hat{m}_4 \hat{m}_5 \hat{m}_6} + \frac{1}{8} \Lambda_{\hat{m}\hat{n}} \overset{\circ}{F}^{\hat{m}\hat{l}_1 \hat{l}_2} G_{\hat{l}_1 \hat{l}_2; \hat{l}_3 \hat{l}_4} \overset{\circ}{F}^{\hat{n}\hat{l}_3 \hat{l}_4} \\ &\quad + \frac{1}{12} \epsilon_{\hat{m}_1 \dots \hat{m}_6} \tilde{\Lambda}_{\hat{n}}^{\hat{m}_1} \overset{\circ}{F}^{\hat{m}_2 \hat{m}_3 \hat{m}_4} \overset{\circ}{F}^{\hat{m}_5 \hat{m}_6 \hat{n}}, \end{aligned} \quad (40)$$

where $\Phi, \Lambda_{\hat{a}\hat{b}}$ are Lagrange multipliers with symmetric traceless tensors $\Lambda_{\hat{a}\hat{b}}$'s. The background metrics with tensor indices are

$$\begin{aligned} G_{\hat{m}_1 \hat{m}_2; \hat{n}_1 \hat{n}_2} &= E_{\hat{m}_1}^{\hat{a}_1} E_{\hat{m}_2}^{\hat{a}_2} \hat{\eta}_{\hat{a}_1 [\hat{b}_1} \hat{\eta}_{\hat{b}_2] \hat{a}_2} E_{\hat{n}_1}^{\hat{b}_1} E_{\hat{n}_2}^{\hat{b}_2}, \\ G_{\hat{m}_1 \hat{m}_2 \hat{m}_3; \hat{n}_1 \hat{n}_2 \hat{n}_3} &= E_{\hat{m}_1}^{\hat{a}_1} E_{\hat{m}_2}^{\hat{a}_2} E_{\hat{m}_3}^{\hat{a}_3} \hat{\eta}_{\hat{a}_1 [\hat{b}_1} \hat{\eta}_{\hat{b}_2 | \hat{a}_2} \hat{\eta}_{\hat{b}_3] \hat{a}_3} E_{\hat{n}_1}^{\hat{b}_1} E_{\hat{n}_2}^{\hat{b}_2} E_{\hat{n}_3}^{\hat{b}_3}. \end{aligned} \quad (41)$$

4 Lagrangian of D-dimensional string from O(D,D) \mathcal{T} -string

In this section we derive the O(D,D) \mathcal{T} -string Lagrangian from the O(D,D) Hamiltonian by the double zweibein method [58, 59]. Then the reduction procedure from the O(D,D) \mathcal{T} -string Lagrangian to the conventional string Lagrangian is presented.

4.1 O(D,D) \mathcal{T} -string

We begin with the sigma model string Lagrangian

$$I = \int d^2\sigma L, \quad (42)$$

$$L = -\frac{1}{2}\partial_\mu x^m (\sqrt{-h}h^{\mu\nu}g_{mn} + \epsilon^{\mu\nu}B_{mn})\partial_\nu x^n,$$

with $\mu = (\tau, \sigma)$. In the conformal gauge the Lagrangian becomes

$$\begin{aligned} L &= \frac{1}{2}(\dot{x}^m g_{mn} \dot{x}^n - x'^m g_{mn} x'^n) - \dot{x}^m B_{mn} x'^n \\ &= \frac{1}{2}(\dot{x}^m x'^m) \begin{pmatrix} g_{mn} & B_{mn} \\ B_{mn} & g_{mn} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} \dot{x}^n \\ x'^n \end{pmatrix} \\ &= \frac{1}{2}\partial_+ x^m (g_{mn} + B_{mn})\partial_- x^n, \end{aligned} \quad (43)$$

with $\dot{x} = \partial_\tau x$, $x' = \partial_\sigma x$ and $\partial_\pm x = \dot{x} \pm x'$.

The Hamiltonian is given by the Legendre transformation where the canonical momentum of x^m is given by $p_m = \partial L / \partial \dot{x}^m$,

$$\begin{aligned} H &= p_m \dot{x}^m - L \\ &= \frac{1}{2}(p_m x'^m) \begin{pmatrix} g^{mn} & g^{ml} B_{ln} \\ -B_{ml} g^{ln} & g_{mn} - B_{ml} g^{lk} B_{kn} \end{pmatrix} \begin{pmatrix} p_n \\ x'^n \end{pmatrix} \\ &= \frac{1}{2}\{(p_m - x'^l B_{lm})g^{mn}(p_n + B_{nk}x'^k) + x'^m g_{mn}x'^n\}. \end{aligned} \quad (44)$$

The background field is the O(D,D) matrix G^{MN} written in terms of the vielbein E_A^M as $E_A^M \rightarrow h_A^B E_B^N g_N^M$, $h \in \text{SO}(D-1,1)$ and $g \in \text{O}(D,D)$

$$E_A^M \eta_{MN} E_B^N = \eta_{AB}. \quad (45)$$

The background metric G^{MN} in the string Hamiltonian (44) is given as

$$\begin{aligned} G^{MN} &= \begin{pmatrix} g^{mn} & g^{ml} B_{ln} \\ -B_{ml} g^{ln} & g_{mn} - B_{ml} g^{lk} B_{kn} \end{pmatrix} = E_A^M \hat{\eta}^{AB} E_B^N, \\ E_A^M &= \begin{pmatrix} e_a^m & e_a^l B_{lm} \\ 0 & e_m^a \end{pmatrix}, \end{aligned} \quad (46)$$

while its inverse is given by

$$\begin{aligned} G_{MN} &= \begin{pmatrix} g_{mn} - B_{ml} g^{lk} B_{kn} & -B_{ml} g^{ln} \\ g^{ml} B_{ln} & g^{mn} \end{pmatrix} = E_M^A \hat{\eta}_{AB} E_N^B, \\ E_M^A &= \begin{pmatrix} e_m^a & -B_{ml} e_a^l \\ 0 & e_a^m \end{pmatrix}. \end{aligned} \quad (47)$$

This $O(D,D)$ background metric is utilized in the Lagrangian with manifest $O(D,D)$ T-duality symmetry.

The $O(D,D)$ covariant space is constructed in such a way that the $O(D,D)$ covariant derivative $\triangleright_M(\sigma)$ algebra satisfies the same algebra of $\triangleright_M = (p_m, x'^m)$ up to the normalization

$$[\triangleright_M(\sigma), \triangleright_N(\sigma')] = 2i\eta_{MN}\partial_\sigma\delta(\sigma - \sigma'), \quad \eta_{MN} = \begin{pmatrix} & \delta_n^m \\ \delta_n^m & \end{pmatrix}. \quad (48)$$

The covariant derivative \triangleright_M is realized in terms of the doubled coordinate X^M and P_M with $[P_M(\sigma), X^N(\sigma')] = -i\delta_M^N\delta(\sigma - \sigma')$ as

$$\overset{\circ}{\triangleright}_M = P_M + \partial_\sigma X^N \eta_{NM}, \quad (49)$$

which is left moving current in the doubled space. The right moving current is also introduced as

$$\tilde{\triangleright}_M = P_M - \partial_\sigma X^N \eta_{NM}, \quad (50)$$

which satisfies the same current algebra (48) with opposite sign. The number of canonical variables of the doubled space are 4D, while the physical one is 2D. The 2D equations $\tilde{\triangleright}_M = 0$ is the usual selfduality condition to suppress 2D unphysical degrees of freedom, so we call $\tilde{\triangleright}_M$ “anti-selfdual current”. Another current $\overset{\circ}{\triangleright}_M$ is selfdual current.

There are two sets of Virasoro operators written in terms of the selfdual current and the anti-selfdual current

$$\begin{cases} \mathcal{H} = \frac{1}{4}\overset{\circ}{\triangleright}_M \hat{\eta}^{MN} \overset{\circ}{\triangleright}_N, \\ \mathcal{S} = \frac{1}{4}\overset{\circ}{\triangleright}_M \eta^{MN} \overset{\circ}{\triangleright}_N, \end{cases} \quad \begin{cases} \tilde{\mathcal{H}} = \frac{1}{4}\tilde{\triangleright}_M \hat{\eta}^{MN} \tilde{\triangleright}_N, \\ \tilde{\mathcal{S}} = \frac{1}{4}\tilde{\triangleright}_M \eta^{MN} \tilde{\triangleright}_N. \end{cases} \quad (51)$$

\mathcal{H} and \mathcal{S} satisfy the Virasoro algebra

$$\begin{aligned} [\mathcal{S}(\sigma), \mathcal{S}(\sigma')] &= i\{\mathcal{S}(\sigma) + \mathcal{S}(\sigma')\}\partial_\sigma\delta(\sigma - \sigma'), \\ [\mathcal{S}(\sigma), \mathcal{H}(\sigma')] &= i\{\mathcal{H}(\sigma) + \mathcal{H}(\sigma')\}\partial_\sigma\delta(\sigma - \sigma'), \\ [\mathcal{H}(\sigma), \mathcal{H}(\sigma')] &= i\{\mathcal{S}(\sigma) + \mathcal{S}(\sigma')\}\partial_\sigma\delta(\sigma - \sigma'), \end{aligned} \quad (52)$$

while $\tilde{\mathcal{H}}$ and $\tilde{\mathcal{S}}$ satisfy the same Virasoro algebra with opposite signs on the right hand side.

As seen in the Hamiltonian in curved background (44) currents \triangleright_M coupled to the vielbein as

$$\triangleright_A = E_A^M \overset{\circ}{\triangleright}_M, \quad \tilde{\triangleright}_A = E_A^M \tilde{\triangleright}_M. \quad (53)$$

In curved background the Virasoro constraints become

$$\begin{cases} \mathcal{H} = \frac{1}{4}\triangleright_A \hat{\eta}^{AB} \triangleright_B = \frac{1}{4}\overset{\circ}{\triangleright}_M G^{MN} \overset{\circ}{\triangleright}_N, \\ \mathcal{S} = \frac{1}{4}\triangleright_A \eta^{AB} \triangleright_B = \frac{1}{4}\overset{\circ}{\triangleright}_M \eta^{MN} \overset{\circ}{\triangleright}_N, \end{cases} \quad \begin{cases} \tilde{\mathcal{H}} = \frac{1}{4}\tilde{\triangleright}_A \hat{\eta}^{AB} \tilde{\triangleright}_B = \frac{1}{4}\tilde{\triangleright}_M G^{MN} \tilde{\triangleright}_N, \\ \tilde{\mathcal{S}} = \frac{1}{4}\tilde{\triangleright}_A \eta^{AB} \tilde{\triangleright}_B = \frac{1}{4}\tilde{\triangleright}_M \eta^{MN} \tilde{\triangleright}_N. \end{cases} \quad (54)$$

The $O(D,D)$ covariant Hamiltonian is given by the sum of all these Virasoro constraints with Lagrange multipliers which are doubled zweibeins [58]

$$\begin{aligned} H &= g\mathcal{H} + s\mathcal{S} + \tilde{g}\tilde{\mathcal{H}} + \tilde{s}\tilde{\mathcal{S}} \\ &= \frac{1}{2}[P_A M^{AB} P_B + 2P_A N^{AC} \eta_{CB} X'^B + X'^A \eta_{AC} M^{CD} \eta_{DB} X'^B], \end{aligned} \quad (55)$$

with $P_A = P_M E_A^M$ and $X'^A \equiv X'^M E_M^A$. We used the fact that the covariant derivatives are rewritten as $\triangleright_A = P_A + X'^B \eta_{BA}$ and $\tilde{\triangleright}_A = P_A - X'^B \eta_{BA}$ by the orthogonal condition (45). Matrices M^{AB} and N^{AB} are given as

$$\begin{aligned} M^{AB} &= \frac{g + \tilde{g}}{2} \hat{\eta}^{AB} + \frac{s + \tilde{s}}{2} \eta^{AB}, \\ N^{AB} &= \frac{g - \tilde{g}}{2} \hat{\eta}^{AB} + \frac{s - \tilde{s}}{2} \eta^{AB}, \end{aligned} \quad (56)$$

with the inverse of M^{AB} as

$$M^{-1}_{AB} = \frac{2}{(g + \tilde{g})^2 - (s + \tilde{s})^2} \{ (g + \tilde{g})\hat{\eta}_{AB} - (s + \tilde{s})\eta_{AB} \}. \quad (57)$$

The Legendre transformation of the Hamiltonian (55) with (56) leads to the following Lagrangian

$$L = P_M \dot{X}^M - H = \frac{1}{2} J_+^A M^{-1}_{AB} J_-^B, \quad (58)$$

$$\begin{cases} J_+^A = \dot{X}^A + (\tilde{g}\hat{\eta}^{AB} + \tilde{s}\eta^{AB})\eta_{BC}X'^C, \\ J_-^A = \dot{X}^A - (g\hat{\eta}^{AB} + s\eta^{AB})\eta_{BC}X'^C, \end{cases} \quad (59)$$

with $\dot{X}^A \equiv \dot{X}^M E_M^A$.

The Lagrangian in (58) can be written in terms of the selfdual current and the anti-selfdual current which is equal to J_- in (59). The selfdual and anti-selfdual currents are given by

$$\begin{cases} J_{SD}^A = (\dot{X}^A - sX'^A) + g\hat{\eta}^{AB}\eta_{BC}X'^C, \\ J_{\overline{SD}}^A = (\dot{X}^A - sX'^A) - g\hat{\eta}^{AB}\eta_{BC}X'^C. \end{cases} \quad (60)$$

The selfdual and anti-selfdual currents in the flat background, $J_{SD/\overline{SD}}^M = J_{SD/\overline{SD}}^A E_A^M$, are written as

$$\begin{cases} \overset{\circ}{J}_{SD}^M = (\dot{X}^M - sX'^M) + g\hat{\eta}^{MN}\eta_{NL}X'^L, \\ \overset{\circ}{J}_{\overline{SD}}^M = (\dot{X}^M - sX'^M) - g\hat{\eta}^{MN}\eta_{NL}X'^L. \end{cases} \quad (61)$$

It is denoted that $\hat{\eta}^{MN}$ becomes G^{MN} in a curved background. The resultant O(D,D) covariant Lagrangian for a \mathcal{T} -string is given [58] as

$$\begin{aligned} I &= \int d\tau d\sigma L, \\ L &= \phi J_{SD}^A \hat{\eta}_{AB} J_{\overline{SD}}^B + \bar{\phi} J_{\overline{SD}}^A \hat{\eta}_{AB} J_{SD}^B + \tilde{\phi} J_{SD}^A \eta_{AB} J_{\overline{SD}}^B \\ &= \phi \overset{\circ}{J}_{SD}^M G_{MN} \overset{\circ}{J}_{\overline{SD}}^N + \bar{\phi} \overset{\circ}{J}_{\overline{SD}}^M G_{MN} \overset{\circ}{J}_{SD}^N + \tilde{\phi} \overset{\circ}{J}_{SD}^M \eta_{MN} \overset{\circ}{J}_{\overline{SD}}^N. \end{aligned} \quad (62)$$

The first term is the kinetic term, while the rest are constraints that are squares of the anti-selfdual currents. The Lagrange multipliers ϕ , $\bar{\phi}$ and $\tilde{\phi}$ are related to the doubled zweibeins as

$$\begin{cases} \phi = \frac{1}{2g}, \\ \bar{\phi} = \frac{1}{2g[(g + \tilde{g})^2 - (s + \tilde{s})^2]} \{ (s + \tilde{s})^2 + g^2 - \tilde{g}^2 \}, \\ \tilde{\phi} = -\frac{s + \tilde{s}}{(g + \tilde{g})^2 - (s + \tilde{s})^2}. \end{cases} \quad (63)$$

4.2 String from O(D,D) \mathcal{T} -string

We break the O(D,D) T-duality symmetry of \mathcal{T} -string into the GL(D) symmetry of the usual string. The background gauge field of \mathcal{T} -string is O(D,D)/O(D-1,1)² coset parameter which includes the D-dimensional metric g_{mn} and B_{mn} field, while the background gauge field of a string is GL(D)/SO(D-1,1) coset parameter which includes only g_{mn} . In this subsection we use the coordinate $X^M = (x^m, y_m)$ with off-diagonal η_{MN} to describe \mathcal{T} -string, while the left/right moving coordinate with diagonal $\eta_{MN} = (\mathbf{1}, -\mathbf{1})$ was used in the reference [58]. The Weyl/Lorentz gauge of the zweibein [58] is given as

$$\varepsilon_{\pm}^{\mu} = \begin{pmatrix} \varepsilon_+^{\tau} & \varepsilon_+^{\sigma} \\ \varepsilon_-^{\tau} & \varepsilon_-^{\sigma} \end{pmatrix} = \begin{pmatrix} 1 & g-s \\ 1 & -g-s \end{pmatrix}. \quad (64)$$

The left/right moving modes with the zweibein is $\varepsilon_{\pm}X \equiv \varepsilon_{\pm}^{\mu}\partial_{\mu}X$. The selfdual and anti-selfdual currents (61) are expressed as

$$\begin{cases} \dot{J}_{SD/\overline{SD}}^m = \dot{x}^m - s x'^m \pm g \hat{\eta}^{mn} y'_n = \varepsilon_{\tau} x^m \pm g \hat{\eta}^{mn} \varepsilon_{\sigma} y_n, \\ \dot{J}_{SD/\overline{SD};m} = \dot{y}_m - s y'_m \pm g \hat{\eta}_{mn} x'^n = \varepsilon_{\tau} y_m \pm g \hat{\eta}_{mn} \varepsilon_{\sigma} x^n, \end{cases} \quad (65)$$

with

$$\varepsilon_{\mu}X \equiv \varepsilon_{\mu}^{\nu}\partial_{\nu}X, \quad \varepsilon_{\mu}^{\nu} = \begin{pmatrix} \varepsilon_{\tau}^{\tau} & \varepsilon_{\tau}^{\sigma} \\ \varepsilon_{\sigma}^{\tau} & \varepsilon_{\sigma}^{\sigma} \end{pmatrix} = \begin{pmatrix} 1 & -s \\ 0 & 1 \end{pmatrix}. \quad (66)$$

The condition of vanishing the anti-selfdual current in $s = 0$ and $g = 1$ gauge leads to the selfduality constraint in flat space as $\partial_{\mu}y = \varepsilon_{\mu\nu}\partial^{\nu}x$ with $\partial^{\tau} = -\partial_{\tau}$. In the gauge $\phi = \frac{1}{2g}$ and $\bar{\phi} = 0 = \tilde{\phi}$, corresponding to $g = \tilde{g}$ and $s + \tilde{s} = 0$, the O(D,D) covariant Lagrangian (62) is written as

$$\begin{aligned} \frac{1}{2g} J_{SD}^A \hat{\eta}_{AB} J_{SD}^B &= \frac{1}{2g} (\dot{X} - s X')^A \hat{\eta}_{AB} (\dot{X} - s X')^B - \frac{g}{2} X'^C \eta_{CD} \hat{\eta}^{DA} \hat{\eta}_{AB} \hat{\eta}^{BF} \eta_{FG} X'^G \\ &= \frac{1}{2g} (\dot{X} - s X')^M E_M^A \hat{\eta}_{AB} E_N^B (\dot{X} - s X')^N - \frac{g}{2} X'^M \eta_{ML} E_A^L \hat{\eta}^{AB} E_B^K \eta_{KN} X'^N \\ &= \frac{1}{2g} \varepsilon_+ X^M G_{MN} \varepsilon_- X^N \\ &= \frac{1}{2g} \varepsilon_+ X^M E_M^A \hat{\eta}_{AB} E_N^B \varepsilon_- X^N. \end{aligned} \quad (67)$$

The orthogonality condition is used in the second equality, $\eta_{ML} G^{LK} \eta_{KN} = G_{MN}$, so $E_M^A \eta_{AB} = \eta_{MN} E_B^N$, is used in the last equality. In terms of x^m, y_m coordinates it is given by

$$\begin{aligned} \frac{1}{2g} J_{SD}^A \hat{\eta}_{AB} J_{SD}^B &= \frac{1}{2g} (\varepsilon_+ x^m \varepsilon_+ y_m) \begin{pmatrix} g_{mn} - B_{ml} g^{lk} B_{kn} & -B_{ml} g^{ln} \\ g^{ml} B_{ln} & g^{mn} \end{pmatrix} \begin{pmatrix} \varepsilon_- x^n \\ \varepsilon_- y_n \end{pmatrix} \\ &= \frac{1}{2g} (\varepsilon_+ x^m \varepsilon_+ y_m) \begin{pmatrix} e_m^a & -B_{ml} e_a^l \\ 0 & e_a^m \end{pmatrix} \begin{pmatrix} \eta_{ab} & 0 \\ 0 & \eta^{ab} \end{pmatrix} \begin{pmatrix} e_n^b & 0 \\ -B_{nk} e_b^k & e_b^n \end{pmatrix} \begin{pmatrix} \varepsilon_- x^n \\ \varepsilon_- y_n \end{pmatrix} \\ &= \frac{1}{2g} [\varepsilon_+ x^m g_{mn} \varepsilon_- x^n + (\varepsilon_+ y_m - \varepsilon_+ x^l B_{lm}) g^{mn} (\varepsilon_- y_n + B_{nk} \varepsilon_- x^k)]. \end{aligned} \quad (68)$$

We break the O(D,D) symmetry into the GL(D) symmetry by the dimensional reduction (2). The resultant Lagrangian is the kinetic term of the usual string with the zweibein field;

$$L_0 = \frac{1}{2g} \varepsilon_+ x^m g_{mn} \varepsilon_- x^n. \quad (69)$$

In order to obtain the Wess-Zumino term we add the total derivative term

$$\partial_{\mu}(\varepsilon^{\mu\nu} x^m \partial_{\nu} y_m) = \dot{x} y' - x' \dot{y} = -\frac{1}{2g} (\varepsilon_+ x \varepsilon_- y - \varepsilon_+ y \varepsilon_- x), \quad (70)$$

to the O(D,D) Lagrangian L (68)

$$\begin{aligned} \frac{1}{2g} J_{SD}^A \hat{\eta}_{AB} J_{SD}^B - \partial_{\mu}(\varepsilon^{\mu\nu} x^m \partial_{\nu} y_m) &= \frac{1}{2g} \{ \varepsilon_+ x^m g_{mn} \varepsilon_- x^n + (\varepsilon_+ y_m - \varepsilon_+ x^l B_{lm} - \varepsilon_+ x^l g_{lm}) g^{mn} (\varepsilon_- y_n + B_{nk} \varepsilon_- x^k + g_{nk} \varepsilon_- x^k) \\ &\quad + \varepsilon_+ x^m g_{mn} \varepsilon_- x^n + 2\varepsilon_+ x^m B_{mn} \varepsilon_- x^n \}. \end{aligned} \quad (71)$$

By the dimensional reduction (2) the Lagrangian with the total derivative term reduces into the string Lagrangian in curved background with the Wess-Zumino term as the curved world-sheet version of (43),

$$L_0 + L_{\text{WZ}} = \frac{1}{g} \varepsilon_+ x^m (g_{mn} + B_{mn}) \varepsilon_- x^n. \quad (72)$$

The zweibeins in (72) and (42) are related as

$$g = -\frac{2}{\sqrt{-h}h^{00}}, \quad s = -\frac{h^{01}}{h^{00}}. \quad (73)$$

5 Lagrangians of string via \mathcal{T} -string from $\mathcal{A}5$ -brane

In this section, we derive the \mathcal{T} -string Lagrangian from the $\mathcal{A}5$ -brane Lagrangian. The resulting \mathcal{T} -string Lagrangian is formulated in terms of an $\text{SL}(4)$ rank-two antisymmetric tensor coordinate, which is coupled to the string background. We then present the reduction procedure from the \mathcal{T} -string Lagrangian to the conventional string Lagrangian.

5.1 \mathcal{T} -string from $\mathcal{A}5$ -brane

The $\text{O}(3,3)$ \mathcal{T} -string from $\mathcal{A}5$ -brane is described by the $\text{SL}(4)$ rank-two anti-symmetric tensor coordinate $X^{\underline{mn}} = (x^{\underline{m}}, y^{\underline{m}\bar{n}})$ with $\underline{m} = 1, \dots, 4$ and $\bar{m} = 1, 2, 3$ as listed in Tab. 1. The $\text{SL}(6)$ rank-two tensor coordinate is decomposed as $\text{SL}(6) \rightarrow \text{SL}(5) \rightarrow \text{SL}(4)$ as $X^{\hat{m}\hat{n}} = (X^{0n} = Y^n, X^{mn}) \rightarrow X^{mn} = (X^{5n} = Y^{\underline{m}}, X^{\underline{mn}}) \rightarrow X^{\underline{mn}} = (X^{4\bar{m}} = x^{\bar{m}}, X^{\bar{m}\bar{n}} = y^{\bar{m}\bar{n}})$ with $\hat{m} = 0, 1, \dots, 5$ and $m = 1, \dots, 5$. The 6-dimensional world-volume derivative is reduced into the string world-sheet derivatives as $\partial^{\hat{m}} = (\partial^0 = \partial_\tau, \partial^5 = \partial_\sigma, \partial^{\underline{m}} = 0)$. The $\text{SL}(6)$ field strength for the \mathcal{T} -string has the following components

$$\overset{\circ}{F}^{0\underline{mn}} = \partial_\tau X^{\underline{mn}}, \quad \overset{\circ}{F}^{5\underline{mn}} = \partial_\sigma X^{\underline{mn}}, \quad \overset{\circ}{F}^{05\underline{m}} = 0 = \overset{\circ}{F}^{\underline{mnl}}. \quad (74)$$

The $\text{SL}(6)$ vielbein for the \mathcal{T} -string has a block diagonal form as

$$E_{\hat{m}}^{\hat{a}} = \begin{pmatrix} E_0^{\hat{0}} & E_0^{\hat{5}} & E_0^{\underline{a}} \\ E_5^{\hat{0}} & E_5^{\hat{5}} & E_5^{\underline{a}} \\ E_{\underline{m}}^{\hat{0}} & E_{\underline{m}}^{\hat{5}} & E_{\underline{m}}^{\underline{a}} \end{pmatrix} = \begin{pmatrix} \frac{1}{g} & 0 & 0 \\ -\frac{s}{g} & 1 & 0 \\ 0 & 0 & g^{1/4} E_{\underline{m}}^{\underline{a}} \end{pmatrix}. \quad (75)$$

The selfdual and the anti-selfdual currents are the following combinations of the $\text{SL}(6)$ field strengths in (74) with (38) as

$$J_{\text{SD}/\overline{\text{SD}}}^{a_1 a_2} = g \left(F^{\hat{0} a_1 a_2} \pm \frac{1}{2} \epsilon^{\hat{0} a_1 a_2}{}_{5 a_3 a_4} F^{\hat{5} a_3 a_4} \right). \quad (76)$$

The zweibein fields g and s are part of the $\text{SL}(6)$ vielbein (75) in the new $\text{SL}(6)$ duality symmetry formulation in (38), contrast to that the world-volume vielbein fields are separated from the $\text{SL}(4)$ spacetime vielbein $E_{\underline{m}}^{\underline{a}}$ in the $\text{SL}(5)$ formulation in (31) and (32) as

$$\begin{aligned} J_{\text{SD}/\overline{\text{SD}}}^{a_1 a_2} &= \overset{\circ}{J}_{\text{SD}/\overline{\text{SD}}}^{m_1 m_2} E_{\underline{m}_1}^{a_1} E_{\underline{m}_2}^{a_2}, \\ \overset{\circ}{J}_{\text{SD}/\overline{\text{SD}}}^{m_1 m_2} &= \varepsilon_\tau X^{m_1 m_2} \pm \frac{1}{2} \hat{\eta}^{m_1 n_1} \hat{\eta}^{m_2 n_2} (-\epsilon_{n_1 \dots n_4}) \varepsilon_\sigma X^{n_3 n_4}, \end{aligned} \quad (77)$$

with (64). The minus sign in the last equation is caused from $\epsilon_{\tilde{m}_1\tilde{m}_2\tilde{m}_3}x^{\tilde{m}_3} = -\epsilon_{\tilde{m}_1\tilde{m}_24\tilde{m}_3}X^{4\tilde{m}_3}$. The $O(3,3)$ invariant metric η_{MN} becomes $SL(4)$ invariant metric $\epsilon_{\underline{m}_1\cdots\underline{m}_4}$. The current in (77) is written in terms of $x^{\tilde{m}}$ and $y^{\tilde{m}\tilde{n}}$ as

$$\begin{cases} \overset{\circ}{J}_{SD/\overline{SD}}^{4\tilde{m}} = \dot{x}^{\tilde{m}} - s x'^{\tilde{m}} \pm g \frac{1}{2} \epsilon^{\tilde{m}}_{\tilde{n}_1\tilde{n}_2} y'^{\tilde{n}_1\tilde{n}_2} = \epsilon_\tau x^{\tilde{m}} \pm \frac{1}{2} \epsilon^{\tilde{m}}_{\hat{n}_1\hat{n}_2} \epsilon_\sigma y^{\hat{n}_1\hat{n}_2}, \\ \overset{\circ}{J}_{SD/\overline{SD}}^{\tilde{m}_1\tilde{m}_2} = \dot{y}^{\tilde{m}_1\tilde{m}_2} - s y'^{\tilde{m}_1\tilde{m}_2} \pm g \epsilon^{\tilde{m}_1\tilde{m}_2}_{\tilde{n}} x'^{\tilde{n}} = \epsilon_\tau y^{\tilde{m}_1\tilde{m}_2} \pm \epsilon^{\tilde{m}_1\tilde{m}_2}_{\hat{n}} \epsilon_\sigma x^{\hat{n}}, \end{cases} \quad (78)$$

which is related to the $O(D,D)$ vector currents (65) with $y^{\tilde{m}\tilde{n}} \equiv \epsilon^{\tilde{m}\tilde{n}\tilde{l}} y_{\tilde{l}}$.

In order to obtain the usual 3-dimensional string Lagrangian we express the spacetime vielbein $E_{\underline{m}}^{\underline{a}} \in SL(4)/SO(4)$ in terms of the 3-dimensional metric $g_{\tilde{m}\tilde{n}}$ and the $B_{\tilde{m}\tilde{n}}$ field. The $O(3,3)$ vector index contraction and the $SL(4)$ tensor index contraction are assumed to be equal up to the normalization as

$$dX^M E_M^A = dx^{\tilde{m}} E_{\tilde{m}}^A + dy_{\tilde{m}} E^{\tilde{m};A} = dX^{4\tilde{x}} E_{4\tilde{m}}^A + \frac{1}{2} dX^{\tilde{m}\tilde{n}} E_{\tilde{m}\tilde{n}}^A = \frac{1}{2} dX^{\underline{mn}} E_{\underline{mn}}^A. \quad (79)$$

We rewrite the $O(D,D)$ vielbein in (47) in terms of tensor indices for $D=3$ case as

$$\begin{aligned} E_M^A &= \begin{pmatrix} e_{\tilde{m}}^{\tilde{a}} & -B_{\tilde{m}\tilde{l}} e_{\tilde{b}}^{\tilde{l}} \epsilon^{\tilde{a}_1\tilde{a}_2\tilde{b}} \\ 0 & \epsilon_{\tilde{m}_1\tilde{m}_2\tilde{m}} e_{\tilde{b}}^{\tilde{m}} \epsilon^{\tilde{a}_1\tilde{a}_2\tilde{b}} \end{pmatrix} \\ &= c \begin{pmatrix} E_{4\tilde{m}}^{4\tilde{a}} & E_{4\tilde{m}}^{\tilde{a}_1\tilde{a}_2} \\ E_{\tilde{m}_1\tilde{m}_2}^{4\tilde{a}} & E_{\tilde{m}_1\tilde{m}_2}^{\tilde{a}_1\tilde{a}_2} \end{pmatrix} = c E_{\underline{m}_1\underline{m}_2}^{\underline{a}_1\underline{a}_2} = c E_{[\underline{m}_1}^{\underline{a}_1} E_{\underline{m}_2]}^{\underline{a}_2}, \end{aligned} \quad (80)$$

with a normalization factor c . The vielbein with the tensor indices can be written as the product of the one with the vector indices

$$\begin{aligned} E_{\underline{m}}^{\underline{a}} &= \begin{pmatrix} E_{\tilde{m}}^{4\tilde{a}} & E_{\tilde{m}}^{\tilde{a}} \\ E_{\tilde{m}}^{4\tilde{a}} & E_{\tilde{m}}^{\tilde{a}} \end{pmatrix} = \mathbf{e}^{-1/4} \begin{pmatrix} 1 & -\tilde{B}^{\tilde{n}} e_{\tilde{n}}^{\tilde{a}} \\ 0 & e_{\tilde{m}}^{\tilde{a}} \end{pmatrix}, \\ \tilde{B}^{\tilde{m}} &= \frac{1}{2} \epsilon^{\tilde{m}\tilde{n}\tilde{l}} B_{\tilde{n}\tilde{l}}, \\ \mathbf{e} &= \det e_{\tilde{m}}^{\tilde{a}}. \end{aligned} \quad (81)$$

The background gauge field in the tensor index is now

$$\begin{aligned} G_{MN} &= G_{\underline{m}_1\underline{m}_2;\underline{n}_1\underline{n}_2} = \frac{1}{22} E_{\underline{m}_1\underline{m}_2}^{\underline{a}_1\underline{a}_2} \hat{\eta}_{\underline{a}_1[\underline{b}_1} \hat{\eta}_{\underline{b}_2]\underline{a}_2} E_{\underline{n}_1\underline{n}_2}^{\underline{b}_1\underline{b}_2} \\ &= \begin{pmatrix} G_{\tilde{m}\tilde{n}} & G_{\tilde{m};\tilde{n}_1\tilde{n}_2} \\ G_{\tilde{m}_1\tilde{m}_2;\tilde{n}} & G_{\tilde{m}_1\tilde{m}_2;\tilde{n}_1\tilde{n}_2} \end{pmatrix} \\ &= \mathbf{e}^{-1} \begin{pmatrix} g_{\tilde{m}\tilde{n}} - \tilde{B}^{\tilde{p}} g_{\tilde{m}[\tilde{p}} g_{\tilde{n}]\tilde{q}} \tilde{B}^{\tilde{q}} & g_{\tilde{m}[\tilde{n}_1} g_{\tilde{n}_2]\tilde{l}} \tilde{B}^{\tilde{l}} \\ -\tilde{B}^{\tilde{p}} g_{\tilde{p}[\tilde{m}_1} g_{\tilde{m}_2]\tilde{n}} & g_{\tilde{m}_1[\tilde{n}_1} g_{\tilde{n}_2]\tilde{m}_2} \end{pmatrix}, \end{aligned} \quad (82)$$

where metric of the stability group is denoted as $\hat{\eta}_{\underline{mn}}$ to distinguish from η_{MN} .

The \mathcal{T} -string Lagrangian is obtained from the world-volume covariant $\mathcal{A}5$ -Lagrangian (40)

$$\begin{aligned} I &= \int d^2\sigma L, \\ L &= \frac{\Phi}{2} \left(-(F^{\hat{0}\underline{a}_1\underline{a}_2})^2 + (F^{\hat{5}\underline{a}_1\underline{a}_2})^2 \right) + \frac{1}{2} \Lambda_{\hat{0}\hat{5}} F^{\hat{0}\underline{a}_1\underline{a}_2} F^{\hat{5}}_{\underline{a}_1\underline{a}_2} + \frac{1}{2} \Lambda_{\underline{ab}} F^{\underline{a}\hat{0}\underline{c}} F^{\underline{b}}_{\hat{0}\underline{c}} + \frac{1}{2} \Lambda_{\underline{ab}} F^{\underline{a}\hat{5}\underline{c}} F^{\underline{b}}_{\hat{5}\underline{c}} \\ &\quad + \frac{\epsilon^{\hat{0}\hat{5}\underline{a}_1\cdots\underline{a}_4}}{4} \left(\tilde{\Lambda}_{\hat{0}}^{\hat{0}} F^{\hat{5}\underline{a}_1\underline{a}_2} F^{\hat{0}\underline{a}_3\underline{a}_4} + \tilde{\Lambda}_{\hat{5}}^{\hat{0}} F^{\hat{5}\underline{a}_1\underline{a}_2} F^{\hat{5}\underline{a}_3\underline{a}_4} - \tilde{\Lambda}_{\hat{0}}^{\hat{5}} F^{\hat{0}\underline{a}_1\underline{a}_2} F^{\hat{0}\underline{a}_3\underline{a}_4} - \tilde{\Lambda}_{\hat{5}}^{\hat{5}} F^{\hat{0}\underline{a}_1\underline{a}_2} F^{\hat{5}\underline{a}_3\underline{a}_4} \right), \end{aligned} \quad (83)$$

with $\eta^{\hat{0}\hat{0}} = -1$ and $\eta^{\hat{5}\hat{5}} = 1$. Although currents are written as field strengths, there is no gauge symmetry of the coordinate δX^{mn} . The \mathcal{T} -string Lagrangian in the SL(4) tensor coordinate is given by

$$\begin{aligned} L &= \phi \frac{1}{2^2} J_{SD}^{a_1 a_2} \hat{\eta}_{a_1 [b_1} \hat{\eta}_{b_2] a_2} J_{SD}^{b_1 b_2} + \bar{\phi} \frac{1}{2^2} J_{SD}^{a_1 a_2} \hat{\eta}_{a_1 [b_1} \hat{\eta}_{b_2] a_2} J_{SD}^{b_1 b_2} \\ &\quad + \tilde{\phi} \frac{1}{2^2} J_{SD}^{a_1 a_2} \epsilon_{a_1 \dots a_4} J_{SD}^{a_3 a_4} \\ &= \phi \frac{1}{2^2} \overset{\circ}{J}_{SD}^{m_1 m_2} G_{m_1 m_2; n_1 n_2} \overset{\circ}{J}_{SD}^{n_1 n_2} + \bar{\phi} \frac{1}{2^2} \overset{\circ}{J}_{SD}^{m_1 m_2} G_{m_1 m_2; n_1 n_2} \overset{\circ}{J}_{SD}^{n_1 n_2} \\ &\quad + \tilde{\phi} \frac{1}{2^2} \overset{\circ}{J}_{SD}^{m_1 m_2} \epsilon_{m_1 \dots m_4} \overset{\circ}{J}_{SD}^{m_3 m_4}, \end{aligned} \quad (84)$$

with the background metric $G_{m_1 m_2; n_1 n_2}$ in (82).

The \mathcal{T} -string Lagrangian in the gauge $\Phi = g^2 \mathbf{e}$ and $\Lambda_{\hat{a}\hat{b}} = 0 = \tilde{\Lambda}_{\hat{a}}^{\hat{b}}$ as

$$L = -\frac{g^2 \mathbf{e}}{2} \left((F^{\hat{0}a_1 a_2})^2 - (F^{\hat{5}a_1 a_2})^2 \right). \quad (85)$$

The SL(4) covariant Lagrangian (84) in the gauge $\phi = \frac{1}{2g}$ and $\bar{\phi} = 0 = \tilde{\phi}$ is given as

$$\begin{aligned} L &= \frac{1}{2g} J_{SD}^A \hat{\eta}_{AB} J_{SD}^B \\ &= \frac{1}{2^3 g} \varepsilon_+ X^{m_1 m_2} G_{m_1 m_2; n_1 n_2} \varepsilon_- X^{n_1 n_2} \\ &= \frac{1}{2g} (\varepsilon_+ x^{\tilde{m}} \varepsilon_+ y^{\tilde{m}_1 \tilde{m}_2}) \begin{pmatrix} g_{\tilde{m}\tilde{n}} - \tilde{B}^{\tilde{p}} g_{\tilde{m}[\tilde{p}} g_{\tilde{n}]\tilde{q}} \tilde{B}^{\tilde{q}} & g_{\tilde{m}[\tilde{n}_1} g_{\tilde{n}_2]\tilde{l}} \tilde{B}^{\tilde{l}} \\ -\tilde{B}^{\tilde{p}} g_{\tilde{p}[\tilde{m}_1} g_{\tilde{m}_2]\tilde{n}} & g_{\tilde{m}_1[\tilde{n}_1} g_{\tilde{n}_2]\tilde{m}_2} \end{pmatrix} \begin{pmatrix} \varepsilon_- x^{\tilde{m}} \\ \varepsilon_- y^{\tilde{m}_1 \tilde{m}_2} \end{pmatrix}. \end{aligned} \quad (86)$$

5.2 String from \mathcal{T} -string

We break SL(4) symmetry of \mathcal{T} -string into GL(3) for the usual string, where the reduction of the spacetime coordinate is performed as $X^{mn} = (X^{4\tilde{m}}, X^{\tilde{m}\tilde{n}}) = (x^{\tilde{m}}, y^{\tilde{m}\tilde{n}}) \rightarrow x^{\tilde{m}}$. We repeat the same procedure of subsection 4.2. The SL(4) Lagrangian (86) is rewritten analogously to (67)

$$\begin{aligned} \frac{1}{2g} J_{SD}^A \hat{\eta}_{AB} J_{SD}^B &= \frac{1}{2g} \varepsilon_+ x^{\tilde{m}} g_{\tilde{m}\tilde{n}} \varepsilon_- x^{\tilde{n}} \\ &\quad + \frac{1}{2^3 g} (\varepsilon_+ y^{\tilde{m}_1 \tilde{m}_2} - \varepsilon_+ x^{[\tilde{m}_1} \tilde{B}^{\tilde{m}_2]}) g_{\tilde{m}_1[\tilde{n}_1} g_{\tilde{n}_2]\tilde{m}_2} (\varepsilon_- y^{\tilde{n}_1 \tilde{n}_2} + \tilde{B}^{[\tilde{n}_1} \varepsilon_- x^{\tilde{n}_2]}) \end{aligned} \quad (87)$$

By the dimensional reduction (2) the Lagrangian (86) reduces to the kinetic term of the string (69).

The total derivative term which is added to obtain the Wess-Zumino term (70) becomes

$$\begin{aligned} -\frac{1}{2^2 g} (\varepsilon_+ x^{\tilde{m}_1} \varepsilon_- y^{\tilde{m}_2 \tilde{m}_3} - \varepsilon_- x^{\tilde{m}_1} \varepsilon_+ y^{\tilde{m}_2 \tilde{m}_3}) \epsilon_{\tilde{m}_1 \tilde{m}_2 \tilde{m}_3} &= -\frac{1}{2} \partial_\mu (\epsilon^{\mu\nu} x^{\tilde{m}_1} \partial_\nu y^{\tilde{m}_2 \tilde{m}_3} \epsilon_{\tilde{m}_1 \tilde{m}_2 \tilde{m}_3}) \\ &= \frac{1}{2^2} \partial_\mu (\epsilon^{\mu\nu} X^{m_1 m_2} \partial_\nu X^{m_3 m_4} \epsilon_{m_1 \dots m_4}). \end{aligned} \quad (88)$$

Adding this term to the SL(4) Lagrangian (86)

$$\begin{aligned} &\frac{1}{2g} J_{SD}^A \hat{\eta}_{AB} J_{SD}^B + \frac{1}{2^2} \partial_\mu (\epsilon^{\mu\nu} X^{m_1 m_2} \partial_\nu X^{m_3 m_4} \epsilon_{m_1 \dots m_4}) \\ &= \frac{1}{2g} \left\{ \varepsilon_+ x^{\tilde{m}} g_{\tilde{m}\tilde{n}} \varepsilon_- x^{\tilde{n}} + \frac{1}{2^2} (\varepsilon_+ y^{\tilde{m}_1 \tilde{m}_2} - \varepsilon_+ x^{[\tilde{m}_1} \tilde{B}^{\tilde{m}_2]}) g_{\tilde{m}_1[\tilde{n}_1} g_{\tilde{n}_2]\tilde{m}_2} \right. \\ &\quad \times (\varepsilon_- y^{\tilde{n}_1 \tilde{n}_2} + \tilde{B}^{[\tilde{n}_1} \varepsilon_- x^{\tilde{n}_2]}) - g^{\tilde{n}_1 \tilde{k}_1} g^{\tilde{n}_2 \tilde{k}_2} \epsilon_{\tilde{k}_1 \tilde{k}_2 \tilde{k}_3} \varepsilon_- x^{\tilde{k}_3} + \varepsilon_+ x^{\tilde{m}} g_{\tilde{m}\tilde{n}} \varepsilon_- x^{\tilde{n}} + 2\varepsilon_+ x^{\tilde{m}} B_{\tilde{m}\tilde{n}} \varepsilon_- x^{\tilde{n}} \left. \right\}. \end{aligned} \quad (89)$$

After the dimensional reduction (2), the Lagrangian with the total derivative term reduces into the usual string Lagrangian with the Wess-Zumino term (72),

$$L_0 + L_{\text{WZ}} = \frac{1}{g} \varepsilon_+ x^{\bar{m}} (g_{\bar{m}\bar{n}} + B_{\bar{m}\bar{n}}) \varepsilon_- x^{\bar{n}}.$$

6 Lagrangians of M2-brane via $\mathcal{M}5$ -brane from $\mathcal{A}5$ -brane

6.1 $\mathcal{M}5$ -brane from $\mathcal{A}5$ -brane

The GL(4) $\mathcal{M}5$ -brane from $\mathcal{A}5$ -brane is described by the GL(4) vector coordinate $X^{5\bar{m}} = x^{\bar{m}}$ [18] as listed in Tab. 1. The SL(6) rank-two tensor coordinate is decomposed as $SL(6) \rightarrow SL(5) \rightarrow GL(4)$ as $X^{\hat{m}\hat{n}} = (X^{0n} = Y^n, X^{mn}) \rightarrow X^{mn} = (X^{5\bar{n}} = x^{\bar{m}}, X^{\bar{m}\bar{n}} = y^{\bar{m}\bar{n}})$ and $Y^m = (Y^5 = Y, Y^{\bar{m}})$. The 6-dimensional world-volume derivative is reduced into the 5-brane world-sheet derivatives as $\partial^{\hat{m}} = (\partial^0 = \partial_\tau, \partial^5 = 0, \partial^{\bar{m}} = \partial_\sigma^{\bar{m}})$. The SL(6) field strength for the $\mathcal{M}5$ -brane has the following components

$$\begin{cases} \overset{\circ}{F}^{05\bar{m}} = \partial_\tau x^{\bar{m}} + \partial^{\bar{m}} Y, \\ \overset{\circ}{F}^{5\bar{m}\bar{n}} = -\partial^{[\bar{m}} x^{\bar{n}]}, \\ \overset{\circ}{F}^{0m\bar{n}} = \partial_\tau y^{\bar{m}\bar{n}} - \partial^{[\bar{m}} Y^{\bar{n}]}, \\ \overset{\circ}{F}^{m\bar{n}\bar{l}} = \frac{1}{2} \partial^{[\bar{m}} y^{\bar{n}\bar{l}]}, \end{cases} \quad (90)$$

where the auxiliary coordinates $y^{\bar{m}\bar{n}}$ and $Y^{\bar{m}}$ are preserved to begin with the SL(5) A -symmetric \mathcal{M} -theory Lagrangian [30].

The SL(6) vielbein for the $\mathcal{M}5$ -brane with SL(5) A -symmetry is given by

$$E_{\hat{m}}^{\hat{a}} = \begin{pmatrix} E_0^{\hat{0}} & E_0^a \\ E_m^{\hat{0}} & E_m^a \end{pmatrix} = \begin{pmatrix} \frac{1}{g} & 0 \\ -\frac{s_m}{g} & g^{1/5} E_m^a \end{pmatrix}. \quad (91)$$

It is stressed that the world-volume vielbein fields g, s_m and the spacetime vielbein E_m^a cannot be in block diagonal form unlike \mathcal{T} -string case (75). The selfdual and anti-selfdual currents in curved background given by (32) based on (31) are the following combination of the SL(6) field strengths in (90) with (38) as

$$F_{\text{SD}/\overline{\text{SD}}}^{a_1 a_2} = g \left(F^{\hat{0} a_1 a_2} \pm \frac{1}{3!} \epsilon^{\hat{0} a_1 a_2}{}_{a_3 a_4 a_5} F^{a_3 a_4 a_5} \right). \quad (92)$$

The GL(4) covariant selfdual and anti-selfdual currents in flat space are derived from the ones of SL(5) (31) given in [1] as

$$\begin{cases} \overset{\circ}{F}_{\text{SD}/\overline{\text{SD}}}^{\bar{m}} &= F_\tau^{\bar{m}} - \frac{1}{2} \epsilon^{mn_1 n_2 n_3} s_{n_1} F_{\sigma; n_2 n_3} \pm g \hat{\eta}^{mn} F_{\sigma; \bar{n}} \\ &= \varepsilon_\tau x^{\bar{m}} \pm g \hat{\eta}^{mn} (\varepsilon_\sigma Y)_{\bar{n}}, \\ \overset{\circ}{F}_{\text{SD}/\overline{\text{SD}}}^{\bar{m}_1 \bar{m}_2} &= F_\tau^{\bar{m}_1 \bar{m}_2} + \epsilon^{\bar{m}_1 \dots \bar{m}_4} s_{\bar{m}_3} F_{\sigma; \bar{m}_4} - \frac{1}{2} \epsilon^{\bar{m}_1 \dots \bar{m}_4} s_5 F_{\sigma; \bar{m}_3 \bar{m}_4} \pm g \hat{\eta}^{\bar{m}_1 \bar{n}_1} \hat{\eta}^{\bar{m}_2 \bar{n}_2} F_{\sigma; \bar{n}_1 \bar{n}_2} \\ &= \varepsilon_\tau y^{\bar{m}_1 \bar{m}_2} \pm g \hat{\eta}^{\bar{m}_1 \bar{n}_1} \hat{\eta}^{\bar{m}_2 \bar{n}_2} (\varepsilon_\sigma x)_{\bar{n}_1 \bar{n}_2}, \end{cases} \quad (93)$$

where $\hat{\eta}^{mn}$ becomes G^{mn} in a curved background. The brane world-volume derivatives are

given as a generalization of the world-sheet zweibein dependence in (64) as

$$\begin{cases} \varepsilon_\tau x^{\underline{m}} & \equiv F_\tau^{\underline{m}} - \frac{1}{2} \epsilon^{\underline{mn}_1 \underline{n}_2 \underline{n}_3} s_{\underline{n}_1} F_{\sigma; \underline{n}_2 \underline{n}_3} \\ & = \dot{x}^{\underline{m}} + \partial^{\underline{m}} Y + s_{\underline{n}} \partial^{[\underline{m}} x^{\underline{n}]}, \\ \varepsilon_\tau y^{\underline{m}_1 \underline{m}_2} & \equiv F_\tau^{\underline{m}_1 \underline{m}_2} + \epsilon^{\underline{m}_1 \dots \underline{m}_4} s_{\underline{m}_3} F_{\sigma; \underline{m}_4} \\ & = \dot{y}^{\underline{m}_1 \underline{m}_2} - \partial^{[\underline{m}_1} Y^{\underline{m}_2]} + \frac{1}{2} s_{\underline{m}_3} \partial^{[\underline{m}_1} y^{\underline{m}_2 \underline{m}_3]} + s_5 \partial^{[\underline{m}_1} x^{\underline{m}_2]}, \\ (\varepsilon_\sigma y)_{\underline{m}} & \equiv F_{\sigma; \underline{m}} \\ & = \frac{1}{2} \epsilon^{\underline{mn}_1 \underline{n}_2 \underline{n}_3} \partial^{\underline{n}_1} y^{\underline{n}_2 \underline{n}_3}, \\ (\varepsilon_\sigma x)_{\underline{m}_1 \underline{m}_2} & \equiv F_{\sigma; \underline{m}_1 \underline{m}_2} \\ & = -\epsilon^{\underline{m}_1 \dots \underline{m}_4} \partial^{\underline{m}_3} x^{\underline{m}_4}. \end{cases} \quad (94)$$

The 11-dimensional supergravity background includes the gravitational metric g_{mn} and the three form gauge field C_{mnl} . We focus on the 4-dimensional subspace of the 11-dimensional space, where the background fields are g_{mn} and C_{mnl} whose number of degrees of freedom is $10 + 4 = 14$. The dimension of the coset $SL(5)/SO(5)$ is also $24 - 10 = 14$. The vector vielbein $E_m^a \in SL(5)/SO(5)$ with $GL(4)$ indices where $m = (5, \underline{m})$ and $\underline{m} = 1, \dots, 4$ is given by [65]

$$\begin{aligned} E_m^a &= \begin{pmatrix} E_5^5 & E_5^{\underline{a}} \\ E_{\underline{m}}^5 & E_{\underline{m}}^{\underline{a}} \end{pmatrix} = \begin{pmatrix} e^{3/5} & e^{-2/5} \tilde{C}^{\underline{n}} e_{\underline{n}}^{\underline{a}} \\ 0 & e^{-2/5} e_{\underline{m}}^{\underline{a}} \end{pmatrix}, \\ \tilde{C}^{\underline{m}} &= \frac{1}{3!} \epsilon^{\underline{m} \underline{m}_1 \underline{m}_2 \underline{m}_3} C_{\underline{m}_1 \underline{m}_2 \underline{m}_3}, \\ e &= \det e_{\underline{m}}^{\underline{a}}, \end{aligned} \quad (95)$$

with $\det E_m^a = 1 = \epsilon^{m_1 \dots m_5} E_{m_1}^{a_1} E_{m_2}^{a_2} E_{m_3}^{a_3} E_{m_4}^{a_4} E_{m_5}^{a_5} = \epsilon^{a_1 \dots a_5}$. The tensor vielbein is the product of the vector vielbein (95) as

$$\begin{aligned} E_{m_1 m_2}^{a_1 a_2} &= E_{[m_1}^{a_1} E_{m_2]}^{a_2} \\ &= \begin{pmatrix} E_{5 \underline{m}}^{5 \underline{a}} & E_{5 \underline{m}}^{\underline{a}_1 \underline{a}_2} \\ E_{\underline{m}_1 \underline{m}_2}^{5 \underline{a}} & E_{\underline{m}_1 \underline{m}_2}^{\underline{a}_1 \underline{a}_2} \end{pmatrix} = \begin{pmatrix} e^{1/5} e_{\underline{m}}^{\underline{a}} & -e^{-4/5} \tilde{C}^{\underline{n}} e_{\underline{m}}^{[\underline{a}_1} e_{\underline{n}}^{\underline{a}_2]} \\ 0 & e^{-4/5} e_{\underline{m}_1}^{[\underline{a}_1} e_{\underline{m}_2}^{\underline{a}_2]} \end{pmatrix}. \end{aligned} \quad (96)$$

The background gauge field in tensor index is now

$$\begin{aligned} G_{MN} &= G_{m_1 m_2; n_1 n_2} = \frac{1}{22} E_{m_1 m_2}^{a_1 a_2} \hat{\eta}_{a_1 [b_1} \hat{\eta}_{b_2] a_2} E_{n_1 n_2}^{b_1 b_2} \\ &= \begin{pmatrix} G_{\underline{mn}} & G_{\underline{m}; \underline{n}_1 \underline{n}_2} \\ G_{\underline{m}_1 \underline{m}_2; \underline{n}} & G_{\underline{m}_1 \underline{m}_2; \underline{n}_1 \underline{n}_2} \end{pmatrix} \\ &= e^{-8/5} \begin{pmatrix} e^2 g_{\underline{mn}} - \tilde{C}^{\underline{p}} g_{\underline{m} [\underline{p}} g_{\underline{n}] \underline{q}} \tilde{C}^{\underline{q}} & \tilde{C}^{\underline{l}} g_{\underline{l} [\underline{n}_1} g_{\underline{n}_2] \underline{m}} \\ g_{\underline{p} [\underline{m}_1} g_{\underline{m}_2] \underline{n}} \tilde{C}^{\underline{p}} & g_{\underline{m}_1 [\underline{n}_1} g_{\underline{n}_2] \underline{m}_2} \end{pmatrix} \\ &= e^{2/5} \begin{pmatrix} g_{\underline{mn}} & 0 \\ 0 & 0 \end{pmatrix} + e^{-8/5} \begin{pmatrix} -\tilde{C}^{\underline{p}} g_{\underline{m} [\underline{p}} g_{\underline{n}] \underline{q}} \tilde{C}^{\underline{q}} & \tilde{C}^{\underline{l}} g_{\underline{l} [\underline{n}_1} g_{\underline{n}_2] \underline{m}} \\ g_{\underline{p} [\underline{m}_1} g_{\underline{m}_2] \underline{n}} \tilde{C}^{\underline{p}} & g_{\underline{m}_1 [\underline{n}_1} g_{\underline{n}_2] \underline{m}_2} \end{pmatrix}. \end{aligned} \quad (97)$$

Inverse of these background gauge fields are given by [65]

$$E_a^m = \begin{pmatrix} E_5^5 & E_5^m \\ E_a^5 & E_a^m \end{pmatrix} = \begin{pmatrix} e^{-3/5} & e^{-3/5} \tilde{C}^m \\ 0 & e^{2/5} e_a^m \end{pmatrix}, \quad (98)$$

$$E_{a_1 a_2}^{m_1 m_2} = E_{[a_1}^{m_1} E_{a_2]}^{m_2} \\ = \begin{pmatrix} E_{5a}^{5m} & E_{5a}^{m_1 m_2} \\ E_{a_1 a_2}^{5m} & E_{a_1 a_2}^{m_1 m_2} \end{pmatrix} = \begin{pmatrix} e_a^m & -\tilde{C}^{[m_1} e_a^{m_2]} \\ 0 & e e_{a_1}^{[m_1} e_{a_2}^{m_2]} \end{pmatrix}, \quad (99)$$

$$G^{MN} = G^{m_1 m_2; n_1 n_2} = \frac{1}{2^2} E_{a_1 a_2}^{m_1 m_2} \hat{\eta}^{a_1 [b_1} \hat{\eta}^{b_2] a_2} E_{b_1 b_2}^{n_1 n_2} \\ = \begin{pmatrix} G^{mn} & G^{m; n_1 n_2} \\ G^{m_1 m_2; n} & G^{m_1 m_2; n_1 n_2} \end{pmatrix} \\ = e^{2/5} \begin{pmatrix} g^{mn} & g^{m[n_1} \tilde{C}^{n_2]} \\ -\tilde{C}^{[m_1} g^{m_2]n} & e^2 g^{m_1 [n_1} g^{n_2] m_2} + \tilde{C}^{[m_1} g^{m_2] [n_1} \tilde{C}^{n_2]} \end{pmatrix} \\ = e^{8/5} \begin{pmatrix} 0 & 0 \\ 0 & g^{m_1 [n_1} g^{n_2] m_2} \end{pmatrix} + e^{-2/5} \begin{pmatrix} g^{mn} & g^{m[n_1} \tilde{C}^{n_2]} \\ -\tilde{C}^{[m_1} g^{m_2]n} & \tilde{C}^{[m_1} g^{m_2] [n_1} \tilde{C}^{n_2]} \end{pmatrix}. \quad (100)$$

The $\mathcal{M}5$ -brane Lagrangian is given by the $SL(5)$ covariant Lagrangian (33) with replacing $GL(4)$ indices as

$$L = \frac{1}{2} \phi \left(F_{SD}^a F_{SDa} + \frac{1}{2} F_{SD}^{ab} F_{SDab} \right) + \frac{1}{2} \bar{\phi} \left((F_{SD}^a)^2 + \frac{1}{2} (F_{SD}^{ab})^2 \right) \\ + \frac{1}{2} \lambda F_{SD}^c F_{SDc} + \lambda_a F_{SD}^{ac} F_{SDc} + \frac{1}{2} \lambda_{ab} \left(F_{SD}^a F_{SD}^b + F_{SD}^{ac} F_{SD}^b{}_{c} \right) \\ - \epsilon_{a_1 \dots a_4} \left(\frac{1}{8} \lambda^5 F_{SD}^{a_1 a_2} F_{SD}^{a_3 a_4} - \frac{1}{2} \lambda^{a_1} F_{SD}^{a_2 a_3} F_{SD}^{a_4}{}_{a_1} \right). \quad (101)$$

The Lagrangian in terms of the curved currents $F_{SD/SD}^{ab}$ is simpler than the one in terms of the flat currents $\overset{\circ}{F}_{SD/SD}^{mn}$. The concrete expression of the Lagrangian of the $\mathcal{M}5$ -brane in a curved background (101) is given as follows. We begin by the $SL(5)$ covariant Lagrangian (33) in the gauge $\phi = \frac{1}{2g}$ and $\bar{\phi} = 0 = \lambda$'s

$$\frac{1}{2g} F_{SD}^A \hat{\eta}_{AB} F_{SD}^B = \frac{1}{8g} \overset{\circ}{F}_{SD}^{m_1 m_2} G_{m_1 m_2; n_1 n_2} \overset{\circ}{F}_{SD}^{n_1 n_2} \\ = \frac{1}{8g} \varepsilon_\tau X^{m_1 m_2} G_{m_1 m_2; n_1 n_2} \varepsilon_\tau X^{n_1 n_2} - \frac{g}{8} F_{\sigma m_1 m_2} G^{m_1 m_2; n_1 n_2} F_{\sigma n_1 n_2} \\ = \frac{1}{2g} (\varepsilon_\tau x^m \varepsilon_\tau y^{m_1 m_2}) \left[e^{2/5} \begin{pmatrix} g^{mn} & 0 \\ 0 & 0 \end{pmatrix} \right. \\ \left. + e^{-8/5} \begin{pmatrix} -\tilde{C}^p g_{m[p} g_{n]q} \tilde{C}^q & \tilde{C}^l g_{l[n_1} g_{n_2]m} \\ g_{p[m_1} g_{m_2]n} \tilde{C}^p & g_{m_1 [n_1} g_{n_2] m_2} \end{pmatrix} \right] \begin{pmatrix} \varepsilon_\tau x^n \\ \varepsilon_\tau y^{n_1 n_2} \end{pmatrix} \\ - \frac{g}{2} ((\varepsilon_\sigma y)_m (\varepsilon_\sigma x)_{m_1 m_2}) \left[e^{8/5} \begin{pmatrix} 0 & 0 \\ 0 & g^{m_1 [n_1} g^{n_2] m_2} \end{pmatrix} \right. \\ \left. + e^{-2/5} \begin{pmatrix} g^{mn} & g^{m[n_1} \tilde{C}^{n_2]} \\ -\tilde{C}^{[m_1} g^{m_2]n} & \tilde{C}^{[m_1} g^{m_2] [n_1} \tilde{C}^{n_2]} \end{pmatrix} \right] \begin{pmatrix} (\varepsilon_\sigma y)_n \\ (\varepsilon_\sigma x)_{n_1 n_2} \end{pmatrix} \\ = L_0 + L_y, \quad (102)$$

with

$$\begin{aligned} L_0 &= \frac{e^{2/5}}{2g} \varepsilon_\tau x^{\underline{m}} g_{\underline{mn}} \varepsilon_\tau x^{\underline{n}} - \frac{g e^{8/5}}{8} (\varepsilon_\sigma x)_{\underline{m}_1 \underline{m}_2} g^{\underline{m}_1 [\underline{n}_1} g^{\underline{n}_2] \underline{m}_2} (\varepsilon_\sigma x)_{\underline{n}_1 \underline{n}_2}, \\ L_y &= \frac{e^{-8/5}}{8g} (\varepsilon_\tau y^{\underline{m}_1 \underline{m}_2} - \varepsilon_\tau x^{[\underline{m}_1} \tilde{C}^{\underline{m}_2]}) g_{\underline{m}_1 [\underline{n}_1} g_{\underline{n}_2] \underline{m}_2} (\varepsilon_\tau y^{\underline{n}_1 \underline{n}_2} + \tilde{C}^{[\underline{n}_1} \varepsilon_\tau x^{\underline{n}_2]}) \\ &\quad - \frac{g e^{-2/5}}{2} ((\varepsilon_\sigma y)_{\underline{m}} + (\varepsilon_\sigma x)_{\underline{ml}} \tilde{C}^{\underline{l}}) g^{\underline{mn}} ((\varepsilon_\sigma y)_{\underline{n}} - \tilde{C}^{\underline{l}} (\varepsilon_\sigma x)_{\underline{ln}}). \end{aligned} \quad (103)$$

In the gauge $g = e^{-3/5}$ Lagrangians take simple form as

$$\begin{aligned} L_0 &= -\frac{e}{2} \left[F^{05a} \hat{\eta}_{ab} F^{05b} - \frac{1}{4} F^{5a_1 a_2} \hat{\eta}_{a_1 [b_1} \hat{\eta}_{b_2] a_2} F^{5b_1 b_2} \right], \\ L_y &= -\frac{1}{2e} \left[\frac{1}{2} F^{0a_1 a_2} \hat{\eta}_{a_1 b_1} \hat{\eta}_{b_2 a_2} F^{0b_1 b_2} - \frac{1}{6} F^{a_1 a_2 a_3} \hat{\eta}_{a_1 b_1} \hat{\eta}_{a_2 b_2} \hat{\eta}_{a_3 b_3} F^{b_1 b_2 b_3} \right], \end{aligned} \quad (104)$$

with $\tilde{m} = (0, \underline{m})$.

The SL(5) U-duality symmetry of the Lagrangian (104) is broken to GL(4) symmetry by the dimensional reduction similarly to (2). Then the kinetic term of the new perturbative Lagrangian for a $\mathcal{M}5$ -brane in the 4-dimensions is given by

$$\begin{aligned} L_0 &= \frac{e}{2} \left[(\dot{x}^{\underline{m}} + \partial^{\underline{m}} Y + s_{\underline{l}} \partial^{[\underline{m}} x^{\underline{l}]}) g_{\underline{mn}} (\dot{x}^{\underline{n}} + \partial^{\underline{n}} Y + s_{\underline{k}} \partial^{[\underline{n}} x^{\underline{k}]}) \right. \\ &\quad \left. - \frac{1}{4} \partial^{[\underline{m}_1} x^{\underline{m}_2]} g_{\underline{m}_1 [\underline{n}_1} g_{\underline{n}_2] \underline{m}_2} \partial^{[\underline{n}_1} x^{\underline{n}_2]} \right]. \end{aligned} \quad (105)$$

The total derivative terms to obtain the Wess-Zumino term for the $\mathcal{M}5$ -brane are given analogously to the string case (70) with the gauge $\partial^{\underline{m}} s_5 = 0$ as

$$\begin{aligned} &\varepsilon_\tau x^{\underline{m}} (\varepsilon_\sigma y)_{\underline{m}} - \frac{1}{2} (\varepsilon_\sigma x)_{\underline{m}_1 \underline{m}_2} \varepsilon_\tau y^{\underline{m}_1 \underline{m}_2} \\ &= \frac{1}{2} \epsilon_{\underline{m}_1 \dots \underline{m}_4} \left\{ \partial_\tau (x^{\underline{m}_1} \partial^{\underline{m}_2} y^{\underline{m}_3 \underline{m}_4}) \right. \\ &\quad \left. + \partial^{\underline{m}_1} (x^{\underline{m}_2} \dot{y}^{\underline{m}_3 \underline{m}_4} + Y \partial^{\underline{m}_2} y^{\underline{m}_3 \underline{m}_4} - 2x^{\underline{m}_2} \partial^{\underline{m}_3} Y^{\underline{m}_4} - 2s_5 x^{\underline{m}_2} \partial^{\underline{m}_3} x^{\underline{m}_4}) \right\}, \end{aligned} \quad (106)$$

where the $s_{\underline{n}}$ dependent terms are cancelled out because of the totally antisymmetry of 5 indices

$$\epsilon_{\underline{m}_1 \dots \underline{m}_4} s_{\underline{n}} \left((\partial^{[\underline{m}_1} x^{\underline{n}]}) \partial^{\underline{m}_2} y^{\underline{m}_3 \underline{m}_4} + \frac{1}{2} (\partial^{\underline{m}_4} x^{\underline{m}_1}) \partial^{[\underline{n}} y^{\underline{m}_2 \underline{m}_3]} \right) = \epsilon_{\underline{m}_1 \dots \underline{m}_4} s_{\underline{n}} \frac{1}{4!} \partial^{[\underline{m}_1} x^{\underline{n}} \partial^{\underline{m}_2} y^{\underline{m}_3 \underline{m}_4]} = 0. \quad (107)$$

Adding the total derivative term (106) to the $\mathcal{M}5$ -brane Lagrangian (101) in gauge $\phi = \frac{1}{2g}$, $g = 2e^{-3/5}$, $\bar{\phi} = 0 = \lambda$'s the Lagrangian for the $\mathcal{M}5$ -brane becomes

$$\begin{aligned} &\frac{1}{2g} F_{\text{SD}}^A \hat{\eta}_{AB} F_{\text{SD}}^B - \varepsilon_\tau x^{\underline{m}} (\varepsilon_\sigma y)_{\underline{m}} + \frac{1}{2} (\varepsilon_\sigma x)_{\underline{m}_1 \underline{m}_2} \varepsilon_\tau y^{\underline{m}_1 \underline{m}_2} = L_0 + L_y + L_{\text{WZ}}, \\ L_0 &= \frac{e}{4} \left[\varepsilon_\tau x^{\underline{m}} g_{\underline{mn}} \varepsilon_\tau x^{\underline{n}} - \frac{1}{4} (\varepsilon_\sigma x)_{\underline{m}_1 \underline{m}_2} g^{\underline{m}_1 [\underline{n}_1} g^{\underline{n}_2] \underline{m}_2} (\varepsilon_\sigma x)_{\underline{n}_1 \underline{n}_2} \right], \\ L_y &= \frac{1}{4e} \left[\frac{1}{4} (\varepsilon_\tau y^{\underline{m}_1 \underline{m}_2} - \varepsilon_\tau x^{[\underline{m}_1} \tilde{C}^{\underline{m}_2]}) + e (\varepsilon_\sigma x)_{\underline{l}_1 \underline{l}_2} g^{\underline{l}_1 \underline{m}_1} g^{\underline{l}_2 \underline{m}_2} \right] g_{\underline{m}_1 [\underline{n}_1} g_{\underline{n}_2] \underline{m}_2} \\ &\quad \times (\varepsilon_\tau y^{\underline{n}_1 \underline{n}_2} + \tilde{C}^{[\underline{n}_1} \varepsilon_\tau x^{\underline{n}_2]} + e g^{\underline{n}_1 \underline{k}_1} g^{\underline{n}_2 \underline{k}_2} (\varepsilon_\sigma x)_{\underline{k}_1 \underline{k}_2}) \\ &\quad - ((\varepsilon_\sigma y)_{\underline{m}} + (\varepsilon_\sigma x)_{\underline{ml}} \tilde{C}^{\underline{l}} + e \varepsilon_\tau x^{\underline{l}} g_{\underline{lm}}) g^{\underline{mn}} ((\varepsilon_\sigma y)_{\underline{n}} - \tilde{C}^{\underline{k}} (\varepsilon_\sigma x)_{\underline{kn}} + e g_{\underline{nk}} \varepsilon_\tau x^{\underline{k}}) \Big], \\ L_{\text{WZ}} &= \varepsilon_\tau x^{\underline{m}_1} \tilde{C}^{\underline{m}_2} (\varepsilon_\sigma x)_{\underline{m}_1 \underline{m}_2}. \end{aligned} \quad (108)$$

Dimensional reduction $L_y \rightarrow 0$ gives the $\mathcal{M}5$ -brane Lagrangian with the Wess-Zumino term. The obtained new $\mathcal{M}5$ -brane Lagrangian in the supergravity background (42) is

$$\begin{aligned} L_{\mathcal{M}5} &= L_0 + L_{\text{WZ}}, \\ L_0 &= \frac{\mathbf{e}}{2} \left[(\dot{x}^{\underline{m}} + \partial^{\underline{m}} Y + s_{\underline{l}} \partial^{[\underline{m}} x^{\underline{l}]}) g_{\underline{mn}} (\dot{x}^{\underline{n}} + \partial^{\underline{n}} Y + s_{\underline{k}} \partial^{[\underline{n}} x^{\underline{k}]}) \right. \\ &\quad \left. - \frac{1}{4} \partial^{[\underline{m}_1} x^{\underline{m}_2]} g_{\underline{m}_1[\underline{n}_1} g_{\underline{n}_2]\underline{m}_2} \partial^{[\underline{n}_1} x^{\underline{n}_2]} \right], \\ L_{\text{WZ}} &= (\dot{x}^{\underline{m}_1} + \partial^{\underline{m}_1} Y) C_{\underline{m}_1 \underline{m}_2 \underline{m}_3} \partial^{\underline{m}_2} x^{\underline{m}_3} + \frac{1}{6} (\partial^{\underline{m}_1} x^{\underline{m}_2}) (\partial^{\underline{m}_3} x^{\underline{m}_4}) s_{[\underline{m}_1} C_{\underline{m}_2 \underline{m}_3 \underline{m}_4]}. \end{aligned} \quad (109)$$

6.2 Non-perturbative M2-brane from $\mathcal{M}5$ -brane

A non-perturbative membrane action in the 11-dimensional supergravity theory is given by [55]

$$\begin{aligned} I &= \int d^3 \sigma L, \quad L = L_0 + L_{\text{WZ}}, \\ \begin{cases} L_0 &= -T \sqrt{-\det \partial_\mu x^m \partial_\nu x^n g_{mn}}, \\ L_{\text{WZ}} &= \frac{T}{3!} \epsilon^{\mu\nu\rho} \partial_\mu x^{m_1} \partial_\nu x^{m_2} \partial_\rho x^{m_3} C_{m_1 m_2 m_3}, \end{cases} \end{aligned} \quad (110)$$

with the spacetime index $m = 0, 1, \dots, 10$ and the world-volume index $\mu = 0, 1, 2$. The canonical coordinates are x^m and p_m , and the spacial world-volume coordinate derivative is ∂_i with $i = 1, 2$. The Hamiltonian is given by [65] where $p_m = \partial L / \partial \dot{x}^m$

$$\begin{aligned} H &= p_m \dot{x}^m - L \\ &= \lambda_0 \mathcal{H}_\tau + \lambda^i \mathcal{H}_i, \\ \begin{cases} \mathcal{H}_\tau &= \frac{1}{2} \triangleright_a \eta^{ab} \triangleright_b + \frac{1}{8} \triangleright^{a_1 a_2} \eta_{a_1 [b_1} \eta_{b_2] a_2} \triangleright^{b_1 b_2}, \\ \mathcal{H}_i &= \partial_i x^m p_m. \end{cases} \end{aligned} \quad (111)$$

Here $\triangleright_A = (\triangleright_a, \triangleright^{ab})$ is related to $\triangleright_M = (\triangleright_m = p_m, \triangleright^{mn} = \epsilon^{ij} \partial_i x^{m_1} \partial_j x^{m_2})$ as $\triangleright_A = E_A^M \triangleright_M$ for the background gauge field E_A^M . E_A^M includes g_{mn} and C_{mnl} . The Virasoro constraint $\mathcal{S}^m = 0$ in (19) is related to the constraint $\mathcal{H}_i = 0$ in (111) which generates σ -diffeomorphism by multiplying the world-volume embedding operator in (114) as $\mathcal{S}^m = \mathcal{H}_i \epsilon^{ij} \partial_j x^m$.

We focus on the 4-dimensional subspace where the supergravity background is a representation of the $\text{SL}(5)$ U-duality symmetry, $E_A^M \in \text{SL}(5)/\text{SO}(5)$. The currents $\triangleright_{\underline{m}}$ and $\triangleright_{\underline{mn}}$ are 4 and 6 components of $\text{SL}(4)$ with $\underline{m} = 1, \dots, 4$, which are unified into a $\text{SL}(5)$ tensor $\triangleright_{\underline{mn}} = (\triangleright_{\underline{m}}, \triangleright_{\underline{mn}})$ with $m = 1, \dots, 5$. The currents for a M2-brane in 4-dimensional space (111) obtained from the membrane Lagrangian (110) are written as

$$\begin{cases} \triangleright_{\underline{m}} &= p_{\underline{m}}, \\ \triangleright_{\underline{m}_1 \underline{m}_2} &= \frac{1}{2} \epsilon_{\underline{m}_1 \dots \underline{m}_4} \epsilon^{ij} \partial_i x^{\underline{m}_3} \partial_j x^{\underline{m}_4}. \end{cases} \quad (112)$$

Commutators of (112) are given as

$$\begin{cases} [\triangleright_{\underline{m}}(\sigma), \triangleright_{\underline{n}}(\sigma')] &= 0, \\ [\triangleright_{\underline{m}_1}(\sigma), \triangleright_{\underline{m}_2 \underline{m}_3}(\sigma')] &= 2i \epsilon_{\underline{m}_1 \dots \underline{m}_4} \epsilon^{ij} \partial_j x^{\underline{m}_4} \partial_i \delta^{(2)}(\sigma - \sigma'), \\ [\triangleright_{\underline{m}_1 \underline{m}_2}(\sigma), \triangleright_{\underline{m}_3 \underline{m}_4}(\sigma')] &= 0. \end{cases} \quad (113)$$

The p -brane current algebras with the non-perturbative winding modes $dx^{m_1} \wedge \dots \wedge dx^{m_p}$ are obtained similarly in [66].

Now let us compare the $SL(5)$ current algebra of the non-perturbative M2-brane (113) with the one of the $\mathcal{M}5$ -brane (18). The perturbative $\mathcal{M}5$ -brane current algebra in (18) reduces into the non-perturbative M2-brane algebra in (113) by reducing the 5-dimensional world-volume of the $\mathcal{M}5$ -brane into the 2-dimensional world-volume of the non-perturbative M2-brane as

$$\partial^m = \epsilon^{ij} \partial_j x^m \partial_i. \quad (114)$$

The operator $\partial_j x^m$ is an embedding of the membrane world-volume to the 5-brane world-volume (where the 5-th brane coordinate is in the internal space). It has the constant form $\partial_j x^m = \delta_j^m$ in the static gauge for the ground state [1].

Now we plug the world-volume projection (114) into the $\mathcal{M}5$ -brane Lagrangian (109). The first term in the $Y = 0$ gauge is given by

$$\begin{aligned} & (\dot{x}^m + s_{\underline{l}} \epsilon^{ij} \partial_j x^m \partial_i x^{\underline{l}}) g_{mn} (\dot{x}^n + s_{\underline{k}} \epsilon^{i'j'} \partial_{j'} x^n \partial_{i'} x^{\underline{k}}) \\ &= (\dot{x}^{\underline{a}})^2 + 2s_{\underline{b}} \dot{x}_{\underline{a}} \epsilon^{ij} \partial_j x^{\underline{a}} \partial_i x^{\underline{b}} + (s_{\underline{b}} \epsilon^{ij} \partial_j x^{\underline{a}} \partial_i x^{\underline{b}})^2 \\ &= h_{00} - 2\lambda^i h_{0i} + \lambda^i \lambda^j h_{ij}, \end{aligned} \quad (115)$$

with

$$h_{ij} = \partial_i x^{\underline{a}} \partial_j x_{\underline{a}} = \partial_i x^m g_{mn} \partial_j x^n, \quad \partial_i x^{\underline{a}} \equiv e_{\underline{m}}^{\underline{a}} \partial_i x^m, \quad (116)$$

and the membrane vielbein λ^i and the 5-brane vielbein $s_{\underline{a}}$

$$\lambda^i = s_{\underline{a}} \epsilon^{ij} \partial_j x^{\underline{a}}, \quad s_{\underline{a}} = e_{\underline{m}}^{\underline{a}} s_{\underline{m}}. \quad (117)$$

The second term is given by

$$\frac{1}{2} \left(\frac{1}{2} \epsilon^{mn} \epsilon^{ij} \partial_j x^{\underline{l}_1} \partial_i x^{\underline{l}_2} \right)^2 = -\frac{1}{2} (\epsilon^{ij} \partial_j x^m \partial_i x^n)^2 = -\det h_{ij}, \quad (118)$$

where the following relation is used in the last equality of (118)

$$\det h_{ij} = \frac{1}{2} \epsilon^{ii'} h_{ij} h_{i'j'} \epsilon^{jj'} = \frac{1}{2} \epsilon^{ii'} \partial_i x^{\underline{a}} \partial_j x_{\underline{a}} \partial_{i'} x^{\underline{b}} \partial_{j'} x_{\underline{b}} \epsilon^{jj'} = \frac{1}{2} (\epsilon^{ij} \partial_j x^{\underline{a}} \partial_i x^{\underline{b}})^2. \quad (119)$$

We choose the following gauge of the membrane world-volume metric

$$\phi = -\frac{\sqrt{-h} h^{00}}{2}, \quad h = \det h_{\mu\nu}, \quad \lambda^i = -\frac{h^{0i}}{h^{00}}, \quad g^2 = \frac{-1}{h(h^{00})^2}. \quad (120)$$

Using with the relation

$$\det h_{ij} = h h^{00}, \quad (121)$$

the kinetic term L_0 in (109) becomes

$$\begin{aligned} L_0 &= \frac{1}{2} \left\{ -\sqrt{-h} - \sqrt{-h} \left(h^{00} h_{00} + 2h^{0i} h_{0i} + \frac{h^{0i} h^{0j}}{(h^{00})^2} h_{ij} \right) \right\} \\ &= -\sqrt{-h}. \end{aligned} \quad (122)$$

This is nothing but the Nambu-Goto Lagrangian for a membrane. The Wess-Zumino term L_{WZ} is obtained by using the world-volume projection (114) into (109) as

$$L_{\text{WZ}} = \dot{x}^{\underline{m}_1} \epsilon^{ij} \partial_j x^{\underline{m}_2} \partial_i x^{\underline{m}_3} C_{\underline{m}_1 \underline{m}_2 \underline{m}_3} = \frac{1}{3!} \epsilon^{\mu\nu\rho} \partial_\mu x^{\underline{m}_1} \partial_\nu x^{\underline{m}_2} \partial_\rho x^{\underline{m}_3} C_{\underline{m}_1 \underline{m}_2 \underline{m}_3}. \quad (123)$$

Together with the Nambu-Goto term (122) the non-perturbative M2-brane Lagrangian is obtained from the perturbative $\mathcal{A}5$ -brane as

$$\begin{aligned} I_{\text{M2}} &= \int d^3\sigma L, \quad L = L_0 + L_{\text{WZ}}, \\ L_0 &= -\sqrt{-\det \partial_\mu x^{\underline{m}} \partial_\nu x^{\underline{n}} g_{\underline{mn}}}, \\ L_{\text{WZ}} &= \frac{1}{3!} \epsilon^{\mu\nu\rho} \partial_\mu x^{\underline{m}_1} \partial_\nu x^{\underline{m}_2} \partial_\rho x^{\underline{m}_3} C_{\underline{m}_1 \underline{m}_2 \underline{m}_3}. \end{aligned} \quad (124)$$

This is the expected M2-brane Lagrangian (110) where we set $T = 1$.

7 Discussion

In this paper we have shown how the conventional strings and membrane are obtained from \mathcal{A} -theory five-brane with the $\text{SL}(5)$ U-duality symmetry.

The following topics are interesting for future problems.

1. From $\mathcal{A}5$ -brane to D-branes: The \mathcal{A} -theory background vielbein field includes the R-R gauge fields which couple to D-branes. The Nambu-Goto Lagrangian will be obtained analogously to the non-perturbative M2-brane Lagrangian as in subsection 6.2 with special care of the B -field. The Wess-Zumino term will be obtained by adding total derivative term with the B -field cloud, in such a way that the gauge transformation rule of the R-R gauge field involves the B -field.
2. From \mathcal{A} -theory branes to the non-perturbative $\mathcal{M}5$ - and NS5-branes: The superstring theories admit the NS5-brane solutions which couple to the B -field magnetically. \mathcal{M} -theory features the $\mathcal{M}5$ -brane whose U-duality symmetry is realized by the current algebra [67], while type IIB superstring theory contains both the NS5-brane and D5-brane related by S-duality. These 5-brane Lagrangians are expected to be derived from $\mathcal{A}5$ -brane and all such 5-branes should be connected via duality transformations. It is interesting to clarify the structure of the 5-brane WEB including $\mathcal{A}5$ -brane for Lagrangians analogous to the one for current algebras [68].
3. From open \mathcal{A} -theory branes to heterotic strings and type I string: The Lagrangians of open \mathcal{A} -theory branes [26], which involve the $\text{SO}(32)$ and $\text{E8} \times \text{E8}$ gauge groups, as well as other half-BPS branes, are of particular interest.
4. Quantization of \mathcal{A} - and \mathcal{M} -branes: The main motivation for constructing the perturbative \mathcal{A} -brane Lagrangian is to facilitate a simpler quantization procedure. Quantum effects in string theory, including winding modes of strings and branes, play a crucial role in understanding Planck-scale physics, such as the resolution of the early-universe singularity. Quantizing \mathcal{A} -theory may provide valuable insights into a unified description of string spectra and S-matrices [69–72].
5. Higher dimensional cases: \mathcal{A} -theories in dimensions $D > 3$ possess U-duality symmetry E_{D+1} [46, 73–75]. In these cases, the spacetime and world-volume dimensions become so large that they necessitate a new interpretation of the unphysical components of spacetime and world-volume. The construction of \mathcal{A} -theory may offer a new perspective on the fundamental description of string theory.

Acknowledgments

We are grateful to Olaf Hohm, Igor Bandos, Martin Roček and Yuqi Li for the fruitful discussions. M.H. would like to thank Yuho Sakatani for useful discussions. We also acknowledge the Simons Center for Geometry and Physics for its hospitality during “The Simons Summer Workshop in Mathematics and Physics 2023 and 2024” where this work has been developed.

Funding information W.S. is supported by NSF award PHY-2210533. M.H. is supported in part by Grant-in-Aid for Scientific Research (C), JSPS KAKENHI Grant Numbers JP22K03603 and JP20K03604.

References

- [1] M. Hatsuda, O. Hulík, W. D. Linch, W. D. Siegel, D. Wang and Y.-P. Wang, *A-theory — A brane world-volume theory with manifest U-duality*, J. High Energy Phys. **10**, 087 (2023), doi:[10.1007/JHEP10\(2023\)087](https://doi.org/10.1007/JHEP10(2023)087) [preprint doi:[10.48550/arXiv.2307.04934](https://doi.org/10.48550/arXiv.2307.04934)].
- [2] E. Witten, *String theory dynamics in various dimensions*, Nucl. Phys. B **443**, 85 (1995), doi:[10.1016/0550-3213\(95\)00158-O](https://doi.org/10.1016/0550-3213(95)00158-O) [preprint doi:[10.48550/arXiv.hep-th/9503124](https://doi.org/10.48550/arXiv.hep-th/9503124)].
- [3] C. Vafa, *Evidence for F-theory*, Nucl. Phys. B **469**, 403 (1996), doi:[10.1016/0550-3213\(96\)00172-1](https://doi.org/10.1016/0550-3213(96)00172-1) [preprint doi:[10.48550/arXiv.hep-th/9602022](https://doi.org/10.48550/arXiv.hep-th/9602022)].
- [4] T. Weigand, *F-theory*, Proc. Sci. **305**, 016 (2018), doi:[10.22323/1.305.0016](https://doi.org/10.22323/1.305.0016).
- [5] W. Siegel, *Superspace duality in low-energy superstrings*, Phys. Rev. D **48**, 2826 (1993), doi:[10.1103/PhysRevD.48.2826](https://doi.org/10.1103/PhysRevD.48.2826) [preprint doi:[10.48550/arXiv.hep-th/9305073](https://doi.org/10.48550/arXiv.hep-th/9305073)].
- [6] W. Siegel, *Two-vierbein formalism for string-inspired axionic gravity*, Phys. Rev. D **47**, 5453 (1993), doi:[10.1103/PhysRevD.47.5453](https://doi.org/10.1103/PhysRevD.47.5453) [preprint doi:[10.48550/arXiv.hep-th/9302036](https://doi.org/10.48550/arXiv.hep-th/9302036)].
- [7] W. Siegel, *Manifest duality in low-energy superstrings*, (arXiv preprint) doi:[10.48550/arXiv.hep-th/9308133](https://doi.org/10.48550/arXiv.hep-th/9308133).
- [8] A. A. Tseytlin, *Duality symmetric closed string theory and interacting chiral scalars*, Nucl. Phys. B **350**, 395 (1991), doi:[10.1016/0550-3213\(91\)90266-Z](https://doi.org/10.1016/0550-3213(91)90266-Z).
- [9] C. Hull and B. Zwiebach, *Double field theory*, J. High Energy Phys. **09**, 099 (2009), doi:[10.1088/1126-6708/2009/09/099](https://doi.org/10.1088/1126-6708/2009/09/099) [preprint doi:[10.48550/arXiv.0904.4664](https://doi.org/10.48550/arXiv.0904.4664)].
- [10] O. Hohm, C. Hull and B. Zwiebach, *Generalized metric formulation of double field theory*, J. High Energy Phys. **08**, 008 (2010), doi:[10.1007/JHEP08\(2010\)008](https://doi.org/10.1007/JHEP08(2010)008) [preprint doi:[10.48550/arXiv.1006.4823](https://doi.org/10.48550/arXiv.1006.4823)].
- [11] G. Aldazabal, D. Marqués and C. Núñez, *Double field theory: A pedagogical review*, Class. Quantum Gravity **30**, 163001 (2013), doi:[10.1088/0264-9381/30/16/163001](https://doi.org/10.1088/0264-9381/30/16/163001) [preprint doi:[10.48550/arXiv.1305.1907](https://doi.org/10.48550/arXiv.1305.1907)].
- [12] O. Hohm, D. Lüst and B. Zwiebach, *The spacetime of double field theory: Review, remarks, and outlook*, Fortschr. Phys. **61**, 926 (2013), doi:[10.1002/prop.201300024](https://doi.org/10.1002/prop.201300024) [preprint doi:[10.48550/arXiv.1309.2977](https://doi.org/10.48550/arXiv.1309.2977)].

- [13] M. Poláček and W. Siegel, *T-duality off shell in 3D type II superspace*, J. High Energy Phys. **06**, 107 (2014), doi:[10.1007/JHEP06\(2014\)107](https://doi.org/10.1007/JHEP06(2014)107) [preprint doi:[10.48550/arXiv.1403.6904](https://doi.org/10.48550/arXiv.1403.6904)].
- [14] M. Poláček and W. Siegel, *Pre-potential in the $AdS_5 \times S^5$ type IIB superspace*, J. High Energy Phys. **01**, 059 (2017), doi:[10.1007/JHEP01\(2017\)059](https://doi.org/10.1007/JHEP01(2017)059) [preprint doi:[10.48550/arXiv.1608.02036](https://doi.org/10.48550/arXiv.1608.02036)].
- [15] M. Hatsuda, K. Kamimura and W. Siegel, *Superspace with manifest T-duality from type II superstring*, J. High Energy Phys. **06**, 039 (2014), doi:[10.1007/JHEP06\(2014\)039](https://doi.org/10.1007/JHEP06(2014)039) [preprint doi:[10.48550/arXiv.1403.3887](https://doi.org/10.48550/arXiv.1403.3887)].
- [16] M. Hatsuda, K. Kamimura and W. Siegel, *Ramond-Ramond gauge fields in superspace with manifest T-duality*, J. High Energy Phys. **02**, 134 (2015), doi:[10.1007/JHEP02\(2015\)134](https://doi.org/10.1007/JHEP02(2015)134) [preprint doi:[10.48550/arXiv.1411.2206](https://doi.org/10.48550/arXiv.1411.2206)].
- [17] M. Hatsuda, K. Kamimura and W. Siegel, *Type II chiral affine Lie algebras and string actions in doubled space*, J. High Energy Phys. **09**, 113 (2015), doi:[10.1007/JHEP09\(2015\)113](https://doi.org/10.1007/JHEP09(2015)113) [preprint doi:[10.48550/arXiv.1507.03061](https://doi.org/10.48550/arXiv.1507.03061)].
- [18] W. D. Linch and W. Siegel, *F-theory from fundamental five-branes*, J. High Energy Phys. **02**, 047 (2021), doi:[10.1007/JHEP02\(2021\)047](https://doi.org/10.1007/JHEP02(2021)047) [preprint doi:[10.48550/arXiv.1502.00510](https://doi.org/10.48550/arXiv.1502.00510)].
- [19] W. D. Linch and W. Siegel, *F-theory with worldvolume sectioning*, J. High Energy Phys. **04**, 022 (2021), doi:[10.1007/JHEP04\(2021\)022](https://doi.org/10.1007/JHEP04(2021)022) [preprint doi:[10.48550/arXiv.1503.00940](https://doi.org/10.48550/arXiv.1503.00940)].
- [20] W. D. Linch III and W. Siegel, *F-brane dynamics*, (arXiv preprint) doi:[10.48550/arXiv.1610.01620](https://doi.org/10.48550/arXiv.1610.01620).
- [21] W. D. Linch III and W. Siegel, *Critical super F-theories*, (arXiv preprint) doi:[10.48550/arXiv.1507.01669](https://doi.org/10.48550/arXiv.1507.01669).
- [22] W. Siegel and D. Wang, *Enlarged exceptional symmetries of first-quantized F-theory*, (arXiv preprint) doi:[10.48550/arXiv.1806.02423](https://doi.org/10.48550/arXiv.1806.02423).
- [23] W. D. Linch and W. Siegel, *F-theory superspace*, J. High Energy Phys. **03**, 059 (2021), doi:[10.1007/JHEP03\(2021\)059](https://doi.org/10.1007/JHEP03(2021)059) [preprint doi:[10.48550/arXiv.1501.02761](https://doi.org/10.48550/arXiv.1501.02761)].
- [24] W. Siegel and D. Wang, *F-theory superspace backgrounds*, (arXiv preprint) doi:[10.48550/arXiv.1910.01710](https://doi.org/10.48550/arXiv.1910.01710).
- [25] M. Hatsuda and W. Siegel, *Perturbative F-theory 10-brane and M-theory 5-brane*, J. High Energy Phys. **11**, 201 (2021), doi:[10.1007/JHEP11\(2021\)201](https://doi.org/10.1007/JHEP11(2021)201) [preprint doi:[10.48550/arXiv.2107.10568](https://doi.org/10.48550/arXiv.2107.10568)].
- [26] M. Hatsuda and W. Siegel, *Open F-branes*, J. High Energy Phys. **04**, 073 (2022), doi:[10.1007/JHEP04\(2022\)073](https://doi.org/10.1007/JHEP04(2022)073) [preprint doi:[10.48550/arXiv.2110.13010](https://doi.org/10.48550/arXiv.2110.13010)].
- [27] W. Siegel, *F-theory with zeroth-quantized ghosts*, (arXiv preprint) doi:[10.48550/arXiv.1601.03953](https://doi.org/10.48550/arXiv.1601.03953).
- [28] W. Siegel and Y.-P. Wang, *F-theory amplitudes*, (arXiv preprint) doi:[10.48550/arXiv.2010.14590](https://doi.org/10.48550/arXiv.2010.14590).

- [29] C.-Y. Ju and W. Siegel, *Gauging unbroken symmetries in F-theory*, Phys. Rev. D **94**, 106004 (2016), doi:[10.1103/PhysRevD.94.106004](https://doi.org/10.1103/PhysRevD.94.106004) [preprint doi:[10.48550/arXiv.1607.03017](https://doi.org/10.48550/arXiv.1607.03017)].
- [30] W. Siegel and D. Wang, *M theory from F theory*, (arXiv preprint) doi:[10.48550/arXiv.2010.09564](https://doi.org/10.48550/arXiv.2010.09564).
- [31] W. Linch and W. Siegel, *F-brane superspace: The new world volume*, (arXiv preprint) doi:[10.48550/arXiv.1709.03536](https://doi.org/10.48550/arXiv.1709.03536).
- [32] D. S. Berman and M. J. Perry, *Generalized geometry and M theory*, J. High Energy Phys. **06**, 074 (2011), doi:[10.1007/JHEP06\(2011\)074](https://doi.org/10.1007/JHEP06(2011)074) [preprint doi:[10.48550/arXiv.1008.1763](https://doi.org/10.48550/arXiv.1008.1763)].
- [33] A. Coimbra, C. Strickland-Constable and D. Waldram, *Supergravity as generalised geometry I: Type II theories*, J. High Energy Phys. **11**, 091 (2011), doi:[10.1007/JHEP11\(2011\)091](https://doi.org/10.1007/JHEP11(2011)091) [preprint doi:[10.48550/arXiv.1107.1733](https://doi.org/10.48550/arXiv.1107.1733)].
- [34] A. Coimbra, C. Strickland-Constable and D. Waldram, *Supergravity as generalised geometry II: $E_{d(d)} \times \mathbb{R}^+$ and M theory*, J. High Energy Phys. **03**, 019 (2014), doi:[10.1007/JHEP03\(2014\)019](https://doi.org/10.1007/JHEP03(2014)019) [preprint doi:[10.48550/arXiv.1212.1586](https://doi.org/10.48550/arXiv.1212.1586)].
- [35] D. S. Berman, M. Cederwall, A. Kleinschmidt and D. C. Thompson, *The gauge structure of generalised diffeomorphisms*, J. High Energy Phys. **01**, 064 (2013), doi:[10.1007/JHEP01\(2013\)064](https://doi.org/10.1007/JHEP01(2013)064) [preprint doi:[10.48550/arXiv.1208.5884](https://doi.org/10.48550/arXiv.1208.5884)].
- [36] O. Hohm and H. Samtleben, *Exceptional form of $D = 11$ supergravity*, Phys. Rev. Lett. **111**, 231601 (2013), doi:[10.1103/PhysRevLett.111.231601](https://doi.org/10.1103/PhysRevLett.111.231601) [preprint doi:[10.48550/arXiv.1308.1673](https://doi.org/10.48550/arXiv.1308.1673)].
- [37] O. Hohm and H. Samtleben, *Exceptional field theory. I. $E_{6(6)}$ -covariant form of M-theory and type IIB*, Phys. Rev. D **89**, 066016 (2014), doi:[10.1103/PhysRevD.89.066016](https://doi.org/10.1103/PhysRevD.89.066016) [preprint doi:[10.48550/arXiv.1312.0614](https://doi.org/10.48550/arXiv.1312.0614)].
- [38] O. Hohm and H. Samtleben, *Exceptional field theory. II. $E_{7(7)}$* , Phys. Rev. D **89**, 066017 (2014), doi:[10.1103/PhysRevD.89.066017](https://doi.org/10.1103/PhysRevD.89.066017) [preprint doi:[10.48550/arXiv.1312.4542](https://doi.org/10.48550/arXiv.1312.4542)].
- [39] O. Hohm and H. Samtleben, *Exceptional field theory. III. $E_{8(8)}$* , Phys. Rev. D **90**, 066002 (2014), doi:[10.1103/PhysRevD.90.066002](https://doi.org/10.1103/PhysRevD.90.066002) [preprint doi:[10.48550/arXiv.1406.3348](https://doi.org/10.48550/arXiv.1406.3348)].
- [40] D. S. Berman, C. D. A. Blair, E. Malek and F. J. Rudolph, *An action for F-theory: $SL(2)\mathbb{R}^+$ exceptional field theory*, Class. Quantum Gravity **33**, 195009 (2016), doi:[10.1088/0264-9381/33/19/195009](https://doi.org/10.1088/0264-9381/33/19/195009) [preprint doi:[10.48550/arXiv.1512.06115](https://doi.org/10.48550/arXiv.1512.06115)].
- [41] M. Cederwall, J. Edlund and A. Karlsson, *Exceptional geometry and tensor fields*, J. High Energy Phys. **07**, 028 (2013), doi:[10.1007/JHEP07\(2013\)028](https://doi.org/10.1007/JHEP07(2013)028) [preprint doi:[10.48550/arXiv.1302.6736](https://doi.org/10.48550/arXiv.1302.6736)].
- [42] M. Cederwall, *Twistors and supertwistors for exceptional field theory*, J. High Energy Phys. **12**, 001 (2015), doi:[10.1007/JHEP12\(2015\)123](https://doi.org/10.1007/JHEP12(2015)123) [preprint doi:[10.48550/arXiv.1510.02298](https://doi.org/10.48550/arXiv.1510.02298)].
- [43] A. S. Arvanitakis and C. D. A. Blair, *The exceptional sigma model*, J. High Energy Phys. **04**, 064 (2018), doi:[10.1007/jhep04\(2018\)064](https://doi.org/10.1007/jhep04(2018)064).
- [44] L. Chabrol, *Geometry of $\mathbb{R}^+ \times E_{3(3)}$ exceptional field theory and F-theory*, J. High Energy Phys. **08**, 073 (2019), doi:[10.1007/JHEP08\(2019\)073](https://doi.org/10.1007/JHEP08(2019)073) [preprint doi:[10.48550/arXiv.1901.08295](https://doi.org/10.48550/arXiv.1901.08295)].

- [45] M. J. Duff, $E_8 \times SO(16)$ symmetry of $d = 11$ supergravity?, in *Nuclear spectroscopy*, Academic Press, New York, USA, ISBN 9781483230641 (1987).
- [46] E. Cremmer and B. Julia, *The $SO(8)$ supergravity*, Nucl. Phys. B **159**, 141 (1979), doi:[10.1016/0550-3213\(79\)90331-6](https://doi.org/10.1016/0550-3213(79)90331-6).
- [47] D. Osten, *Currents, charges and algebras in exceptional generalised geometry*, J. High Energy Phys. **06**, 070 (2021), doi:[10.1007/JHEP06\(2021\)070](https://doi.org/10.1007/JHEP06(2021)070) [preprint doi:[10.48550/arXiv.2103.03267](https://doi.org/10.48550/arXiv.2103.03267)].
- [48] D. Osten, *On exceptional QP-manifolds*, J. High Energy Phys. **01**, 028 (2024), doi:[10.1007/JHEP01\(2024\)028](https://doi.org/10.1007/JHEP01(2024)028) [preprint doi:[10.48550/arXiv.2306.11093](https://doi.org/10.48550/arXiv.2306.11093)].
- [49] D. Osten, *On the universal exceptional structure of world-volume theories in string and M-theory*, Phys. Lett. B **855**, 138814 (2024), doi:[10.1016/j.physletb.2024.138814](https://doi.org/10.1016/j.physletb.2024.138814) [preprint doi:[10.48550/arXiv.2402.10269](https://doi.org/10.48550/arXiv.2402.10269)].
- [50] Y. Sakatani and S. Uehara, *Branes in extended spacetime: Brane worldvolume theory based on duality symmetry*, Phys. Rev. Lett. **117**, 191601 (2016), doi:[10.1103/PhysRevLett.117.191601](https://doi.org/10.1103/PhysRevLett.117.191601) [preprint doi:[10.48550/arXiv.1607.04265](https://doi.org/10.48550/arXiv.1607.04265)].
- [51] C. D. A. Blair and E. T. Musaev, *Five-brane actions in double field theory*, J. High Energy Phys. **03**, 111 (2018), doi:[10.1007/JHEP03\(2018\)111](https://doi.org/10.1007/JHEP03(2018)111) [preprint doi:[10.48550/arXiv.1712.01739](https://doi.org/10.48550/arXiv.1712.01739)].
- [52] Y. Sakatani and S. Uehara, *Exceptional M-brane sigma models and η -symbols*, Prog. Theor. Exp. Phys. 033B05 (2018), doi:[10.1093/ptep/pty021](https://doi.org/10.1093/ptep/pty021) [preprint doi:[10.48550/arXiv.1712.10316](https://doi.org/10.48550/arXiv.1712.10316)].
- [53] C. D. A. Blair, *Open exceptional strings and D-branes*, J. High Energy Phys. **07**, 083 (2019), doi:[10.1007/JHEP07\(2019\)083](https://doi.org/10.1007/JHEP07(2019)083) [preprint doi:[10.48550/arXiv.1904.06714](https://doi.org/10.48550/arXiv.1904.06714)].
- [54] Y. Sakatani and S. Uehara, *Born sigma model for branes in exceptional geometry*, Prog. Theor. Exp. Phys. 073B05 (2020), doi:[10.1093/ptep/ptaa081](https://doi.org/10.1093/ptep/ptaa081) [preprint doi:[10.48550/arXiv.2004.09486](https://doi.org/10.48550/arXiv.2004.09486)].
- [55] E. Bergshoeff, E. Sezgin and P. K. Townsend, *Supermembranes and eleven-dimensional supergravity*, Phys. Lett. B **189**, 75 (1987), doi:[10.1016/0370-2693\(87\)91272-X](https://doi.org/10.1016/0370-2693(87)91272-X).
- [56] M. J. Duff and J. X. Lu, *Duality rotations in membrane theory*, Nucl. Phys. B **347**, 394 (1990), doi:[10.1016/0550-3213\(90\)90565-U](https://doi.org/10.1016/0550-3213(90)90565-U).
- [57] M. J. Duff, J. X. Lu, R. Percacci, C. N. Pope, H. Samtleben and E. Sezgin, *Membrane duality revisited*, Nucl. Phys. B **901**, 1 (2015), doi:[10.1016/j.nuclphysb.2015.10.003](https://doi.org/10.1016/j.nuclphysb.2015.10.003) [preprint doi:[10.48550/arXiv.1509.02915](https://doi.org/10.48550/arXiv.1509.02915)].
- [58] M. Hatsuda and W. Siegel, *$O(D,D)$ gauge fields in the T-dual string Lagrangian*, J. High Energy Phys. **02**, 010 (2019), doi:[10.1007/JHEP02\(2019\)010](https://doi.org/10.1007/JHEP02(2019)010) [preprint doi:[10.48550/arXiv.1810.04761](https://doi.org/10.48550/arXiv.1810.04761)].
- [59] M. Hatsuda and W. Siegel, *T-dual superstring Lagrangian with double zweibeins*, J. High Energy Phys. **03**, 058 (2020), doi:[10.1007/JHEP03\(2020\)058](https://doi.org/10.1007/JHEP03(2020)058) [preprint doi:[10.48550/arXiv.1912.05092](https://doi.org/10.48550/arXiv.1912.05092)].
- [60] M. Hatsuda, K. Kamimura and W. Siegel, *Type II chiral affine Lie algebras and string actions in doubled space*, J. High Energy Phys. **09**, 113 (2015), doi:[10.1007/jhep09\(2015\)113](https://doi.org/10.1007/jhep09(2015)113).

- [61] E. Cremmer, B. Julia, H. Lü and C. N. Pope, *Dualisation of dualities*, Nucl. Phys. B **523**, 73 (1998), doi:[10.1016/s0550-3213\(98\)00136-9](https://doi.org/10.1016/s0550-3213(98)00136-9).
- [62] C. M. Hull, *A geometry for non-geometric string backgrounds*, J. High Energy Phys. **10**, 065 (2005), doi:[10.1088/1126-6708/2005/10/065](https://doi.org/10.1088/1126-6708/2005/10/065).
- [63] C. M. Hull, *Doubled geometry and T-folds*, J. High Energy Phys. **07**, 080 (2007), doi:[10.1088/1126-6708/2007/07/080](https://doi.org/10.1088/1126-6708/2007/07/080).
- [64] K. Lee and J.-H. Park, *Covariant action for a string in doubled yet gauged spacetime*, Nucl. Phys. B **880**, 134 (2014), doi:[10.1016/j.nuclphysb.2014.01.003](https://doi.org/10.1016/j.nuclphysb.2014.01.003).
- [65] M. Hatsuda and K. Kamimura, *SL(5) duality from canonical M2-brane*, J. High Energy Phys. **11**, 001 (2012), doi:[10.1007/JHEP11\(2012\)001](https://doi.org/10.1007/JHEP11(2012)001) [preprint doi:[10.48550/arXiv.1208.1232](https://doi.org/10.48550/arXiv.1208.1232)].
- [66] M. Hatsuda and T. Kimura, *Canonical approach to Courant brackets for D-branes*, J. High Energy Phys. **06**, 034 (2012), doi:[10.1007/JHEP06\(2012\)034](https://doi.org/10.1007/JHEP06(2012)034) [preprint doi:[10.48550/arXiv.1203.5499](https://doi.org/10.48550/arXiv.1203.5499)].
- [67] M. Hatsuda and K. Kamimura, *M5 algebra and SO(5,5) duality*, J. High Energy Phys. **06**, 095 (2013), doi:[10.1007/JHEP06\(2013\)095](https://doi.org/10.1007/JHEP06(2013)095) [preprint doi:[10.48550/arXiv.1305.2258](https://doi.org/10.48550/arXiv.1305.2258)].
- [68] M. Hatsuda, S. Sasaki and M. Yata, *Five-brane current algebras in type II string theories*, J. High Energy Phys. **03**, 298 (2021), doi:[10.1007/JHEP03\(2021\)298](https://doi.org/10.1007/JHEP03(2021)298) [preprint doi:[10.48550/arXiv.2011.13145](https://doi.org/10.48550/arXiv.2011.13145)].
- [69] H. Feng and W. Siegel, *Gauge-covariant S-matrices for field theory and strings*, Phys. Rev. D **71**, 106001 (2005), doi:[10.1103/PhysRevD.71.106001](https://doi.org/10.1103/PhysRevD.71.106001) [preprint doi:[10.48550/arXiv.hep-th/0409187](https://doi.org/10.48550/arXiv.hep-th/0409187)].
- [70] H. Elvang, D. Z. Freedman and M. Kiermaier, *Solution to the Ward identities for superamplitudes*, J. High Energy Phys. **10**, 103 (2010), doi:[10.1007/JHEP10\(2010\)103](https://doi.org/10.1007/JHEP10(2010)103) [preprint doi:[10.48550/arXiv.0911.3169](https://doi.org/10.48550/arXiv.0911.3169)].
- [71] H. Elvang, D. Z. Freedman and M. Kiermaier, *SUSY Ward identities, superamplitudes and counterterms*, J. Phys. A: Math. Theor. **44**, 454009 (2011), doi:[10.1088/1751-8113/44/45/454009](https://doi.org/10.1088/1751-8113/44/45/454009) [preprint doi:[10.48550/arXiv.1012.3401](https://doi.org/10.48550/arXiv.1012.3401)].
- [72] W. Siegel, *S-matrices from 4d worldvolume*, (arXiv preprint) doi:[10.48550/arXiv.2012.12938](https://doi.org/10.48550/arXiv.2012.12938).
- [73] E. Cremmer and B. Julia, *The N = 8 supergravity theory. I. The Lagrangian*, Phys. Lett. B **80**, 48 (1978), doi:[10.1016/0370-2693\(78\)90303-9](https://doi.org/10.1016/0370-2693(78)90303-9).
- [74] G. Bossard, F. Ciceri, G. Inverso, A. Kleinschmidt and H. Samtleben, *E₉ exceptional field theory. Part I. The potential*, J. High Energy Phys. **03**, 089 (2019), doi:[10.1007/JHEP03\(2019\)089](https://doi.org/10.1007/JHEP03(2019)089) [preprint doi:[10.48550/arXiv.1811.04088](https://doi.org/10.48550/arXiv.1811.04088)].
- [75] G. Bossard, F. Ciceri, G. Inverso, A. Kleinschmidt and H. Samtleben, *E₉ exceptional field theory. Part II. The complete dynamics*, J. High Energy Phys. **05**, 107 (2021), doi:[10.1007/JHEP05\(2021\)107](https://doi.org/10.1007/JHEP05(2021)107) [preprint doi:[10.48550/arXiv.2103.12118](https://doi.org/10.48550/arXiv.2103.12118)].