

Hopf exceptional points

 Tsuneya Yoshida^{1,2*},  Emil J. Bergholtz^{3†} and  Tomáš Bzdušek^{4‡}

¹ Department of Physics, Kyoto University, Kyoto 606-8502, Japan 

² Institute for Theoretical Physics, ETH Zurich, 8093 Zurich, Switzerland 

³ Department of Physics, Stockholm University,
AlbaNova University Center, 10691 Stockholm, Sweden 

⁴ Department of Physics, University of Zürich,
Winterthurerstrasse 190, 8057 Zürich, Switzerland 

* yoshida.tsuneya.2z@kyoto-u.ac.jp, † emil.bergholtz@fysik.su.se, ‡ tomas.bzdusek@uzh.ch

Abstract

Exceptional points at which eigenvalues and eigenvectors of non-Hermitian matrices coalesce are ubiquitous in the description of a wide range of platforms from photonic or mechanical metamaterials to open quantum systems. Here, we introduce a class of *Hopf exceptional points* (HEPs) that are protected by the Hopf invariants (including the higher-dimensional generalizations) and which exhibit phenomenology sharply distinct from conventional exceptional points. Saliently, owing to their $\mathbb{Z}_2$ topological invariant related to the Witten anomaly, three-fold HEPs and symmetry-protected five-fold HEPs act as their own “antiparticles”. Furthermore, based on higher homotopy groups of spheres, we predict the existence of multifold HEPs and symmetry-protected HEPs with non-Hermitian topology captured by a range of finite groups (such as $\mathbb{Z}_3$, $\mathbb{Z}_{12}$, or $\mathbb{Z}_{24}$) beyond the periodic table of Bernard-LeClair symmetry classes.



Copyright T. Yoshida *et al.*

This work is licensed under the Creative Commons
Attribution 4.0 International License.

Published by the SciPost Foundation.

Received 2025-08-01

Accepted 2025-12-09

Published 2026-01-07

doi:[10.21468/SciPostPhys.20.1.001](https://doi.org/10.21468/SciPostPhys.20.1.001)



Check for
updates

Contents

1	Introduction	2
2	HEP3s with $\mathbb{Z}_2$ topology	3
3	Symmetry-protected HEP5s with $\mathbb{Z}_2$ topology	4
4	Symmetry-protected HEP4s with $\mathbb{Z}$ topology	7
5	General characteristics of HEPs	8
5.1	Multiply-charged aspect of HEPs	8
5.2	Additional bands	8
5.3	Higher dimensions	9
6	Conclusion and outlook	9

A	Defining topology from resultants	10
A.1	Resultant of a pair of polynomials	10
A.2	Resultant winding number	11
A.3	Berry connection and curvature of resultant Hamiltonians for Hopf topology	12
B	Details of the presented Hamiltonians	12
B.1	Model in Eq. (4) and computation of the $\mathbb{Z}_2$ invariant	12
B.2	Model in Eq. (8)	13
B.3	Model in Eq. (13)	13
C	Fake EP3s	14
	References	14

1 Introduction

The notion of topology plays a pivotal role in modern condensed matter physics of both quantum [1–5] and classical systems [6–10]. Saliiently, topological physics implies the possibility of various exotic quasiparticles.¹ One of the prime examples is a Majorana zero mode [12–15] whose antiparticle is itself. In addition, a Weyl fermion [16–22], protected by Chern number in topological semimetals, is a source of negative magnetoresistance [23] which is a signal of chiral anomaly. The periodic table for the ten Altland-Zirnbauer symmetry classes provides a systematic understanding of the topological obstructions inducing these exotic excitations in Hermitian systems [24–30].

Notably, open systems coupled to environments host topological excitations for which non-Hermiticity is essential [31, 32], such as exceptional points [33–37]. At exceptional points, two energy bands touch in both the real and the imaginary parts. Such band touchings are protected by the winding topology of energy eigenvalues [38]. In sharp contrast to Hermitian topological excitations, exceptional points exhibit a dispersion with a fractional exponent. Exceptional points and their variants [38–52] are reported for a wide range of platforms from quantum systems [53–62] to metamaterials [63–76], indicating the ubiquity of these non-Hermitian excitations. In particular, the high controllability of synthetic systems allows for the realization of exceptional points in dimensions larger than three [63, 64, 77, 78].

Among the various exceptional points, multifold exceptional points exhibit an n -fold band touching [37, 79–83] whose stability due to topology and symmetry has recently been revealed in Refs. [84–89]. These unique excitations are beyond the existing classification table for Bernard-LeClair symmetry classes [90–94] and discussed in interdisciplinary fields [68, 77, 95–110]. However, topology of the formerly reported n -fold exceptional points (EPns) is generally characterized by $\mathbb{Z}$ invariants (i.e., winding topology) for $n \geq 3$ [84, 88]. This, in particular, implies that an exceptional point cannot be its own “antiparticle”. If such n -fold band touching exists, it should be distinguished from the formerly reported EPn.

In this work, we report novel non-Hermitian topological excitations, dubbed n -fold Hopf exceptional points (HEPns, $n = 3, \dots, 7$),² which are topologically stabilized by higher homo-

¹Here, the term “quasiparticle” denotes a topologically protected band singularity, as is widely used in the literature [1, 2, 11].

²To achieve concise terminology, we refer to *all* such EPs protected by topologically nontrivial maps between higher-dimensional spheres with $d > m \geq 2$ collectively as *Hopf exceptional points*, although not all of these maps are termed as Hopf maps in the mathematical literature.




Figure 1: Illustration of $\mathbb{Z}_2$ topology protecting a three-fold Hopf exceptional point (HEP3). The resultant vector $\mathbf{R}(\mathbf{k})$ defines a map from a 4-sphere in the momentum (or parameter) space to a 3-sphere in the space of the resultant vector [see Eqs. (1) and (2)]. If the map is topologically nontrivial, an HEP3 with $\mathbb{Z}_2$ topology exists inside the 4-sphere.

topology groups of spheres, labeled $\pi_d(S^m)$, with integers $d > m \geq 2$. Saliiently, an HEP3 and a symmetry-protected HEP5 exhibit unusual $\mathbb{Z}_2$ topology, meaning that they act as their own antiparticle. We trace this striking feature to the homotopy group $\pi_4(S^3) = \mathbb{Z}_2$ (see Fig. 1) classifying the map of resultants. We further discover symmetry-protected HEP4 whose topology is classified by $\pi_3(S^2) = \mathbb{Z}$. By systematically leveraging higher homotopy groups of spheres, we elucidate the potential presence of HEPns characterized by abundant finite groups (e.g., $\mathbb{Z}_3$, $\mathbb{Z}_{12}$, and $\mathbb{Z}_{24}$) beyond the existing classification table.

2 HEP3s with $\mathbb{Z}_2$ topology

As a first example, we consider a generic three-band non-Hermitian Hamiltonian $H(\mathbf{k})$ in a five-dimensional momentum (or parameter) space denoted by $\mathbf{k} = (k_1, \dots, k_5)$. Such a momentum space whose dimensionality is higher than three can be accessed in metamaterials [63, 64, 77, 78, 88]. The formation of an EP3 is captured by vanishing resultants $r_j \in \mathbb{C}$

$$r_j(\mathbf{k}) = \text{Res}[\partial_E^{n-1-j} P(E, \mathbf{k}), \partial_E^{n-1} P(E, \mathbf{k})], \quad (1)$$

with $j = 1, 2, \dots, n-1$ and $n = 3$ (see Appendix A.1).^{3,4} Here, $P(E, \mathbf{k}) = \det[H(\mathbf{k}) - E\mathbb{1}] \in \mathbb{C}$ is the characteristic polynomial, and ∂_E denotes derivative with respect to the polynomial variable $E \in \mathbb{C}$.

To expose whether the EP3 is in fact an HEP3, we consider a 4-sphere $S^4 \subset \mathbb{R}^5$ enclosing the band touching. Assuming $(r_1, r_2) \neq (0, 0)$ on the 4-sphere, we introduce a normalized vector $\mathbf{n} = \mathbf{R}/\|\mathbf{R}\|$ ($\|\mathbf{R}\| = \sqrt{\mathbf{R} \cdot \mathbf{R}}$) with the resultant vector

$$\mathbf{R}(\mathbf{k}) = (\text{Re}[r_1], \text{Im}[r_1], \text{Re}[r_2], \text{Im}[r_2]). \quad (2)$$

Vector $\mathbf{n}(\mathbf{k})$ defines a map from S^4 to S^3 whose topology is classified by an element of the homotopy group $\pi_4(S^3) = \mathbb{Z}_2$. When the map of $\mathbf{n}(\mathbf{k})$ possesses nontrivial $\mathbb{Z}_2$ topology, the enclosed band touching is an HEP3. In passing, we note that the resultant topology is neither point-gap nor line-gap topology [84, 88] (see also Appendix A.2 for a brief overview of the resultant winding topology).

³The vanishing resultants $r_j = 0$ ($j = 1, 2, \dots, n-1$) indicate the n -tuple root; it captures the algebraic multiplicity rather than geometric multiplicity. We note, however, that these two coincide with each other because breaking the coincidence requires fine-tuning.

⁴For $n = 2$, Eq. (1) reduces to the discriminant whose winding topology protects EP2 for systems without symmetry [38, 46].

The $\mathbb{Z}_2$ invariant ν_F of maps from S^4 to S^3 , originally discovered by Freudenthal [111], was considered in physics in the context of the Witten anomaly in $SU(2)$ gauge theory [112] and to describe topological defects in superfluid ^3He [113]. Numerical computation of this $\mathbb{Z}_2$ invariant is carried out by the following representation [114]

$$\nu_F = \frac{1}{4\pi} \oint d^4p \epsilon^{\mu\nu\rho\lambda} [\partial_\mu \Delta\varphi(\mathbf{p})] A_\nu F_{\rho\lambda}, \quad (3)$$

with $\mu, \nu, \rho, \lambda = 1, \dots, 4$ and anti-symmetric tensor $\epsilon^{\mu\nu\rho\lambda}$ taking $\epsilon^{1234} = 1$. The Berry connection A_μ and the Berry curvature $F_{\mu\nu}$ are obtained from the *resultant Hamiltonian*, and the phase $\Delta\varphi(\mathbf{p})$ ($0 \leq \Delta\varphi \leq 2\pi$) is obtained from the resultant vector. For the precise definitions of A_μ , $F_{\mu\nu}$ and $\Delta\varphi$, see Appendix A.3. Vector $\mathbf{p}$ parametrizes the 4-sphere in the momentum space. While the integral in Eq. (3) can take an arbitrary integer value, gauge transformations can change ν_F by multiples of two [112].

We demonstrate the emergence of an HEP3 in five dimensions by analyzing a toy model. The Hamiltonian reads

$$H = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ \frac{\zeta_2}{6} & \zeta_1 & 0 \end{pmatrix}, \quad (4)$$

where the functions $\zeta_{1,2}(\mathbf{k})$ are parameterized by δ , m_0 , and $f(k_5)$ (for the explicit form, see Appendix B.1). Here the function $f(k_5)$ is either even [$f(k_5) = 1$] or odd [$f(k_5) = 2\sin(k_5/2)$].

To motivate the Hamiltonian form in Eq. (4), recall that a three-fold exceptional point is generally captured by a 3×3 Jordan block form [86]. The Hamiltonian including the perturbations ζ 's in the bottom row ensures that the resultants [recall Eq. (1)] are expressed as $r_j \propto \zeta_j$ for $j \in \{1, 2\}$ [88], which provides clear correspondence between the perturbations ζ 's and the Hopf topology of $\mathbf{R}$. The sought higher Hopf topology is then ensured by imposing the appropriate dependence [114] of the functions $\zeta_{1,2}(\mathbf{k})$ on momenta (parameters) $\mathbf{k}$.⁵ The vanishing resultant vector determines where the HEP3s emerge in the momentum space. The explicit form of the resultant vector [see Appendix B.1] indicates the emergence of HEP3s at $\mathbf{k} = (0, 0, 0, \pi/2, \pm\pi/3)$ for $\delta = 0.5$ and $m_0 = 1.5$. Figure 2(a,b) displays the emergence of the HEP3 at $\mathbf{k} = (0, 0, 0, \pi/2, \pi/3)$.

Notably, the HEP acts as its own antiparticle as a direct consequence of its $\mathbb{Z}_2$ topology.⁶ This fact is elucidated by examining pair annihilation of HEP3s in two cases: $f(k_5) = 1$ and $f(k_5) = 2\sin(k_5/2)$ [see Fig. 2(c,d)]. For $f(k_5) = 1$ and $\delta = 0.5$, the system hosts two HEP3s demarcated by the planes at $k_5 = \pi/2, 0$ and $-\pi/2$ where numerically computed ν_F is equal to 1, 0, and 1, respectively [for computation of ν_F , see Appendix B.1]. As δ decreases, the two HEP3s approach and annihilate each other [see Fig. 2(c)]. Changing the parity of $f(k_5)$ flips the sign of the numerically computed ν_F at $k_5 = -\pi/2$. Even in this case, pair annihilation occurs [see Fig. 2(d)]. The occurrence of pair annihilation in both arrangements manifests that HEP3s are indeed protected by $\mathbb{Z}_2$ topology, implying that an HEP3 is its own antiparticle.

3 Symmetry-protected HEP5s with $\mathbb{Z}_2$ topology

Symmetry further enriches Hopf exceptional points, as exemplified by the emergence of symmetry-protected HEP5 in five dimensions. We consider a five-band non-Hermitian Hamiltonian which preserves parity-time (PT -) symmetry

$$U_{PT} H^*(\mathbf{k}) U_{PT}^{-1} = H(\mathbf{k}), \quad (5)$$

⁵Another choice of ζ 's yields an EP3 with winding topology [88] (see Appendix A.2).

⁶The pair annihilation of topological nodes with the same charge is commonly observed for $\mathbb{Z}_2$ topology; for instance, Ref. [48] demonstrates pair annihilation of EP2s with $\mathbb{Z}_2$ topology.




Figure 2: Energy bands of Hamiltonian (4) for $m_0 = 1.5$. The real (imaginary) part is represented as height (color). In panel (a), the complex conjugate of the upper band is omitted. Panels (a) and (b) are obtained for $k_5 = \pi/3$, $\delta = 0.5$, and $f(k_5) = 1$. Here, Δk_4 is defined as $\Delta k_4 = k_4 - \pi/2$. Panel (c) [(d)] displays pair annihilation of HEP3s for $(k_2, k_3, k_4) = (0, 0, \pi/2)$, and $f(k_5) = 1$ [$f(k_5) = 2 \sin(k_5/2)$]. In these panels, numerically computed ν_F for $k_5 = -\pi/2, 0$, and $\pi/2$ at $\delta = 0.5$ is represented by numbers highlighted in yellow. We used a mesh of 40^4 points to evaluate the integrals with momenta $k_{1,2,3} \in [-\pi, \pi]$ and $k_4 \in [0, 2\pi]$.

with a unitary matrix satisfying $U_{PT} U_{PT}^* = \mathbb{1}$. Here, $\mathbb{1}$ is the identity matrix, and asterisks denote complex conjugation. The PT -symmetry imposes the constraint

$$P(E) = P^*(E^*), \quad (6)$$

indicating that all coefficients of the characteristic polynomial $P(E)$ are real at each $\mathbf{k}$. For $n = 5$, the formation of an EP5 is captured by vanishing resultants $r_{1,\dots,4} = 0$ [see Eq. (1)]. Due to PT -symmetry [see Eqs. (5) and (6)] these resultants are real: $r_{1,\dots,4} \in \mathbb{R}$.

To expose whether the EP5 is in fact an HEP5, we consider a 4-sphere $S^4 \subset \mathbb{R}$ enclosing the band touching. Assuming $r_{1,\dots,4} \neq 0$ on the 4-sphere, we introduce a normalized vector $\mathbf{n} = \mathbf{R}/\|\mathbf{R}\|$ with

$$\mathbf{R} = (r_1, r_2, r_3, r_4)^T. \quad (7)$$

The normalized vector $\mathbf{n}$ defines a map from S^4 to S^3 whose topology is classified by $\pi_4(S^3) = \mathbb{Z}_2$. When the map of $\mathbf{n}$ possesses nontrivial $\mathbb{Z}_2$ topology, the enclosed band touching is a symmetry-protected HEP5. The topological invariant is introduced in a similar way as




Figure 3: Energy bands of Hamiltonian (8) for $f(k_5)=1$, $k_5=\pi/3$, $m_0=1.5$ and $\delta=0.5$. The real (imaginary) part is represented as height (color). Panel (a) [(b)] displays the data for $(k_3, k_4)=(0, \pi/2)$ [$(k_1, k_2)=(0, 0)$]. In panel (a), the complex conjugates of the upper and the lower bands are omitted. In panel (b), Δk_4 is defined as $\Delta k_4=k_4-\pi/2$.

Eq. (3) with the only difference that we compute the $\mathbb{Z}_2$ invariant from the resultant vector in Eq. (7) instead of the one in Eq. (2).

We demonstrate the emergence of a symmetry-protected HEP5 by analyzing a toy model whose Hamiltonian reads

$$H = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ \frac{\zeta_4}{(5!)^3} & \frac{\zeta_3}{(5!)^2} & \frac{\zeta_2}{5!2!} & \frac{\zeta_1}{3!} & 0 \end{pmatrix}, \quad (8)$$

with real functions $\zeta_{1,\dots,4}(\mathbf{k})$ parameterized by δ , m_0 , and $f(k_5)$ [for the explicit form, see Appendix B.2]. Here, $f(k_5)$ is either $f(k_5)=1$ or $f(k_5)=2\sin(k_5/2)$. This Hamiltonian preserves PT -symmetry [Eq. (5)] with $U_{PT}=\mathbb{1}$. In analogy with the discussion of the Hamiltonian in Eq. (4), the perturbation of the 5×5 Jordan form in Eq. (8) is chosen so that the resultants obey $r_j \propto \zeta_j$ for each $j \in \{1, \dots, 4\}$. The explicit form of $\zeta_j(\mathbf{k})$ follows from a known representative of the nontrivial class in $\pi_4(S^3)$ [114].

The vanishing resultant vector $\mathbf{R} \propto (\zeta_1, \zeta_2, \zeta_3, \zeta_4)^T = \mathbf{0}$ specifies where the symmetry-protected HEP5s emerge. The explicit form of the resultant vector [see Appendix B.2] indicates that the model in Eq. (8) hosts symmetry-protected $\mathbb{Z}_2$ HEP5s at $\mathbf{k}=(0, 0, 0, \pi/2, \pm\pi/3)$ which possesses $\mathbb{Z}_2$ topological charge [see Fig. 3].

The argument of PT -symmetry protected HEPs can be applied to other cases of symmetry: pseudo-Hermiticity, CP -, and chiral symmetry. We consider a Hamiltonian preserving pseudo-Hermiticity

$$U_{\text{pH}} H^\dagger(\mathbf{k}) U_{\text{pH}}^\dagger = H(\mathbf{k}), \quad (9)$$

with $U_{\text{pH}}^\dagger$ being a unitary matrix and dagger denotes Hermitian conjugation. Because the transposition does not affect the determinant, Eq. (9) leads to Eq. (6), implying that the resultants [Eq. (1)] are real.

For CP - and chiral symmetry, the Hamiltonian obeys

$$U_{\text{CP}} H^*(\mathbf{k}) U_{\text{CP}}^\dagger = -H(\mathbf{k}), \quad (10)$$

$$U_\Gamma H^\dagger(\mathbf{k}) U_\Gamma^\dagger = -H(\mathbf{k}), \quad (11)$$

with U_{CP} and U_Γ being unitary matrices. In these cases, replacing H to $H' = iH$ reduces to the case of PT -symmetry or pseudo-Hermiticity. Therefore, symmetry-protected HEPs may emerge when systems preserve pseudo-Hermiticity, CP -, or chiral symmetry.




Figure 4: (a) and (b): Energy eigenvalues of Hamiltonian in Eq. (13) for $m_0 = 3$ and $\delta = 0$. The real (imaginary) part is represented as height (color). In panel (b), complex conjugates of the upper and lower bands are omitted. (c) [(d)]: Lines in the momentum space (k_1, k_2, k_3) for $m_0 = 3$, $\delta = 0$ and $k_4 = -\pi/2$ [$k_4 = \pi/2$], where red and blue lines denote the momenta satisfying $\mathbf{R} \propto (0, 0, 1)^T$ and $\mathbf{R} \propto (0, 0, -1)^T$, respectively. The linking of these lines determines the value of the Hopf invariant ν_H . (e): Momenta satisfying $\mathbf{R}(\mathbf{k}) = \mathbf{0}$ for $k_1 = 0$ and $\delta = 0$ resp. $\delta = 0.2$ (orange manifolds). As δ is introduced, the symmetry-protected HEP4 inflates into a loop. The gray oval and the blue loop illustrate the S^3 resp. S^2 , both extending in the fourth dimension k_1 (not shown), on which one computes the Hopf invariant ν_H resp. the resultant winding number W_2 (see Appendix A.2) [88].

4 Symmetry-protected HEP4s with $\mathbb{Z}$ topology

Symmetry protection also enables HEP4s with $\mathbb{Z}$ topology. To illustrate such a possibility, we consider a four-band non-Hermitian Hamiltonian with PT -symmetry [see Eq. (5)] in a four-dimensional momentum space described by $\mathbf{k} = (k_1, k_2, k_3, k_4)$. For $n = 4$, the formation of an EP4 is captured by vanishing resultants $r_{1,\dots,3} = 0$ [see Eq. (1)].

To expose whether the EP4 is in fact an HEP4, we consider a 3-sphere $S^3 \subset \mathbb{R}^4$ enclosing the band touching. The homotopy group $\pi_3(S^2) = \mathbb{Z}$ implies the existence of nontrivial maps $\mathbf{n}(\mathbf{k}) = \mathbf{R}/\|\mathbf{R}\|$. When the map of $\mathbf{n}$ possesses nontrivial $\mathbb{Z}$ topology, the enclosed band touching is a symmetry-protected HEP4.

The topology of such HEP4s is characterized by the Hopf invariant, expressed as [115–117]

$$\nu_H = \oint \frac{d^3 \mathbf{p}}{2} \epsilon^{\mu\nu\rho} A_\mu F_{\nu\rho}, \quad (12)$$

with $\mu, \nu, \rho = 1, 2, 3$. Here, A_μ and $F_{\mu\nu}$ are obtained from the resultant Hamiltonian (for the definitions, see Appendix A.3). Vector $\mathbf{p}$ parametrizes the 3-sphere in the momentum space.

We demonstrate the emergence of a symmetry-protected HEP4 by analyzing a toy model whose Hamiltonian reads

$$H = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ \frac{\zeta_3}{24^2} & -\frac{\zeta_2}{24} & \frac{\zeta_1}{2} & 0 \end{pmatrix}, \quad (13)$$

with real functions $\zeta_{1,\dots,3}(\mathbf{k})$ parametrized by δ and m_0 [for the explicit form, see Appendix B.3]. As is the case with Hamiltonians in Eqs. (4) and (8), the Hopf topology [115] of ζ 's in the perturbed model [Eq. (13)] is equivalent to that of the resultant vector.

The vanishing resultant vector $\mathbf{R} \propto (\zeta_1, \zeta_2, \zeta_3)^T = \mathbf{0}$ specifies where the symmetry-protected HEP4 emerges in the momentum space. The explicit form of the resultant [see Appendix B.3] indicates the emergence of a symmetry-protected HEP4 at $\mathbf{k} = \mathbf{0}$ for $m_0 = 3$ [see Fig. 4(a,b)]. Here, we characterize the symmetry-protected HEP4 enclosed by two planes at $k_4 = -\pi/2$ and $k_4 = \pi/2$. Numerically evaluating the Hopf invariant [Eq. (12)], we obtain $\nu_H = 0$ [$\nu_H = 1$] for $k_4 = -\pi/2$ [$k_4 = \pi/2$] which is consistent with linking of the inverse maps in the momentum space [see Fig. 4(c,d)] [118, 119]. These results indicate that the symmetry-protected HEP4 is characterized by $\nu_H = 1$.

5 General characteristics of HEPs

5.1 Multiply-charged aspect of HEPs

Reference 84 has pointed out that the codimension of symmetry-protected EP4s is three, whereas the codimension of the HEP4 in our model is four [see Fig. 4(a,b)]. This mismatch in codimension implies that the HEP4 inflates into a loop of EP4s under a generic perturbation of the Hamiltonian matrix [see Fig. 4(e)]. Such a perturbation does not trivialize the Hopf topology but enriches it [30]: the loop of EP4s carries both the Hopf invariant on S^3 and the resultant winding number on S^2 .⁷ In the same spirit, a similar assignment of multiple topological invariants applies to all HEPs introduced in this work.

5.2 Additional bands

So far, we have discussed HEPns in n -band models. For systems with more than n bands, EPns are still associated with winding of the resultant vector; however, the converse does not hold. Specifically, as discussed in Appendix C, it is possible to find situations where a finite value of the winding number leads to the vanishing resultant vector $\mathbf{R} = \mathbf{0}$ without a symmetry-protected EP3.

Nevertheless, topological invariants of resultants are applicable when we focus around an HEPn (or an EPn). Specifically, if we know that an HEPn emerges at momenta $\mathbf{k} = \mathbf{k}_0$ and energy $E = E_0$, we may apply the Taylor expansion to the characteristic polynomial $P(E) = \det[H(\mathbf{k}) - E\mathbb{I}]$, which leads to a polynomial $\tilde{P}(\tilde{E})$ with degree n

$$\tilde{P}(\tilde{E}) = \sum_{j=0}^n a_j(\mathbf{k}) \tilde{E}^j, \quad (14)$$

with $a_j(\mathbf{k})$ ($j = 0, 1, \dots, n$) being complex functions and $\tilde{E} = E - E_0$. Computing the topological invariant [e.g., Eq. (3)] of $\tilde{P}(\tilde{E})$ around $\mathbf{k} = \mathbf{k}_0$, we can characterize the HEPn for systems with more than n bands.

⁷In contrast to the EP4s with the Hopf topology, a loop of formerly reported EP4s can be contracted and annihilated.

Table 1: Hopf exceptional points (HEPs) are classified by higher homotopy groups of spheres. The topology explicitly analyzed in this manuscript corresponds to the blue entries. HEPns [symmetry-protected HEPns] of codimension $c = 4, 5, \dots$ are classified by $\pi_{c-1}(S^{2n-3})$ [$\pi_{c-1}(S^{n-2})$].

c	HEP3	SP-HEP6	HEP4
	SP-HEP4, SP-HEP5		SP-HEP7
4	$\mathbb{Z}$	0	0
5	$\mathbb{Z}_2$	$\mathbb{Z}$	0
6	$\mathbb{Z}_2$	$\mathbb{Z}_2$	$\mathbb{Z}$
7	$\mathbb{Z}_{12}$	$\mathbb{Z}_2$	$\mathbb{Z}_2$
8	$\mathbb{Z}_2$	$\mathbb{Z} \times \mathbb{Z}_{12}$	$\mathbb{Z}_2$
9	$\mathbb{Z}_2$	$\mathbb{Z}_2^2$	$\mathbb{Z}_{24}$
10	$\mathbb{Z}_3$	$\mathbb{Z}_2^2$	$\mathbb{Z}_2$
11	$\mathbb{Z}_{15}$	$\mathbb{Z}_{24} \times \mathbb{Z}_3$	$\mathbb{Z}_2$
12	$\mathbb{Z}_2$	$\mathbb{Z}_{15}$	$\mathbb{Z}_2$
13	$\mathbb{Z}_2^2$	$\mathbb{Z}_2$	$\mathbb{Z}_{30}$
14	$\mathbb{Z}_{12} \times \mathbb{Z}_2$	$\mathbb{Z}_2^3$	$\mathbb{Z}_2$
15	$\mathbb{Z}_{84} \times \mathbb{Z}_2^2$	$\mathbb{Z}_{120} \times \mathbb{Z}_{12} \times \mathbb{Z}_2$	$\mathbb{Z}_2^3$
16	$\mathbb{Z}_2^2$	$\mathbb{Z}_{84} \times \mathbb{Z}_2^5$	$\mathbb{Z}_{72} \times \mathbb{Z}_2$

5.3 Higher dimensions

Higher homotopy groups of spheres indicate abundant topology of HEPns. HEPns are captured by the $(2n-2)$ -component resultant vector, and thus are classified by $\pi_{c-1}(S^{2n-3})$ with codimension $c \geq 4$. Symmetry-protected HEPns are captured by the $(n-1)$ -component resultant vector, and thus are classified by $\pi_{c-1}(S^{n-2})$. The classification results for HEPns ($n=3, 4$) and symmetry-protected HEPns ($n=4, 5, 6, 7$) are summarized in Table 1 (see also Sec. 4.1 of Ref. [120] and Chapter XIV of Ref. [121]). This table predicts the presence of various HEPns following exotic fusion rules. For instance, $\mathbb{Z}_3$ topology of codimension $c=10$ implies an HEP3 annihilate with two copies of itself, which is reminiscent of parafermions [122–125]. In addition, there exist HEP3s with $\mathbb{Z}_{12}$ topology of codimension $c=7$ and HEP4s with $\mathbb{Z}_{24}$ topology of codimension $c=9$. Topological invariants for these HEPns remain to be formulated.

6 Conclusion and outlook

We have discovered novel non-Hermitian topological excitations, dubbed Hopf exceptional points. Saliiently, HEP3s and symmetry-protected HEP5s exhibit an unusual $\mathbb{Z}_2$ topology, meaning that they act as their own antiparticle. This striking feature arises from the homotopy group $\pi_4(S^3) = \mathbb{Z}_2$. We have further discovered symmetry-protected HEP4s whose topology is classified by $\pi_3(S^2) = \mathbb{Z}$. Leveraging higher homotopy groups of spheres, we elucidate the potential presence of HEPns characterized by abundant finite groups (e.g., $\mathbb{Z}_3$, $\mathbb{Z}_{12}$, and $\mathbb{Z}_{24}$) beyond the classification table of Bernard-LeClair symmetry classes.

Our work on non-Hermitian multiband systems opens up a new direction of topological physics. Here, we outline several concrete open questions motivated by our findings. On one hand, from the implementation perspective, we anticipate that HEPns can be realized in a variety of experimental settings. Notably they may be realized in terms of non-unitary photon dynamics where both the eigenvalues and eigenstates have been measured accurately at and around multifold EPs [99], hence suggesting a path to a particularly comprehensive simula-

tion of HEPns. Further promising platforms include nitrogen-vacancy spin systems [109] and coupled micro-resonators [100] in which multifold EPs have also been realized. Quite generally, the high controllability of metamaterials allows access to momentum (or parameter) spaces with dimensions larger than three [63, 64, 77, 78, 88], thus inviting various realistic experimental verifications of the unusual topology of HEPns.

On the other hand, our work also indicates several concrete theoretical aspects that deserve a deeper mathematical analysis. First, Table 1 implies the presence of novel HEPs following a unique fusion rule due to their topology. Explicit analysis based on topological invariants and concrete models is an interesting issue. Next, unless an explicit band structure is provided, one-to-one correspondence between EPns and the resultant winding numbers is lost for many-band systems. This observation calls for a more general topological characterization of Hopf exceptional points. Finally, it is interesting to consider non-Hermitian topological bands, captured by suitably adapted Hopf invariants [119], that arise in models with a hopfion texture [126] of the resultant vector over the Brillouin zone torus. We postpone the analysis of these questions to future studies.

Acknowledgments

T. Y. is grateful for the support from the ETH Pauli Center for Theoretical Studies.

Funding information T. Y. is supported by JSPS KAKENHI Grant Nos. JP21K13850, JP23KK0247, JP25K07152, and JP25H02136 as well as JSPS Bilateral Program No. JPJSBP120249925 and Yamada Science Foundation. E. J. B. is supported by the Wallenberg Scholars program (2023.0256) and the Göran Gustafsson Foundation for Research in Natural Sciences and Medicine. T. B. was supported by the Starting Grant No. 211310 by the Swiss National Science Foundation (SNSF).

A Defining topology from resultants

A.1 Resultant of a pair of polynomials

For given two polynomials

$$f(x) = a_n x^n + \dots + a_1 x + a_0, \quad (\text{A.1})$$

$$g(x) = b_m x^m + \dots + b_1 x + b_0, \quad (\text{A.2})$$

with complex coefficients a 's and b 's, the resultant is defined as

$$\text{Res}[f(x), g(x)] = a_n^m b_m^n \prod_{i,j} (\alpha_i - \beta_j). \quad (\text{A.3})$$

Here, α 's and β 's are roots of polynomials $f(x)$ and $g(x)$, respectively. Symbol $\prod_{i,j}$ denotes the product over all pairs of the roots. The resultant vanishes when the two polynomials have a common root.

The resultant can be computed from the Sylvester matrix

$$\text{Res}[f(x), g(x)] = \det \begin{bmatrix} a_n & \cdots & a_0 & & & \\ & a_n & \cdots & a_0 & & \\ & & \ddots & & \ddots & \\ & & & a_n & \cdots & a_0 \\ b_m & \cdots & b_0 & & & \\ & b_m & \cdots & b_0 & & \\ & & \ddots & & \ddots & \\ & & & b_m & \cdots & b_0 \end{bmatrix}. \quad (\text{A.4})$$

Here, the empty elements are zero. The size of the matrix is $n + m$ and the first n rows are composed of the coefficients a 's and the remaining m rows are composed of the coefficients b 's.

A.2 Resultant winding number

We next consider an N -component resultant vector $\mathbf{R}(\mathbf{k})$, such as the one in Eq. (2), defined on an N -dimensional momentum space $\mathbb{R}^N$. When the norm is finite $\|\mathbf{R}\| \neq 0$ on an $(N-1)$ -sphere $S^{N-1} \subset \mathbb{R}^N$, one can consider the normalized vector $\mathbf{n}(\mathbf{k}) = \mathbf{R}(\mathbf{k})/\|\mathbf{R}(\mathbf{k})\|$ whose topology is classified by $\pi_{N-1}(S^{N-1}) = \mathbb{Z}$.

The topology of such a map is characterized by the resultant winding number W_{N-1} [84,88]

$$W_{N-1} = \frac{\epsilon^{i_1 \cdots i_N}}{A_{N-1}} \int d^{N-1} \mathbf{p} f_{i_1 \cdots i_N}(\mathbf{p}), \quad (\text{A.5})$$

$$f_{i_1 \cdots i_N}(\mathbf{p}) = n_{i_1} \partial_1 n_{i_2} \partial_2 n_{i_3} \cdots \partial_{N-1} n_{i_N}, \quad (\text{A.6})$$

where the integral is taken over S^{N-1} in the momentum space parameterized by vector $\mathbf{p}$. Here, the area of the $(N-1)$ -dimensional sphere A_{N-1} is expressed by

$$A_{2l-1} = \frac{2\pi^l}{(l-1)!}, \quad (\text{A.7})$$

$$A_{2l-2} = \frac{2^{2l-1} \pi^{l-1} (l-1)!}{(2l-2)!}, \quad (\text{A.8})$$

with a positive integer l . We note that the resultant topology protecting the EP n is neither point-gap nor line-gap topology. In general, the EP n is accompanied by manifolds of EP m s with $2 \leq m < n$ (e.g., a symmetry-protected EP3 emerges on lines of symmetry-protected EP2s). As EP m s require the point-gap closing on the sphere enclosing the EP n , the point-gap topology [90, 91, 93, 94] cannot protect the EP n . In addition, the line-gap topology does not protect band touchings for both real- and imaginary parts [93], simultaneously.

The invariants in Eq. (A.5) are rewritten as Chern numbers or winding numbers of the resultant Hamiltonian [84]. Specifically, for $N = 3$, Eq. (A.5) is rewritten as the first Chern number of the resultant Hamiltonian $H_R(\mathbf{k}) = \mathbf{R} \cdot \boldsymbol{\sigma}$. For $N = 4$, Eq. (A.5) is rewritten as the winding number of three-dimensional chiral symmetric Hamiltonian $H_R = \sum_{i=1, \dots, 4} R_i \gamma_i$ satisfying $\gamma_5 H_R \gamma_5 = -H_R$ with $\boldsymbol{\gamma} = (\sigma_1 \tau_0, \sigma_2 \tau_0, \sigma_3 \tau_1, \sigma_3 \tau_2, \sigma_3 \tau_3)^T$. Here, $\tau_{1, \dots, 3}$ are Pauli matrices, and τ_0 is the 2×2 identity matrix. We note that the presence of the gap of H_R on the sphere is reduced to $\|\mathbf{R}\| \neq 0$. This is because the eigenvalues are given by $E_R = \pm \|\mathbf{R}\|$ which arises from the anti-commutation relation $\{\gamma_i, \gamma_j\} = 2\delta_{ij}$.

A.3 Berry connection and curvature of resultant Hamiltonians for Hopf topology

The $\mathbb{Z}_2$ invariant in Eq. (3) is obtained from the resultant vector $\mathbf{R}=(R_1, R_2, R_3, R_4)^T$ ($R_1, \dots, R_4 \in \mathbb{R}$). Specifically, the Berry connection A_μ and the Berry curvature $F_{\mu\nu}$ are defined as

$$A_\mu = \frac{1}{2\pi i} \langle z | \partial_\mu z \rangle, \quad (\text{A.9})$$

$$F_{\mu\nu} = \frac{1}{2\pi i} \left(\langle \partial_\mu z | \partial_\nu z \rangle - \langle \partial_\nu z | \partial_\mu z \rangle \right), \quad (\text{A.10})$$

with the negative eigenstate $|z\rangle$ of the resultant Hamiltonian $\tilde{\mathbf{n}} \cdot \boldsymbol{\sigma}$ with $\tilde{\mathbf{n}} = \tilde{\mathbf{R}}/\|\tilde{\mathbf{R}}\|$ and $\tilde{\mathbf{R}}=(R_1, R_2, R_3)$. Here, $\sigma_{1, \dots, 3}$ denote Pauli matrices. The ratio $\|\tilde{\mathbf{R}}\|/R_4$ defines the phase $\Delta\varphi(\mathbf{p})=2\varphi(\mathbf{p})$ with $\varphi = \arctan\|\tilde{\mathbf{R}}\|/R_4$ ($0 \leq \varphi \leq \pi$). Substituting $\Delta\varphi$, A_μ , and $F_{\mu\nu}$ into Eq. (3) yields the $\mathbb{Z}_2$ invariant ν_F .

The $\mathbb{Z}$ invariant in Eq. (12) is also obtained from the resultant vector $\mathbf{R}=(R_1, R_2, R_3)^T$ ($R_1, \dots, R_3 \in \mathbb{R}$). Specifically, the Berry connection A_μ and the Berry curvature $F_{\mu\nu}$ are defined in the same way as Eqs. (A.9) and (A.10) except for the definition of $|z\rangle$. In this case, $|z\rangle$ is defined as the negative eigenstate of the resultant Hamiltonian $\mathbf{n} \cdot \boldsymbol{\sigma}$ with $\mathbf{n} = \mathbf{R}/\|\mathbf{R}\|$. Substituting the specific form of the Berry curvature A_μ and the Berry connection $F_{\mu\nu}$ into Eq. (12) yields $\mathbb{Z}$ invariant ν_H .

B Details of the presented Hamiltonians

B.1 Model in Eq. (4) and computation of the $\mathbb{Z}_2$ invariant

The explicit form of ζ 's is

$$\zeta_1 = -2 \sin \phi \left(\eta_\uparrow^* \eta_\downarrow \right), \quad (\text{B.1a})$$

$$\zeta_2 = -\sin \phi \left(|\eta_\uparrow|^2 - |\eta_\downarrow|^2 \right) + i \cos \phi. \quad (\text{B.1b})$$

Here $\eta_\uparrow$, $\eta_\downarrow$ and ϕ are defined as

$$\eta_\uparrow = \sin k_1 + i f(k_5) \sin k_2, \quad (\text{B.2a})$$

$$\eta_\downarrow = \sin k_3 + i \left[\xi(\mathbf{k}) + \frac{3}{2} \sin k_4 - 3(\cos k_5 + \delta) \right], \quad (\text{B.2b})$$

$$\phi = \frac{\pi}{2} (1 - \cos k_4), \quad (\text{B.2c})$$

with $\xi(\mathbf{k}) = \sum_{j=1,2,3} \cos k_j - m_0$ and $f(k_5)$ being $f(k_5)=1$ or $f(k_5)=2 \sin(k_5/2)$.

The resultant vector $\mathbf{R}$ is obtained as

$$\mathbf{R} = (3!)^2 \left(-2 \sin \phi \operatorname{Re}[\eta_\uparrow \eta_\downarrow], -2 \sin \phi \operatorname{Im}[\eta_\uparrow \eta_\downarrow], -\sin \phi (|\eta_\uparrow|^2 - |\eta_\downarrow|^2), \cos \phi \right)^T. \quad (\text{B.3})$$

The above equation indicates that the resultant vector vanishes when $\eta_\uparrow = \eta_\downarrow = 0$ and $\phi = \pi/2$ both hold.

From the given resultant vector [see Eq. (B.3)], the $\mathbb{Z}_2$ invariant ν_F is obtained as follows. With $\tilde{\mathbf{n}} = \tilde{\mathbf{R}}/\|\tilde{\mathbf{R}}\|$ and Eq. (B.3), the resultant Hamiltonian is obtained as

$$\tilde{\mathbf{n}} \cdot \boldsymbol{\sigma} = -\frac{1}{\sqrt{|\eta_\uparrow|^2 + |\eta_\downarrow|^2}} \begin{pmatrix} |\eta_\uparrow|^2 - |\eta_\downarrow|^2 & 2\eta_\uparrow \eta_\downarrow^* \\ 2\eta_\uparrow^* \eta_\downarrow & -|\eta_\uparrow|^2 + |\eta_\downarrow|^2 \end{pmatrix}. \quad (\text{B.4})$$

Thus, the negative eigenstate of $\tilde{\mathbf{n}} \cdot \boldsymbol{\sigma}$ is obtained as

$$|z(\mathbf{k})\rangle = \frac{1}{\sqrt{|\eta_{\uparrow}|^2 + |\eta_{\downarrow}|^2}} \begin{pmatrix} \eta_{\uparrow} \\ \eta_{\downarrow} \end{pmatrix}. \quad (\text{B.5})$$

In addition, from the resultant vector in Eq. (B.3), we obtain

$$\Delta\varphi = 2 \arctan\|\tilde{\mathbf{R}}\|/R_4 = 2\phi. \quad (\text{B.6})$$

Substituting the obtained $|z\rangle$ and $\Delta\varphi$ into Eq. (3), we can numerically compute the $\mathbb{Z}_2$ invariant.

B.2 Model in Eq. (8)

The explicit form of ζ 's are

$$\zeta_1 + i\zeta_2 = -2 \sin \phi (\eta_{\uparrow} \eta_{\downarrow}), \quad (\text{B.7a})$$

$$\zeta_3 = -\sin \phi (|\eta_{\uparrow}|^2 - |\eta_{\downarrow}|^2), \quad (\text{B.7b})$$

$$\zeta_4 = \cos \phi, \quad (\text{B.7c})$$

with η 's and ϕ defined in Eq. (B.2).

The resultant vector of this Hamiltonian is obtained as

$$\mathbf{R} = (5!)^2 \left(-2 \sin \phi \operatorname{Re}[\eta_{\uparrow} \eta_{\downarrow}], -2 \sin \phi \operatorname{Im}[\eta_{\uparrow} \eta_{\downarrow}], -\sin \phi (|\eta_{\uparrow}|^2 - |\eta_{\downarrow}|^2), \cos \phi \right)^T, \quad (\text{B.8})$$

which is proportional to the resultant vector in Eq. (B.3). The above equation indicates that the resultant vector vanishes when $\eta_{\uparrow} = \eta_{\downarrow} = 0$ and $\phi = \pi/2$ both hold.

B.3 Model in Eq. (13)

The explicit form of ζ 's are

$$\zeta_1 + i\zeta_2 = 2(\eta_{\uparrow}^* \eta_{\downarrow}), \quad (\text{B.9a})$$

$$\zeta_3 = |\eta_{\uparrow}|^2 - |\eta_{\downarrow}|^2 + \delta. \quad (\text{B.9b})$$

Here $\eta_{\uparrow}$ and $\eta_{\downarrow}$ are

$$\eta_{\uparrow} = \sin k_1 + i \sin k_2, \quad (\text{B.10a})$$

$$\eta_{\downarrow} = \sin k_3 + i[\xi(\mathbf{k}) + \sin k_4], \quad (\text{B.10b})$$

with $\xi(\mathbf{k}) = \sum_{j=1,2,3} \cos k_j - m_0$.

The resultant vector $\mathbf{R}$ is obtained as

$$\mathbf{R} = -(24)^2 \left(2\operatorname{Re}[\eta_{\uparrow}^* \eta_{\downarrow}], 2\operatorname{Im}[\eta_{\uparrow}^* \eta_{\downarrow}], |\eta_{\uparrow}|^2 - |\eta_{\downarrow}|^2 + \delta \right)^T. \quad (\text{B.11})$$

The above equation indicates that the resultant vector vanishes when $\eta_{\uparrow} = 0$ and $|\eta_{\downarrow}| = \sqrt{\delta}$ both hold for $\delta > 0$ or when $\eta_{\downarrow} = 0$ and $|\eta_{\uparrow}| = \sqrt{|\delta|}$ both hold for $\delta < 0$.




Figure 5: (a) [(b)]: Argument of $R_1 + iR_2$ [band structure] of Hamiltonian (C.1). In panel (b), the complex conjugate of the top band is omitted.

C Fake EP3s

For three-band systems, EP3s with PT -symmetry are characterized by the resultant winding number [84, 88] of the resultant Hamiltonian $R_1\sigma_1 + R_2\sigma_2$ with $R = (r_1, r_2)$ which satisfies chiral symmetry. However, for systems with four or more bands, the resultant winding number may additionally capture “fake EP3s” unless Taylor expansion is applied. This is because the resultant vector may vanish without a triple root in the characteristic polynomial $P(E)$ whose degree is four or higher.

As an example, we consider a non-Hermitian Hamiltonian

$$H = \begin{pmatrix} 1 + ik_1 & 0 & 0 & 0 \\ 0 & 1 - ik_1 & 0 & 0 \\ 0 & 0 & ik_2 & 1 \\ 0 & 0 & 1 & -ik_2 \end{pmatrix}, \quad (\text{C.1})$$

satisfying PT -symmetry [Eq. (5)] with

$$U_{PT} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}. \quad (\text{C.2})$$

The resultant winding number is defined as

$$W_1 = \oint \frac{dp}{2\pi i} \partial_p \log[R_1(p) + iR_2(p)], \quad (\text{C.3})$$

where p parametrizes the circle in the momentum space. The winding number W_1 (see Appendix A.2) is finite for the path illustrated by the black arrow in Fig. 5(a). However, the system does not host EP3s [see Fig. 5(b)].

References

- [1] M. Z. Hasan and C. L. Kane, *Colloquium: Topological insulators*, Rev. Mod. Phys. **82**, 3045 (2010), doi:[10.1103/RevModPhys.82.3045](https://doi.org/10.1103/RevModPhys.82.3045) [preprint doi:[10.48550/arXiv.1002.3895](https://doi.org/10.48550/arXiv.1002.3895)].

- [2] X.-L. Qi and S.-C. Zhang, *Topological insulators and superconductors*, Rev. Mod. Phys. **83**, 1057 (2011), doi:[10.1103/RevModPhys.83.1057](https://doi.org/10.1103/RevModPhys.83.1057) [preprint doi:[10.48550/arXiv.1008.2026](https://doi.org/10.48550/arXiv.1008.2026)].
- [3] C. L. Kane and E. J. Mele, *Z_2 topological order and the quantum spin Hall effect*, Phys. Rev. Lett. **95**, 146802 (2005), doi:[10.1103/PhysRevLett.95.146802](https://doi.org/10.1103/PhysRevLett.95.146802) [preprint doi:[10.48550/arXiv.cond-mat/0506581](https://doi.org/10.48550/arXiv.cond-mat/0506581)].
- [4] C. L. Kane and E. J. Mele, *Quantum spin Hall effect in graphene*, Phys. Rev. Lett. **95**, 226801 (2005), doi:[10.1103/PhysRevLett.95.226801](https://doi.org/10.1103/PhysRevLett.95.226801) [preprint doi:[10.48550/arXiv.cond-mat/0411737](https://doi.org/10.48550/arXiv.cond-mat/0411737)].
- [5] X.-L. Qi, T. L. Hughes and S.-C. Zhang, *Topological field theory of time-reversal invariant insulators*, Phys. Rev. B **78**, 195424 (2008), doi:[10.1103/PhysRevB.78.195424](https://doi.org/10.1103/PhysRevB.78.195424) [preprint doi:[10.48550/arXiv.0802.3537](https://doi.org/10.48550/arXiv.0802.3537)].
- [6] F. D. M. Haldane and S. Raghu, *Possible realization of directional optical waveguides in photonic crystals with broken time-reversal symmetry*, Phys. Rev. Lett. **100**, 013904 (2008), doi:[10.1103/PhysRevLett.100.013904](https://doi.org/10.1103/PhysRevLett.100.013904) [preprint doi:[10.48550/arXiv.cond-mat/0503588](https://doi.org/10.48550/arXiv.cond-mat/0503588)].
- [7] L. Lu, J. D. Joannopoulos and M. Soljačić, *Topological photonics*, Nat. Photonics **8**, 821 (2014), doi:[10.1038/nphoton.2014.248](https://doi.org/10.1038/nphoton.2014.248) [preprint doi:[10.48550/arXiv.1408.6730](https://doi.org/10.48550/arXiv.1408.6730)].
- [8] E. Prodan and C. Prodan, *Topological phonon modes and their role in dynamic instability of microtubules*, Phys. Rev. Lett. **103**, 248101 (2009), doi:[10.1103/PhysRevLett.103.248101](https://doi.org/10.1103/PhysRevLett.103.248101) [preprint doi:[10.48550/arXiv.0909.3492](https://doi.org/10.48550/arXiv.0909.3492)].
- [9] V. V. Albert, L. I. Glazman and L. Jiang, *Topological properties of linear circuit lattices*, Phys. Rev. Lett. **114**, 173902 (2015), doi:[10.1103/PhysRevLett.114.173902](https://doi.org/10.1103/PhysRevLett.114.173902) [preprint doi:[10.48550/arXiv.1410.1243](https://doi.org/10.48550/arXiv.1410.1243)].
- [10] C. H. Lee, S. Imhof, C. Berger, F. Bayer, J. Brehm, L. W. Molenkamp, T. Kiessling and R. Thomale, *Topoelectrical circuits*, Commun. Phys. **1**, 39 (2018), doi:[10.1038/s42005-018-0035-2](https://doi.org/10.1038/s42005-018-0035-2) [preprint doi:[10.48550/arXiv.1705.01077](https://doi.org/10.48550/arXiv.1705.01077)].
- [11] B. Bradlyn, J. Cano, Z. Wang, M. G. Vergniory, C. Felser, R. J. Cava and B. A. Bernevig, *Beyond Dirac and Weyl fermions: Unconventional quasiparticles in conventional crystals*, Science **353**, aaf5037 (2016), doi:[10.1126/science.aaf5037](https://doi.org/10.1126/science.aaf5037) [preprint doi:[10.48550/arXiv.1603.03093](https://doi.org/10.48550/arXiv.1603.03093)].
- [12] A. Y. Kitaev, *Unpaired Majorana fermions in quantum wires*, Phys.-Uspekhi **44**, 131 (2001), doi:[10.1070/1063-7869/44/10S/S29](https://doi.org/10.1070/1063-7869/44/10S/S29) [preprint doi:[10.48550/arXiv.cond-mat/0010440](https://doi.org/10.48550/arXiv.cond-mat/0010440)].
- [13] G. Moore and N. Read, *Nonabelions in the fractional quantum Hall effect*, Nucl. Phys. B **360**, 362 (1991), doi:[10.1016/0550-3213\(91\)90407-O](https://doi.org/10.1016/0550-3213(91)90407-O).
- [14] N. Read and D. Green, *Paired states of fermions in two dimensions with breaking of parity and time-reversal symmetries and the fractional quantum Hall effect*, Phys. Rev. B **61**, 10267 (2000), doi:[10.1103/PhysRevB.61.10267](https://doi.org/10.1103/PhysRevB.61.10267) [preprint doi:[10.48550/arXiv.cond-mat/9906453](https://doi.org/10.48550/arXiv.cond-mat/9906453)].
- [15] L. Fu and C. L. Kane, *Superconducting proximity effect and Majorana fermions at the surface of a topological insulator*, Phys. Rev. Lett. **100**, 096407 (2008), doi:[10.1103/PhysRevLett.100.096407](https://doi.org/10.1103/PhysRevLett.100.096407) [preprint doi:[10.48550/arXiv.0707.1692](https://doi.org/10.48550/arXiv.0707.1692)].

- [16] H. Weyl, *Elektron und Gravitation. I*, Z. Phys. **56**, 330 (1929), doi:[10.1007/BF01339504](https://doi.org/10.1007/BF01339504).
- [17] S. Murakami, *Phase transition between the quantum spin Hall and insulator phases in 3D: Emergence of a topological gapless phase*, New J. Phys. **9**, 356 (2007), doi:[10.1088/1367-2630/9/9/356](https://doi.org/10.1088/1367-2630/9/9/356) [preprint doi:[10.48550/arXiv.0710.0930](https://doi.org/10.48550/arXiv.0710.0930)].
- [18] X. Wan, A. M. Turner, A. Vishwanath and S. Y. Savrasov, *Topological semimetal and Fermi-arc surface states in the electronic structure of pyrochlore iridates*, Phys. Rev. B **83**, 205101 (2011), doi:[10.1103/PhysRevB.83.205101](https://doi.org/10.1103/PhysRevB.83.205101) [preprint doi:[10.48550/arXiv.1007.0016](https://doi.org/10.48550/arXiv.1007.0016)].
- [19] A. A. Burkov and L. Balents, *Weyl semimetal in a topological insulator multilayer*, Phys. Rev. Lett. **107**, 127205 (2011), doi:[10.1103/PhysRevLett.107.127205](https://doi.org/10.1103/PhysRevLett.107.127205) [preprint doi:[10.48550/arXiv.1105.5138](https://doi.org/10.48550/arXiv.1105.5138)].
- [20] B. Yan and C. Felser, *Topological materials: Weyl semimetals*, Annu. Rev. Condens. Matter Phys. **8**, 337 (2017), doi:[10.1146/annurev-conmatphys-031016-025458](https://doi.org/10.1146/annurev-conmatphys-031016-025458) [preprint doi:[10.48550/arXiv.1611.04182](https://doi.org/10.48550/arXiv.1611.04182)].
- [21] S.-Y. Xu et al., *Discovery of a Weyl fermion semimetal and topological Fermi arcs*, Science **349**, 613 (2015), doi:[10.1126/science.aaa9297](https://doi.org/10.1126/science.aaa9297) [preprint doi:[10.48550/arXiv.1502.03807](https://doi.org/10.48550/arXiv.1502.03807)].
- [22] B. Q. Lv et al., *Experimental discovery of Weyl semimetal TaAs*, Phys. Rev. X **5**, 031013 (2015), doi:[10.1103/PhysRevX.5.031013](https://doi.org/10.1103/PhysRevX.5.031013) [preprint doi:[10.48550/arXiv.1502.04684](https://doi.org/10.48550/arXiv.1502.04684)].
- [23] X. Huang et al., *Observation of the chiral-anomaly-induced negative magnetoresistance in 3D Weyl semimetal TaAs*, Phys. Rev. X **5**, 031023 (2015), doi:[10.1103/PhysRevX.5.031023](https://doi.org/10.1103/PhysRevX.5.031023) [preprint doi:[10.48550/arXiv.1503.01304](https://doi.org/10.48550/arXiv.1503.01304)].
- [24] A. P. Schnyder, S. Ryu, A. Furusaki and A. W. W. Ludwig, *Classification of topological insulators and superconductors in three spatial dimensions*, Phys. Rev. B **78**, 195125 (2008), doi:[10.1103/PhysRevB.78.195125](https://doi.org/10.1103/PhysRevB.78.195125) [preprint doi:[10.48550/arXiv.0803.2786](https://doi.org/10.48550/arXiv.0803.2786)].
- [25] A. Kitaev, V. Lebedev and M. Feigel'man, *Periodic table for topological insulators and superconductors*, AIP Conf. Proc. **1134**, 22 (2009), doi:[10.1063/1.3149495](https://doi.org/10.1063/1.3149495) [preprint doi:[10.48550/arXiv.0901.2686](https://doi.org/10.48550/arXiv.0901.2686)].
- [26] S. Ryu, A. P. Schnyder, A. Furusaki and A. W. W. Ludwig, *Topological insulators and superconductors: Tenfold way and dimensional hierarchy*, New J. Phys. **12**, 065010 (2010), doi:[10.1088/1367-2630/12/6/065010](https://doi.org/10.1088/1367-2630/12/6/065010) [preprint doi:[10.48550/arXiv.0912.2157](https://doi.org/10.48550/arXiv.0912.2157)].
- [27] K. Shiozaki and M. Sato, *Topology of crystalline insulators and superconductors*, Phys. Rev. B **90**, 165114 (2014), doi:[10.1103/PhysRevB.90.165114](https://doi.org/10.1103/PhysRevB.90.165114) [preprint doi:[10.48550/arXiv.1403.3331](https://doi.org/10.48550/arXiv.1403.3331)].
- [28] T. Morimoto and A. Furusaki, *Weyl and Dirac semimetals with $\mathbb{Z}_2$ topological charge*, Phys. Rev. B **89**, 235127 (2014), doi:[10.1103/PhysRevB.89.235127](https://doi.org/10.1103/PhysRevB.89.235127) [preprint doi:[10.48550/arXiv.1403.7962](https://doi.org/10.48550/arXiv.1403.7962)].
- [29] C.-K. Chiu and A. P. Schnyder, *Classification of reflection-symmetry-protected topological semimetals and nodal superconductors*, Phys. Rev. B **90**, 205136 (2014), doi:[10.1103/PhysRevB.90.205136](https://doi.org/10.1103/PhysRevB.90.205136) [preprint doi:[10.48550/arXiv.1408.4642](https://doi.org/10.48550/arXiv.1408.4642)].
- [30] T. Bzdušek and M. Sigrist, *Robust doubly charged nodal lines and nodal surfaces in centrosymmetric systems*, Phys. Rev. B **96**, 155105 (2017), doi:[10.1103/PhysRevB.96.155105](https://doi.org/10.1103/PhysRevB.96.155105) [preprint doi:[10.48550/arXiv.1705.07126](https://doi.org/10.48550/arXiv.1705.07126)].

- [31] E. J. Bergholtz, J. C. Budich and F. K. Kunst, *Exceptional topology of non-Hermitian systems*, Rev. Mod. Phys. **93**, 015005 (2021), doi:[10.1103/RevModPhys.93.015005](https://doi.org/10.1103/RevModPhys.93.015005) [preprint doi:[10.48550/arXiv.1912.10048](https://doi.org/10.48550/arXiv.1912.10048)].
- [32] Y. Ashida, Z. Gong and M. Ueda, *Non-Hermitian physics*, Adv. Phys. **69**, 249 (2020), doi:[10.1080/00018732.2021.1876991](https://doi.org/10.1080/00018732.2021.1876991) [preprint doi:[10.48550/arXiv.2006.01837](https://doi.org/10.48550/arXiv.2006.01837)].
- [33] T. Kato, *Perturbation theory for linear operators*, Springer, Berlin, Heidelberg, Germany, ISBN 9783540586616 (1966), doi:[10.1007/978-3-662-12678-3](https://doi.org/10.1007/978-3-662-12678-3).
- [34] W. D. Heiss, M. Müller and I. Rotter, *Collectivity, phase transitions, and exceptional points in open quantum systems*, Phys. Rev. E **58**, 2894 (1998), doi:[10.1103/PhysRevE.58.2894](https://doi.org/10.1103/PhysRevE.58.2894) [preprint doi:[10.48550/arXiv.quant-ph/9805038](https://doi.org/10.48550/arXiv.quant-ph/9805038)].
- [35] I. Rotter, *A non-Hermitian Hamilton operator and the physics of open quantum systems*, J. Phys. A: Math. Theor. **42**, 153001 (2009), doi:[10.1088/1751-8113/42/15/153001](https://doi.org/10.1088/1751-8113/42/15/153001).
- [36] M. V. Berry, *Physics of nonhermitian degeneracies*, Czechoslov. J. Phys. **54**, 1039 (2004), doi:[10.1023/B:CJOP0000044002.05657.04](https://doi.org/10.1023/B:CJOP0000044002.05657.04).
- [37] W. D. Heiss, *The physics of exceptional points*, J. Phys. A: Math. Theor. **45**, 444016 (2012), doi:[10.1088/1751-8113/45/44/444016](https://doi.org/10.1088/1751-8113/45/44/444016) [preprint doi:[10.48550/arXiv.1210.7536](https://doi.org/10.48550/arXiv.1210.7536)].
- [38] H. Shen, B. Zhen and L. Fu, *Topological band theory for non-Hermitian Hamiltonians*, Phys. Rev. Lett. **120**, 146402 (2018), doi:[10.1103/PhysRevLett.120.146402](https://doi.org/10.1103/PhysRevLett.120.146402) [preprint doi:[10.48550/arXiv.1706.07435](https://doi.org/10.48550/arXiv.1706.07435)].
- [39] Y. Xu, S.-T. Wang and L.-M. Duan, *Weyl exceptional rings in a three-dimensional dissipative cold atomic gas*, Phys. Rev. Lett. **118**, 045701 (2017), doi:[10.1103/PhysRevLett.118.045701](https://doi.org/10.1103/PhysRevLett.118.045701) [preprint doi:[10.48550/arXiv.1611.02239](https://doi.org/10.48550/arXiv.1611.02239)].
- [40] J. Carlström and E. J. Bergholtz, *Exceptional links and twisted Fermi ribbons in non-Hermitian systems*, Phys. Rev. A **98**, 042114 (2018), doi:[10.1103/PhysRevA.98.042114](https://doi.org/10.1103/PhysRevA.98.042114) [preprint doi:[10.48550/arXiv.1807.03330](https://doi.org/10.48550/arXiv.1807.03330)].
- [41] J. C. Budich, J. Carlström, F. K. Kunst and E. J. Bergholtz, *Symmetry-protected nodal phases in non-Hermitian systems*, Phys. Rev. B **99**, 041406 (2019), doi:[10.1103/PhysRevB.99.041406](https://doi.org/10.1103/PhysRevB.99.041406) [preprint doi:[10.48550/arXiv.1810.00914](https://doi.org/10.48550/arXiv.1810.00914)].
- [42] R. Okugawa and T. Yokoyama, *Topological exceptional surfaces in non-Hermitian systems with parity-time and parity-particle-hole symmetries*, Phys. Rev. B **99**, 041202 (2019), doi:[10.1103/PhysRevB.99.041202](https://doi.org/10.1103/PhysRevB.99.041202) [preprint doi:[10.48550/arXiv.1810.03376](https://doi.org/10.48550/arXiv.1810.03376)].
- [43] H. Zhou, J. Y. Lee, S. Liu and B. Zhen, *Exceptional surfaces in PT-symmetric non-Hermitian photonic systems*, Optica **6**, 190 (2019), doi:[10.1364/OPTICA.6.000190](https://doi.org/10.1364/OPTICA.6.000190) [preprint doi:[10.48550/arXiv.1810.06549](https://doi.org/10.48550/arXiv.1810.06549)].
- [44] T. Yoshida, R. Peters, N. Kawakami and Y. Hatsugai, *Symmetry-protected exceptional rings in two-dimensional correlated systems with chiral symmetry*, Phys. Rev. B **99**, 121101 (2019), doi:[10.1103/PhysRevB.99.121101](https://doi.org/10.1103/PhysRevB.99.121101) [preprint doi:[10.48550/arXiv.1810.06297](https://doi.org/10.48550/arXiv.1810.06297)].
- [45] K. Kimura, T. Yoshida and N. Kawakami, *Chiral-symmetry protected exceptional torus in correlated nodal-line semimetals*, Phys. Rev. B **100**, 115124 (2019), doi:[10.1103/PhysRevB.100.115124](https://doi.org/10.1103/PhysRevB.100.115124) [preprint doi:[10.48550/arXiv.1905.11761](https://doi.org/10.48550/arXiv.1905.11761)].

- [46] Z. Yang, A. P. Schnyder, J. Hu and C.-K. Chiu, *Fermion doubling theorems in two-dimensional non-Hermitian systems for Fermi points and exceptional points*, Phys. Rev. Lett. **126**, 086401 (2021), doi:[10.1103/PhysRevLett.126.086401](https://doi.org/10.1103/PhysRevLett.126.086401) [preprint doi:[10.48550/arXiv.1912.02788](https://doi.org/10.48550/arXiv.1912.02788)].
- [47] M. M. Denner, A. Skurativska, F. Schindler, M. H. Fischer, R. Thomale, T. Bzdušek and T. Neupert, *Exceptional topological insulators*, Nat. Commun. **12**, 5681 (2021), doi:[10.1038/s41467-021-25947-z](https://doi.org/10.1038/s41467-021-25947-z) [preprint doi:[10.48550/arXiv.2008.01090](https://doi.org/10.48550/arXiv.2008.01090)].
- [48] T. Yoshida, R. Okugawa and Y. Hatsugai, *Discriminant indicators with generalized inversion symmetry*, Phys. Rev. B **105**, 085109 (2022), doi:[10.1103/PhysRevB.105.085109](https://doi.org/10.1103/PhysRevB.105.085109) [preprint doi:[10.48550/arXiv.2111.07077](https://doi.org/10.48550/arXiv.2111.07077)].
- [49] X.-Q. Sun, C. C. Wojcik, S. Fan and T. Bzdušek, *Alice strings in non-Hermitian systems*, Phys. Rev. Res. **2**, 023226 (2020), doi:[10.1103/PhysRevResearch.2.023226](https://doi.org/10.1103/PhysRevResearch.2.023226) [preprint doi:[10.48550/arXiv.1905.04338](https://doi.org/10.48550/arXiv.1905.04338)].
- [50] C. C. Wojcik, X.-Q. Sun, T. Bzdušek and S. Fan, *Homotopy characterization of non-Hermitian Hamiltonians*, Phys. Rev. B **101**, 205417 (2020), doi:[10.1103/PhysRevB.101.205417](https://doi.org/10.1103/PhysRevB.101.205417) [preprint doi:[10.48550/arXiv.1911.12748](https://doi.org/10.48550/arXiv.1911.12748)].
- [51] Z. Li and R. S. K. Mong, *Homotopical characterization of non-Hermitian band structures*, Phys. Rev. B **103**, 155129 (2021), doi:[10.1103/PhysRevB.103.155129](https://doi.org/10.1103/PhysRevB.103.155129) [preprint doi:[10.48550/arXiv.1911.02697](https://doi.org/10.48550/arXiv.1911.02697)].
- [52] K. Yang, Z. Li, J. L. K. König, L. Rødland, M. Stålhammar and E. J. Bergholtz, *Homotopy, symmetry, and non-Hermitian band topology*, Rep. Prog. Phys. **87**, 078002 (2024), doi:[10.1088/1361-6633/ad4e64](https://doi.org/10.1088/1361-6633/ad4e64) [preprint doi:[10.48550/arXiv.2309.14416](https://doi.org/10.48550/arXiv.2309.14416)].
- [53] V. Kozii and L. Fu, *Non-Hermitian topological theory of finite-lifetime quasiparticles: Prediction of bulk Fermi arc due to exceptional point*, Phys. Rev. B **109**, 235139 (2024), doi:[10.1103/PhysRevB.109.235139](https://doi.org/10.1103/PhysRevB.109.235139).
- [54] A. A. Zyuzin and A. Y. Zyuzin, *Flat band in disorder-driven non-Hermitian Weyl semimetals*, Phys. Rev. B **97**, 041203 (2018), doi:[10.1103/PhysRevB.97.041203](https://doi.org/10.1103/PhysRevB.97.041203) [preprint doi:[10.48550/arXiv.1710.05344](https://doi.org/10.48550/arXiv.1710.05344)].
- [55] T. Yoshida, R. Peters and N. Kawakami, *Non-Hermitian perspective of the band structure in heavy-fermion systems*, Phys. Rev. B **98**, 035141 (2018), doi:[10.1103/PhysRevB.98.035141](https://doi.org/10.1103/PhysRevB.98.035141) [preprint doi:[10.48550/arXiv.1805.01172](https://doi.org/10.48550/arXiv.1805.01172)].
- [56] T. Yoshida, R. Peters, N. Kawakami and Y. Hatsugai, *Exceptional band touching for strongly correlated systems in equilibrium*, Prog. Theor. Exp. Phys. **12A**109 (2020), doi:[10.1093/ptep/ptaa059](https://doi.org/10.1093/ptep/ptaa059) [preprint doi:[10.48550/arXiv.2002.11265](https://doi.org/10.48550/arXiv.2002.11265)].
- [57] H. Shen and L. Fu, *Quantum oscillation from in-gap states and a non-Hermitian Landau level problem*, Phys. Rev. Lett. **121**, 026403 (2018), doi:[10.1103/PhysRevLett.121.026403](https://doi.org/10.1103/PhysRevLett.121.026403) [preprint doi:[10.48550/arXiv.1802.03023](https://doi.org/10.48550/arXiv.1802.03023)].
- [58] Y. Nagai, Y. Qi, H. Isobe, V. Kozii and L. Fu, *DMFT reveals the non-Hermitian topology and Fermi arcs in heavy-fermion systems*, Phys. Rev. Lett. **125**, 227204 (2020), doi:[10.1103/PhysRevLett.125.227204](https://doi.org/10.1103/PhysRevLett.125.227204) [preprint doi:[10.48550/arXiv.2005.06729](https://doi.org/10.48550/arXiv.2005.06729)].
- [59] M. Am-Shallem, R. Kosloff and N. Moiseyev, *Exceptional points for parameter estimation in open quantum systems: Analysis of the Bloch equations*, New J. Phys. **17**, 113036 (2015), doi:[10.1088/1367-2630/17/11/113036](https://doi.org/10.1088/1367-2630/17/11/113036) [preprint doi:[10.48550/arXiv.1411.6364](https://doi.org/10.48550/arXiv.1411.6364)].

- [60] F. Minganti, A. Miranowicz, R. W. Chhajlany and F. Nori, *Quantum exceptional points of non-Hermitian Hamiltonians and Liouvillians: The effects of quantum jumps*, Phys. Rev. A **100**, 062131 (2019), doi:[10.1103/PhysRevA.100.062131](https://doi.org/10.1103/PhysRevA.100.062131) [preprint doi:[10.48550/arXiv.1909.12276](https://doi.org/10.48550/arXiv.1909.12276)].
- [61] M. Nakagawa, N. Kawakami and M. Ueda, *Exact Liouvillian spectrum of a one-dimensional dissipative Hubbard model*, Phys. Rev. Lett. **126**, 110404 (2021), doi:[10.1103/PhysRevLett.126.110404](https://doi.org/10.1103/PhysRevLett.126.110404) [preprint doi:[10.48550/arXiv.2003.14202](https://doi.org/10.48550/arXiv.2003.14202)].
- [62] W. Chen, M. Abbasi, B. Ha, S. Erdamar, Y. N. Joglekar and K. W. Murch, *Decoherence-induced exceptional points in a dissipative superconducting Qubit*, Phys. Rev. Lett. **128**, 110402 (2022), doi:[10.1103/PhysRevLett.128.110402](https://doi.org/10.1103/PhysRevLett.128.110402) [preprint doi:[10.48550/arXiv.2111.04754](https://doi.org/10.48550/arXiv.2111.04754)].
- [63] C. Dembowski, H.-D. Gräf, H. L. Harney, A. Heine, W. D. Heiss, H. Rehfeld and A. Richter, *Experimental observation of the topological structure of exceptional points*, Phys. Rev. Lett. **86**, 787 (2001), doi:[10.1103/PhysRevLett.86.787](https://doi.org/10.1103/PhysRevLett.86.787).
- [64] S.-B. Lee, J. Yang, S. Moon, S.-Y. Lee, J.-B. Shim, S. W. Kim, J.-H. Lee and K. An, *Observation of an exceptional point in a chaotic optical microcavity*, Phys. Rev. Lett. **103**, 134101 (2009), doi:[10.1103/PhysRevLett.103.134101](https://doi.org/10.1103/PhysRevLett.103.134101) [preprint doi:[10.48550/arXiv.0905.4478](https://doi.org/10.48550/arXiv.0905.4478)].
- [65] J. Doppler et al., *Dynamically encircling an exceptional point for asymmetric mode switching*, Nature **537**, 76 (2016), doi:[10.1038/nature18605](https://doi.org/10.1038/nature18605) [preprint doi:[10.48550/arXiv.1603.02325](https://doi.org/10.48550/arXiv.1603.02325)].
- [66] B. Zhen, C. W. Hsu, Y. Igarashi, L. Lu, I. Kaminer, A. Pick, S.-L. Chua, J. D. Joannopoulos and M. Soljačić, *Spawning rings of exceptional points out of Dirac cones*, Nature **525**, 354 (2015), doi:[10.1038/nature14889](https://doi.org/10.1038/nature14889) [preprint doi:[10.48550/arXiv.1504.00734](https://doi.org/10.48550/arXiv.1504.00734)].
- [67] K. Takata and M. Notomi, *Photonic topological insulating phase induced solely by gain and loss*, Phys. Rev. Lett. **121**, 213902 (2018), doi:[10.1103/PhysRevLett.121.213902](https://doi.org/10.1103/PhysRevLett.121.213902) [preprint doi:[10.48550/arXiv.1710.09169](https://doi.org/10.48550/arXiv.1710.09169)].
- [68] H. Zhou, C. Peng, Y. Yoon, C. W. Hsu, K. A. Nelson, L. Fu, J. D. Joannopoulos, M. Soljačić and B. Zhen, *Observation of bulk Fermi arc and polarization half charge from paired exceptional points*, Science **359**, 1009 (2018), doi:[10.1126/science.aap9859](https://doi.org/10.1126/science.aap9859) [preprint doi:[10.48550/arXiv.1709.03044](https://doi.org/10.48550/arXiv.1709.03044)].
- [69] H. Meng, Y. S. Ang and C. H. Lee, *Exceptional points in non-Hermitian systems: Applications and recent developments*, Appl. Phys. Lett. **124**, 060502 (2024), doi:[10.1063/5.0183826](https://doi.org/10.1063/5.0183826).
- [70] T. Ozawa et al., *Topological photonics*, Rev. Mod. Phys. **91**, 015006 (2019), doi:[10.1103/RevModPhys.91.015006](https://doi.org/10.1103/RevModPhys.91.015006) [preprint doi:[10.48550/arXiv.1802.04173](https://doi.org/10.48550/arXiv.1802.04173)].
- [71] M.-A. Miri and A. Alù, *Exceptional points in optics and photonics*, Science **363**, eaar7709 (2019), doi:[10.1126/science.aar7709](https://doi.org/10.1126/science.aar7709).
- [72] X. Zhu, H. Ramezani, C. Shi, J. Zhu and X. Zhang, *$\mathcal{PT}$ -symmetric acoustics*, Phys. Rev. X **4**, 031042 (2014), doi:[10.1103/PhysRevX.4.031042](https://doi.org/10.1103/PhysRevX.4.031042).
- [73] C. Shi, M. Dubois, Y. Chen, L. Cheng, H. Ramezani, Y. Wang and X. Zhang, *Accessing the exceptional points of parity-time symmetric acoustics*, Nat. Commun. **7**, 11110 (2016), doi:[10.1038/ncomms11110](https://doi.org/10.1038/ncomms11110).

- [74] T. Yoshida and Y. Hatsugai, *Exceptional rings protected by emergent symmetry for mechanical systems*, Phys. Rev. B **100**, 054109 (2019), doi:[10.1103/PhysRevB.100.054109](https://doi.org/10.1103/PhysRevB.100.054109) [preprint doi:[10.48550/arXiv.1904.10764](https://doi.org/10.48550/arXiv.1904.10764)].
- [75] T. Hofmann et al., *Reciprocal skin effect and its realization in a topoelectrical circuit*, Phys. Rev. Res. **2**, 023265 (2020), doi:[10.1103/PhysRevResearch.2.023265](https://doi.org/10.1103/PhysRevResearch.2.023265) [preprint doi:[10.48550/arXiv.1908.02759](https://doi.org/10.48550/arXiv.1908.02759)].
- [76] Y. Li et al., *Anti-parity-time symmetry in diffusive systems*, Science **364**, 170 (2019), doi:[10.1126/science.aaw6259](https://doi.org/10.1126/science.aaw6259).
- [77] W. Tang, X. Jiang, K. Ding, Y.-X. Xiao, Z.-Q. Zhang, C. T. Chan and G. Ma, *Exceptional nexus with a hybrid topological invariant*, Science **370**, 1077 (2020), doi:[10.1126/science.abd8872](https://doi.org/10.1126/science.abd8872).
- [78] W. Tang, K. Ding and G. Ma, *Realization and topological properties of third-order exceptional lines embedded in exceptional surfaces*, Nat. Commun. **14**, 6660 (2023), doi:[10.1038/s41467-023-42414-z](https://doi.org/10.1038/s41467-023-42414-z) [preprint doi:[10.48550/arXiv.2211.15921](https://doi.org/10.48550/arXiv.2211.15921)].
- [79] W. D. Heiss, *Chirality of wavefunctions for three coalescing levels*, J. Phys. A: Math. Theor. **41**, 244010 (2008), doi:[10.1088/1751-8113/41/24/244010](https://doi.org/10.1088/1751-8113/41/24/244010) [preprint doi:[10.48550/arXiv.0708.1392](https://doi.org/10.48550/arXiv.0708.1392)].
- [80] E. M. Graefe, U. Günther, H. J. Korsch and A. E. Niederle, *A non-Hermitian $\mathcal{PT}$ symmetric Bose-Hubbard model: Eigenvalue rings from unfolding higher-order exceptional points*, J. Phys. A: Math. Theor. **41**, 255206 (2008), doi:[10.1088/1751-8113/41/25/255206](https://doi.org/10.1088/1751-8113/41/25/255206) [preprint doi:[10.48550/arXiv.0802.3164](https://doi.org/10.48550/arXiv.0802.3164)].
- [81] G. Demange and E.-M. Graefe, *Signatures of three coalescing eigenfunctions*, J. Phys. A: Math. Theor. **45**, 025303 (2011), doi:[10.1088/1751-8113/45/2/025303](https://doi.org/10.1088/1751-8113/45/2/025303) [preprint doi:[10.48550/arXiv.1110.1489](https://doi.org/10.48550/arXiv.1110.1489)].
- [82] J.-W. Ryu, J.-H. Han and C.-H. Yi, *Classification of multiple arbitrary-order non-Hermitian singularities*, Phys. Rev. A **106**, 012218 (2022), doi:[10.1103/PhysRevA.106.012218](https://doi.org/10.1103/PhysRevA.106.012218) [preprint doi:[10.48550/arXiv.2112.02547](https://doi.org/10.48550/arXiv.2112.02547)].
- [83] K. Yang and I. Mandal, *Enhanced eigenvector sensitivity and algebraic classification of sublattice-symmetric exceptional points*, Phys. Rev. B **107**, 144304 (2023), doi:[10.1103/PhysRevB.107.144304](https://doi.org/10.1103/PhysRevB.107.144304) [preprint doi:[10.48550/arXiv.2211.08449](https://doi.org/10.48550/arXiv.2211.08449)].
- [84] P. Delplace, T. Yoshida and Y. Hatsugai, *Symmetry-protected multifold exceptional points and their topological characterization*, Phys. Rev. Lett. **127**, 186602 (2021), doi:[10.1103/PhysRevLett.127.186602](https://doi.org/10.1103/PhysRevLett.127.186602) [preprint doi:[10.48550/arXiv.2103.08232](https://doi.org/10.48550/arXiv.2103.08232)].
- [85] I. Mandal and E. J. Bergholtz, *Symmetry and higher-order exceptional points*, Phys. Rev. Lett. **127**, 186601 (2021), doi:[10.1103/PhysRevLett.127.186601](https://doi.org/10.1103/PhysRevLett.127.186601) [preprint doi:[10.48550/arXiv.2103.15729](https://doi.org/10.48550/arXiv.2103.15729)].
- [86] S. Sayyad and F. K. Kunst, *Realizing exceptional points of any order in the presence of symmetry*, Phys. Rev. Res. **4**, 023130 (2022), doi:[10.1103/PhysRevResearch.4.023130](https://doi.org/10.1103/PhysRevResearch.4.023130) [preprint doi:[10.48550/arXiv.2202.07009](https://doi.org/10.48550/arXiv.2202.07009)].
- [87] A. Montag and F. K. Kunst, *Symmetry-induced higher-order exceptional points in two dimensions*, Phys. Rev. Res. **6**, 023205 (2024), doi:[10.1103/PhysRevResearch.6.023205](https://doi.org/10.1103/PhysRevResearch.6.023205) [preprint doi:[10.48550/arXiv.2401.10913](https://doi.org/10.48550/arXiv.2401.10913)].

- [88] T. Yoshida, J. L. K. König, L. Rødland, E. J. Bergholtz and M. Stålhammar, *Winding topology of multifold exceptional points*, Phys. Rev. Res. **7**, L012021 (2025), doi:[10.1103/PhysRevResearch.7.L012021](https://doi.org/10.1103/PhysRevResearch.7.L012021) [preprint doi:[10.48550/arXiv.2409.09153](https://doi.org/10.48550/arXiv.2409.09153)].
- [89] M. Stålhammar and L. Rødland, *Abelian spectral topology of multifold exceptional points*, Phys. Rev. Res. **7**, 033246 (2025), doi:[10.1103/8lyx-vtnw](https://doi.org/10.1103/8lyx-vtnw).
- [90] K. Kawabata, T. Bessho and M. Sato, *Classification of exceptional points and non-Hermitian topological semimetals*, Phys. Rev. Lett. **123**, 066405 (2019), doi:[10.1103/PhysRevLett.123.066405](https://doi.org/10.1103/PhysRevLett.123.066405) [preprint doi:[10.48550/arXiv.1902.08479](https://doi.org/10.48550/arXiv.1902.08479)].
- [91] Z. Gong, Y. Ashida, K. Kawabata, K. Takasan, S. Higashikawa and M. Ueda, *Topological phases of non-Hermitian systems*, Phys. Rev. X **8**, 031079 (2018), doi:[10.1103/PhysRevX.8.031079](https://doi.org/10.1103/PhysRevX.8.031079) [preprint doi:[10.48550/arXiv.1802.07964](https://doi.org/10.48550/arXiv.1802.07964)].
- [92] K. Kawabata, S. Higashikawa, Z. Gong, Y. Ashida and M. Ueda, *Topological unification of time-reversal and particle-hole symmetries in non-Hermitian physics*, Nat. Commun. **10**, 297 (2019), doi:[10.1038/s41467-018-08254-y](https://doi.org/10.1038/s41467-018-08254-y) [preprint doi:[10.48550/arXiv.1804.04676](https://doi.org/10.48550/arXiv.1804.04676)].
- [93] K. Kawabata, K. Shiozaki, M. Ueda and M. Sato, *Symmetry and topology in non-Hermitian physics*, Phys. Rev. X **9**, 041015 (2019), doi:[10.1103/PhysRevX.9.041015](https://doi.org/10.1103/PhysRevX.9.041015) [preprint doi:[10.48550/arXiv.1812.09133](https://doi.org/10.48550/arXiv.1812.09133)].
- [94] H. Zhou and J. Y. Lee, *Periodic table for topological bands with non-Hermitian symmetries*, Phys. Rev. B **99**, 235112 (2019), doi:[10.1103/PhysRevB.99.235112](https://doi.org/10.1103/PhysRevB.99.235112) [preprint doi:[10.48550/arXiv.1812.10490](https://doi.org/10.48550/arXiv.1812.10490)].
- [95] Z. Lin, A. Pick, M. Lončar and A. W. Rodriguez, *Enhanced spontaneous emission at third-order Dirac exceptional points in inverse-designed photonic crystals*, Phys. Rev. Lett. **117**, 107402 (2016), doi:[10.1103/PhysRevLett.117.107402](https://doi.org/10.1103/PhysRevLett.117.107402) [preprint doi:[10.48550/arXiv.1605.06175](https://doi.org/10.48550/arXiv.1605.06175)].
- [96] J. Schnabel, H. Cartarius, J. Main, G. Wunner and W. D. Heiss, *$\mathcal{PT}$ -symmetric waveguide system with evidence of a third-order exceptional point*, Phys. Rev. A **95**, 053868 (2017), doi:[10.1103/PhysRevA.95.053868](https://doi.org/10.1103/PhysRevA.95.053868) [preprint doi:[10.48550/arXiv.1703.05091](https://doi.org/10.48550/arXiv.1703.05091)].
- [97] J. Wiersig, *Review of exceptional point-based sensors*, Photonics Res. **8**, 1457 (2020), doi:[10.1364/PRJ.396115](https://doi.org/10.1364/PRJ.396115).
- [98] J. Wiersig, *Revisiting the hierarchical construction of higher-order exceptional points*, Phys. Rev. A **106**, 063526 (2022), doi:[10.1103/PhysRevA.106.063526](https://doi.org/10.1103/PhysRevA.106.063526) [preprint doi:[10.48550/arXiv.2210.04459](https://doi.org/10.48550/arXiv.2210.04459)].
- [99] K. Wang, L. Xiao, H. Lin, W. Yi, E. J. Bergholtz and P. Xue, *Experimental simulation of symmetry-protected higher-order exceptional points with single photons*, Sci. Adv. **9**, eadi0732 (2023), doi:[10.1126/sciadv.adi0732](https://doi.org/10.1126/sciadv.adi0732) [preprint doi:[10.48550/arXiv.2303.11834](https://doi.org/10.48550/arXiv.2303.11834)].
- [100] H. Hodaei, A. U. Hassan, S. Wittek, H. Garcia-Gracia, R. El-Ganainy, D. N. Christodoulides and M. Khajavikhan, *Enhanced sensitivity at higher-order exceptional points*, Nature **548**, 187 (2017), doi:[10.1038/nature23280](https://doi.org/10.1038/nature23280).
- [101] K. Bai, J.-Z. Li, T.-R. Liu, L. Fang, D. Wan and M. Xiao, *Nonlinear exceptional points with a complete basis in dynamics*, Phys. Rev. Lett. **130**, 266901 (2023), doi:[10.1103/PhysRevLett.130.266901](https://doi.org/10.1103/PhysRevLett.130.266901).

- [102] Q. Liu, D. A. Kessler and E. Barkai, *Designing exceptional-point-based graphs yielding topologically guaranteed quantum search*, Phys. Rev. Res. **5**, 023141 (2023), doi:[10.1103/PhysRevResearch.5.023141](https://doi.org/10.1103/PhysRevResearch.5.023141).
- [103] Y. S. S. Patil, J. Höller, P. A. Henry, C. Guria, Y. Zhang, L. Jiang, N. Kralj, N. Read and J. G. E. Harris, *Measuring the knot of non-Hermitian degeneracies and non-commuting braids*, Nature **607**, 271 (2022), doi:[10.1038/s41586-022-04796-w](https://doi.org/10.1038/s41586-022-04796-w) [preprint doi:[10.48550/arXiv.2112.00157](https://doi.org/10.48550/arXiv.2112.00157)].
- [104] A. Leclerc, L. Jezequel, N. Perez, A. Bhandare, G. Laibe and P. Delplace, *Exceptional ring of the buoyancy instability in stars*, Phys. Rev. Res. **6**, L012055 (2024), doi:[10.1103/PhysRevResearch.6.L012055](https://doi.org/10.1103/PhysRevResearch.6.L012055).
- [105] L. Crippa, J. C. Budich and G. Sangiovanni, *Fourth-order exceptional points in correlated quantum many-body systems*, Phys. Rev. B **104**, L121109 (2021), doi:[10.1103/PhysRevB.104.L121109](https://doi.org/10.1103/PhysRevB.104.L121109) [preprint doi:[10.48550/arXiv.2106.11987](https://doi.org/10.48550/arXiv.2106.11987)].
- [106] M. Stålhammar and E. J. Bergholtz, *Classification of exceptional nodal topologies protected by $\mathcal{PT}$ symmetry*, Phys. Rev. B **104**, L201104 (2021), doi:[10.1103/PhysRevB.104.L201104](https://doi.org/10.1103/PhysRevB.104.L201104) [preprint doi:[10.48550/arXiv.2106.04582](https://doi.org/10.48550/arXiv.2106.04582)].
- [107] N. Hatano, *Exceptional points of the Lindblad operator of a two-level system*, Mol. Phys. **117**, 2121 (2019), doi:[10.1080/00268976.2019.1593535](https://doi.org/10.1080/00268976.2019.1593535) [preprint doi:[10.48550/arXiv.1903.04676](https://doi.org/10.48550/arXiv.1903.04676)].
- [108] S. Khandelwal, N. Brunner and G. Haack, *Signatures of Liouvillian exceptional points in a quantum thermal machine*, PRX Quantum **2**, 040346 (2021), doi:[10.1103/PRXQuantum.2.040346](https://doi.org/10.1103/PRXQuantum.2.040346) [preprint doi:[10.48550/arXiv.2101.11553](https://doi.org/10.48550/arXiv.2101.11553)].
- [109] Y. Wu, Y. Wang, X. Ye, W. Liu, Z. Niu, C.-K. Duan, Y. Wang, X. Rong and J. Du, *Third-order exceptional line in a nitrogen-vacancy spin system*, Nat. Nanotechnol. **19**, 160 (2024), doi:[10.1038/s41565-023-01583-0](https://doi.org/10.1038/s41565-023-01583-0).
- [110] J.-W. Ryu, J.-H. Han, C.-H. Yi, M. J. Park and H. C. Park, *Exceptional classifications of non-Hermitian systems*, Commun. Phys. **7**, 109 (2024), doi:[10.1038/s42005-024-01595-9](https://doi.org/10.1038/s42005-024-01595-9).
- [111] H. Freudenthal, *Über die Klassen der Sphärenabbildungen I. Große Dimensionen*, Compos. Math. **5**, 299 (1938).
- [112] E. Witten, *Current algebra, baryons, and quark confinement*, Nucl. Phys. B **223**, 433 (1983), doi:[10.1016/0550-3213\(83\)90064-0](https://doi.org/10.1016/0550-3213(83)90064-0).
- [113] P. G. Grinevich and G. E. Volovik, *Topology of gap nodes in superfluid³He: π_4 homotopy group for³He-B disclination*, J. Low Temp. Phys. **72**, 371 (1988), doi:[10.1007/BF00682148](https://doi.org/10.1007/BF00682148).
- [114] T. Schuster, S. Gazit, J. E. Moore and N. Y. Yao, *Floquet Hopf insulators*, Phys. Rev. Lett. **123**, 266803 (2019), doi:[10.1103/PhysRevLett.123.266803](https://doi.org/10.1103/PhysRevLett.123.266803) [preprint doi:[10.48550/arXiv.1903.02558](https://doi.org/10.48550/arXiv.1903.02558)].
- [115] J. E. Moore, Y. Ran and X.-G. Wen, *Topological surface states in three-dimensional magnetic insulators*, Phys. Rev. Lett. **101**, 186805 (2008), doi:[10.1103/PhysRevLett.101.186805](https://doi.org/10.1103/PhysRevLett.101.186805) [preprint doi:[10.48550/arXiv.0804.4527](https://doi.org/10.48550/arXiv.0804.4527)].

- [116] D.-L. Deng, S.-T. Wang, C. Shen and L.-M. Duan, *Hopf insulators and their topologically protected surface states*, Phys. Rev. B **88**, 201105 (2013), doi:[10.1103/PhysRevB.88.201105](https://doi.org/10.1103/PhysRevB.88.201105) [preprint doi:[10.48550/arXiv.1307.7206](https://doi.org/10.48550/arXiv.1307.7206)].
- [117] Z. Yang and J. Hu, *Non-Hermitian Hopf-link exceptional line semimetals*, Phys. Rev. B **99**, 081102 (2019), doi:[10.1103/PhysRevB.99.081102](https://doi.org/10.1103/PhysRevB.99.081102) [preprint doi:[10.48550/arXiv.1807.05661](https://doi.org/10.48550/arXiv.1807.05661)].
- [118] F. Wilczek and A. Zee, *Linking numbers, spin, and statistics of solitons*, Phys. Rev. Lett. **51**, 2250 (1983), doi:[10.1103/PhysRevLett.51.2250](https://doi.org/10.1103/PhysRevLett.51.2250).
- [119] R. Kennedy, *Topological Hopf-Chern insulators and the Hopf superconductor*, Phys. Rev. B **94**, 035137 (2016), doi:[10.1103/PhysRevB.94.035137](https://doi.org/10.1103/PhysRevB.94.035137) [preprint doi:[10.48550/arXiv.1604.02840](https://doi.org/10.48550/arXiv.1604.02840)].
- [120] A. Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, UK, ISBN 9780521795401 (2001).
- [121] H. Toda, *Composition methods in homotopy groups of spheres*, Princeton University Press, Princeton, USA, ISBN 9780691095868 (1963).
- [122] N. Read and E. Rezayi, *Beyond paired quantum Hall states: Parafermions and incompressible states in the first excited Landau level*, Phys. Rev. B **59**, 8084 (1999), doi:[10.1103/PhysRevB.59.8084](https://doi.org/10.1103/PhysRevB.59.8084) [preprint doi:[10.48550/arXiv.cond-mat/9809384](https://doi.org/10.48550/arXiv.cond-mat/9809384)].
- [123] P. Fendley, *Parafermionic edge zero modes in Z_n -invariant spin chains*, J. Stat. Mech.: Theory Exp. P11020 (2012), doi:[10.1088/1742-5468/2012/11/P11020](https://doi.org/10.1088/1742-5468/2012/11/P11020) [preprint doi:[10.48550/arXiv.1209.0472](https://doi.org/10.48550/arXiv.1209.0472)].
- [124] R. S. K. Mong et al., *Universal topological quantum computation from a superconductor-Abelian quantum Hall heterostructure*, Phys. Rev. X **4**, 011036 (2014), doi:[10.1103/PhysRevX.4.011036](https://doi.org/10.1103/PhysRevX.4.011036) [preprint doi:[10.48550/arXiv.1307.4403](https://doi.org/10.48550/arXiv.1307.4403)].
- [125] H. Liu, R. Perea-Causin and E. J. Bergholtz, *Parafermions in moiré minibands*, Nat. Commun. **16**, 1770 (2025), doi:[10.1038/s41467-025-57035-x](https://doi.org/10.1038/s41467-025-57035-x) [preprint doi:[10.48550/arXiv.2406.08546](https://doi.org/10.48550/arXiv.2406.08546)].
- [126] F. N. Rybakov, N. S. Kiselev, A. B. Borisov, L. Döring, C. Melcher and S. Blügel, *Magnetic hopfions in solids*, APL Mater. **10**, 111113 (2022), doi:[10.1063/5.0099942](https://doi.org/10.1063/5.0099942) [preprint doi:[10.48550/arXiv.1904.00250](https://doi.org/10.48550/arXiv.1904.00250)].