

A brief note on the G_2 Affleck-Kennedy-Lieb-Tasaki chain

Hosho Katsura^{1,2,3*} and Dirk Schuricht^{4†}

¹ Department of Physics, Graduate School of Science, The University of Tokyo,
7-3-1, Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan

² Institute for Physics of Intelligence, The University of Tokyo,
7-3-1, Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan

³ Trans-scale Quantum Science Institute, The University of Tokyo,
7-3-1, Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan

⁴ Institute for Theoretical Physics, Center for Extreme Matter and Emergent Phenomena,
Utrecht University, Princetonplein 5, 3584 CE Utrecht, The Netherlands

★ katsura@phys.s.u-tokyo.ac.jp, † d.schuricht@uu.nl

Part of the *In Memoriam: Ian Affleck Collection*

Abstract

We consider the valence bond solid (VBS) state built of singlet pairs of fundamental representations and projected onto adjoint representations of the exceptional Lie group G_2 . The two-point correlation function in the VBS state is non-vanishing only for nearest neighbours, but possesses finite string order. We construct a parent Hamiltonian for the VBS state, which constitutes the G_2 analog of the famous AKLT chain.



Copyright H. Katsura and D. Schuricht.
This work is licensed under the Creative Commons
[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).
Published by the SciPost Foundation.

Received 2025-03-28

Accepted 2025-07-17

Published 2025-08-15

doi:[10.21468/SciPostPhysCore.8.3.052](https://doi.org/10.21468/SciPostPhysCore.8.3.052)



Check for
updates

Contents

1	Introduction	2
2	Valence bond solid state	2
3	Construction of a parent Hamiltonian	5
4	Static correlation functions	6
5	String order	7
6	Conclusion	7
A	Explicit representation matrices	8
B	On the tensor product $7 \otimes 7$	9

C Matrix product representation of the VBS state	9
D Family of parent Hamiltonians	11
References	12

In memory of Ian Affleck.

1 Introduction

In 1987 Ian Affleck together with Tom Kennedy, Elliot Lieb and Hal Tasaki introduced [1, 2] a class of exactly solvable Hamiltonians. These Hamiltonians possess unique ground states constructed out of valence bonds and were thus named valence bond solids (VBSs). It was shown that there exists an energy gap and that the VBS states have short-range correlations and unbroken symmetry. Its simplest variant, now generally known as the Affleck–Kennedy–Lieb–Tasaki (AKLT) chain, corresponds to a specific point in the phase diagram [3, 4] of the bilinear-biquadratic spin-1 isotropic quantum chain. In fact, the AKLT chain is located in the phase of the isotropic spin-1 Heisenberg chain (also known as the Haldane chain). Thus the physics of the AKLT chain provided a very intuitive framework for the appearance of the ‘Haldane gap’ [5–7], which was originally predicted using sophisticated field-theoretical arguments. More generally, VBS states helped to understand the fundamental distinction between integer and half-integer quantum spin chains [8, 9], provided examples for quantum number fractionalisation and edge modes [10–12], and, in their matrix product form, constitute a natural framework for numerical approaches like the density matrix renormalisation group method [13, 14]. The very intuitive construction of the AKLT chain initiated numerous generalisations, including to higher spins and higher-dimensional models (see for example [15]), but also extensions of the underlying spin symmetry to q -deformed [16–21] and supersymmetric [22, 23] models as well as higher-dimensional Lie groups like $SU(n)$ [24–30], $SO(n)$ [31, 32] and $SP(n)$ [33]. Here we will extend this list with G_2 , the simplest exceptional Lie group.

In this brief note we consider an N -site spin chain, where the local degrees of freedom at each lattice site transform under the 14-dimensional, adjoint representation of the exceptional Lie group G_2 . We construct a valence bond solid (VBS) state and calculate static correlation functions as well as a string order parameter. Furthermore, we derive a parent Hamiltonian of which the VBS state is a zero-energy ground state. Our results can be regarded as the G_2 version of the famous AKLT construction.

2 Valence bond solid state

We consider the exceptional Lie group G_2 [34, 35], whose Lie algebra $\mathfrak{g}_2$ is 14-dimensional. We denote the generators by Λ^a , $a = 1, \dots, 14$. Furthermore we use bold symbols $\Lambda = (\Lambda^1, \dots, \Lambda^{14})$ to denote the vector formed by the generators. The algebra $\mathfrak{g}_2$ has rank two, thus two of the generators can be chosen to be diagonal, with the states in a given representation being labeled by their eigenvalues, see Figures 2 and 3 for examples.

The fundamental representation of G_2 is denoted by its dimension 7, meaning the 14 generators Λ^a can be represented as 7×7 matrices. A possible explicit realisation is given in Appendix A. We fix the normalisation of the generators via $\text{tr}[(\Lambda^a)^\dagger \Lambda^b] = 2\delta_{ab}$. The quadratic Casimir operator takes the value $\Lambda^2 = \Lambda \cdot \Lambda = \sum_{a=1}^{14} \Lambda^a \Lambda^a = 4 \text{id} \equiv 4$. Here and in the following we assume the generators to be Hermitian.




Figure 1: Graphical representation of the VBS state $|\Psi_{\text{VBS}}\rangle$. Each circle stands for a fundamental representation **7**, each line joining two circles for a singlet bond, i.e., the singlet **1** in the tensor product (**1**), and each oval for a lattice site on which we project onto the adjoint representation **14**.

For the construction of the physical Hilbert space we consider a chain with N lattice sites and periodic boundary conditions. At each lattice site we place two fundamental representations **7**, see Figure 1 for a sketch. On every lattice site we thus have the tensor product (see Appendix B for more details)

$$\mathbf{7} \otimes \mathbf{7} = \mathbf{1} \oplus \mathbf{7} \oplus \mathbf{14} \oplus \mathbf{27}, \quad (1)$$

where on the right-hand side we see its decomposition into the irreducible representations **1** (singlet), **7** (fundamental), **14** (adjoint) and **27**. We recall that the appearing representations are labeled by their dimensions. Under exchange of the factors on the left-hand side, i.e., under the action of the permutation operator on $\mathbf{7} \otimes \mathbf{7}$, the representations **1** and **27** are even, while **7** and **14** are odd. Furthermore the eigenvalues of the quadratic Casimir operator $(\Lambda_1 + \Lambda_2)^2$ are 0, 4, 8 and $28/3$, respectively. Finally, the physical Hilbert space is obtained by projecting¹ onto the factor **14** on the right-hand side, thus obtaining a tensor product of N local Hilbert spaces each transforming in the 14-dimensional, adjoint representation of G_2 .

In this Hilbert space, the VBS state is constructed as follows: We form a singlet between one of the fundamental representations **7** on lattice site j with one on the neighbouring site $j - 1$, while we form another singlet with the second representation **7** on lattice site j with one on the neighbouring site $j + 1$. We stress that the formation of these singlets is imposed in addition to the already implemented projection onto the adjoint representation **14** at each lattice site. If we further impose periodic boundary conditions this yields a unique² VBS state $|\Psi_{\text{VBS}}\rangle$, which is translationally invariant and can be represented graphically as shown in Figure 1. The entanglement entropy of the VBS state can be straightforwardly obtained using its Schmidt decomposition [38]. The result in the thermodynamic limit is $S_{\text{EE}} = 2 \ln 7$, where the factor 2 stems from the fact that for the system with periodic boundary conditions two singlet bonds have to be cut, each contributing $\ln 7$.

As we show in Appendix C one can derive an explicit matrix product representation for the VBS state, which takes the form

$$|\Psi_{\text{VBS}}\rangle = \sum_{a_1, a_2, \dots, a_N=1}^{14} \text{tr}_7[M^{a_1} M^{a_2} \cdots M^{a_N}] |\psi_{a_1}\rangle_1 |\psi_{a_2}\rangle_2 \cdots |\psi_{a_N}\rangle_N, \quad (2)$$

where $M^a = \Lambda^a / \sqrt{2}$ with Λ^a , $a = 1, \dots, 14$, defined in (A.2) and the trace is taken over the fundamental representation **7**. Here we introduced the local states $|\psi_a\rangle_j$, $a = 1, \dots, 14$, forming an orthonormal basis of the adjoint representation **14** at lattice site j . Explicitly, these

¹We note that since the representation **14** is odd under the exchange of the factors in $\mathbf{7} \otimes \mathbf{7}$, the projection cannot be implemented using Schwinger bosons creating the particles in the fundamental representations, as can be done in the original AKLT chain [36, 37].

²In the case of open boundary conditions on the lattice sites 1 and N one unpaired fundamental representation **7** would remain. Thus the construction would yield $7 \cdot 7 = 49$ VBS states.

basis states are given by

$$\begin{aligned}
 |\psi_1\rangle &= \frac{|10\rangle + |13\rangle}{\sqrt{2}}, & |\psi_2\rangle &= -i \frac{|10\rangle - |13\rangle}{\sqrt{2}}, & |\psi_3\rangle &= -\frac{|3\rangle + \sqrt{3}|9\rangle}{2}, \\
 |\psi_4\rangle &= \frac{|5\rangle + |12\rangle}{\sqrt{2}}, & |\psi_5\rangle &= -i \frac{|5\rangle - |12\rangle}{\sqrt{2}}, & |\psi_6\rangle &= \frac{|4\rangle + |8\rangle}{\sqrt{2}}, \\
 |\psi_7\rangle &= -i \frac{|4\rangle - |8\rangle}{\sqrt{2}}, & |\psi_8\rangle &= \frac{-\sqrt{3}|3\rangle + |9\rangle}{2}, & |\psi_9\rangle &= i \frac{|6\rangle + i|7\rangle}{\sqrt{2}}, \\
 |\psi_{10}\rangle &= \frac{|6\rangle - i|7\rangle}{\sqrt{2}}, & |\psi_{11}\rangle &= i \frac{|2\rangle - i|11\rangle}{\sqrt{2}}, & |\psi_{12}\rangle &= -\frac{|2\rangle + i|11\rangle}{\sqrt{2}}, \\
 |\psi_{13}\rangle &= -i \frac{|1\rangle + i|14\rangle}{\sqrt{2}}, & |\psi_{14}\rangle &= \frac{|1\rangle - i|14\rangle}{\sqrt{2}}, & &
 \end{aligned} \tag{3}$$

where $\{|a\rangle : a = 1, \dots, 14\}$ is another basis of the adjoint representation that diagonalises the elements of the Cartan subalgebra, i.e., Λ^3 and Λ^8 ; we show the corresponding weight diagram in Figure 2. In Appendix C we provide an explicit basis in terms of the basis states of the underlying fundamental representations in the tensor product (1). We note that the matrices M^a , $a = 1, \dots, 14$, are injective, and thus the state (2) is an injective matrix product state (see for example Theorem 7.5 in Reference [15] for the definition). The representation (2) will be the starting point for the calculation of the static correlation functions and string order parameters below.




Figure 2: Weight diagram of the adjoint representation **14**. The state with $\Lambda^3 = \Lambda^8 = 0$ is doubly degenerate. An explicit expansion of the basis states in terms of the basis states of the underlying fundamental representations in the tensor product (1) is provided in Appendix C. The states appearing in the matrix product state (2) are given in (3).

3 Construction of a parent Hamiltonian

Having constructed the VBS state on the N -site chain, we now proceed in deriving a parent Hamiltonian. Specifically we construct a Hamiltonian H_{AKLT} which possesses the VBS state $|\Psi_{\text{VBS}}\rangle$ as its zero-energy ground state.

We start by considering two neighbouring sites of the chain. We denote the local generators by Λ_j^a , $a = 1, \dots, 14$, where the subindex indicates that they act non-trivially at site j only.³ These local generators act in the adjoint representation **14**, thus on the two neighbouring sites we have to consider the tensor product

$$\mathbf{14} \otimes \mathbf{14} = \mathbf{1} \oplus \mathbf{14} \oplus \mathbf{27} \oplus \mathbf{77} \oplus \mathbf{77}'. \quad (4)$$

We note that there exist two non-isomorphic 77-dimensional representations in the decomposition. The eigenvalues of the quadratic Casimir operator $(\Lambda_j + \Lambda_{j+1})^2$ on the representations on the right-hand side of (4) are 0, 8, 28/3, 16 and 20, respectively. The representations **1**, **27** and **77'** are even under exchange of the factors on the left-hand side, the representations **14** and **77** are odd.

The parent Hamiltonian for the VBS state is constructed by noting that on each two neighbouring sites in the VBS state we find one singlet and two uncoupled fundamental representations. Hence, the total G_2 contents on two neighbouring sites is given by the tensor product (1). If we construct an operator which annihilates (1) but is strictly positive on the complement of (1) in (4), we will obtain the VBS state as zero-energy ground state. Such an operator is conveniently implemented using the quadratic Casimir operator $(\Lambda_j + \Lambda_{j+1})^2$ on the two sites by

$$H_{j,j+1} = \frac{1}{1152} (\Lambda_j + \Lambda_{j+1})^2 \left[(\Lambda_j + \Lambda_{j+1})^2 - 8 \right] \left[(\Lambda_j + \Lambda_{j+1})^2 - \frac{28}{3} \right], \quad (5)$$

where the subindices indicate that the operator is acting locally on sites j and $j+1$. We note that the operators $H_{j,j+1}$ are not projectors⁴ as they take the different values 20/27 and 20/9 on the representations **77** and **77'**. The prefactor in (5) is chosen for convenience.

As the operators (5) annihilate the VBS state (2), a translationally invariant parent Hamiltonian H_{AKLT} is obtained by simply summing them up for each bond on the chain. Furthermore using $\Lambda_j^2 = 8$ and thus $(\Lambda_j + \Lambda_{j+1})^2 = 16 + 2\Lambda_j \cdot \Lambda_{j+1}$ we find⁵

$$H_{\text{AKLT}} = \sum_{j=1}^N \left[\frac{1}{2} \Lambda_j \cdot \Lambda_{j+1} + \frac{23}{216} (\Lambda_j \cdot \Lambda_{j+1})^2 + \frac{1}{144} (\Lambda_j \cdot \Lambda_{j+1})^3 + \frac{20}{27} \right]. \quad (6)$$

By construction we have $H_{\text{AKLT}} |\Psi_{\text{VBS}}\rangle = 0$, and as all terms in H_{AKLT} are non-negative we conclude that $|\Psi_{\text{VBS}}\rangle$ is indeed a zero-energy ground state. Full numerical diagonalisation of the Hamiltonian on $N = 4$ lattice sites shows that the ground state is unique for periodic boundary conditions and 49th fold degenerate for open ones.⁶ We stress that the generators appearing in the Hamiltonian act in the 14-dimensional, adjoint representation of G_2 . The original AKLT chain is formulated in terms of spin-1 operators, which form the adjoint representation of $SU(2)$. In this sense the model (6) can be regarded as the G_2 analog of the AKLT chain.

³More precisely, they are defined as $\Lambda_j^a = \text{id}^{\otimes j-1} \otimes \Lambda^a \otimes \text{id}^{\otimes N-j}$, where id is the identity matrix of dimension 14.

⁴In fact, using the projectors onto **77** and **77'** one can construct a whole family of parent Hamiltonians. We present some details in Appendix D.

⁵We note that the Hamiltonian is manifestly interacting, in line with a general argument that G_2 invariance is incompatible with non-interacting systems [39].

⁶Since the VBS state (2) is injective, following the procedure by Perez-Garcia et al. [40] one can construct a parent Hamiltonian such that the VBS state is its unique gapped ground state. However, this procedure typically yields more than two-body interactions, and thus there is no guarantee that the resulting parent Hamiltonian coincides with H_{AKLT} .

4 Static correlation functions

The explicit matrix product form (2) of the VBS state allows us to exactly calculate the static correlation functions. We follow the procedure presented for the original AKLT chain in Reference [15]. First we introduce a matrix $\tilde{M}^a$ obtained by conjugation of M^a , $\tilde{M}_{\sigma,\sigma'}^a = (M_{\sigma,\sigma'}^a)^*$, i.e., without taking the transpose. Using this we define a 49×49 transfer matrix

$$R_{\mu,\nu} = R_{(\sigma,\tau),(\sigma',\tau')} = \sum_{a=1}^{14} \tilde{M}_{\sigma,\sigma'}^a M_{\tau,\tau'}^a, \quad (7)$$

where the indices $\mu, \nu = 1, 2, \dots, 49$ correspond to the double indices $(1, 1), (1, 2), \dots, (7, 7)$. The transfer matrix R is introduced for each lattice site, but given the translational invariance we suppress the index j . Then the squared norm of the VBS state is given by

$$\langle \Psi_{\text{VBS}} | \Psi_{\text{VBS}} \rangle = \text{tr}(R^N) = 2^N + 7 + 27 \left(-\frac{1}{3}\right)^N, \quad (8)$$

where we used that the eigenvalues of R are given by 2 (unique), 1 (7-fold), $-1/3$ (27-fold) and 0 (14-fold). Note that the eigenvector corresponding to the leading eigenvalue 2 is given by $|\phi_0\rangle = \frac{1}{\sqrt{7}} \sum_{\sigma=1}^7 |\sigma\rangle_7 |\sigma\rangle_7$, which is a maximally entangled state⁷ between the two fundamental representations. The uniqueness of the leading eigenvalue of R implies the injectivity of the VBS state (2).

Next we introduce a similar transfer matrix including the local action of a G_2 generator, namely

$$T_{\mu,\nu}^a = T_{(\sigma,\tau),(\sigma',\tau')}^a = \sum_{b,c=1}^{14} \tilde{M}_{\sigma,\sigma'}^b \langle \psi_b | \Lambda^a | \psi_c \rangle M_{\tau,\tau'}^c, \quad (9)$$

where the generator Λ^a acts on the adjoint representation 14. Using this we obtain

$$\langle \Psi_{\text{VBS}} | \Lambda_1^a \Lambda_j^b | \Psi_{\text{VBS}} \rangle = \text{tr}(T^a R^{j-2} T^b R^{N-j}). \quad (10)$$

Note that we have again suppressed the site indices, keeping the necessary information by the number of matrices R between T^a and T^b . The trace can be evaluated in the eigenbasis of R . As can be checked⁸ using the explicit matrices given in Appendix A, we have $RT^b |\phi_0\rangle = 0$, thus leading to

$$\langle \Lambda_1^a \Lambda_{j \geq 3}^b \rangle \equiv \frac{\langle \Psi_{\text{VBS}} | \Lambda_1^a \Lambda_{j \geq 3}^b | \Psi_{\text{VBS}} \rangle}{\langle \Psi_{\text{VBS}} | \Psi_{\text{VBS}} \rangle} \sim 2^{-N} \rightarrow 0, \quad \text{for } N \rightarrow \infty. \quad (11)$$

For nearest neighbours, however, the result remains finite in the thermodynamic limit and is given by

$$\langle \Lambda_j^a \Lambda_{j+1}^b \rangle \equiv \frac{\langle \Psi_{\text{VBS}} | \Lambda_j^a \Lambda_{j+1}^b | \Psi_{\text{VBS}} \rangle}{\langle \Psi_{\text{VBS}} | \Psi_{\text{VBS}} \rangle} = -\left(\frac{2}{7} + \frac{2}{3} 2^{-N} + \dots\right) \delta_{ab}. \quad (12)$$

Thus we conclude that the static correlations in the VBS state are nearest neighbour only, in contrast to the exponentially decaying correlations in the original AKLT chain [1, 2] or various of its generalisations [17, 27, 29, 31, 33].

⁷Here we used the following property of the maximally entangled state. Let id_7 be the 7×7 identity matrix. For any 7×7 matrix A , we have $(A \otimes \text{id}_7) |\phi_0\rangle = (\text{id}_7 \otimes A^T) |\phi_0\rangle$. Noting that $R = \sum_{a=1}^{14} \tilde{M}^a \otimes M^a$, we find $R |\phi_0\rangle = \frac{1}{2} (\text{id}_7 \otimes \Lambda^2) |\phi_0\rangle = 2 |\phi_0\rangle$, where we used the fact that $\Lambda^2 |\sigma\rangle_7 = 4 |\sigma\rangle_7$ for all $\sigma = 1, \dots, 7$.

⁸More specifically we need the following properties: (i) $\langle \psi_b | \Lambda^a | \psi_c \rangle = -\langle \psi_c | \Lambda^a | \psi_b \rangle$, $a = 1, \dots, 14$, for the generators of G_2 in 14 and (ii) $\sum_{b=1}^{14} \Lambda^b \Lambda^a \Lambda^b = 0$, $a = 1, \dots, 14$, for the generators of G_2 in 7.

5 String order

While the correlation functions in the AKLT chain are short-ranged, hidden antiferromagnetic order exists. This can be seen by considering a non-local string operator [41, 42] which possesses a finite expectation value. Here, the matrix product form (2) allows the calculation of a similar string order. Specifically we define the string operators

$$O_{ij}^{ab}(\alpha, \beta) = -\Lambda_i^a \exp\left(i\pi\sqrt{2}\alpha \sum_{k=i+1}^{j-1} \Lambda_k^3 + i\pi\sqrt{6}\beta \sum_{k=i+1}^{j-1} \Lambda_k^8\right) \Lambda_j^b, \quad (13)$$

where the string between the sites i and j is over the diagonal generators, which are Λ^3 and Λ^8 in our convention, see (C.3). Introducing a transfer matrix representing the exponential at site k via

$$T(\alpha, \beta)_{\mu, \nu} = T(\alpha, \beta)_{(\sigma, \tau), (\sigma', \tau')} = \sum_{c, d=1}^{14} \tilde{M}_{\sigma, \sigma'}^c \langle \psi_c | \exp\left(i\pi\sqrt{2}\alpha\Lambda_k^3 + i\pi\sqrt{6}\beta\Lambda_k^8\right) | \psi_d \rangle M_{\tau, \tau'}^d, \quad (14)$$

we obtain

$$\langle \Psi_{\text{VBS}} | O_{1j}^{ab}(\alpha, \beta) | \Psi_{\text{VBS}} \rangle = -\text{tr}(T^a T(\alpha, \beta)^{j-2} T^b R^{N-j}). \quad (15)$$

A finite result in the thermodynamic limit requires the non-vanishing matrix element of $T^a T(\alpha, \beta)^{j-2} T^b$ multiplying the 2^{N-j} coming from R^{N-j} . We have found that as $N \rightarrow \infty$

$$\langle O_{1j}^{33}(\frac{1}{2}, 0) \rangle \equiv \frac{\langle \Psi_{\text{VBS}} | O_{1j}^{33}(\frac{1}{2}, 0) | \Psi_{\text{VBS}} \rangle}{\langle \Psi_{\text{VBS}} | \Psi_{\text{VBS}} \rangle} = \frac{8}{49} \left(1 + 27 \frac{(-1)^j}{6^j}\right) \xrightarrow{j \rightarrow \infty} \frac{8}{49}, \quad (16)$$

while for example $\langle O_{1j}^{33}(\frac{1}{2}, \frac{1}{2}) \rangle \sim 2^{-j}$. Thus we observe finite string order (16). It is unclear whether there exists a non-local Kennedy–Tasaki transformation [43, 44] that maps the VBS state to a classical product state. We note that on general grounds a family of models including H_{AKLT} as a special case does not allow for symmetry protected topological phases with respect to G_2 [45]. Thus we expect that the VBS state (2) can be smoothly connected to a trivial state, in analogy to the situation for the spin-2 AKLT state [12].

6 Conclusion

We considered a spin chain where the local Hilbert spaces transform under the adjoint representation 14 of G_2 , constructed a VBS state in this chain as well as a parent Hamiltonian for which the VBS state is a zero-energy ground state. Furthermore we showed that in the thermodynamic limit the static correlation functions are non-vanishing for nearest neighbours only, but that a non-local string order exists. In this sense our results can be seen as a G_2 analog of the famous AKLT chain.

Acknowledgments

We would like to thank Philippe Lecheminant and Thomas Quella for useful comments on the manuscript. DS thanks the organisers of the 2025 Banff workshop “Exact Solutions in Quantum Information: Entanglement, Topology, and Quantum Circuits”, where this work has been initiated.

Funding information HK was supported by JSPS KAKENHI Grants No. JP23K25783, No. JP23K25790, and MEXT KAKENHI Grant-in-Aid for Transformative Research Areas A “Extreme Universe” (KAKENHI Grant No. JP21H05191).

A Explicit representation matrices

An explicit representation for the 14 generators Λ^a , $a = 1, \dots, 14$, in the fundamental representation **7** can be chosen as (see for example Eq. (14) in Reference [46])

$$\sum_{a=1}^{14} f_a \Lambda^a = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & f_3 & f_1 - if_2 & f_4 - if_5 & 0 & 0 & 0 \\ 0 & f_1 + if_2 & -f_3 & f_6 - if_7 & 0 & 0 & 0 \\ 0 & f_4 + if_5 & f_6 + if_7 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -f_3 & -f_1 - if_2 & -f_4 - if_5 \\ 0 & 0 & 0 & 0 & -f_1 + if_2 & f_3 & -f_6 - if_7 \\ 0 & 0 & 0 & 0 & -f_4 + if_5 & -f_6 + if_7 & 0 \end{pmatrix} \quad (\text{A.1})$$

$$+ \frac{1}{\sqrt{3}} \begin{pmatrix} 0 & f_{13} - if_{14} & f_{11} - if_{12} & if_9 + f_{10} & -if_{13} + f_{14} & -if_{11} + f_{12} & -f_9 - if_{10} \\ f_{13} + if_{14} & \frac{f_8}{\sqrt{2}} & 0 & 0 & 0 & \frac{f_9 - if_{10}}{\sqrt{2}} & \frac{if_{11} + f_{12}}{\sqrt{2}} \\ f_{11} + if_{12} & 0 & \frac{f_8}{\sqrt{2}} & 0 & -\frac{f_9 - if_{10}}{\sqrt{2}} & 0 & -\frac{if_{13} + f_{14}}{\sqrt{2}} \\ -if_9 + f_{10} & 0 & 0 & -\sqrt{2}f_8 & -\frac{if_{11} + f_{12}}{\sqrt{2}} & \frac{if_{13} + f_{14}}{\sqrt{2}} & 0 \\ if_{13} + f_{14} & 0 & -\frac{f_9 + if_{10}}{\sqrt{2}} & \frac{if_{11} - f_{12}}{\sqrt{2}} & -\frac{f_8}{\sqrt{2}} & 0 & 0 \\ if_{11} + f_{12} & \frac{f_9 + if_{10}}{\sqrt{2}} & 0 & -\frac{if_{13} - f_{14}}{\sqrt{2}} & 0 & -\frac{f_8}{\sqrt{2}} & 0 \\ -f_9 + if_{10} & -\frac{if_{11} - f_{12}}{\sqrt{2}} & \frac{if_{13} - f_{14}}{\sqrt{2}} & 0 & 0 & 0 & \sqrt{2}f_8 \end{pmatrix}.$$

The so defined generators Λ^a are Hermitian and satisfy

$$\text{tr}[(\Lambda^a)^\dagger \Lambda^b] = 2\delta_{ab}, \quad (\text{A.2})$$

$$[\Lambda^a, \Lambda^b] = \sum_{c=1}^{14} f^{abc} \Lambda^c, \quad \text{with} \quad f^{abc} = \frac{1}{2} \text{tr}([\Lambda^a, \Lambda^b] \Lambda^c), \quad (\text{A.3})$$

$$\Lambda^2 = \sum_{a=1}^{14} \Lambda^a \Lambda^a = 4. \quad (\text{A.4})$$

The last equation states that the eigenvalue of the quadratic Casimir operator Λ^2 is 4 in the fundamental representation. We note that the basis defined above exhibits the appearance of the Gell-Mann matrices, showing that $\text{su}(3)$ appears as a sub-algebra [34, 46]. The weight diagram of the fundamental representation is shown in Figure 3.

The adjoint representation is 14-dimensional. Corresponding generators can be defined via $(\Lambda^a)_{bc} = f^{acb}$, which act on the basis $\{|\psi_a\rangle : a = 1, \dots, 14\}$ defined in (3). We note that none of the generators Λ^a are diagonal in this basis. The quadratic Casimir operator takes the value $\Lambda^2 = 8$. Alternatively one can define the generators of **14** from the projection in the tensor product $\mathbf{7} \otimes \mathbf{7}$, see below. The weight diagram is shown in Figure 2.




Figure 3: Weight diagram of the fundamental representation $\mathbf{7}$. The dots represent eigenstates $|\sigma\rangle$, $\sigma = 1, \dots, 7$, of the two diagonal generators Λ^3 and Λ^8 , as fixed by the explicit matrix representation (A.2).

B On the tensor product $\mathbf{7} \otimes \mathbf{7}$

The projectors onto the irreducible representations in the decomposition of the tensor product (1) are explicitly given by

$$P_1 = -\frac{3}{896} [(\Lambda_1 + \Lambda_2)^2 - 4] [(\Lambda_1 + \Lambda_2)^2 - 8] [(\Lambda_1 + \Lambda_2)^2 - \frac{28}{3}], \quad (\text{B.1})$$

$$P_7 = \frac{3}{256} (\Lambda_1 + \Lambda_2)^2 [(\Lambda_1 + \Lambda_2)^2 - 8] [(\Lambda_1 + \Lambda_2)^2 - \frac{28}{3}], \quad (\text{B.2})$$

$$P_{14} = -\frac{3}{128} (\Lambda_1 + \Lambda_2)^2 [(\Lambda_1 + \Lambda_2)^2 - 4] [(\Lambda_1 + \Lambda_2)^2 - \frac{28}{3}], \quad (\text{B.3})$$

$$P_{27} = \frac{27}{1792} (\Lambda_1 + \Lambda_2)^2 [(\Lambda_1 + \Lambda_2)^2 - 4] [(\Lambda_1 + \Lambda_2)^2 - 8]. \quad (\text{B.4})$$

Here $\Lambda_{1,2}$ denote the generators in the first and second fundamental representation $\mathbf{7}$ respectively. In the following we use the explicit matrices given in Appendix A, i.e., we fix the basis $\{|\sigma\rangle_{1,2} : \sigma = 1, \dots, 7\}$. Then the unique singlet state can be written as

$$\begin{aligned} |\circ-\circ\rangle &= \frac{1}{\sqrt{7}} (i|1\rangle_1|1\rangle_2 + |2\rangle_1|5\rangle_2 + |3\rangle_1|6\rangle_2 + |4\rangle_1|7\rangle_2 + |5\rangle_1|2\rangle_2 + |6\rangle_1|3\rangle_2 + |7\rangle_1|4\rangle_2) \\ &= \frac{1}{\sqrt{7}} (i|1\rangle_1, |2\rangle_1, |3\rangle_1, \dots, |7\rangle_1) (|1\rangle_2, |5\rangle_2, |6\rangle_2, |7\rangle_2, |2\rangle_2, |3\rangle_2, |4\rangle_2)^t \\ &= \frac{1}{\sqrt{7}} \vec{v}_1 \vec{w}_2^t, \end{aligned} \quad (\text{B.5})$$

where in the last line we have introduced auxiliary vectors $\vec{v}_1$ and $\vec{w}_2$ with elements in the local Hilbert spaces of $\mathbf{7}_{1,2}$. We note that the singlet state is even under exchange of the factors. Similarly, a basis of the adjoint representation can be obtained from the eigenspace of the projector P_{14} , see Appendix C below.

C Matrix product representation of the VBS state

In this appendix we derive the explicit matrices M^a in the matrix product representation (2). We start with a singlet formed by two fundamental representations at neighbouring lattices

j and $j + 1$, which according to (B.5) is given by $\vec{v}_j \vec{w}_{j+1}^t$ and we dropped the normalisation. Thus considering two singlets on the bonds $(j - 1, j)$ and $(j, j + 1)$ we find

$$|\circ - \circ \quad \circ - \circ\rangle \propto \vec{v}_{j-1} \vec{w}_j^t \vec{v}_j \vec{w}_{j+1}^t = \vec{v}_{j-1} \mathcal{M}_j^{7 \otimes 7} \vec{w}_{j+1}^t, \quad (\text{C.1})$$

where we introduced the matrix $\mathcal{M}_j^{7 \otimes 7} = \vec{w}_j^t \vec{v}_j$ which takes values in the tensor product $7 \otimes 7$ at lattice site j . Explicitly we find

$$\mathcal{M}_j^{7 \otimes 7} = \begin{pmatrix} i|1\rangle_7|1\rangle_7 & |1\rangle_7|2\rangle_7 & |1\rangle_7|3\rangle_7 & |1\rangle_7|4\rangle_7 & |1\rangle_7|5\rangle_7 & |1\rangle_7|6\rangle_7 & |1\rangle_7|7\rangle_7 \\ i|5\rangle_7|1\rangle_7 & |5\rangle_7|2\rangle_7 & |5\rangle_7|3\rangle_7 & |5\rangle_7|4\rangle_7 & |5\rangle_7|5\rangle_7 & |5\rangle_7|6\rangle_7 & |5\rangle_7|7\rangle_7 \\ i|6\rangle_7|1\rangle_7 & |6\rangle_7|2\rangle_7 & |6\rangle_7|3\rangle_7 & |6\rangle_7|4\rangle_7 & |6\rangle_7|5\rangle_7 & |6\rangle_7|6\rangle_7 & |6\rangle_7|7\rangle_7 \\ i|7\rangle_7|1\rangle_7 & |7\rangle_7|2\rangle_7 & |7\rangle_7|3\rangle_7 & |7\rangle_7|4\rangle_7 & |7\rangle_7|5\rangle_7 & |7\rangle_7|6\rangle_7 & |7\rangle_7|7\rangle_7 \\ i|2\rangle_7|1\rangle_7 & |2\rangle_7|2\rangle_7 & |2\rangle_7|3\rangle_7 & |2\rangle_7|4\rangle_7 & |2\rangle_7|5\rangle_7 & |2\rangle_7|6\rangle_7 & |2\rangle_7|7\rangle_7 \\ i|3\rangle_7|1\rangle_7 & |3\rangle_7|2\rangle_7 & |3\rangle_7|3\rangle_7 & |3\rangle_7|4\rangle_7 & |3\rangle_7|5\rangle_7 & |3\rangle_7|6\rangle_7 & |3\rangle_7|7\rangle_7 \\ i|4\rangle_7|1\rangle_7 & |4\rangle_7|2\rangle_7 & |4\rangle_7|3\rangle_7 & |4\rangle_7|4\rangle_7 & |4\rangle_7|5\rangle_7 & |4\rangle_7|6\rangle_7 & |4\rangle_7|7\rangle_7 \end{pmatrix}. \quad (\text{C.2})$$

Here the subscripts indicate that the states belong to the fundamental representations **7**. The next step is now to project the states in the matrix $\mathcal{M}_j^{7 \otimes 7}$ onto the adjoint representation **14**. To this end we obtain the representation space as the eigenspace of the projector P_{14} and within this choose the orthonormal basis (with the subscripts indicating the respective representations)

$$\begin{aligned} |1\rangle_{14} &= -\frac{1}{\sqrt{3}}(|1\rangle_7|2\rangle_7 - |2\rangle_7|1\rangle_7) - \frac{1}{\sqrt{6}}(|6\rangle_7|7\rangle_7 - |7\rangle_7|6\rangle_7), \\ |2\rangle_{14} &= \frac{1}{\sqrt{3}}(|1\rangle_7|3\rangle_7 - |3\rangle_7|1\rangle_7) - \frac{1}{\sqrt{6}}(|5\rangle_7|7\rangle_7 - |7\rangle_7|5\rangle_7), \\ |3\rangle_{14} &= \frac{1}{2}(|2\rangle_7|5\rangle_7 - |5\rangle_7|2\rangle_7) - \frac{1}{2}(|4\rangle_7|7\rangle_7 - |7\rangle_7|4\rangle_7), \\ |4\rangle_{14} &= -\frac{1}{\sqrt{2}}(|3\rangle_7|7\rangle_7 - |7\rangle_7|3\rangle_7), \\ |5\rangle_{14} &= -\frac{1}{\sqrt{2}}(|2\rangle_7|7\rangle_7 - |7\rangle_7|2\rangle_7), \\ |6\rangle_{14} &= -\frac{1}{\sqrt{3}}(|1\rangle_7|7\rangle_7 - |7\rangle_7|1\rangle_7) + \frac{i}{\sqrt{6}}(|2\rangle_7|3\rangle_7 - |3\rangle_7|2\rangle_7), \\ |7\rangle_{14} &= -\frac{1}{\sqrt{3}}(|1\rangle_7|4\rangle_7 - |4\rangle_7|1\rangle_7) - \frac{1}{\sqrt{6}}(|5\rangle_7|6\rangle_7 - |6\rangle_7|5\rangle_7), \\ |8\rangle_{14} &= -\frac{1}{\sqrt{2}}(|4\rangle_7|6\rangle_7 - |6\rangle_7|4\rangle_7), \\ |9\rangle_{14} &= \frac{1}{2\sqrt{3}}(|2\rangle_7|5\rangle_7 - |5\rangle_7|2\rangle_7) - \frac{1}{\sqrt{3}}(|3\rangle_7|6\rangle_7 - |6\rangle_7|3\rangle_7) \\ &\quad + \frac{1}{2\sqrt{3}}(|4\rangle_7|7\rangle_7 - |7\rangle_7|4\rangle_7), \\ |10\rangle_{14} &= -\frac{1}{\sqrt{2}}(|2\rangle_7|6\rangle_7 - |6\rangle_7|2\rangle_7), \\ |11\rangle_{14} &= -\frac{1}{\sqrt{3}}(|1\rangle_7|6\rangle_7 - |6\rangle_7|1\rangle_7) - \frac{i}{\sqrt{6}}(|2\rangle_7|4\rangle_7 - |4\rangle_7|2\rangle_7), \\ |12\rangle_{14} &= -\frac{1}{\sqrt{2}}(|4\rangle_7|5\rangle_7 - |5\rangle_7|4\rangle_7), \\ |13\rangle_{14} &= -\frac{1}{\sqrt{2}}(|3\rangle_7|5\rangle_7 - |5\rangle_7|3\rangle_7), \\ |14\rangle_{14} &= -\frac{1}{\sqrt{3}}(|1\rangle_7|5\rangle_7 - |5\rangle_7|1\rangle_7) + \frac{i}{\sqrt{6}}(|3\rangle_7|4\rangle_7 - |4\rangle_7|3\rangle_7). \end{aligned}$$

The states are shown in the weight diagram of the representation **14** in Figure 2. We note that the basis vectors are indeed odd under exchange of the factors. One can straightforwardly check that indeed ${}_{14}\langle a|b\rangle_{14} = \delta_{ab}$. The action of the generators Λ on the basis states $|a\rangle_{14}$ is obtained from their action on the respective factors. For example,

$$\Lambda^1 |10\rangle_{14} = (\Lambda_1^1 + \Lambda_2^1) \left[-\frac{1}{\sqrt{2}} (|2\rangle_7 |6\rangle_7 - |6\rangle_7 |2\rangle_7) \right] = \frac{1}{2} |3\rangle_{14} + \frac{\sqrt{3}}{2} |9\rangle_{14}. \quad (\text{C.3})$$

In this basis the generators Λ^3 and Λ^8 are indeed diagonal. Using this the action of the generators on the matrix $\mathcal{M}_j$ required in the calculation of the static correlation functions and string order parameter can be worked out.

Finally, to obtain the explicit expressions of M^a we project the matrix $\mathcal{M}_j^{7\otimes 7}$ in (C.2) onto the basis $\{|a\rangle_{14} : a = 1, \dots, 14\}$ and dropping the subscript **14**. As a result we have

$$\mathcal{M}_j^{14} = \begin{pmatrix} 0 & -\frac{1}{\sqrt{3}} |1\rangle & \frac{1}{\sqrt{3}} |2\rangle & -\frac{1}{\sqrt{3}} |7\rangle & -\frac{1}{\sqrt{3}} |14\rangle & -\frac{1}{\sqrt{3}} |11\rangle & -\frac{1}{\sqrt{3}} |6\rangle \\ \frac{i}{\sqrt{3}} |14\rangle & -\frac{1}{2} |3\rangle - \frac{1}{2\sqrt{3}} |9\rangle & \frac{1}{\sqrt{2}} |13\rangle & \frac{1}{\sqrt{2}} |12\rangle & 0 & -\frac{1}{\sqrt{6}} |7\rangle & -\frac{1}{\sqrt{6}} |2\rangle \\ \frac{i}{\sqrt{3}} |11\rangle & \frac{1}{\sqrt{2}} |10\rangle & \frac{1}{\sqrt{3}} |9\rangle & \frac{1}{\sqrt{2}} |8\rangle & \frac{1}{\sqrt{6}} |7\rangle & 0 & -\frac{1}{\sqrt{6}} |1\rangle \\ \frac{i}{\sqrt{3}} |6\rangle & \frac{1}{\sqrt{2}} |5\rangle & \frac{1}{\sqrt{3}} |4\rangle & \frac{1}{2} |3\rangle - \frac{1}{2\sqrt{3}} |9\rangle & \frac{1}{\sqrt{6}} |2\rangle & \frac{1}{\sqrt{6}} |1\rangle & 0 \\ \frac{i}{\sqrt{3}} |1\rangle & 0 & -\frac{1}{\sqrt{6}} |6\rangle & \frac{i}{\sqrt{6}} |11\rangle & \frac{1}{2} |3\rangle + \frac{1}{2\sqrt{3}} |9\rangle & -\frac{1}{\sqrt{2}} |10\rangle & -\frac{1}{\sqrt{2}} |5\rangle \\ -\frac{i}{\sqrt{3}} |2\rangle & \frac{i}{\sqrt{6}} |6\rangle & 0 & -\frac{1}{\sqrt{6}} |14\rangle & -\frac{1}{\sqrt{2}} |13\rangle & -\frac{1}{\sqrt{3}} |9\rangle & -\frac{1}{\sqrt{2}} |4\rangle \\ \frac{i}{\sqrt{3}} |7\rangle & -\frac{i}{\sqrt{6}} |11\rangle & \frac{1}{\sqrt{6}} |14\rangle & 0 & -\frac{1}{\sqrt{2}} |12\rangle & -\frac{1}{\sqrt{2}} |8\rangle & -\frac{1}{2} |3\rangle + \frac{1}{2\sqrt{3}} |9\rangle \end{pmatrix}. \quad (\text{C.4})$$

This matrix can be rewritten in terms of the basis states $|\psi_a\rangle$ in (3) as

$$\mathcal{M}_j^{14} = \frac{1}{\sqrt{2}} \sum_{a=1}^{14} \check{\Lambda}^a |\psi_a\rangle_j, \quad (\text{C.5})$$

where $\check{\Lambda}^a = V \Lambda^a V^\dagger$ with $V = \text{diag}(-i, 1, 1, 1, 1, 1, 1)$ and the index j on the right-hand side was added to indicate that $|\psi_a\rangle$ is a local state at site j . Using this we obtain

$$\begin{aligned} |\Psi_{\text{VBS}}\rangle &= \text{tr}_7 [\mathcal{M}_1 \mathcal{M}_2 \cdots \mathcal{M}_N] \\ &= \frac{1}{2^{N/2}} \sum_{a_1, a_2, \dots, a_N=1}^{14} \text{tr}_7 [\check{\Lambda}^{a_1} \check{\Lambda}^{a_2} \cdots \check{\Lambda}^{a_N}] |\psi_{a_1}\rangle_1 |\psi_{a_2}\rangle_2 \cdots |\psi_{a_N}\rangle_N, \end{aligned} \quad (\text{C.6})$$

where the trace is taken over the 7-dimensional auxiliary space. In the trace the matrices V and $V^\dagger$ cancel out in pairs, and we are left with

$$|\Psi_{\text{VBS}}\rangle = \frac{1}{2^{N/2}} \sum_{a_1, a_2, \dots, a_N=1}^{14} \text{tr}_7 [\Lambda^{a_1} \Lambda^{a_2} \cdots \Lambda^{a_N}] |\psi_{a_1}\rangle_1 |\psi_{a_2}\rangle_2 \cdots |\psi_{a_N}\rangle_N, \quad (\text{C.7})$$

which is the desired (2).

D Family of parent Hamiltonians

Using the projectors onto the representations **77** and **77'** in (4) one can construct a family of parent Hamiltonians for the VBS state (2). We start from the projectors

$$\begin{aligned} P_{77} &= -\frac{3}{10240} (\Lambda_1 + \Lambda_2)^2 \left[(\Lambda_1 + \Lambda_2)^2 - 8 \right] \left[(\Lambda_1 + \Lambda_2)^2 - \frac{28}{3} \right] \left[(\Lambda_1 + \Lambda_2)^2 - 20 \right], \\ P_{77'} &= \frac{1}{10240} (\Lambda_1 + \Lambda_2)^2 \left[(\Lambda_1 + \Lambda_2)^2 - 8 \right] \left[(\Lambda_1 + \Lambda_2)^2 - \frac{28}{3} \right] \left[(\Lambda_1 + \Lambda_2)^2 - 16 \right], \end{aligned}$$

where $\Lambda_{1,2}$ denote the generators on the respective factors in $\mathbf{14} \otimes \mathbf{14}$. Using these an operator which annihilates (1) and is strictly positive on the complement of (1) in (4) is given by

$$\tilde{H}_{j,j+1} = a P_{77} + b P_{77'} = \sum_{n=0}^4 \mu_n (\Lambda_j \cdot \Lambda_{j+1})^n, \quad (\text{D.1})$$

where $a, b > 0$ and

$$\mu_0 = a, \quad \mu_1 = \frac{7a}{40} + \frac{b}{6}, \quad \mu_2 = -\frac{31a}{160} + \frac{9b}{80}, \quad \mu_3 = -\frac{a}{16} + \frac{23b}{960}, \quad \mu_4 = -\frac{3a}{640} + \frac{b}{640}. \quad (\text{D.2})$$

The parameters a and b can be tuned such that one of the coefficients μ_n , $n = 2, 3, 4$, vanishes. For example, choosing $b = 3a$ implies $\mu_4 = 0$ [with $a = 20/27$ giving (5)]; in this sense the Hamiltonian (6) is the simplest parent Hamiltonian for the VBS state.

References

- [1] I. Affleck, T. Kennedy, E. H. Lieb and H. Tasaki, *Rigorous results on valence-bond ground states in antiferromagnets*, Phys. Rev. Lett. **59**, 799 (1987), doi:[10.1103/PhysRevLett.59.799](https://doi.org/10.1103/PhysRevLett.59.799).
- [2] I. Affleck, T. Kennedy, E. H. Lieb and H. Tasaki, *Valence bond ground states in isotropic quantum antiferromagnets*, Commun. Math. Phys. **115**, 477 (1988), doi:[10.1007/BF01218021](https://doi.org/10.1007/BF01218021).
- [3] U. Schollwöck, T. Jolicœur and T. Garel, *Onset of incommensurability at the valence-bond-solid point in the $S = 1$ quantum spin chain*, Phys. Rev. B **53**, 3304 (1996), doi:[10.1103/PhysRevB.53.3304](https://doi.org/10.1103/PhysRevB.53.3304) [preprint doi:[10.48550/arXiv.cond-mat/9505086](https://doi.org/10.48550/arXiv.cond-mat/9505086)].
- [4] A. Läuchli, G. Schmid and S. Trebst, *Spin nematics correlations in bilinear-biquadratic $S = 1$ spin chains*, Phys. Rev. B **74**, 144426 (2006), doi:[10.1103/PhysRevB.74.144426](https://doi.org/10.1103/PhysRevB.74.144426) [preprint doi:[10.48550/arXiv.cond-mat/0311082](https://doi.org/10.48550/arXiv.cond-mat/0311082)].
- [5] F. D. M. Haldane, *Continuum dynamics of the 1-D Heisenberg antiferromagnet: Identification with the $O(3)$ nonlinear sigma model*, Phys. Lett. A **93**, 464 (1983), doi:[10.1016/0375-9601\(83\)90631-X](https://doi.org/10.1016/0375-9601(83)90631-X).
- [6] F. D. M. Haldane, *Nonlinear field theory of large-spin Heisenberg antiferromagnets: Semi-classically quantized solitons of the one-dimensional easy-axis Néel state*, Phys. Rev. Lett. **50**, 1153 (1983), doi:[10.1103/PhysRevLett.50.1153](https://doi.org/10.1103/PhysRevLett.50.1153).
- [7] I. Affleck, *Quantum spin chains and the Haldane gap*, J. Phys.: Condens. Matter **1**, 3047 (1989), doi:[10.1088/0953-8984/1/19/001](https://doi.org/10.1088/0953-8984/1/19/001).
- [8] M. Oshikawa, *Hidden $Z_2 \times Z_2$ symmetry in quantum spin chains with arbitrary integer spin*, J. Phys.: Condens. Matter **4**, 7469 (1992), doi:[10.1088/0953-8984/4/36/019](https://doi.org/10.1088/0953-8984/4/36/019).
- [9] D. Maekawa and H. Tasaki, *The asymmetric valence-bond-solid states in quantum spin chains: The difference between odd and even spins*, J. Math. Phys. **64**, 031901 (2023), doi:[10.1063/5.0123743](https://doi.org/10.1063/5.0123743) [preprint doi:[10.48550/arXiv.2205.00653](https://doi.org/10.48550/arXiv.2205.00653)].
- [10] T. Kennedy, *Exact diagonalisations of open spin-1 chains*, J. Phys.: Condens. Matter **2**, 5737 (1990), doi:[10.1088/0953-8984/2/26/010](https://doi.org/10.1088/0953-8984/2/26/010).

- [11] M. Hagiwara, K. Katsumata, I. Affleck, B. I. Halperin and J. P. Renard, *Observation of $S = 1/2$ degrees of freedom in an $S = 1$ linear-chain Heisenberg antiferromagnet*, Phys. Rev. Lett. **65**, 3181 (1990), doi:[10.1103/PhysRevLett.65.3181](https://doi.org/10.1103/PhysRevLett.65.3181).
- [12] F. Pollmann, E. Berg, A. M. Turner and M. Oshikawa, *Symmetry protection of topological phases in one-dimensional quantum spin systems*, Phys. Rev. B **85**, 075125 (2012), doi:[10.1103/PhysRevB.85.075125](https://doi.org/10.1103/PhysRevB.85.075125) [preprint doi:[10.48550/arXiv.0909.4059](https://doi.org/10.48550/arXiv.0909.4059)].
- [13] S. Östlund and S. Rommer, *Thermodynamic limit of density matrix renormalization*, Phys. Rev. Lett. **75**, 3537 (1995), doi:[10.1103/PhysRevLett.75.3537](https://doi.org/10.1103/PhysRevLett.75.3537) [preprint doi:[10.48550/arXiv.cond-mat/9503107](https://doi.org/10.48550/arXiv.cond-mat/9503107)].
- [14] U. Schollwöck, *The density-matrix renormalization group in the age of matrix product states*, Ann. Phys. **326**, 96 (2011), doi:[10.1016/j.aop.2010.09.012](https://doi.org/10.1016/j.aop.2010.09.012) [preprint doi:[10.48550/arXiv.1008.3477](https://doi.org/10.48550/arXiv.1008.3477)].
- [15] H. Tasaki, *Physics and mathematics of quantum many-body systems*, Springer, Cham, Switzerland, ISBN 9783030412647 (2020), doi:[10.1007/978-3-030-41265-4](https://doi.org/10.1007/978-3-030-41265-4).
- [16] A. Klümper, A. Schadschneider and J. Zittartz, *Equivalence and solution of anisotropic spin-1 models and generalized t-J fermion models in one dimension*, J. Phys. A: Math. Gen. **24**, L955 (1991), doi:[10.1088/0305-4470/24/16/012](https://doi.org/10.1088/0305-4470/24/16/012).
- [17] A. Klümper, A. Schadschneider and J. Zittartz, *Groundstate properties of a generalized VBS-model*, Z. Phys. B Condens. Matter **87**, 281 (1992), doi:[10.1007/BF01309281](https://doi.org/10.1007/BF01309281).
- [18] M. T. Batchelor and C. M. Yung, *q-deformations of quantum spin chains with exact valence-bond ground states*, Int. J. Mod. Phys. B **08**, 3645 (1994), doi:[10.1142/S021797929400155X](https://doi.org/10.1142/S021797929400155X) [preprint doi:[10.48550/arXiv.cond-mat/9403080](https://doi.org/10.48550/arXiv.cond-mat/9403080)].
- [19] M. Fannes, B. Nachtergaele and R. F. Werner, *Quantum spin chains with quantum group symmetry*, Commun. Math. Phys. **174**, 477 (1996), doi:[10.1007/BF02101525](https://doi.org/10.1007/BF02101525) [preprint doi:[10.48550/arXiv.cond-mat/9504002](https://doi.org/10.48550/arXiv.cond-mat/9504002)].
- [20] T. Quella, *Symmetry-protected topological phases beyond groups: The q-deformed Affleck-Kennedy-Lieb-Tasaki model*, Phys. Rev. B **102**, 081120 (2020), doi:[10.1103/PhysRevB.102.081120](https://doi.org/10.1103/PhysRevB.102.081120) [preprint doi:[10.48550/arXiv.2005.09072](https://doi.org/10.48550/arXiv.2005.09072)].
- [21] T. Franke and T. Quella, *Duality and hidden symmetry breaking in the q-deformed Affleck-Kennedy-Lieb-Tasaki model*, SciPost Phys. Core **7**, 078 (2024), doi:[10.21468/SciPostPhysCore.7.4.078](https://doi.org/10.21468/SciPostPhysCore.7.4.078) [preprint doi:[10.48550/arXiv.2403.09287](https://doi.org/10.48550/arXiv.2403.09287)].
- [22] D. P. Arovas, K. Hasebe, X.-L. Qi and S.-C. Zhang, *Supersymmetric valence bond solid states*, Phys. Rev. B **79**, 224404 (2009), doi:[10.1103/PhysRevB.79.224404](https://doi.org/10.1103/PhysRevB.79.224404) [preprint doi:[10.48550/arXiv.0901.1498](https://doi.org/10.48550/arXiv.0901.1498)].
- [23] K. Hasebe and K. Totsuka, *Hidden order and dynamics in supersymmetric valence-bond solid states: Super-matrix product state formalism*, Phys. Rev. B **84**, 104426 (2011), doi:[10.1103/PhysRevB.84.104426](https://doi.org/10.1103/PhysRevB.84.104426) [preprint doi:[10.48550/arXiv.1105.3529](https://doi.org/10.48550/arXiv.1105.3529)].
- [24] I. Affleck, D. P. Arovas, J. B. Marston and D. A. Rabson, *$SU(2n)$ quantum antiferromagnets with exact C-breaking ground states*, Nucl. Phys. B **366**, 467 (1991), doi:[10.1016/0550-3213\(91\)90027-U](https://doi.org/10.1016/0550-3213(91)90027-U).

- [25] M. Greiter, S. Rachel and D. Schuricht, *Exact results for $SU(3)$ spin chains: Trimer states, valence bond solids, and their parent Hamiltonians*, Phys. Rev. B **75**, 060401 (2007), doi:[10.1103/PhysRevB.75.060401](https://doi.org/10.1103/PhysRevB.75.060401) [preprint doi:[10.48550/arXiv.cond-mat/0701354](https://doi.org/10.48550/arXiv.cond-mat/0701354)].
- [26] M. Greiter and S. Rachel, *Valence bond solids for $SU(n)$ spin chains: Exact models, spinon confinement, and the Haldane gap*, Phys. Rev. B **75**, 184441 (2007), doi:[10.1103/PhysRevB.75.184441](https://doi.org/10.1103/PhysRevB.75.184441) [preprint doi:[10.48550/arXiv.cond-mat/0702443](https://doi.org/10.48550/arXiv.cond-mat/0702443)].
- [27] D. P. Arovas, *Simplex solid states of $SU(N)$ quantum antiferromagnets*, Phys. Rev. B **77**, 104404 (2008), doi:[10.1103/PhysRevB.77.104404](https://doi.org/10.1103/PhysRevB.77.104404) [preprint doi:[10.48550/arXiv.0711.3921](https://doi.org/10.48550/arXiv.0711.3921)].
- [28] H. Katsura, T. Hirano and V. E. Korepin, *Entanglement in an $SU(n)$ valence-bond-solid state*, J. Phys. A: Math. Theor. **41**, 135304 (2008), doi:[10.1088/1751-8113/41/13/135304](https://doi.org/10.1088/1751-8113/41/13/135304) [preprint doi:[10.48550/arXiv.0711.3882](https://doi.org/10.48550/arXiv.0711.3882)].
- [29] S. Rachel, D. Schuricht, B. Scharfenberger, R. Thomale and M. Greiter, *Spontaneous parity violation in a quantum spin chain*, J. Phys.: Conf. Ser. **200**, 022049 (2010), doi:[10.1088/1742-6596/200/2/022049](https://doi.org/10.1088/1742-6596/200/2/022049) [preprint doi:[10.48550/arXiv.0905.4895](https://doi.org/10.48550/arXiv.0905.4895)].
- [30] S. Gozel, D. Poilblanc, I. Affleck and F. Mila, *Novel families of $SU(N)$ AKLT states with arbitrary self-conjugate edge states*, Nucl. Phys. B **945**, 114663 (2019), doi:[10.1016/j.nuclphysb.2019.114663](https://doi.org/10.1016/j.nuclphysb.2019.114663) [preprint doi:[10.48550/arXiv.1904.01822](https://doi.org/10.48550/arXiv.1904.01822)].
- [31] H.-H. Tu, G.-M. Zhang and T. Xiang, *Class of exactly solvable $SO(n)$ symmetric spin chains with matrix product ground states*, Phys. Rev. B **78**, 094404 (2008), doi:[10.1103/PhysRevB.78.094404](https://doi.org/10.1103/PhysRevB.78.094404) [preprint doi:[10.48550/arXiv.0806.1839](https://doi.org/10.48550/arXiv.0806.1839)].
- [32] H.-H. Tu, G.-M. Zhang and T. Xiang, *String order and hidden topological symmetry in the $SO(2n+1)$ symmetric matrix product states*, J. Phys. A: Math. Theor. **41**, 415201 (2008), doi:[10.1088/1751-8113/41/41/415201](https://doi.org/10.1088/1751-8113/41/41/415201) [preprint doi:[10.48550/arXiv.0804.1685](https://doi.org/10.48550/arXiv.0804.1685)].
- [33] D. Schuricht and S. Rachel, *Valence bond solid states with symplectic symmetry*, Phys. Rev. B **78**, 014430 (2008), doi:[10.1103/PhysRevB.78.014430](https://doi.org/10.1103/PhysRevB.78.014430) [preprint doi:[10.48550/arXiv.0805.3918](https://doi.org/10.48550/arXiv.0805.3918)].
- [34] W. Fulton and J. Harris, *Representation theory: A first course*, Springer, New York, USA, ISBN 9780387975276 (1991), doi:[10.1007/978-1-4612-0979-9](https://doi.org/10.1007/978-1-4612-0979-9).
- [35] B. C. Hall, *Lie groups, Lie algebras, and representations*, Springer, Cham, Switzerland, ISBN 9783319134666 (2015), doi:[10.1007/978-3-319-13467-3](https://doi.org/10.1007/978-3-319-13467-3).
- [36] D. P. Arovas and A. Auerbach, *Extended Heisenberg models of antiferromagnetism: Analogies to the fractional quantum Hall effect*, Phys. Rev. Lett. **60**, 531 (1988), doi:[10.1103/PhysRevLett.60.531](https://doi.org/10.1103/PhysRevLett.60.531).
- [37] A. Auerbach, *Interacting electrons and quantum magnetism*, Springer, New York, USA, ISBN 9780387942865 (1994), doi:[10.1007/978-1-4612-0869-3](https://doi.org/10.1007/978-1-4612-0869-3).
- [38] H. Katsura, N. Kawashima, A. N. Kirillov, V. E. Korepin and S. Tanaka, *Entanglement in valence-bond-solid states on symmetric graphs*, J. Phys. A: Math. Theor. **43**, 255303 (2010), doi:[10.1088/1751-8113/43/25/255303](https://doi.org/10.1088/1751-8113/43/25/255303) [preprint doi:[10.48550/arXiv.1003.2007](https://doi.org/10.48550/arXiv.1003.2007)].
- [39] Z.-Q. Gao and C. Wu, *Construction of G_2 symmetry in a Hubbard-type model*, (arXiv preprint) doi:[10.48550/arXiv.2010.14126](https://doi.org/10.48550/arXiv.2010.14126).

- [40] D. Perez-Garcia, F. Verstraete, M. M. Wolf and J. I. Cirac, *Matrix product state representations*, Quantum Inf. Comput. **7**, 401 (2007), doi:[10.26421/QIC7.5-6-1](https://doi.org/10.26421/QIC7.5-6-1) [preprint doi:[10.48550/arXiv.quant-ph/0608197](https://doi.org/10.48550/arXiv.quant-ph/0608197)].
- [41] M. den Nijs and K. Rommelse, *Preroughening transitions in crystal surfaces and valence-bond phases in quantum spin chains*, Phys. Rev. B **40**, 4709 (1989), doi:[10.1103/PhysRevB.40.4709](https://doi.org/10.1103/PhysRevB.40.4709).
- [42] S. M. Girvin and D. P. Arovas, *Hidden topological order in integer quantum spin chains*, Phys. Scr. **T27**, 156 (1989), doi:[10.1088/0031-8949/1989/T27/027](https://doi.org/10.1088/0031-8949/1989/T27/027).
- [43] T. Kennedy and H. Tasaki, *Hidden $Z_2 \times Z_2$ symmetry breaking in Haldane-gap antiferromagnets*, Phys. Rev. B **45**, 304 (1992), doi:[10.1103/PhysRevB.45.304](https://doi.org/10.1103/PhysRevB.45.304).
- [44] T. Kennedy and H. Tasaki, *Hidden symmetry breaking and the Haldane phase in $S = 1$ quantum spin chains*, Commun. Math. Phys. **147**, 431 (1992), doi:[10.1007/BF02097239](https://doi.org/10.1007/BF02097239).
- [45] K. Duivenvoorden and T. Quella, *Topological phases of spin chains*, Phys. Rev. B **87**, 125145 (2013), doi:[10.1103/PhysRevB.87.125145](https://doi.org/10.1103/PhysRevB.87.125145) [preprint doi:[10.48550/arXiv.1206.2462](https://doi.org/10.48550/arXiv.1206.2462)].
- [46] C. McRae, *Exploring triality explicitly: Convenient bases for $SO(8)$, $Spin(1,7)$, and G_2* , (arXiv preprint) doi:[10.48550/arXiv.2502.14016](https://doi.org/10.48550/arXiv.2502.14016).