

# Intermittency in $pp$ collisions at $\sqrt{s} = 0.9, 7$ and $8$ TeV from the CMS collaboration

Z. Ong\*, P. Agarwal, H. W. Ang, A. H. Chan and C. H. Oh

Department of Physics, National University of Singapore

\* ongzungjin@u.nus.edu

50th International Symposium on Multiparticle Dynamics



(ISMD2021)

12-16 July 2021

doi:[10.21468/SciPostPhysProc.10](https://doi.org/10.21468/SciPostPhysProc.10)

## Abstract

The intermittency-type fluctuations as outlined by Bialas and Peschanski in the 1980s is analysed in  $pp$  collisions at  $\sqrt{s} = 0.9, 7$  and  $8$  TeV from the CMS collaboration at CERN. Our preliminary analysis shows that the intermittency exponents in the bin-averaged scaled factorial moments decrease in magnitude with increasing collision energy at the TeV scale, which suggests that the cascading nature of multiparticle production described by the  $\alpha$ -model is weakening. We outline possible areas planned for future studies.



Copyright Z. Ong *et al.*

This work is licensed under the Creative Commons

[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Published by the SciPost Foundation.

Received 26-10-2021

Accepted 25-03-2022

Published 11-08-2022



doi:[10.21468/SciPostPhysProc.10.032](https://doi.org/10.21468/SciPostPhysProc.10.032)

## 1 Introduction

Intermittency in multiparticle production, where fluctuations in particle multiplicity can be observed at all scales, can be quantified via the bin-averaged scaled factorial moments [1, 2]:

$$F_q(\delta\eta) = M^{q-1} \sum_{m=1}^M \frac{\langle n_m(n_m - 1)\dots(n_m - q + 1) \rangle}{\langle N \rangle^q}, \quad (1)$$

where  $F_q$  is the  $q^{\text{th}}$  bin-averaged moment,  $M$  is the number of bins that the (pseudo)rapidity space is divided into (each with size  $\delta\eta$ ),  $n_m$  is the multiplicity in bin  $m$ ,  $N$  is the total event multiplicity and  $\langle \dots \rangle$  represents an average over events. Equation (1) is sometimes known as the “horizontal moments” in intermittency literature.

In Bialas and Peschanski’s formulation [1, 2], a system is said to be intermittent if a power-law relation  $F_q \sim (\delta\eta)^{-\phi_q}$  exists. We expect an intermittent system to produce a straight line in a  $\ln F_q$  vs.  $\ln(1/\delta\eta)$  plot towards  $\delta\eta \rightarrow 0$ , with some positive gradient  $\phi_q$  called the “slope parameter” or “intermittency exponent”. The magnitude of  $\phi_q$  reflects the strength of intermittency of the system.

## 2 Motivation and Objectives

With the LHC at CERN ushering in a new era of TeV-scale high energy physics, it is worthwhile to reinvestigate this phenomenon in a new energy regime. In our previous work [3], we found that intermittent phenomenon exists in  $pp$  collisions at  $\sqrt{s} = 7$  TeV. That motivates our current investigation, where the analysis is extended to include the other available energies in Run 1 from the CERN Open Data Portal – 0.9 and 8 TeV from the CMS collaboration.

Hence we aim to: (1) compute the bin-averaged scaled factorial moments and produce log-log plots of  $F_q$  vs.  $(1/\delta\eta)$  for  $pp$  collisions at  $\sqrt{s} = 0.9, 7$  and  $8$  TeV data from the CERN Open Data Portal, and (2) obtain the intermittency exponents for  $q = 2, 3, 4$  and  $5$  from the data and make observations about any trends that may appear.

## 3 Data Processing

MinimumBias datasets from the CERN Open Data Portal are used for this analysis – 0.9 TeV data from the Commissioning run of 2010 [4], 7 TeV data from RunA of 2010 [5] and 8 TeV data from RunB of 2012 [6]. About one million events for each energy were included in this analysis.

### Event Selection

Vertices are required to have number of degrees of freedom greater than 4, and be within 15 cm of the beamspot in the  $z$ -direction. If more than one primary vertex is present, the one with the highest number of associated tracks is chosen to be the primary vertex.

To remove secondaries from background and pileup, tracks are required to have impact parameter significances  $d_0/\sigma_{d_0}$  and  $d_z/\sigma_{d_z}$  to be smaller than 5. The relative uncertainty in the momentum measurement  $\sigma_{p_T}/p_T$  is required to be less than 10%. To ensure good reconstruction efficiency, only highPurity tracks with  $p_T > 0.5$  GeV/c are selected.

## 4 Results

Figure 1 shows the log-log plots in  $F_q$  vs  $(1/\delta\eta)$  for  $\sqrt{s} = 0.9, 7$  and  $8$  TeV with error bars plotted (only statistical; most are too small to be seen). Table 1 summarises the intermittency exponents, which are obtained via linear regression on the rightmost 10 data points of each set of data<sup>1</sup>.

Table 1: Summary of intermittency exponents

$q$	0.9 TeV	7 TeV	8 TeV
2	$0.889 \pm 0.003$	$0.787 \pm 0.002$	$0.560 \pm 0.003$
3	$0.89 \pm 0.02$	$0.795 \pm 0.007$	$0.612 \pm 0.004$
4	$0.87 \pm 0.08$	$0.82 \pm 0.03$	$0.72 \pm 0.01$
5	$0.9 \pm 0.3$	$0.87 \pm 0.09$	$0.90 \pm 0.03$

<sup>1</sup>The results in this section have been updated from what was presented at ISMD2021.




Figure 1: Horizontal moments for  $\sqrt{s} = 0.9, 7$  and  $8$  TeV

## 5 Conclusion and Discussion

From the non-zero values in Table 1, we can immediately conclude that intermittency still persists in  $pp$  collisions up to  $\sqrt{s} = 8$  TeV.

With the exception of  $q = 5$ , the intermittency exponents generally appear to show a decreasing trend as  $\sqrt{s}$  increases from 0.9 TeV to 8 TeV. For  $q = 5$ , the intermittency exponents appear to be equal within errors. These findings are made with data treatment kept uniform across all three energies.

### Scaling, Universality and Branching

The results clearly show that the scaling relation  $F_q \sim (\delta\eta)^{-\phi_q}$  still holds at the energies being investigated. However, universality, as noted by Bialas [7] where the intermittency exponents are the same in all experiments, does not seem to hold. More analysis needs to be done before further conclusions can be made about this.

The concept of intermittency is closely related to parton branching processes in multiparticle production [7]. The decreasing intermittency exponents lead us to surmise that as the centre-of-mass energy increases into the TeV scale, other competing processes in multiparticle production might become increasingly important. Still, parton branching ideas remain valid and are a present area of study (see for example, [8]).

## 6 Forthcoming Research

This analysis can be further extended in the following areas:

1. **Unfolding (corrections):** In this analysis, the particle multiplicities are derived by

counting charged tracks from the tracker in CMS. The multiplicity distributions need to be unfolded back to the charged hadron level, so that we can study the intermittency effects from a QCD perspective. Unfolding will also correct for detector-related inefficiencies.

2. **Higher energies:** With the expected release of Run 2 data to the public, we can look forward to extending this study up to  $\sqrt{s} = 13$  TeV, which will shed more light into intermittent behaviour at the TeV scale.

## Acknowledgements

The authors would like to thank the National University of Singapore, and the support and helpful discussions with colleagues. We would also like to thank the CMS collaboration at CERN for the well-organised release of collision data to the public on the CERN Open Data Portal. This work is also supported by the NUS Research Scholarship.

## References

- [1] A. Bialas and R. Peschanski, *Moments of rapidity distributions as a measure of short-range fluctuations in high-energy collisions*, Nucl. Phys. B **273**, 703 (1986), doi:[10.1016/0550-3213\(86\)90386-X](https://doi.org/10.1016/0550-3213(86)90386-X).
- [2] A. Bialas and R. Peschanski, *Intermittency in multiparticle production at high energy*, Nucl. Phys. B **308**, 857 (1988), doi:[10.1016/0550-3213\(88\)90131-9](https://doi.org/10.1016/0550-3213(88)90131-9).
- [3] Z. Ong, E. Yuen, H. W. Ang, A. H. Chan and C. H. Oh, *Intermittency in pseudorapidity space of pp collisions at = 7 TeV*, EPJ Web Conf. **206**, 09004 (2019), doi:[10.1051/epjconf/201920609004](https://doi.org/10.1051/epjconf/201920609004).
- [4] CMS collaboration, *MinimumBias primary dataset in RECO format from the 0.9 TeV Commissioning run of 2010* ([/MinimumBias/Commissioning10-07JunReReco\\_900GeV/RECO](https://opendata.cern.ch/record/260000)), CERN Open Data Portal (2019), doi:[10.7483/OPENDATA.CMS.1R58.OMBD](https://doi.org/10.7483/OPENDATA.CMS.1R58.OMBD).
- [5] CMS collaboration, *MinimumBias primary dataset in AOD format from RunA of 2010* ([/MinimumBias/Run2010A-Apr21ReReco-v1/AOD](https://opendata.cern.ch/record/260000)), CERN Open Data Portal (2019), doi:[10.7483/OPENDATA.CMS.6B3H.TR6Z](https://doi.org/10.7483/OPENDATA.CMS.6B3H.TR6Z).
- [6] CMS collaboration, *MinimumBias primary dataset in AOD format from Run of 2012* ([/MinimumBias/Run2012B-22Jan2013-v1/AOD](https://opendata.cern.ch/record/260000)), CERN Open Data Portal (2017), doi:[10.7483/OPENDATA.CMS.HU6U.DRLD](https://doi.org/10.7483/OPENDATA.CMS.HU6U.DRLD).
- [7] A. Bialas, *Intermittency in high energy collisions*, Nucl. Phys. A **545**, 285 (1992), doi:[10.1016/0375-9474\(92\)90469-Z](https://doi.org/10.1016/0375-9474(92)90469-Z).
- [8] Z. Ong, P. Agarwal, H. W. Ang, A. H. Chan and C. H. Oh, *Numerical solutions to Giovannini's parton branching equation up to TeV energies at the LHC*, Mod. Phys. Lett. A **35**, 2050325 (2020), doi:[10.1142/S0217732320503253](https://doi.org/10.1142/S0217732320503253).