

# Ruling out QCD phase transition as a PBH origin of LIGO/Virgo events

Joaquim Iguaz Juan\*

LAPTh, CNRS, Université Savoie Mont Blanc, F-74940 Annecy, France

\* [iguaz@lapth.cnrs.fr](mailto:iguaz@lapth.cnrs.fr)



14th International Conference on Identification of Dark Matter  
Vienna, Austria, 18-22 July 2022  
doi:[10.21468/SciPostPhysProc.12](https://doi.org/10.21468/SciPostPhysProc.12)

## Abstract

The best-motivated scenario for a sizable primordial black hole (PBH) contribution to the LIGO/Virgo binary black hole mergers invokes the QCD phase transition, which naturally enhances the probability to form PBH with masses of stellar scale. We reconsider the expected mass function assuming a CMB-like primordial spectrum and associated not only to the QCD phase transition proper, but also the  $e^+e^-$  annihilation process, and analyze the constraints on this scenario from a number of observations. We find that the scenario is not viable, unless an ad hoc mass evolution for the PBH mass function and a cutoff in power-spectrum very close to the QCD scale are introduced by hand.



Copyright J. Iguaz Juan.

This work is licensed under the Creative Commons

[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Published by the SciPost Foundation.

Received 21-11-2022

Accepted 28-04-2023

Published 05-07-2023

doi:[10.21468/SciPostPhysProc.12.052](https://doi.org/10.21468/SciPostPhysProc.12.052)



Check for updates

## 1 Introduction

The detection of heavy black hole merger events, see for instance [1], provides a strong motivation for primordial black holes (PBHs) as the candidates responsible for the bulk of these events (see e.g. [2–5]). PBHs are theoretical objects that were firstly discussed in the 60s and 70s by Zeldovich & Novikov [6] and Hawking [7] and are typically assumed to be formed in the early universe from the collapse of large overdensities. PBHs are a well-studied non-particle dark matter (DM) candidate. Indeed, while there is no shortage of DM particle candidates in extensions of the SM, there is no guarantee nor observational indication that DM is made of microscopic fundamental particles. For a review of PBHs as DM and current bounds see [8]. In addition, PBHs also are very interesting objects in the context of supermassive black holes (SMBHs), LIGO/Virgo coalescing events, inflation etc.

In this work, we will focus on PBHs in the mass range  $M_{\text{PBH}} \sim 10^{-2}M_{\odot} - 10^9M_{\odot}$ . This mass window is interesting at least for a couple of reasons. On the heavy end, PBHs whose mass is above  $M_{\text{PBH}} \sim 10^6M_{\odot}$  provide a possible explanation for the most massive BHs observed in

the universe and, in particular, those at high redshift, which are difficult to explain through standard astrophysical processes otherwise. On the light end, PBHs falling within the stellar mass range, namely  $M_{\text{PBH}} \sim 1M_{\odot} - 10^2M_{\odot}$ , are particularly interesting in light of LIGO/Virgo merger events observations. Even if the abundance of PBHs in the the stellar mass range is pretty constrained, some authors explored the possibility of PBHs constituting a large fraction of the events detected by LIGO/Virgo. In particular, as reported in [9], PBHs with masses  $M_{\text{PBH}} \sim \mathcal{O}(10)M_{\odot}$  contributing a fraction  $f_{\text{PBH}} \simeq \mathcal{O}(10^{-3})$  could explain a significant fraction of the events, improving fits to the inferred mass distribution with respect to the simplest astrophysical source templates.

The question now is: do we have any PBH production model that can yield such abundance in this specific mass range? Generally speaking, PBH models are hardly predictive on its mass distribution. However, it turns out that there is at least one physically motivated model amenable to observational tests based on physics in the early universe, in particular the QCD phase transition. Such model [10–16], when including other early universe phenomena (like  $e^+e^-$  annihilation) yields a peculiar mass function with physically motivated features extending up to  $M_{\text{PBH}} \sim 10^7M_{\odot}$ . In this work, we revisit this “best motivated” scenario to assess its viability in the light of current constraints from cosmic microwave background (CMB) anisotropies associated to accretion onto PBH [17], from CMB spectral distortions [18], as well as null searches of sub-solar PBHs [19] and a stochastic gravitational wave background [20] in LIGO/Virgo. To do so, we compute the expected mass function associated not only to the QCD phase transition proper, but also the following particle antiparticle annihilation processes, down to the electron-positron annihilation.

The material included in this paper summarizes the work presented in the 14th conference on the identification of dark matter (IDM2022) organized by HEPHY in Vienna and closely follows reference [21], where we will constantly refer the reader to for more detailed calculations and further discussion.

## 2 Physics in the early universe

The PBH mass distribution adopts a very characteristic shape due to physical phenomena such as the QCD phase transition and electron-positron annihilation. In particular, an enhancement of PBH production is induced at those particular times, associated to a specific mass scale. A simple picture to understand how the mass function is shaped is the following: essentially, it all boils down to the decrease of relativistic d.o.f. which take place as a consequence of the drop of the temperature of the primordial plasma due to the expansion of the universe and the disappearance of species from it when the temperature falls roughly below its mass. This phenomena induces a decrease of the E.o.S. parameter which can be translated into a decrease of the overdensity threshold above which a PBH is formed. Therefore, whenever this drop in d.o.f. happens, the value an overdensity has to reach in order to collapse decreases and as a result PBH production is enhanced. This is summarized in Figure 1, where one can see the drops induced in the E.o.S. parameter.

Indeed, as we explained before, a decrease in the number of d.o.f. induces the dip structure observed in the right plot in Fig. 1. In particular, the most prominent dip at around  $\sim 1M_{\odot}$  is due to the QCD phase transition, the one at  $\sim 100M_{\odot}$  to the pion and muon annihilation and the third one at  $\sim 10^7M_{\odot}$  is caused by the electron-positron annihilation.




Figure 1: (Left) Effective number of relativistic degrees of freedom,  $g_{\text{eff}}$  and  $h_{\text{eff}}$ , as a function of the temperature (upper x-axis) and amount of mass enclosed in a Hubble patch (lower x-axis). (Right) Equation of state parameter  $w$  as a function of the temperature of the universe (top scale) or Hubble mass  $M_H$  (bottom scale). The gray horizontal line corresponds to the value during radiation domination  $w = 1/3$ .

### 3 The power spectrum

The early universe phenomena we just revisited in the previous section turns out to not be enough to obtain a significant production of PBHs. There is a second ingredient we need to deal with in order to account for a non-negligible amount of PBHs: the power spectrum. This object is well constrained at large scales, namely at CMB scales. However, PBHs are associated to the smallest scales, where constraints on the power spectrum still allow for a large variety of options. Naively, a first attempt to provide an expression for the power spectrum at such small scales would be to extrapolate it from the CMB scale. Nonetheless, one quickly realizes that such scenario leads to negligible production of PBHs. Therefore, in order to derive interesting scenarios, we need to introduce an enhancement of the power spectrum to larger values at the scales relevant for PBH production.

A couple of considerations regarding the scale of enhancement are in order. Firstly, it should not be placed too close to CMB scales since the power spectrum is already well constrained in this range and we don't want to mess it up. And secondly, it should not be placed too close to the QCD scale either, since we are trying to evaluate the scenario where the QCD phase transition is shaping the mass function in a very characteristic way and we don't want to spoil the natural appeal of it. All in all, the power spectrum should ideally be enhanced at a given scaled fulfilling condition 1.

$$k_{\text{QCD}} \gg k_{\text{cut}} \gg k_{\text{CMB}} \iff M_{\text{QCD}} \ll M_{\text{cut}} \ll M_{\text{CMB}}, \tag{1}$$

where  $M_{\text{cut}} = (\frac{k_{\text{cut}}}{10^6 \text{Mpc}^{-1}} (\frac{g_s}{10.75})^{1/12} 17^{-1/2})^{-2} M_{\odot}$ . For a particular parametrization of the power spectrum fulfilling condition 1, we refer the reader to [21], where one can see a phenomenological expression used to obtain some of the results in the next section.

### 4 Results and Conclusion

We first derive a mass distribution by requiring that the fraction of PBHs in the stellar mass range amounts to  $10^{-3}$ , as this is the value that seems to be preferred from the statistical fits




Figure 2: (Left) PBH mass distribution for a quasi-flat spectrum with a spectral index  $n_M = 0.025$ . The thin black line corresponds to the scenario without QCD/ $e^+e^-$  enhancement. It corresponds to Figure 5 in [13]. We also plot excluded regions from microlensing [25] [26] [27] in light green, GW production [19] for two different two-point delta mass distributions in blue and purple, accretion effects on CMB anisotropies [17] in pink/red and inferred SMBH population at high redshift [17] in gray. (Right) Mass functions consistent with three different sets of bounds for fixed values of  $M_c$ . We show the results for  $M_c = 10^8 M_\odot$  and SMBH counting (gray),  $M_c = 10^4 M_\odot$  and spherical accretion (red) and  $M_c = 10^2 M_\odot$  and GW production (blue and purple).

of LIGO/Virgo data. Therefore, we impose condition 2

$$f_{GW} \equiv \int_{5M_\odot}^{160M_\odot} \psi_p(M) dM \sim 10^{-3}, \tag{2}$$

where  $\psi_p$  is the mass function that ultimately depends on the power spectrum and all its parameters  $p$ . For a more detailed definition of the mass function and its derivation we again refer the reader to [21]. For the moment, we will not study any particular parametrization of the power spectrum in the whole wavenumber range (from CMB to PBHs) so we just assume condition 1 is implicitly fulfilled and take a CMB-like expression valid on CMB scales only of the form

$$\sigma^2 = 0.0033 \left( \frac{M}{10M_\odot} \right)^{n_M}, \tag{3}$$

where  $n_M = 0.025$  ( $n_M = 0$  corresponds to the scale invariant limit) and the numerical factor is obtained from 2. The resulting mass distribution is displayed in the left plot of Figure 2.

Before making any assessment on the validity of such scenario let us note the following remark. In Fig. 2 we are overplotting an extended mass distribution on top of a set of monochromatic bounds. In order to check the agreement among them, one cannot compare them directly as it is shown in the plot but instead one should first translate the monochromatic bounds into their extended version. Under some linear assumptions, one can derive the constraints imposed by a monochromatic bound  $f_{\text{mono}}^{\text{max}}$  on an extended mass function by imposing equation 4 as discussed in [28].

$$\int_{M_{\text{min}}}^{M_{\text{max}}} dM \frac{\psi_p(M)}{f_{\text{mono}}^{\text{max}}(M)} = 1, \tag{4}$$

where  $M_{\text{min}}$  ( $M_{\text{max}}$ ) is taken as the minimum (maximum) value for which the monochromatic bound has support.




Figure 3: Upper bounds on  $f_{GW}$  vs the cutoff mass  $M_c$  from CMB anisotropies (pink/red excluded regions) and non-observations of mergers with a BH whose mass is sub-solar; these bounds mildly depend on the heavier partner mass  $M_p$ , and the two blue bands in the plot bracket the extremes; see [19] for more details.

With this in mind, one can easily check that this particular scenario is in tension with most of the upper bounds. Is there a way to still get a considerable amount of PBHs in the stellar range and avoid the upper bounds at the same time? One option is to play with the enhancement scale. Clearly, depending on where we set this scale, we can easily avoid some of the bounds. Some allowed models for different values of the enhancement scale ( $M_{cut}$ ) are shown in the right plot of Figure 2. For scales such that  $M_{cut} \gtrsim 10^4 M_\odot$ , the mass function gets in tension with the CMB anisotropies bound and even for the SMBH counting bound at larger values. On the other hand, for  $M_{cut} \lesssim \mathcal{O}(10^2 M_\odot)$ , the cut is just above the QCD scale and, as discussed, cutting below means renouncing the idea of a QCD-inspired scenario.

The last issue we need to assess now is whether any of the models in the right plot of Fig. 2 can actually account for a fraction of  $f_{GW} \sim 10^{-3}$ . As it can be seen in Figure 3, current bounds on  $f_{PBH}$  lead to an upper limit of  $f_{GW} \lesssim 10^{-5}$ , which is well below (about two orders of magnitude!) the amount required in phenomenological fits. Therefore, in QCD-inspired scenarios, PBHs have at most a tiny contribution to LIGO/Virgo events.

Clearly, the results displayed in the previous plots are only valid under certain assumptions. In particular, we implicitly assumed a fixed mass function, that is the primordial mass distribution of PBHs at formation time is the same as the one today. This assumption might seem quite strong since we expect a significant evolution of the mass function, most notably due to accretion phenomena and PBH mergers. However, as discussed in [21], it does not seem plausible that such phenomena can modify the mass function in such a way that  $f_{GW} \sim 10^{-3}$  is attained and all the bounds avoided.

In conclusion, the most appealing scenario to explain the required mass function to significantly contribute to LIGO/Virgo merger events, invoking the physics of the early universe between the QCD phase transition and the  $e^\pm$  annihilation era does not appear viable. Of course, one could always tailor an alternative model leading to a prominent enough peak in the stellar mass range amounting to  $f_{GW} \sim 10^{-3}$  and avoiding all the bounds, although that would be at the expense of its predictability power.

## References

- [1] B. P. Abbott et al., *Observation of gravitational waves from a binary black hole merger*, Phys. Rev. Lett. **116**, 061102 (2016), doi:[10.1103/PhysRevLett.116.061102](https://doi.org/10.1103/PhysRevLett.116.061102).
- [2] M. Sasaki, T. Suyama, T. Tanaka and S. Yokoyama, *Primordial black hole scenario for the gravitational-wave event GW150914*, Phys. Rev. Lett. **117**, 061101 (2016), doi:[10.1103/PhysRevLett.117.061101](https://doi.org/10.1103/PhysRevLett.117.061101).
- [3] S. Bird et al., *Did LIGO detect dark matter?*, Phys. Rev. Lett. **116**, 201301 (2016), doi:[10.1103/PhysRevLett.116.201301](https://doi.org/10.1103/PhysRevLett.116.201301).
- [4] Y. Ali-Haïmoud, E. D. Kovetz and M. Kamionkowski, *Merger rate of primordial black-hole binaries*, Phys. Rev. D **96**, 123523 (2017), doi:[10.1103/PhysRevD.96.123523](https://doi.org/10.1103/PhysRevD.96.123523).
- [5] B. J. Kavanagh, D. Gaggero and G. Bertone, *Merger rate of a subdominant population of primordial black holes*, Phys. Rev. D **98**, 023536 (2018), doi:[10.1103/PhysRevD.98.023536](https://doi.org/10.1103/PhysRevD.98.023536).
- [6] Y. B. Zel'dovich and I. D. Novikov, *The hypothesis of cores retarded during expansion and the hot cosmological model*, Sov. Astron. **10**, 602 (1967).
- [7] S. Hawking, *Gravitationally collapsed objects of very low mass*, Mon. Not. R. Astron. Soc. **152**, 75 (1971), doi:[10.1093/mnras/152.1.75](https://doi.org/10.1093/mnras/152.1.75).
- [8] B. Carr and F. Kühnel, *Primordial black holes as dark matter: Recent developments*, Annu. Rev. Nucl. Part. Sci. **70**, 355 (2020), doi:[10.1146/annurev-nucl-050520-125911](https://doi.org/10.1146/annurev-nucl-050520-125911).
- [9] G. Franciolini et al., *Searching for a subpopulation of primordial black holes in LIGO-Virgo gravitational-wave data*, Phys. Rev. D **105**, 083526 (2022), doi:[10.1103/PhysRevD.105.083526](https://doi.org/10.1103/PhysRevD.105.083526).
- [10] K. Jedamzik, *Could MACHOS be primordial black holes formed during the QCD epoch?*, Phys. Rep. **307**, 155 (1998), doi:[10.1016/S0370-1573\(98\)00067-2](https://doi.org/10.1016/S0370-1573(98)00067-2).
- [11] K. Jedamzik and J. C. Niemeyer, *Primordial black hole formation during first-order phase transitions*, Phys. Rev. D **59**, 124014 (1999), doi:[10.1103/PhysRevD.59.124014](https://doi.org/10.1103/PhysRevD.59.124014).
- [12] J. L. G. Sobrinho, P. Augusto and A. L. Gonçalves, *New thresholds for primordial black hole formation during the QCD phase transition*, Mon. Not. R. Astron. Soc. **463**, 2348 (2016), doi:[10.1093/mnras/stw2138](https://doi.org/10.1093/mnras/stw2138).
- [13] C. T. Byrnes, M. Hindmarsh, S. Young and M. R. S. Hawkins, *Primordial black holes with an accurate QCD equation of state*, J. Cosmol. Astropart. Phys. **2018**, 041 (2018), doi:[10.1088/1475-7516/2018/08/041](https://doi.org/10.1088/1475-7516/2018/08/041).
- [14] B. Carr, S. Clesse and J. García-Bellido, *Primordial black holes from the QCD epoch: Linking dark matter, baryogenesis, and anthropic selection*, Mon. Not. R. Astron. Soc. **501**, 1426 (2020), doi:[10.1093/mnras/staa3726](https://doi.org/10.1093/mnras/staa3726).
- [15] B. Carr, S. Clesse, J. García-Bellido and F. Kühnel, *Cosmic conundra explained by thermal history and primordial black holes*, Phys. Dark Universe **31**, 100755 (2021), doi:[10.1016/j.dark.2020.100755](https://doi.org/10.1016/j.dark.2020.100755).
- [16] K. Jedamzik, *Consistency of primordial black hole dark matter with LIGO/Virgo merger rates*, Phys. Rev. Lett. **126**, 051302 (2021), doi:[10.1103/PhysRevLett.126.051302](https://doi.org/10.1103/PhysRevLett.126.051302).

- [17] V. Poulin, P. D. Serpico, F. Calore, S. Clesse and K. Kohri, *CMB bounds on disk-accreting massive primordial black holes*, Phys. Rev. D **96**, 083524 (2017), doi:[10.1103/PhysRevD.96.083524](https://doi.org/10.1103/PhysRevD.96.083524).
- [18] J. Chluba, A. L. Erickcek and I. Ben-Dayan, *Probing the inflation: Small-scale power spectrum constraints from measurements of the cosmic microwave background energy spectrum*, Astrophys. J. **758**, 76 (2012), doi:[10.1088/0004-637x/758/2/76](https://doi.org/10.1088/0004-637x/758/2/76).
- [19] A. H. Nitz and Y.-F. Wang, *Broad search for gravitational waves from subsolar-mass binaries through LIGO and Virgo's third observing run*, Phys. Rev. D **106**, 023024 (2022), doi:[10.1103/PhysRevD.106.023024](https://doi.org/10.1103/PhysRevD.106.023024).
- [20] R. Abbott et al., *Upper limits on the isotropic gravitational-wave background from advanced LIGO and advanced Virgo's third observing run*, Phys. Rev. D **104**, 022004 (2021), doi:[10.1103/PhysRevD.104.022004](https://doi.org/10.1103/PhysRevD.104.022004).
- [21] J. Iguaz, P. D. Serpico and G. Franco-Abellán, *The QCD phase transition behind a PBH origin of LIGO/Virgo events?*, J. Cosmol. Astropart. Phys. **2022**, 009 (2022), doi:[10.1088/1475-7516/2022/07/009](https://doi.org/10.1088/1475-7516/2022/07/009).
- [22] E. W. Kolb and M. S. Turner, *The early universe*, CRC Press, Boca Raton, USA, ISBN 9780201626742 (1994), doi:[10.1201/9780429492860](https://doi.org/10.1201/9780429492860).
- [23] L. Husdal, *On effective degrees of freedom in the early universe*, Galaxies **4**, 78 (2016), doi:[10.3390/galaxies4040078](https://doi.org/10.3390/galaxies4040078).
- [24] S. Borsanyi et al., *Calculation of the axion mass based on high-temperature lattice quantum chromodynamics*, Nature **539**, 69 (2016), doi:[10.1038/nature20115](https://doi.org/10.1038/nature20115).
- [25] H. Niikura, M. Takada, S. Yokoyama, T. Sumi and S. Masaki, *Constraints on Earth-mass primordial black holes from OGLE 5-year microlensing events*, Phys. Rev. D **99**, 083503 (2019), doi:[10.1103/PhysRevD.99.083503](https://doi.org/10.1103/PhysRevD.99.083503).
- [26] P. Tisserand et al., *Limits on the macho content of the galactic halo from the EROS-2 Survey of the Magellanic clouds*, Astron. Astrophys. **469**, 387 (2007), doi:[10.1051/0004-6361:20066017](https://doi.org/10.1051/0004-6361:20066017).
- [27] C. Alcock et al., *MACHO project limits on black hole dark matter in the 1 – 30  $M_{\odot}$  range*, Astrophys. J. **550**, L169 (2001), doi:[10.1086/319636](https://doi.org/10.1086/319636).
- [28] B. Carr, M. Raidal, T. Tenkanen, V. Vaskonen and H. Veermäe, *Primordial black hole constraints for extended mass functions*, Phys. Rev. D **96**, 023514 (2017), doi:[10.1103/PhysRevD.96.023514](https://doi.org/10.1103/PhysRevD.96.023514).