

Neutrino phenomenology via Type-(I+II) seesaw in the $U(1)_{L_e-L_\mu}$ model

Priya Mishra*, Mitesh Kumar Behera, Papia Panda,
Shivaramakrishna Singirala and Rukmani Mohanta

University of Hyderabad, Hyderabad, Telangana, India, 500046

* mishpriya99@gmail.com

16th International Workshop on Tau Lepton Physics (TAU2021)
Online, September 27 - October 1, 2021
doi:[10.21468/SciPostPhysProc.16](https://doi.org/10.21468/SciPostPhysProc.16)

Abstract

We show how, $L_e - L_\mu$ gauged symmetry is used as the Standard Model (SM) extension with the type (I+II) seesaw mechanism for neutrino mass generation. The particle content of this model involves three extra right handed neutrinos, one scalar triplet and one singlet with the SM spectrum. We discuss about lepton flavour violating decays $\mu \rightarrow e\gamma$, $\tau \rightarrow \mu\bar{\mu}\mu$ and anomalous magnetic moment of muon.



Copyright P. Mishra *et al.*

This work is licensed under the Creative Commons

[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Published by the SciPost Foundation.

Received 2021-11-23

Accepted 2024-12-13

Published 2025-07-15

doi:[10.21468/SciPostPhysProc.16.023](https://doi.org/10.21468/SciPostPhysProc.16.023)



Check for
updates

1 Introduction

Though successful, SM is not able to explain many observed phenomena in particle physics, astrophysics and cosmology. For instance, SM is unable to resolve the puzzle of neutrino physics. These days neutrinos are the most talked about topic in the scientific community as they are believed to contribute both in microscopic and macroscopic world. SM does not include right-handed neutrinos; as a result, neutrinos become unfavorable to have Dirac mass. Therefore to explain neutrino mass, we need to extend the boundary of the Standard Model. Here in this work, we use $U(1)_{L_e-L_\mu}$ gauge symmetry to go beyond SM and add three right handed neutrinos ν_{jR} , where $j = e, \mu, \tau$; one scalar singlet S and one scalar triplet Δ in framework of type (I+II) mechanism [1]. The neutrino mass matrix comes out to be two-zero A_2 texture. We constrain the model parameters consistent with current neutrino oscillation data. Furthermore, we obtain new contributions to muon $g-2$ and also charged lepton flavor violating decays such as $\mu \rightarrow e\gamma$, $\tau \rightarrow \mu\bar{\mu}\mu$.

2 The main ingredients of the model

As we extend the Standard model through Gauge ($U(1)_{L_e-L_\mu}$), fermion ($\nu_{eR}, \nu_{\mu R}, \nu_{\tau R}$) and scalar sector (S, Δ), the particle spectrum of the new model is:

The Lagrangian for leptonic sector, which is invariant under the $SU(2)_L \times U(1)_Y \times U(1)_{L_e-L_\mu}$ symmetry,

$$\begin{aligned} \mathcal{L}_{\text{lepton}} = & -y_l^\alpha \bar{\ell}_{\alpha L} H \alpha_R - \frac{1}{2} y_\Delta \left(\bar{\ell}_{\tau L} \Delta i \sigma_2 \ell_{\mu L}^C + \bar{\ell}_{\mu L} \Delta i \sigma_2 \ell_{\tau L}^C \right) - y_\nu^\alpha \bar{\ell}_{\alpha L} \tilde{H} \nu_{\alpha R} \\ & - \frac{1}{2} y_S^{e\tau} \left(\bar{\nu}_{eR}^C \nu_{\tau R} + \bar{\nu}_{\tau R}^C \nu_{eR} \right) S^\dagger - \frac{1}{2} y_S^{\mu\tau} \left(\bar{\nu}_{\mu R}^C \nu_{\tau R} + \bar{\nu}_{\tau R}^C \nu_{\mu R} \right) S \\ & - \frac{1}{2} \left[m_R^{\tau\tau} \bar{\nu}_{\tau R}^C \nu_{\tau R} + m_R^{e\mu} \left(\bar{\nu}_{\mu R}^C \nu_{eR} + \bar{\nu}_{eR}^C \nu_{\mu R} \right) \right] + \text{h.c.} \end{aligned} \quad (1)$$

3 Neutrino mass matrix

For the active neutrino mass matrix we implement type (I+II) seesaw mechanics,

$$M_\nu = M_L - M_D M_R^{-1} M_D^T. \quad (2)$$

From the Lagrangian given above we can get the forms of M_D , M_R and M_L ,

$$\begin{aligned} M_D &= \frac{v_H}{\sqrt{2}} \begin{pmatrix} y_\nu^e & 0 & 0 \\ 0 & y_\nu^\mu & 0 \\ 0 & 0 & y_\nu^\tau \end{pmatrix}, \\ M_L &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & y_\Delta v_\Delta \\ 0 & y_\Delta v_\Delta & 0 \end{pmatrix}, \\ M_R &= \begin{pmatrix} 0 & m_R^{e\mu} & |y_S^{e\tau}| \frac{v_S}{\sqrt{2}} e^{i\phi} \\ m_R^{e\mu} & 0 & y_S^{\mu\tau} \frac{v_S}{\sqrt{2}} \\ |y_S^{e\tau}| \frac{v_S}{\sqrt{2}} e^{i\phi} & y_S^{\mu\tau} \frac{v_S}{\sqrt{2}} & m_R^{\tau\tau} \end{pmatrix}, \end{aligned} \quad (3)$$

where M_R is the mass matrix of right-handed neutrinos.

Assuming $y_S^{e\tau} \gg y_S^{\mu\tau}$ in M_R to have correlations between mixing parameters and observables, the active neutrino mass matrix is given by,

$$M_\nu = \begin{pmatrix} 0 & -\frac{v_H^2 y_\nu^e y_\nu^\mu}{2m_R^{e\mu}} & 0 \\ -\frac{v_H^2 y_\nu^e y_\nu^\mu}{2m_R^{e\mu}} & -\frac{v_S^2 |y_S^{e\tau}|^2 v_H^2 (y_\nu^\mu)^2}{(m_R^{e\mu})^2 m_R^{\tau\tau}} e^{2i\phi} & y_\Delta v_\Delta + \frac{v_H^2 |y_S^{e\tau}| v_S y_\nu^\mu y_\nu^\tau}{2\sqrt{2} m_R^{e\mu} m_R^{\tau\tau}} e^{i\phi} \\ 0 & y_\Delta v_\Delta + \frac{v_H^2 |y_S^{e\tau}| v_S y_\nu^\mu y_\nu^\tau}{2\sqrt{2} m_R^{e\mu} m_R^{\tau\tau}} e^{i\phi} & -\frac{v_H^2 (y_\nu^\tau)^2}{2m_R^{\tau\tau}} \end{pmatrix}, \quad (4)$$

where $(M_\nu)_{ee} = (M_\nu)_{e\tau} = 0$, so M_ν has two-zero A_2 texture.

Table 1: Particle contents in $U(1)_{L_e-L_\mu}$ model.

Particles	$SU(2)_L$	$U(1)_Y$	$U(1)_{L_e-L_\mu}$
$l_{eL}, l_{\mu L}, l_{\tau L}$	2	-1	1, 0, -1
e_R, μ_R, τ_R	1	-2	1, 0, -1
$\nu_{eR}, \nu_{\mu R}, \nu_{\tau R}$	1	0	1, 0, -1
H	2	1	0
S	1	0	1
Δ	3	-2	-1




Figure 1: Left (right) plot shows the variation of $\sum_i m_i$ with $\sin^2\theta_{13}$ ($\sin^2\theta_{12}, \sin^2\theta_{23}$)




Figure 2: Left (right) plot shows the variation of $\sin^2\theta_{12}$ ($\sin^2\theta_{23}$) with respect to $\sin^2\theta_{13}$.

4 Numerical analysis

We have diagonalised the neutrino mass matrix which came out to be two-zero A_2 texture and hence supporting the normal hierarchy in neutrino masses [2]. We assume model parameters in the following region,

$$\begin{aligned} y_\nu^\alpha &\in [10^{-6}, 10^{-7}], & y_\Delta &\in [0.01, 1], \\ M_\Delta &\in [1, 2] \text{ TeV}, & M_1, M_2, M_3 &\in [1, 10] \text{ TeV}, \\ \nu_\Delta &\in [10^{-2}, 10^{-1}] \text{ eV}, \end{aligned} \quad (5)$$

which can show that the neutrino oscillation parameters are in good agreement with experimental data from NuFit [3] at 3σ range.

We show the plots between cosmic bound ($\sum m_i < 0.12 \text{ eV}$) for neutrino mass and mixing angle for the allowed range of parameters, correlation between mixing angles, CP violating phase and Jarlskog invariant with $\sin^2\theta_{13}$.

5 Lepton flavor violation

After the discovery of neutrino oscillations, it is evident that neutrino oscillates from one flavour to the other and if so in charged lepton sector, that lead to go beyond Standard Model.




Figure 3: Left (right) plot shows the variation of J_{CP} (δ_{CP}) with $\sin^2 \theta_{13}$




Figure 4: Left plot shows the correlation between the BR of $\mu \rightarrow e\gamma$ with respect to $m_{e\mu}$ and right one shows BR for $\tau \rightarrow \mu\bar{\mu}\mu$ with the mass element $m_{\mu\mu}$ and $m_{\mu\tau}$.

Here in our present model we show branching ratios (BR) for $\mu \rightarrow e\gamma$, $\tau \rightarrow \mu\bar{\mu}\mu$ decays are in good agreement with their current experimental limits. The BR for the lepton flavor violating decay modes are [4], [5]

$$\text{Br}(\mu \rightarrow e\gamma) = \frac{27\alpha_{em} |\langle m^2 \rangle_{e\tau}|^2}{256\pi G_F^2 v_\Delta^4 M_\Delta^4} < 4.2 \times 10^{-13}, \quad (6)$$

$$\text{Br}(\tau \rightarrow \mu\bar{\mu}\mu) = \frac{|m_{\tau\mu}|^2 |m_{\mu\mu}|^2}{16G_F^2 v_\Delta^4 M_\Delta^4} < 2.1 \times 10^{-8}, \quad (7)$$

where fine structure constant α_{em} is $\frac{1}{137}$ and fermi coupling G_F is $1.17 \times 10^{-5} \text{ GeV}^2$.

6 Muon g-2

The muon g-2 data reported by BNL has a discrepancy at 3.7σ level [6] and the data from E989 experiment at Fermilab [7] has deviation at 4.2σ level from the prediction of SM. In our present model, $Z'_{e\mu}$ couples to muon and hence the expression for Δa_μ in presence of $Z'_{e\mu}$ is,

$$\begin{aligned} \Delta a_\mu &= \frac{g'^2}{4\pi^2} \int_0^1 \frac{x^2(1-x)}{x^2 + (M_{Z'}^2/m_\mu^2)(1-x)} dx \\ &\approx \frac{m_\mu^2}{12\pi^2} \frac{g'^2_{e\mu}}{M_{Z'_{e\mu}}^2}. \end{aligned} \quad (8)$$




Figure 5: Projected limits on gauge couplings and mass of $Z'_{e\mu}$.

The value of $\frac{M_{Z'_{e\mu}}}{g'_{e\mu}}$ in terms of Δa_μ ,

$$\frac{M_{Z'_{e\mu}}}{g'_{e\mu}} = \sqrt{\frac{m_\mu^2}{12\pi^2 \Delta a_\mu^{\text{FNAL}}}},$$

and its required value to have muon magnetic moment in 3σ range is $148 < \frac{M_{Z'_{e\mu}}}{g'_{e\mu}} < 357$ in GeV. The plot is showing the gauge parameter space of $g-2$ of muon in the $(M_{Z'_{e\mu}} - g'_{e\mu})$ plane. Green region is excluded by neutrino trident production.

7 Conclusion

We have formulated two-zero A_2 texture neutrino mass matrix and shown the correlation plots with mixing parameters for allowed range of model parameter space and upper bound for LFV decays $\mu \rightarrow e\gamma, \tau \rightarrow \mu\bar{\mu}\mu$. Also, we discussed muon $g-2$ and the results are in favor of E989 experiment of Fermilab. Hence we demonstrated Neutrino phenomenology in extended SM with type (I+II) seesaw mechanism.

References

- [1] P. Panda, P. Mishra, M. Kumar Behera, S. Singirala and R. Mohanta, *Exploring neutrino masses, $(g-2)_{\mu,e}$ in type I+II seesaw in $L_e - L_\alpha$ -gauge extended model*, Universe **10**, 387 (2024), doi:[10.3390/universe10100387](https://doi.org/10.3390/universe10100387).
- [2] H. Fritzsch, Z.-z. Xing and S. Zhou, *Two-zero textures of the Majorana neutrino mass matrix and current experimental tests*, J. High Energy Phys. **09**, 083 (2011), doi:[10.1007/JHEP09\(2011\)083](https://doi.org/10.1007/JHEP09(2011)083).
- [3] M. C. Gonzalez-Garcia, M. Maltoni, J. Salvado and T. Schwetz, *Global fit to three neutrino mixing: Critical look at present precision*, J. High Energy Phys. **12**, 123 (2012), doi:[10.1007/JHEP12\(2012\)123](https://doi.org/10.1007/JHEP12(2012)123).

- [4] T. Fukuyama, H. Sugiyama and K. Tsumura, *Constraints from LFV processes in the Higgs triplet model*, J. High Energy Phys. **03**, 044 (2010), doi:[10.1007/JHEP03\(2010\)044](https://doi.org/10.1007/JHEP03(2010)044).
- [5] A. G. Akeroyd, M. Aoki and H. Sugiyama, *Probing Majorana phases and the neutrino mass spectrum in the Higgs triplet model at the CERN LHC*, Phys. Rev. D **77**, 075010 (2008), doi:[10.1103/PhysRevD.77.075010](https://doi.org/10.1103/PhysRevD.77.075010).
- [6] R. Prigl et al., *Status of the BNL Muon $g - 2$ Experiment*, IEEE Trans. Instrum. Meas. **48**, 182 (1999), doi:[10.1109/19.769558](https://doi.org/10.1109/19.769558).
- [7] T. Albahri et al., *Measurement of the anomalous precession frequency of the muon in the Fermilab muon $g - 2$ experiment*, Phys. Rev. D **103**, 072002 (2021), doi:[10.1103/PhysRevD.103.072002](https://doi.org/10.1103/PhysRevD.103.072002).