

New τ -based evaluation of the hadronic vacuum polarization contribution to the muon anomalous magnetic moment

Jesús Alejandro Miranda Hernández* and Pablo Roig Garcés

Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN,
Apdo. Postal 14-740, 07000 Ciudad de México, México

* jmiranda@fis.cinvestav.mx

16th International Workshop on Tau Lepton Physics (TAU2021)
Online, September 27 - October 1, 2021
doi:[10.21468/SciPostPhysProc.16](https://doi.org/10.21468/SciPostPhysProc.16)

Abstract

We revisit the radiative corrections to the $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau(\gamma)$ decays using Resonance Chiral Theory for the structure-dependent part. This channel is the most important one to understand the hadronic vacuum polarization piece of the muon g-2, which dominates the uncertainty of its data-driven prediction in the Standard Model. Using our results, the discrepancy between theory and experiment for this observable is reduced to $\sim 2.2\sigma$.



Copyright J. A. Miranda Hernández
and P. Roig Garcés.

This work is licensed under the Creative Commons
[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Published by the SciPost Foundation.

Received 2021-11-04

Accepted 2024-12-09

Published 2025-07-15

doi:[10.21468/SciPostPhysProc.16.032](https://doi.org/10.21468/SciPostPhysProc.16.032)



Check for
updates

1 Introduction

The combination of the first measurement from the new muon g-2 experiment at FNAL [1] with the final average from the muon g-2 measurements at BNL [2] yields

$$a_\mu^{\text{Exp}} = 116592061(41) \times 10^{-11}. \quad (1)$$

The Muon g-2 Theory Initiative quotes [3]

$$a_\mu^{\text{SM}} = 116591810(43) \times 10^{-11}, \quad (2)$$

as the Standard Model prediction for a_μ ,¹ which deviates 4.2σ from a_μ^{Exp} . However, the very precise lattice evaluation of the (leading order) hadronic vacuum polarization contribution to the muon g-2 ($a_\mu^{\text{HVP, LO}}$) by the BMW Coll. reduces this difference to only 1.6σ [39]. While this result is corroborated or refuted by another lattice calculation with commensurate accuracy (see Aida El-Khadra and Gilberto Colangelo, these proceedings), it is timely to reexamine the dominant contribution to $a_\mu^{\text{HVP, LO}}$ using hadronic tau decay data (instead of

¹This result is based on Refs. [4–38].

$\sigma(e^+e^- \rightarrow \text{hadrons})$, as in [3]). We will present here our results [40], for an updated evaluation of the radiative corrections needed for such purpose and our value of $a_\mu^{\text{HVP, LO}}$, based on tau data (see also Ref. [41]). Our study extends Refs. [42, 43], which first used Resonance Chiral Theory ($R\chi T$) [44, 45] in this setting,² by going one order further in the chiral expansion.

In section 2 we introduce $a_\mu^{\text{HVP, LO}}$. In section 3 we summarize the computation of the radiative corrections to di-pion tau decays and in section 4 we present our results. Finally, we conclude in section 5.

2 Hadronic vacuum polarization

Based on analyticity and unitarity, loop integrals containing HVP insertions in photon propagators can be expressed as dispersive integrals over the cross-section of a virtual photon decaying into hadrons

$$a_\mu^{\text{HVP, LO}} = \frac{\alpha^2}{3\pi^2} \int_{m_\pi^2}^{\infty} ds \frac{K(s)}{s} R(s), \quad (3)$$

where $R(s) = \sigma(e^+e^- \rightarrow \text{hadrons}(\gamma))/\sigma(e^+e^- \rightarrow \mu^+\mu^-)|_{\text{LO}}$ and $K(s)$ is a kernel function behaving as $\sim 1/s$, which enhances the weight of the low-energy contributions. Indeed, about 73% of these (and 58% of their uncertainty) comes from the $\pi^+\pi^-(\gamma)$ final states for $4m_\pi^2 \leq s \leq 0.8 \text{ GeV}^2$, in which we will focus here. Use of hadronic tau decay data in $a_\mu^{\text{HVP, LO}}$ requires to account for isospin-breaking (IB) corrections. At LO in them [42]

$$\sigma(e^+e^- \rightarrow \pi^+\pi^-(\gamma))(s) = \frac{K_\sigma(s)}{K_\Gamma(s)} \frac{d\Gamma(\tau^- \rightarrow \pi^-\pi^0\nu_\tau)}{ds} \frac{R_{\text{IB}}(s)}{S_{\text{EW}}}, \quad (4)$$

where $s = (p_\pi + p'_\pi)^2$, and

$$\begin{aligned} R_{\text{IB}}(s) &= \frac{FSR(s)}{G_{\text{EM}}(s)} \frac{\beta_{\pi^+\pi^-}^3(s)}{\beta_{\pi^-\pi^0}^3(s)} \frac{F_V^{\pi^+\pi^-}(s)}{f_+^{\pi^-\pi^0}(s)}, \\ K_\sigma(s) &= \frac{\pi\alpha^2}{3s}, \\ K_\Gamma(s) &= \frac{G_F^2 |V_{ud}|^2 M_\tau^3}{384\pi^3} \left(1 - \frac{s}{M_\tau^2}\right)^2 \left(1 + \frac{2s}{M_\tau^2}\right). \end{aligned} \quad (5)$$

FSR stands for final-state radiation, the ratio of β functions corresponds to the slightly different phase space (PS) for $\pi^{0/+}$, and that of the form factors accounts for ρ^- , ... or ρ^0/ω , ... resonances exchanged in the charged/neutral channel. G_{EM} includes the long-distance (real and virtual) radiative corrections and S_{EW} the universal short-distance ones. $K_{\sigma/\Gamma}$ collect the remaining differing factors between these processes. $\pi\pi$ upper/lower indices will be omitted in what follows.

For years, evaluations of $a_\mu^{\text{HVP, LO}}$ using tau data [29, 42, 43, 48–53] for the $\pi\pi$ cut have been closer to the world average for a_μ^{exp} than those employing e^+e^- data. We must also note that the $\rho^0 - \gamma$ mixing was proposed [54] to bring both sets of data into agreement (although the ρ meson was treated as an elementary boson). These differences have been interpreted in the context of effective Lagrangians for non-standard interactions [55–57], yielding constraints on new physics at the TeV level (see P. Roig, these proceedings).

² $R\chi T$ has also been employed in calculating the lightest pseudoscalar's pole contribution to a_μ [46, 47], in agreement with Ref. [3].

3 $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau(\gamma)$ decays

These decays were studied within $R\chi T$ [42, 43] including the contributions saturating the next-to-leading (NLO) order chiral low-energy constants (LECs).³ Here, we go [40] one order further in this expansion (NNLO). Refs. [60–62] analyzed these processes within a vector meson dominance model and put forward the importance of the $\rho - \omega - \pi$ couplings (these arise at chiral NNLO and thus were absent in Refs. [42, 43]) on the $G_{EM}(s)$. Our work [40] (see also [63] and Z. H. Guo, these proceedings) wanted to verify the importance of the $\rho - \omega - \pi$ vertices on $G_{EM}(s)$ and, further, to estimate an uncertainty for the original $R\chi T$ computation [42, 43] (and for ours, at the next chiral order), which is mandatory to assess the e^+e^- vs. τ data discrepancy in the two-pion channels, and its impact on $a_\mu^{\text{HVP, LO}}$.

The matrix element for the process with a real photon reads [64]

$$T = eG_F V_{ud}^* \epsilon^\mu(K)^* \left\{ F_\nu \bar{u}(q) \gamma^\nu (1 - \gamma_5) (M_\tau + \not{p} - \not{k}) \gamma_\mu u(P) + (V_{\mu\nu} - A_{\mu\nu}) \bar{u}(q) \gamma^\nu (1 - \gamma_5) u(P) \right\}, \quad (6)$$

where $F_\nu = (p_0 - p_-)_\nu f_+(s)/(2p \cdot k)$.⁴ The $V_{\mu\nu} - A_{\mu\nu}$ term can be split into structure-independent (SI) and dependent (SD) parts:

$$V^{\mu\nu} = V_{SI}^{\mu\nu} + V_{SD}^{\mu\nu}, \quad A^{\mu\nu} = A_{SD}^{\mu\nu}, \quad (7)$$

with the SI pieces fulfilling low-energy theorems [67, 68]. The evaluation of the SD part in $R\chi T$ up to NLO in the chiral expansion [43] only depends -keeping just the lightest spin-one resonance multiplets- on three resonance couplings (F_V, G_V, F_A) and the $U(3)$ $M_{V,A}$ masses. All of them can be determined employing a consistent set of short-distance constraints on the two-point Green functions, according to the operator product expansion of QCD, leading to [44, 45, 69]

$$F_V = \sqrt{2}F, \quad F_A = F, \quad G_V = \frac{F}{\sqrt{2}}, \quad M_A = \sqrt{2}M_V, \quad (8)$$

with $M_V \sim M_\rho(770)$ and $F \sim F_\pi \sim 92$ MeV. These were employed in Ref. [43], without estimating any uncertainty associated to the previous high-energy restrictions or to the missing higher-order contributions in the chiral expansion.

Including the $R\chi T$ Lagrangian which saturates the chiral LECs at NNLO [70, 71] increases the number of couplings constants (now also in the odd-intrinsic parity sector) entering $(V/A)_{SD}^{\mu\nu}$ to 55. Short-distance QCD constraints on both sectors [70–72] reduce them to 41 undetermined couplings,⁵ challenging predictive power. Different phenomenological determinations have bounded six of them [58, 74–80]. To estimate the remaining 35, we rely on chiral counting [81–83]. For the indeed most important $\rho - \omega - \pi$ couplings we also fit the corresponding LECs [84]. This yields results compatible with the chiral counting and allows to reduce the error on these couplings by a half, approximately. Still, our uncertainty estimation based on this procedure appears to be extremely conservative when we float Gaussianly all parameters within their corresponding range (with so many of them, it is likely that a few are close to their maxima, producing thus an unrealistically big error). Lacking a better way to take this uncertainty into account, we report the errors obtained in this way [40]. We have estimated the errors of Ref. [43], stemming from uncomputed higher-order chiral corrections, by adding, at NNLO, only couplings which are predicted using short-distance QCD constraints (this should give a lower bound on their uncertainty).

³Radiative corrections for the one-meson tau decays have also been computed within $R\chi T$ [58, 59].

⁴For our numerical evaluations we use the dispersive $f_+(s)$ form factor from Refs. [65, 66].

⁵These give rise to one-, two- and three-resonance mediated contributions, in both intrinsic parity sectors, to $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau \gamma$ decays, see Figs. 1-7 in Ref. [40]. Among these, the strength of the $\rho - \omega - \pi$ vertices is encoded in the κ_i^V couplings [71], see also Ref. [73].




Figure 1: $G_{\text{EM}}(s)$ function (see main text for details).

For different photon energy cuts, we predict [40] the di-pion invariant mass distribution, the branching ratio and the photon energy distribution in $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$ decays (see Figs. 10-14 and tables 2 and 3 in this reference). For definiteness, when cutting $E_\gamma < 100$ MeV, we obtain a branching ratio of $(1.9 \pm 0.3) \cdot 10^{-3}$, while this is $(9.5^{+3.5}_{-0.5}) \cdot 10^{-4}$ for Ref. [43] input. We emphasize the importance of measuring these observables at Belle-II. This will immediately shrink the errors of $a_\mu^{\text{HVP, LO}}$ using tau data and increase the new physics reach of other tests using them (see P. Roig, these proceedings).

The $G_{\text{EM}}(s)$ function is defined via

$$\frac{d\Gamma(\tau^- \rightarrow \pi^- \pi^0 \nu_\tau(\gamma))}{ds} = \frac{G_F^2 |V_{ud}|^2 M_\tau^3 S_{\text{EW}}}{384\pi^3} |f_+(s)|^2 \left(1 - \frac{s}{M_\tau^2}\right)^2 \left(1 - \frac{4m_\pi^2}{s}\right)^{3/2} \left(1 + \frac{2s}{M_\tau^2}\right) G_{\text{EM}}(s). \quad (9)$$

In Fig. 1 the solid line shows the $G_{\text{EM}}(s)$ function neglecting the structure-dependent part (only SI), the dashed and dotted lines are the NLO $G_{\text{EM}}(s)$ function (with either the constraints (8), or with the ones applying to two- and three-point Green functions, respectively). The blue shaded region is the full NNLO contribution, including (overestimated) uncertainties.⁶

4 Isospin-breaking corrections for the τ -based evaluation of $a_\mu^{\text{HVP, LO}}$

The effect of each IB correction can be computed using

$$\Delta a_\mu^{\text{HVP, LO}} = \frac{1}{4\pi^3} \int_{4m_\pi^2}^{s_{\text{cut}}} ds K(s) \frac{K_\sigma(s)}{K_\Gamma(s)} \frac{d\Gamma(\tau^- \rightarrow \pi^- \pi^0 \nu_\tau(\gamma))}{ds} \left(\frac{R_{\text{IB}}(s)}{S_{\text{EW}}^{\pi\pi}} - 1 \right), \quad (10)$$

and is summarized in table 1. Di-pion tau decay data from ALEPH [85], Belle [86], CLEO [87] and OPAL [88] was used.

As a test of these radiative corrections, the branching ratio ($\mathcal{B}$) of di-pion tau decays can be estimated from the measured $\sigma(e^+e^- \rightarrow \pi^+\pi^-(\gamma))$, by means of

$$\mathcal{B}(\tau^- \rightarrow \pi^- \pi^0 \nu_\tau(\gamma)) = \mathcal{B}(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau(\gamma)) \int_{4m_\pi^2}^{M_\tau^2} ds \sigma(e^+e^- \rightarrow \pi^+\pi^-(\gamma)) \mathcal{N}(s) \frac{S_{\text{EW}}}{R_{\text{IB}}(s)}, \quad (11)$$

⁶The $G_{\text{EM}}^0(s)$ function was introduced in Ref. [42], and corresponds to the leading contribution in a low-energy photon expansion [67].

Table 1: Different IB contributions to $a_\mu^{\text{HVP, LO}}$, obtained using tau data. For the form factors (FF) effect we follow the two approaches proposed in Refs. [43] ([50]) (see Ref. [40] for details).

Source	$\Delta a_\mu^{\text{HVP, LO}} (\times 10^{11})$
S_{EW}	−103.1
PS	−74.5
FSR	−45.6 ± 4.6
FF	+40.9 ± 48.9 (+77.6 ± 24.0)
EM	−15.9 ^{+5.7} _{−16.0}
Total	−107.1 ^{+49.4} _{−51.7} (−70.4 ^{+25.1} _{−29.2})




Figure 2: Comparison between the different data sets from BaBar (left) and KLOE (right) with our prediction using tau data and IB.

where $\mathcal{N}(s) = \frac{3|V_{ud}|^2}{2\pi\alpha^2 M_\tau^3} s \left(1 - \frac{s}{M_\tau^2}\right)^2 \left(1 + \frac{2s}{M_\tau^2}\right)$. Using BaBar $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ data [89] and the required IB corrections, we get

$$\mathcal{B}(\tau^- \rightarrow \pi^- \pi^0 \nu_\tau(\gamma)) = (24.68 \pm 0.15)\%, (24.70^{+0.26}_{-0.15})\%, \quad (12)$$

where the first (second) result includes NLO (NNLO) chiral contributions to $G_{\text{EM}}(s)$.⁷ NNLO results agree at the one σ level with the ALEPH [85] and Belle [86] measurements: $(25.24 \pm 0.39)\%$ and $(25.47 \pm 0.13)\%$, respectively (a 4σ tension appears at NLO with respect to ALEPH). A comparison between the BaBar [89] and KLOE [90] $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ data and the prediction obtained from $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau(\gamma)$ including IB corrections is seen in Fig. 2. We are only showing it for the difference between the charged and neutral di-pion vector form factor called ‘FF1’ in Ref. [40]. Results for ‘FF2’ are very similar (although the comparison is slightly worse). Our prediction is in better agreement with BaBar data. The dashed and solid lines represent the contributions at NLO using (8) or its generalization, respectively. The dotted line is the SI contribution. The red line depicts the envelope of $G_{\text{EM}}(s)$ at NNLO, with unoverestimated uncertainty. The blue dotdashed line is the NNLO prediction using only those couplings which are determined by asymptotic QCD.

Accounting for all di-pion tau decay data [85–88], we get

$$10^{10} \times a_\mu^{\text{HVP, LO}} = 519.6 \pm 2.8_{\text{spectra}+\mathcal{B}}^{+1.9}_{-2.1\text{IB}}, \quad \text{at NLO}, \quad (13)$$

and

$$10^{10} \times a_\mu^{\text{HVP, LO}} = 514.6 \pm 2.8_{\text{spectra}+\mathcal{B}}^{+5.0}_{-3.9\text{IB}}, \quad \text{at NNLO}. \quad (14)$$

⁷Contributions to the uncertainty are given in Ref. [40]. At NLO, errors are dominated by statistical uncertainty of BaBar data and by the error on $\Gamma_{\rho^+} - \Gamma_{\rho^0}$. At NNLO, these are overcome by the (asymmetric) error on $G_{\text{EM}}(s)$.




Figure 3: SM prediction [3], BMW value [39] and our results [40] compared to the a_μ measurements [1,2].

When these results are supplemented with the four-pion tau decay measurements and with e^+e^- data for the other channels and energies higher than the tau mass [8], we find the overall contribution

$$10^{10} \times a_\mu^{\text{HVP, LO}|\tau \text{ data}} = 705.7^{+4.0}_{-4.1}, \quad \text{at NLO}, \quad (15)$$

and

$$10^{10} \times a_\mu^{\text{HVP, LO}|\tau \text{ data}} = 700.7^{+6.1}_{-5.2}, \quad \text{at NNLO}. \quad (16)$$

When all other (QED, EW and subleading hadronic) contributions are added, the 4.2σ deficit of the SM prediction with respect to the FNAL+BNL average is reduced to

$$\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{th}} = (12.5 \pm 6.0) \times 10^{-10}, \quad \text{at NLO}, \quad (17)$$

and

$$\Delta a_\mu = (17.5^{+6.8}_{-7.5}) \times 10^{-10}, \quad \text{at NNLO}, \quad (18)$$

2.1 and 2.3σ deviations, respectively. Our results [40] are compared to the BNL [2] and FNAL [1] measurements, and to the White Paper SM prediction [3] and the BMW lattice result [39] in Fig. 3.

5 Conclusion

There is a global effort towards improving the Standard Model predictions of the hadronic contributions to a_μ . Specifically, there are dedicated studies to ameliorate lattice evaluations, dispersive data-driven computations, measurements of hadronic e^+e^- cross-section and τ decays, and related Monte Carlos.

The measurement of several observables in $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau \gamma$ decays will reduce drastically the model-dependent errors of the isospin-breaking corrections needed to use tau input in $a_\mu^{\text{HVP, LO}}$. Our computation of these improves agreement (also in spectra) between the measurements of e^+e^- and tau di-pion channels.

Our tau-based evaluation yields $\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{th}} = (12.5 \pm 6.0) \times 10^{-10}$, at NLO, and $\Delta a_\mu = (17.5^{+6.8}_{-7.5}) \times 10^{-10}$ at NNLO, corresponding to 2.1 and 2.3σ deviations, respectively.

Acknowledgments

The authors thank the organizing committees for the very nice TAU2021 Workshop.

Funding information The authors acknowledge Conacyt (México) and Cátedras Marcos Moshinsky (Fundación Marcos Moshinsky) for financial support.

References

- [1] B. Abi et al., *Measurement of the positive muon anomalous magnetic moment to 0.46 ppm*, Phys. Rev. Lett. **126**, 141801 (2021), doi:[10.1103/PhysRevLett.126.141801](https://doi.org/10.1103/PhysRevLett.126.141801).
- [2] G. W. Bennett et al., *Final report of the E821 muon anomalous magnetic moment measurement at BNL*, Phys. Rev. D **73**, 072003 (2006), doi:[10.1103/PhysRevD.73.072003](https://doi.org/10.1103/PhysRevD.73.072003).
- [3] T. Aoyama et al., *The anomalous magnetic moment of the muon in the Standard Model*, Phys. Rep. **887**, 1 (2020), doi:[10.1016/j.physrep.2020.07.006](https://doi.org/10.1016/j.physrep.2020.07.006).
- [4] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, *Reevaluation of the hadronic vacuum polarisation contributions to the Standard Model predictions of the muon $g-2$ and $\alpha(m_Z^2)$ using newest hadronic cross-section data*, Eur. Phys. J. C **77**, 827 (2017), doi:[10.1140/epjc/s10052-017-5161-6](https://doi.org/10.1140/epjc/s10052-017-5161-6).
- [5] A. Keshavarzi, D. Nomura and T. Teubner, *Muon $g-2$ and $\alpha(m_Z^2)$: A new data-based analysis*, Phys. Rev. D **97**, 114025 (2018), doi:[10.1103/PhysRevD.97.114025](https://doi.org/10.1103/PhysRevD.97.114025).
- [6] G. Colangelo, M. Hoferichter and P. Stoffer, *Two-pion contribution to hadronic vacuum polarization*, J. High Energy Phys. **02**, 006 (2019), doi:[10.1007/JHEP02\(2019\)006](https://doi.org/10.1007/JHEP02(2019)006).
- [7] M. Hoferichter, B.-L. Hoid and B. Kubis, *Three-pion contribution to hadronic vacuum polarization*, J. High Energy Phys. **08**, 137 (2019), doi:[10.1007/JHEP08\(2019\)137](https://doi.org/10.1007/JHEP08(2019)137).
- [8] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, *A new evaluation of the hadronic vacuum polarisation contributions to the muon anomalous magnetic moment and to $\alpha(m_Z^2)$* , Eur. Phys. J. C **80**, 241 (2020), doi:[10.1140/epjc/s10052-020-7792-2](https://doi.org/10.1140/epjc/s10052-020-7792-2).
- [9] A. Keshavarzi, D. Nomura and T. Teubner, *$g-2$ of charged leptons, $\alpha(m_Z^2)$, and the hyperfine splitting of muonium*, Phys. Rev. D **101**, 014029 (2020), doi:[10.1103/PhysRevD.101.014029](https://doi.org/10.1103/PhysRevD.101.014029).
- [10] A. Kurz, T. Liu, P. Marquard and M. Steinhauser, *Hadronic contribution to the muon anomalous magnetic moment to next-to-next-to-leading order*, Phys. Lett. B **734**, 144 (2014), doi:[10.1016/j.physletb.2014.05.043](https://doi.org/10.1016/j.physletb.2014.05.043).
- [11] B. Chakraborty et al., *Strong-isospin-breaking correction to the muon anomalous magnetic moment from lattice QCD at the physical point*, Phys. Rev. Lett. **120**, 152001 (2018), doi:[10.1103/PhysRevLett.120.152001](https://doi.org/10.1103/PhysRevLett.120.152001).
- [12] S. Borsanyi et al., *Hadronic vacuum polarization contribution to the anomalous magnetic moments of leptons from first principles*, Phys. Rev. Lett. **121**, 022002 (2018), doi:[10.1103/PhysRevLett.121.022002](https://doi.org/10.1103/PhysRevLett.121.022002).
- [13] T. Blum et al., *Calculation of the hadronic vacuum polarization contribution to the muon anomalous magnetic moment*, Phys. Rev. Lett. **121**, 022003 (2018), doi:[10.1103/PhysRevLett.121.022003](https://doi.org/10.1103/PhysRevLett.121.022003).

- [14] D. Giusti, V. Lubicz, G. Martinelli, F. Sanfilippo and S. Simula, *Electromagnetic and strong isospin-breaking corrections to the muon $g - 2$ from lattice QCD+QED*, Phys. Rev. D **99**, 114502 (2019), doi:[10.1103/PhysRevD.99.114502](https://doi.org/10.1103/PhysRevD.99.114502).
- [15] E. Shintani et al., *Hadronic vacuum polarization contribution to the muon $g - 2$ with $2 + 1$ flavor lattice QCD on a larger than $(10 \text{ fm})^4$ lattice at the physical point*, Phys. Rev. D **100**, 034517 (2019), doi:[10.1103/PhysRevD.100.034517](https://doi.org/10.1103/PhysRevD.100.034517).
- [16] C. T. H. Davies et al., *Hadronic-vacuum-polarization contribution to the muon's anomalous magnetic moment from four-flavor lattice QCD*, Phys. Rev. D **101**, 034512 (2020), doi:[10.1103/PhysRevD.101.034512](https://doi.org/10.1103/PhysRevD.101.034512).
- [17] A. Gérardin, M. Cè, G. von Hippel, B. Hörz, H. B. Meyer, D. Mohler, K. Ottnad, J. Wilhelm and H. Wittig, *Leading hadronic contribution to $(g - 2)_\mu$ from lattice QCD with $N_f = 2 + 1$ flavours of $O(a)$ improved Wilson quarks*, Phys. Rev. D **100**, 014510 (2019), doi:[10.1103/PhysRevD.100.014510](https://doi.org/10.1103/PhysRevD.100.014510).
- [18] C. Aubin, T. Blum, C. Tu, M. Golterman, C. Jung and S. Peris, *Light quark vacuum polarization at the physical point and contribution to the muon $g - 2$* , Phys. Rev. D **101**, 014503 (2020), doi:[10.1103/PhysRevD.101.014503](https://doi.org/10.1103/PhysRevD.101.014503).
- [19] D. Giusti and S. Simula, *Lepton anomalous magnetic moments in Lattice QCD+QED*, Proc. Sci. **363**, 104 (2019), doi:[10.22323/1.363.0104](https://doi.org/10.22323/1.363.0104).
- [20] K. Melnikov and A. Vainshtein, *Hadronic light-by-light scattering contribution to the muon anomalous magnetic moment reexamined*, Phys. Rev. D **70**, 113006 (2004), doi:[10.1103/PhysRevD.70.113006](https://doi.org/10.1103/PhysRevD.70.113006).
- [21] P. Masjuan and P. Sanchez-Puertas, *Pseudoscalar-pole contribution to the $(g_\mu - 2)$: A rational approach*, Phys. Rev. D **95**, 054026 (2017), doi:[10.1103/PhysRevD.95.054026](https://doi.org/10.1103/PhysRevD.95.054026).
- [22] G. Colangelo, M. Hoferichter, M. Procura and P. Stoffer, *Dispersion relation for hadronic light-by-light scattering: Two-pion contributions*, J. High Energy Phys. **04**, 161 (2017), doi:[10.1007/JHEP04\(2017\)161](https://doi.org/10.1007/JHEP04(2017)161).
- [23] M. Hoferichter, B.-L. Hoid, B. Kubis, S. Leupold and S. P. Schneider, *Dispersion relation for hadronic light-by-light scattering: Pion pole*, J. High Energy Phys. **10**, 141 (2018), doi:[10.1007/JHEP10\(2018\)141](https://doi.org/10.1007/JHEP10(2018)141).
- [24] A. Gérardin, H. B. Meyer and A. Nyffeler, *Lattice calculation of the pion transition form factor with $N_f = 2 + 1$ Wilson quarks*, Phys. Rev. D **100**, 034520 (2019), doi:[10.1103/PhysRevD.100.034520](https://doi.org/10.1103/PhysRevD.100.034520).
- [25] J. Bijnens, N. Hermansson-Truedsson and A. Rodríguez-Sánchez, *Short-distance constraints for the HLbL contribution to the muon anomalous magnetic moment*, Phys. Lett. B **798**, 134994 (2019), doi:[10.1016/j.physletb.2019.134994](https://doi.org/10.1016/j.physletb.2019.134994).
- [26] G. Colangelo, F. Hagelstein, M. Hoferichter, L. Laub and P. Stoffer, *Longitudinal short-distance constraints for the hadronic light-by-light contribution to $(g - 2)_\mu$ with large- N_c Regge models*, J. High Energy Phys. **03**, 101 (2020), doi:[10.1007/JHEP03\(2020\)101](https://doi.org/10.1007/JHEP03(2020)101).
- [27] V. Pauk and M. Vanderhaeghen, *Single meson contributions to the muon's anomalous magnetic moment*, Eur. Phys. J. C **74**, 3008 (2014), doi:[10.1140/epjc/s10052-014-3008-y](https://doi.org/10.1140/epjc/s10052-014-3008-y).
- [28] I. Danilkin and M. Vanderhaeghen, *Light-by-light scattering sum rules in light of new data*, Phys. Rev. D **95**, 014019 (2017), doi:[10.1103/PhysRevD.95.014019](https://doi.org/10.1103/PhysRevD.95.014019).

- [29] F. Jegerlehner, *The anomalous magnetic moment of the muon*, Springer, Cham, Switzerland, ISBN 9783319635750 (2017), doi:[10.1007/978-3-319-63577-4](https://doi.org/10.1007/978-3-319-63577-4).
- [30] M. Knecht, S. Narison, A. Rabemananjara and D. Rabetiariivony, *Scalar meson contributions to a_μ from hadronic light-by-light scattering*, Phys. Lett. B **787**, 111 (2018), doi:[10.1016/j.physletb.2018.10.048](https://doi.org/10.1016/j.physletb.2018.10.048).
- [31] G. Eichmann, C. S. Fischer and R. Williams, *Kaon-box contribution to the anomalous magnetic moment of the muon*, Phys. Rev. D **101**, 054015 (2020), doi:[10.1103/PhysRevD.101.054015](https://doi.org/10.1103/PhysRevD.101.054015).
- [32] P. Roig and P. Sanchez-Puertas, *Axial-vector exchange contribution to the hadronic light-by-light piece of the muon anomalous magnetic moment*, Phys. Rev. D **101**, 074019 (2020), doi:[10.1103/PhysRevD.101.074019](https://doi.org/10.1103/PhysRevD.101.074019).
- [33] G. Colangelo, M. Hoferichter, A. Nyffeler, M. Passera and P. Stoffer, *Remarks on higher-order hadronic corrections to the muon $g-2$* , Phys. Lett. B **735**, 90 (2014), doi:[10.1016/j.physletb.2014.06.012](https://doi.org/10.1016/j.physletb.2014.06.012).
- [34] T. Blum, N. Christ, M. Hayakawa, T. Izubuchi, L. Jin, C. Jung and C. Lehner, *Hadronic light-by-light scattering contribution to the muon anomalous magnetic moment from lattice QCD*, Phys. Rev. Lett. **124**, 132002 (2020), doi:[10.1103/PhysRevLett.124.132002](https://doi.org/10.1103/PhysRevLett.124.132002).
- [35] T. Aoyama, M. Hayakawa, T. Kinoshita and M. Nio, *Complete tenth-order QED contribution to the muon $g-2$* , Phys. Rev. Lett. **109**, 111808 (2012), doi:[10.1103/PhysRevLett.109.111808](https://doi.org/10.1103/PhysRevLett.109.111808).
- [36] T. Aoyama, T. Kinoshita and M. Nio, *Theory of the anomalous magnetic moment of the electron*, Atoms **7**, 28 (2019), doi:[10.3390/atoms7010028](https://doi.org/10.3390/atoms7010028).
- [37] A. Czarnecki, W. J. Marciano and A. Vainshtein, *Refinements in electroweak contributions to the muon anomalous magnetic moment*, Phys. Rev. D **67**, 073006 (2003), doi:[10.1103/PhysRevD.67.073006](https://doi.org/10.1103/PhysRevD.67.073006).
- [38] C. Gnendiger, D. Stöckinger and H. Stöckinger-Kim, *The electroweak contributions to $(g-2)_\mu$ after the Higgs-boson mass measurement*, Phys. Rev. D **88**, 053005 (2013), doi:[10.1103/PhysRevD.88.053005](https://doi.org/10.1103/PhysRevD.88.053005).
- [39] S. Borsanyi et al., *Leading hadronic contribution to the muon magnetic moment from lattice QCD*, Nature **593**, 51 (2021), doi:[10.1038/s41586-021-03418-1](https://doi.org/10.1038/s41586-021-03418-1).
- [40] J. A. Miranda and P. Roig, *New τ -based evaluation of the hadronic contribution to the vacuum polarization piece of the muon anomalous magnetic moment*, Phys. Rev. D **102**, 114017 (2020), doi:[10.1103/PhysRevD.102.114017](https://doi.org/10.1103/PhysRevD.102.114017).
- [41] A. Miranda, *Isospin-breaking corrections to $\tau \rightarrow \pi\pi\nu_\tau$ decays and the muon $g-2$* , Proc. Sci. **385**, 043 (2021), doi:[10.22323/1.385.0043](https://doi.org/10.22323/1.385.0043).
- [42] V. Cirigliano, G. Ecker and H. Neufeld, *Isospin violation and the magnetic moment of the muon*, Phys. Lett. B **513**, 361 (2001), doi:[10.1016/S0370-2693\(01\)00764-X](https://doi.org/10.1016/S0370-2693(01)00764-X).
- [43] V. Cirigliano, G. Ecker and H. Neufeld, *Radiative τ decay and the magnetic moment of the muon*, J. High Energy Phys. **08**, 002 (2002), doi:[10.1088/1126-6708/2002/08/002](https://doi.org/10.1088/1126-6708/2002/08/002).
- [44] G. Ecker, J. Gasser, A. Pich and E. De Rafael, *The role of resonances in chiral perturbation theory*, Nucl. Phys. B **321**, 311 (1989), doi:[10.1016/0550-3213\(89\)90346-5](https://doi.org/10.1016/0550-3213(89)90346-5).

- [45] G. Ecker, J. Gasser, H. Leutwyler, A. Pich and E. De Rafael, *Chiral Lagrangians for massive spin-1 fields*, Phys. Lett. B **223**, 425 (1989), doi:[10.1016/0370-2693\(89\)91627-4](https://doi.org/10.1016/0370-2693(89)91627-4).
- [46] P. Roig, A. Guevara and G. L. Castro, *VV'P form factors in resonance chiral theory and the $\pi - \eta - \eta'$ light-by-light contribution to the muon $g - 2$* , Phys. Rev. D **89**, 073016 (2014), doi:[10.1103/PhysRevD.89.073016](https://doi.org/10.1103/PhysRevD.89.073016).
- [47] A. Guevara, P. Roig and J. J. Sanz-Cillero, *Pseudoscalar pole light-by-light contributions to the muon $(g - 2)$ in resonance chiral theory*, J. High Energy Phys. **06**, 160 (2018), doi:[10.1007/JHEP06\(2018\)160](https://doi.org/10.1007/JHEP06(2018)160).
- [48] R. Alemany, M. Davier and A. Höcker, *Improved determination of the hadronic contribution to the muon $(g - 2)$ and to $\alpha(M_Z^2)$ using new data from hadronic τ decays*, Eur. Phys. J. C **2**, 123 (1998), doi:[10.1007/s100520050127](https://doi.org/10.1007/s100520050127).
- [49] M. Davier, S. Eidelman, A. Höcker and Z. Zhang, *Confronting spectral functions from e^+e^- annihilation and τ decays: Consequences for the muon magnetic moment*, Eur. Phys. J. C **27**, 497 (2003), doi:[10.1140/epjc/s2003-01136-2](https://doi.org/10.1140/epjc/s2003-01136-2).
- [50] M. Davier, A. Hoecker, G. L. Castro, B. Malaescu, X. H. Mo, G. T. Sánchez, P. Wang, C. Z. Yuan and Z. Zhang, *The discrepancy between τ and e^+e^- spectral functions revisited and the consequences for the muon magnetic anomaly*, Eur. Phys. J. C **66**, 127 (2010), doi:[10.1140/epjc/s10052-009-1219-4](https://doi.org/10.1140/epjc/s10052-009-1219-4).
- [51] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, *Reevaluation of the hadronic contributions to the muon $g - 2$ and to $\alpha(M_Z^2)$* , Eur. Phys. J. C **71**, 1515 (2011), doi:[10.1140/epjc/s10052-010-1515-z](https://doi.org/10.1140/epjc/s10052-010-1515-z).
- [52] M. Benayoun, P. David, L. DelBuono and F. Jegerlehner, *Upgraded breaking of the HLS model: A full solution to the $\tau^-e^+e^-$ and ϕ decay issues and its consequences on $g - 2$ VMD estimates*, Eur. Phys. J. C **72**, 1848 (2012), doi:[10.1140/epjc/s10052-011-1848-2](https://doi.org/10.1140/epjc/s10052-011-1848-2).
- [53] M. Davier, A. Höcker, B. Malaescu, C. Z. Yuan and Z. Zhang, *Update of the ALEPH non-strange spectral functions from hadronic τ decays*, Eur. Phys. J. C **74**, 2803 (2014), doi:[10.1140/epjc/s10052-014-2803-9](https://doi.org/10.1140/epjc/s10052-014-2803-9).
- [54] F. Jegerlehner and R. Szafron, *$\rho^0 - \gamma$ mixing in the neutral channel pion form factor $F_\pi^e(s)$ and its role in comparing $e^+ \rightarrow e^-$ with τ spectral functions*, Eur. Phys. J. C **71**, 1632 (2011), doi:[10.1140/epjc/s10052-011-1632-3](https://doi.org/10.1140/epjc/s10052-011-1632-3).
- [55] J. A. Miranda and P. Roig, *Effective-field theory analysis of the $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$ decays*, J. High Energy Phys. **11**, 038 (2018), doi:[10.1007/JHEP11\(2018\)038](https://doi.org/10.1007/JHEP11(2018)038).
- [56] V. Cirigliano, A. Falkowski, M. González-Alonso and A. Rodríguez-Sánchez, *Hadronic τ decays as new physics probes in the LHC era*, Phys. Rev. Lett. **122**, 221801 (2019), doi:[10.1103/PhysRevLett.122.221801](https://doi.org/10.1103/PhysRevLett.122.221801).
- [57] S. González-Solís, A. Miranda, J. Rendón and P. Roig, *Exclusive hadronic tau decays as probes of non-SM interactions*, Phys. Lett. B **804**, 135371 (2020), doi:[10.1016/j.physletb.2020.135371](https://doi.org/10.1016/j.physletb.2020.135371).
- [58] Z.-H. Guo and P. Roig, *One-meson radiative tau decays*, Phys. Rev. D **82**, 113016 (2010), doi:[10.1103/PhysRevD.82.113016](https://doi.org/10.1103/PhysRevD.82.113016).

- [59] M. A. Arroyo-Ureña, G. Hernández-Tomé, G. López-Castro, P. Roig and I. Rosell, *Radiative corrections to $\tau \rightarrow \pi(K) \nu_\tau [\gamma]$: A reliable new physics test*, Phys. Rev. D **104**, L091502 (2021), doi:[10.1103/PhysRevD.104.L091502](https://doi.org/10.1103/PhysRevD.104.L091502).
- [60] A. Flores-Tlalpa, G. L. Castro and G. T. Sánchez, *Radiative two-pion decay of the tau lepton*, Phys. Rev. D **72**, 113003 (2005), doi:[10.1103/PhysRevD.72.113003](https://doi.org/10.1103/PhysRevD.72.113003).
- [61] F. Flores-Báez, A. Flores-Tlalpa, G. L. Castro and G. T. Sánchez, *Long distance radiative corrections to the decay $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$* , Phys. Rev. D **74**, 071301 (2006), doi:[10.1103/PhysRevD.74.071301](https://doi.org/10.1103/PhysRevD.74.071301).
- [62] A. Flores-Tlalpa, F. Flores-Báez, G. L. Castro and G. T. Sánchez, *Model-dependent radiative corrections to $\tau^- \rightarrow \pi^- \pi^0 \nu$ revisited*, Nucl. Phys. B - Proc. Suppl. **169**, 250 (2007), doi:[10.1016/j.nuclphysbps.2007.03.011](https://doi.org/10.1016/j.nuclphysbps.2007.03.011).
- [63] J. L. G. Santiago, G. L. Castro and P. Roig, *Lepton-pair production in dipion τ lepton decays*, Phys. Rev. D **103**, 014027 (2021), doi:[10.1103/PhysRevD.103.014027](https://doi.org/10.1103/PhysRevD.103.014027).
- [64] J. Bijnens, G. Ecker and J. Gasser, *Radiative semileptonic kaon decays*, Nucl. Phys. B **396**, 81 (1993), doi:[10.1016/0550-3213\(93\)90259-R](https://doi.org/10.1016/0550-3213(93)90259-R).
- [65] D. G. Dumm and P. Roig, *Dispersive representation of the pion vector form factor in $\tau \rightarrow \pi \pi \nu_\tau$ decays*, Eur. Phys. J. C **73**, 2528 (2013), doi:[10.1140/epjc/s10052-013-2528-1](https://doi.org/10.1140/epjc/s10052-013-2528-1).
- [66] S. González-Solís and P. Roig, *A dispersive analysis of the pion vector form factor and $\tau^- \rightarrow K^- K_S \nu_\tau$ decay*, Eur. Phys. J. C **79**, 436 (2019), doi:[10.1140/epjc/s10052-019-6943-9](https://doi.org/10.1140/epjc/s10052-019-6943-9).
- [67] F. E. Low, *Bremsstrahlung of very low-energy quanta in elementary particle collisions*, Phys. Rev. **110**, 974 (1958), doi:[10.1103/PhysRev.110.974](https://doi.org/10.1103/PhysRev.110.974).
- [68] T. H. Burnett and N. M. Kroll, *Extension of the low soft-photon theorem*, Phys. Rev. Lett. **20**, 86 (1968), doi:[10.1103/PhysRevLett.20.86](https://doi.org/10.1103/PhysRevLett.20.86).
- [69] S. Weinberg, *Precise relations between the spectra of vector and axial-vector mesons*, Phys. Rev. Lett. **18**, 507 (1967), doi:[10.1103/PhysRevLett.18.507](https://doi.org/10.1103/PhysRevLett.18.507).
- [70] V. Cirigliano, G. Ecker, M. Eidemüller, R. Kaiser, A. Pich and J. Portolés, *Towards a consistent estimate of the chiral low-energy constants*, Nucl. Phys. B **753**, 139 (2006), doi:[10.1016/j.nuclphysb.2006.07.010](https://doi.org/10.1016/j.nuclphysb.2006.07.010).
- [71] K. Kampf and J. Novotný, *Resonance saturation in the odd-intrinsic parity sector of low-energy QCD*, Phys. Rev. D **84**, 014036 (2011), doi:[10.1103/PhysRevD.84.014036](https://doi.org/10.1103/PhysRevD.84.014036).
- [72] P. Roig and J. J. S. Cillero, *Consistent high-energy constraints in the anomalous QCD sector*, Phys. Lett. B **733**, 158 (2014), doi:[10.1016/j.physletb.2014.04.034](https://doi.org/10.1016/j.physletb.2014.04.034).
- [73] P. D. Ruiz-Femenía, A. Pich and J. Portolés, *Odd-intrinsic-parity processes within the resonance effective theory of QCD*, J. High Energy Phys. **07**, 003 (2003), doi:[10.1088/1126-6708/2003/07/003](https://doi.org/10.1088/1126-6708/2003/07/003).
- [74] D. G. Dumm, P. Roig, A. Pich and J. Portolés, *Hadron structure in $\tau \rightarrow KK \pi \nu_\tau$ decays*, Phys. Rev. D **81**, 034031 (2010), doi:[10.1103/PhysRevD.81.034031](https://doi.org/10.1103/PhysRevD.81.034031).
- [75] D. G. Dumm, P. Roig, A. Pich and J. Portolés, *$\tau \rightarrow \pi \pi \pi \nu_\tau$ decays and the $a_1(1260)$ off-shell width revisited*, Phys. Lett. B **685**, 158 (2010), doi:[10.1016/j.physletb.2010.01.059](https://doi.org/10.1016/j.physletb.2010.01.059).

- [76] D. G. Dumm and P. Roig, *Resonance chiral Lagrangian analysis of $\tau^- \rightarrow \eta^{(\prime)} \pi^- \pi^0 \nu_\tau$* , Phys. Rev. D **86**, 076009 (2012), doi:[10.1103/PhysRevD.86.076009](https://doi.org/10.1103/PhysRevD.86.076009).
- [77] Y.-H. Chen, Z.-H. Guo and H.-Q. Zheng, *Study of $\eta - \eta'$ mixing from radiative decay processes*, Phys. Rev. D **85**, 054018 (2012), doi:[10.1103/PhysRevD.85.054018](https://doi.org/10.1103/PhysRevD.85.054018).
- [78] L. Y. Dai, J. Portolés and O. Shekhovtsova, *Three pseudoscalar meson production in e^+e^- annihilation*, Phys. Rev. D **88**, 056001 (2013), doi:[10.1103/PhysRevD.88.056001](https://doi.org/10.1103/PhysRevD.88.056001).
- [79] P. Roig, A. Guevara and G. L. Castro, *Weak radiative pion vertex in $\tau^- \rightarrow \pi^- \nu_\tau \ell^+ \ell^-$ decays*, Phys. Rev. D **88**, 033007 (2013), doi:[10.1103/PhysRevD.88.033007](https://doi.org/10.1103/PhysRevD.88.033007).
- [80] Y.-H. Chen, Z.-H. Guo and H.-Q. Zheng, *Radiative transition processes of light vector resonances in a chiral framework*, Phys. Rev. D **90**, 034013 (2014), doi:[10.1103/PhysRevD.90.034013](https://doi.org/10.1103/PhysRevD.90.034013).
- [81] S. Weinberg, *Phenomenological Lagrangians*, Phys. A: Stat. Mech. Appl. **96**, 327 (1979), doi:[10.1016/0378-4371\(79\)90223-1](https://doi.org/10.1016/0378-4371(79)90223-1).
- [82] J. Gasser and H. Leutwyler, *Chiral perturbation theory to one loop*, Ann. Phys. **158**, 142 (1984), doi:[10.1016/0003-4916\(84\)90242-2](https://doi.org/10.1016/0003-4916(84)90242-2).
- [83] J. Gasser and H. Leutwyler, *Chiral perturbation theory: Expansions in the mass of the strange quark*, Nucl. Phys. B **250**, 465 (1985), doi:[10.1016/0550-3213\(85\)90492-4](https://doi.org/10.1016/0550-3213(85)90492-4).
- [84] S.-Z. Jiang, Z.-L. Wei, Q.-S. Chen and Q. Wang, *Computation of the $O(p^6)$ order low-energy constants: An update*, Phys. Rev. D **92**, 025014 (2015), doi:[10.1103/PhysRevD.92.025014](https://doi.org/10.1103/PhysRevD.92.025014).
- [85] S. Schael et al., *Branching ratios and spectral functions of τ decays: Final ALEPH measurements and physics implications*, Phys. Rep. **421**, 191 (2005), doi:[10.1016/j.physrep.2005.06.007](https://doi.org/10.1016/j.physrep.2005.06.007).
- [86] M. Fujikawa et al., *High-statistics study of the $\tau^- \rightarrow \pi^0 \pi^0 \nu_\tau$ decay*, Phys. Rev. D **78**, 072006 (2008), doi:[10.1103/PhysRevD.78.072006](https://doi.org/10.1103/PhysRevD.78.072006).
- [87] S. Anderson et al., *Hadronic structure in the decay $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$* , Phys. Rev. D **61**, 112002 (2000), doi:[10.1103/PhysRevD.61.112002](https://doi.org/10.1103/PhysRevD.61.112002).
- [88] K. Ackerstaff et al., *Measurement of the strong coupling constant α_s and the vector and axial-vector spectral functions in hadronic tau decays*, Eur. Phys. J. C **7**, 571 (1999), doi:[10.1007/s100529901061](https://doi.org/10.1007/s100529901061).
- [89] J. P. Lees, et al., *Precise measurement of the $e^+e^- \rightarrow \pi^+ \pi^- (\gamma)$ cross section with the initial-state radiation method at BABAR*, Phys. Rev. D **86**, 032013 (2012), doi:[10.1103/PhysRevD.86.032013](https://doi.org/10.1103/PhysRevD.86.032013).
- [90] A. Anastasi et al., *Combination of KLOE $\sigma(e^+e^- \rightarrow \pi^+ \pi^- \gamma(\gamma))$ measurements and determination of $a_\mu^{\pi^+ \pi^-}$ in the energy range $0.10 < 0.95 \text{ GeV}^2$* , J. High Energy Phys. **03**, 173 (2018), doi:[10.1007/JHEP03\(2018\)173](https://doi.org/10.1007/JHEP03(2018)173).