


Applications of the Perturbative Gradient Flow

Fabian Lange^{1,2*}

1 Institut für Theoretische Teilchenphysik, Karlsruhe Institute of Technology (KIT), Wolfgang-Gaede Straße 1, 76128 Karlsruhe, Germany

2 Institut für Astroteilchenphysik, Karlsruhe Institute of Technology (KIT), Hermann-von-Helmholtz-Platz 1, 76344 Eggenstein-Leopoldshafen, Germany

* fabian.lange@kit.edu



Abstract

Over the last decade the gradient flow formalism became an important tool for lattice simulations of Quantum Chromodynamics. It offers remarkable renormalization properties which pave the way for cross-fertilization between perturbative and lattice calculations. In this contribution we discuss the perturbative approach. As first application we compute vacuum expectation values of flowed operators which could help to extract parameters like the strong coupling constant from lattice simulations. Afterwards, we apply the flowed operator product expansion to the time-ordered product of two currents which could be employed for an alternative first-principle evaluation of vacuum polarization functions on the lattice.



Copyright F. Lange.

This work is licensed under the Creative Commons

[Attribution 4.0 International License](#).

Published by the SciPost Foundation.

Received 15-10-2021

Accepted 28-02-2022

Published 21-06-2022



[doi:10.21468/SciPostPhysProc.7.033](#)

1 Introduction

The gradient flow formalism introduced in Refs. [1–3] became an important tool for simulations of *Quantum Chromodynamics* (QCD) on the lattice over the last decade. Most prominently, it led to new strategies to set the scale of the lattice, see e.g. Refs. [3–5]. Moreover, it provides strong renormalization properties. Especially, *flowed*¹ composite operators constructed from *flowed* fields do not require renormalization [6]. Therefore, they do not mix under *renormalization group* (RG) running which allows one to match results of lattice and perturbative calculations without scheme transformation. One prominent application is the

¹We use the terms *flowed* and *regular* to distinguish quantities defined at flow time $t > 0$ from those defined at $t = 0$.

extraction of the strong coupling α_s from lattice simulations which, however, did not yield competitive results yet, see Ref. [7] for a recent review.

A powerful tool is the so-called *small-flow-time expansion* which leads to a relation between flowed and regular operators related by a flow-time dependent mixing matrix [6]. By inverting the mixing matrix, one obtains a flowed *operator product expansion* (OPE) which expresses the regular operators through the corresponding flowed operators [8–10]. This was first utilized to construct a regularization independent formula for the *energy-momentum tensor* (EMT) of QCD [8, 9].

In this contribution we briefly introduce the perturbative treatment of the gradient flow at infinite volume² in Section 2 which allows us to compute *vacuum expectation values* (VEVs) of flowed operators through *next-to-next-to-leading order* (NNLO) in Section 3. In Section 4, we discuss the flowed OPE and apply it to *vacuum polarization functions* (VPFs) which might pave the way for an alternative determination of the hadronic corrections to the anomalous magnetic moment and other observables.

2 Perturbative Gradient Flow

The gradient flow formalism continues the gluon and quark fields $A_\mu^a(x)$ and $\psi(x)$ of regular QCD from $D = 4 - 2\epsilon$ Euclidean dimensions to the fields $B_\mu^a(t, x)$ and $\chi(t, x)$ additionally depending on the *flow time* $t > 0$ through the boundary conditions

$$B_\mu^a(t=0, x) = A_\mu^a(x), \quad \chi(t=0, x) = \psi(x) \quad (1)$$

and the flow equations [3, 12]

$$\partial_t B_\mu^a = \mathcal{D}_\nu^{ab} G_{\nu\mu}^b + \kappa \mathcal{D}_\mu^{ab} \partial_\nu B_\nu^b, \quad \partial_t \chi = \Delta \chi - \kappa \partial_\mu B_\mu^a T^a \chi, \quad \partial_t \bar{\chi} = \bar{\chi} \overleftarrow{\Delta} + \kappa \bar{\chi} \partial_\mu B_\mu^a T^a, \quad (2)$$

where

$$\begin{aligned} G_{\mu\nu}^a &= \partial_\mu B_\nu^a - \partial_\nu B_\mu^a + f^{abc} B_\mu^b B_\nu^c, & \mathcal{D}_\mu^{ab} &= \delta^{ab} \partial_\mu - f^{abc} B_\mu^c, \\ \Delta &= \mathcal{D}_\mu^F \mathcal{D}_\mu^F, & \overleftarrow{\Delta} &= \overleftarrow{\mathcal{D}}_\mu^F \overleftarrow{\mathcal{D}}_\mu^F, & \mathcal{D}_\mu^F &= \partial_\mu + B_\mu^a T^a, & \overleftarrow{\mathcal{D}}_\mu^F &= \overleftarrow{\partial}_\mu - B_\mu^a T^a. \end{aligned} \quad (3)$$

The flow time t is a parameter of mass dimension minus two and we use the short-hand notation $\partial_t \equiv \frac{\partial}{\partial t}$. The symmetry generators T^a in the fundamental representation and the structure constants f^{abc} are defined through

$$[T^a, T^b] = f^{abc} T^c, \quad \text{Tr}(T^a T^b) = -T_R \delta^{ab}. \quad (4)$$

κ is an additional gauge parameter and all physical observables should be independent of it [3]. In perturbative calculations it is usually most convenient to set $\kappa = 1$.

The flow equations (2) can be incorporated into a Lagrangian formalism by defining

$$\mathcal{L} = \mathcal{L}_{\text{QCD}} + \mathcal{L}_{\text{gauge-fixing}} + \mathcal{L}_{\text{ghost}} + \mathcal{L}_B + \mathcal{L}_\chi. \quad (5)$$

The first three terms constituting the regular Yang-Mills Lagrangian are given by

$$\begin{aligned} \mathcal{L}_{\text{QCD}} &= \frac{1}{4g_B^2} F_{\mu\nu}^a F_{\mu\nu}^a + \sum_{f=1}^{n_f} \bar{\psi}_f (\not{p}^F + m_{f,B}) \psi_f, \\ \mathcal{L}_{\text{gauge-fixing}} &= \frac{1}{2g_B^2 \xi} (\partial_\mu A_\mu^a)^2, \quad \mathcal{L}_{\text{ghost}} = \frac{1}{g_B^2} \partial_\mu \bar{c}^a D_\mu^{ab} c^b, \end{aligned} \quad (6)$$

²At finite volume different techniques are required, see e.g. Ref. [11].

where

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + f^{abc} A_\mu^b A_\nu^c, \quad D_\mu^F = \partial_\mu + A_\mu^a T^a, \quad D_\mu^{ab} = \delta^{ab} \partial_\mu - f^{abc} A_\mu^c. \quad (7)$$

The flow equations are incorporated through

$$\begin{aligned} \mathcal{L}_B &= -2 \int_0^\infty dt \text{Tr} \left[L_\mu^a T^a \left(\partial_t B_\mu^b T^b - \mathcal{D}_\nu^{bc} G_{\nu\mu}^c T^b - \kappa \mathcal{D}_\mu^{bc} \partial_\nu B_\nu^c T^b \right) \right], \\ \mathcal{L}_\chi &= \sum_{f=1}^{n_f} \int_0^\infty dt \left[\bar{\lambda}_f \left(\partial_t - \Delta + \kappa \left(\partial_\mu B_\mu^a \right) T^a \right) \chi_f + \bar{\chi}_f \left(\overleftarrow{\partial}_t - \overleftarrow{\Delta} - \kappa \left(\partial_\mu B_\mu^a \right) T^a \right) \lambda_f \right], \end{aligned} \quad (8)$$

where $L_\mu^a(t, x)$, $\lambda_f(t, x)$, and $\bar{\lambda}_f(t, x)$ are Lagrange multiplier fields [6, 12]. Their Euler-Lagrange equations lead to Eq. (2).

The Feynman rules for perturbative calculations can be derived from the Lagrangian employing standard techniques [6, 12] and the complete list can be found in Ref. [13].

3 Vacuum Expectation Values

VEVs of gauge-invariant operators at finite flow time are among the simplest quantities one can consider within the gradient flow formalism. As mentioned before, these operators do not require any *ultra-violet* (UV) renormalization beyond that of regular QCD and that of the involved flowed fields [6]. This means that the operators do not mix under RG running, which makes it particularly simple to combine results from different regularization schemes.

The renormalization of the coupling and the masses follows the usual prescription with the known QCD renormalization constants. Throughout this contribution we employ the $\overline{\text{MS}}$ scheme and refer to Refs. [13, 14] for details.

The flowed gauge field $B_\mu^a(t, x)$ does not require renormalization so that matrix elements of the gluon action density

$$E(t, x) \equiv \frac{1}{4} G_{\mu\nu}^a(t, x) G_{\mu\nu}^a(t, x) \quad (9)$$

are finite after just the renormalization of g and m_f [3, 6]. Hence, a direct comparison of results obtained in different regularization schemes is possible.

In contrast, flowed quark fields require a renormalization factor $Z_\chi^{1/2}(\alpha_s)$ in order to render Green's functions finite. In the $\overline{\text{MS}}$ scheme it reads

$$Z_\chi^{-1}(\alpha_s) = 1 - \frac{\alpha_s}{4\pi} \frac{\gamma_{\chi,0}}{\epsilon} + \left(\frac{\alpha_s}{4\pi} \right)^2 \left[\frac{\gamma_{\chi,0}}{2\epsilon^2} (\gamma_{\chi,0} + \beta_0) - \frac{\gamma_{\chi,1}}{2\epsilon} \right] + \mathcal{O}(\alpha_s^3), \quad (10)$$

with

$$\gamma_{\chi,0} = 3 C_F, \quad \gamma_{\chi,1} = \left(\frac{223}{6} - 8 \ln 2 \right) C_A C_F - \left(\frac{3}{2} + 8 \ln 2 \right) C_F^2 - \frac{22}{3} C_F T_R n_f. \quad (11)$$

$\gamma_{\chi,0}$ has been computed in Ref. [12], whereas $\gamma_{\chi,1}$ has been obtained by requiring that the NNLO calculations in Refs. [13, 15] become finite.

The scalar quark density

$$S(t, x) \equiv Z_\chi \sum_{f=1}^{n_f} \bar{\chi}_f(t, x) \chi_f(t, x) \quad (12)$$

thus acquires an anomalous dimension, which prevents a direct comparison of results from different regularization schemes. This can be cured by working with *ringed quark fields* [9], which amounts to renormalizing the flowed quark fields with

$$\dot{Z}_\chi(t, \mu) = -\frac{2N_C n_f}{(4\pi t)^2} \cdot \frac{1}{\langle R(t) \rangle|_{m=0}} \quad (13)$$




Figure 1: Sample diagrams for the VEVs through NNLO. Produced with TikZ-Feynman [16].

instead of Z_χ , where

$$R(t, x) = \sum_{f=1}^{n_f} \bar{\chi}_f(t, x) \overleftrightarrow{D}^F \chi_f(t, x) \quad \text{with} \quad \overleftrightarrow{D}_\mu^F = D_\mu^F - \overleftarrow{D}_\mu^F \quad (14)$$

is the quark kinetic operator. This corresponds to a “physical” renormalization scheme, which means that the anomalous dimension of the operator

$$\dot{S}(t, x) = \zeta_\chi(t, \mu) S(t, x) \quad \text{with} \quad \zeta_\chi(t, \mu) \equiv Z_\chi^{-1} \dot{Z}_\chi(t, \mu) \quad (15)$$

vanishes.

The Feynman rules for the operators $E(t, x)$, $S(t, x)$, and $R(t, x)$ can again be derived by standard techniques and are listed in Ref. [13]. They result in Feynman diagrams like the samples shown in Fig. 1. In Ref. [13] we set up a program chain to automatically generate [17, 18] and process [19–23] these diagrams as well as to perform a reduction to master integrals [24–30] and to solve those subsequently [31–41].

For the gluon action density we then find

$$\begin{aligned} \langle E(t) \rangle|_{m=0} &= \frac{3\alpha_s}{4\pi t^2} \frac{N_A}{8} \left[1 + \frac{\alpha_s}{4\pi} e_1(\mu^2 t) + \left(\frac{\alpha_s}{4\pi} \right)^2 e_2(\mu^2 t) + \mathcal{O}(\alpha_s^3) \right], \\ e_1(z) &= e_{1,0} + \beta_0 L(z), \quad e_2(z) = e_{2,0} + (2\beta_0 e_{1,0} + \beta_1) L(z) + \beta_0^2 L^2(z), \end{aligned} \quad (16)$$

where $\alpha_s = \alpha_s(\mu)$, with μ the renormalization scale,

$$L(z) \equiv \ln(2z) + \gamma_E, \quad (17)$$

and with the $\overline{\text{MS}}$ coefficients β_0 , β_1 . For the non-logarithmic coefficients $e_{i,j}$ we find

$$\begin{aligned} e_{0,0} &= 1, \quad e_{1,0} = \left(\frac{52}{9} + \frac{22}{3} \ln 2 - 3 \ln 3 \right) C_A - \frac{8}{9} T_R n_f \\ e_{2,0} &= 27.9784 C_A^2 - (31.5652 \dots) C_A T_R n_f + \left(16\zeta(3) - \frac{43}{3} \right) C_F T_R n_f + \left(\frac{8\pi^2}{27} - \frac{80}{81} \right) T_R^2 n_f^2, \end{aligned} \quad (18)$$

where $\zeta(3) = 1.20206\dots$. The *next-to-leading order* (NLO) coefficient e_1 was first evaluated in Ref. [3] and the NNLO coefficient in Ref. [31]. The three dots in the coefficient of $C_A T_R n_f$

indicate that we were able to obtain the expression in analytical form in Ref. [13]. Our estimate of the numerical accuracy for the C_A^2 coefficient is at least six digits beyond the four decimal places shown here.

Since these VEVs are formally independent of the renormalization scale μ , the residual scale dependence can be used to study the behavior of the perturbative expansion. As shown in Fig. 2 for $\langle E(t) \rangle$, it is well behaved at high energies and still decent around a central scale of 3 GeV. For a detailed discussion and results for $\langle \dot{S}(t) \rangle$ and $\zeta_\chi(t, \mu)$ we refer to Ref. [13].




Figure 2: Renormalization scale dependence of $t^2\langle E(t) \rangle$ in QCD for two different central scales $\mu_0 = e^{-\gamma_E/2}/\sqrt{2t}$. See Ref. [13] for details.

The proportionality of $\langle E(t) \rangle$ to α_s suggests to define a *gradient flow coupling*

$$\alpha_{\text{GF}}(t) \equiv \frac{1}{\mathcal{N}} \frac{32\pi}{3N_A} t^2 \langle E(t) \rangle_{\text{lattice}}, \quad (19)$$

based on the determination of $\langle E(t) \rangle_{\text{lattice}}$ through lattice simulations, where \mathcal{N} accounts for boundary conditions [42–45]. Our perturbative result in Eq. (16) then might help in the extrapolation to infinite volume.

4 Flowed Operator Product Expansion

A powerful concept in the gradient flow formalism is the flowed OPE [8–10]. Consider a set of operators $\mathcal{O}_i(x)$ and a corresponding set of flowed operators $\tilde{\mathcal{O}}_i(t, x)$ which are constructed from flowed fields. They are related by the *small-flow-time expansion*

$$\tilde{\mathcal{O}}_i(t, x) = \sum_j \zeta_{ij}(t) \mathcal{O}_j(x) + O(t) \quad (20)$$

with the flow-time dependent mixing matrix $\zeta_{ij}(t)$ [6]. By inverting Eq. (20) one can then express any linear combination of the $\mathcal{O}_i(x)$ through their flowed counterparts:

$$T = \sum_i C_i \mathcal{O}_i = \sum_{i,j} C_i \zeta_{ij}^{-1} \tilde{\mathcal{O}}_j + O(t) \equiv \sum_j \tilde{C}_j \tilde{\mathcal{O}}_j + O(t), \quad (21)$$

where C_i are the Wilson coefficients for the object T . Since the flowed operators do not require renormalization beyond field and coupling renormalization [6], the r.h.s. is scheme independent. Thus, one can directly relate T in different schemes, for example lattice and perturbative schemes, by employing the flowed OPE. First, it was applied to the EMT [8, 9, 15] which led

to promising thermodynamical results [46–54]. Other applications include charge conjugation parity violating operators for the nucleon electric dipole moment [55] or the electroweak Hamiltonian [56]. We now apply it to *vacuum polarization functions* (VPFs).

5 Hadronic Vacuum Polarization using Gradient Flow

VPFs for (axial-)vector and (pseudo-)scalar particles are important objects in QCD. Through the optical theorem, their imaginary part is directly related to physical observables such as the decay rates of the Z - or the Higgs boson, or the hadronic R-ratio. Moreover, VPFs also contribute indirectly to physical observables such as anomalous magnetic moments [57, 58], the definition of short-distance quark masses [59], or hadronic contributions to the coupling of Quantum Electrodynamics [60, 61]. However, the latter applications involve an integration of the VPFs over the non-perturbative regime. They are typically computed from experimental data with the help of dispersion relations. First-principle lattice calculations have started to become competitive with these dispersive approaches only very recently. However, for the prominent topic of the hadronic vacuum polarization contribution to the muon's anomalous magnetic moment, the two approaches lead to incompatible results [62].

The perturbative and non-perturbative regimes of VPFs can explicitly be demonstrated through the OPE (see, e.g., Ref. [63]):

$$T(Q) \equiv \int d^4x e^{iQx} \langle T j(x) j(0) \rangle \stackrel{Q^2 \rightarrow \infty}{\sim} \sum_{k,n} C_n^{(k),B}(Q) \langle \mathcal{O}_n^{(k)}(x=0) \rangle, \quad (22)$$

where $j(x)$ generically stands for a scalar, pseudo-scalar, vector, axial-vector, or tensor current, and k labels the mass dimension. In principle, the coefficients $C_n^{(k),B}$ on the r.h.s. of Eq. (22) depend on the quantum numbers of the currents, but we suppress such indices in the following. We furthermore assume that possible global divergences are subtracted off of $T(Q)$.

Up to mass dimension two, only operators proportional to unity contribute to physical matrix elements of QCD. Explicitly, they read

$$\mathcal{O}_1^{(0)} \equiv \mathcal{O}^{(0)} = \mathbb{1}, \quad \mathcal{O}_1^{(2)} \equiv \mathcal{O}^{(2)} = m_B^2 \mathbb{1}, \quad (23)$$

where m_B is the bare mass of the n_h degenerate massive quarks. Therefore, the Wilson coefficients

$$C_1^{(0)} \equiv C^{(0)} \equiv C^{(0),B}, \quad C_1^{(2)} \equiv C^{(2)} \equiv Z_m^2 C^{(2),B} \quad (24)$$

are UV-finite, where Z_m is the $\overline{\text{MS}}$ renormalization constant of the quark mass. At mass dimension four we choose

$$\mathcal{O}_1^{(4)} \equiv \mathcal{O}_1 = \frac{1}{g_B^2} F_{\mu\nu}^a F_{\mu\nu}^a, \quad \mathcal{O}_2^{(4)} \equiv \mathcal{O}_2 = \sum_{f=1}^{n_f} \bar{\psi}_f \not{\partial}^F \psi_f, \quad \mathcal{O}_3^{(4)} \equiv \mathcal{O}_3 = m_B^4 \mathbb{1} \quad (25)$$

as basis of operators. Higher dimensional operators are neglected in the following.

Matrix elements of the dimension-four operators are divergent in general. However, by defining renormalized operators \mathcal{O}_n^R as linear combinations among them, physical matrix elements as well as the Wilson coefficients become finite, i.e.

$$\mathcal{O}_n^R = \sum_k Z_{nk} \mathcal{O}_k, \quad C_n = \sum_m C_m^B (Z^{-1})_{mn}, \quad (26)$$

where $C_n^B \equiv C_n^{(4),B}$, cf. Eqs. (22) and (25). Since the operators of Eq. (25) are part of the QCD Lagrangian, the renormalization matrix Z can be expressed in terms of the anomalous dimensions of QCD [64, 65].

To derive the flowed OPE, we introduce the flowed operators as

$$\begin{aligned}\tilde{\mathcal{O}}_1(t, x) &= \frac{Z_s}{g_B^2} G_{\mu\nu}^a(t, x) G_{\mu\nu}^a(t, x) = \frac{4}{\hat{\mu}^{2\epsilon} g^2} E(t, x), \\ \tilde{\mathcal{O}}_2(t, x) &= \dot{Z}_\chi \sum_{f=1}^{n_f} \bar{\chi}_f(t, x) \overleftrightarrow{\not{D}}^F(t, x) \chi_f(t, x) = \dot{R}(t, x), \quad \tilde{\mathcal{O}}_3(t, x) = m^4 \mathbb{1},\end{aligned}\tag{27}$$

where $E(t, x)$ and $\dot{R}(t, x)$ are the composite operators already introduced in Section 3. The small-flow-time expansion in Eq. (20) allows us to relate the regular QCD operators and coefficients with their flowed counterparts through

$$\tilde{\mathcal{O}}_n(t) = \zeta_n^{(0)}(t) \mathbb{1} + \zeta_n^{(2)}(t) m^2 \mathbb{1} + \sum_k \zeta_{nk}(t) \mathcal{O}_k^R + O(t),\tag{28}$$

where $\zeta_n^{(2)}(t)$ and $\zeta_{nk}(t)$ are the renormalized, finite mixing coefficients. Inverting Eq. (28) yields

$$\mathcal{O}_n^R = \sum_k \zeta_{nk}^{-1}(t) \tilde{\mathcal{O}}_k(t) + O(t), \quad \tilde{\mathcal{O}}_n(t) \equiv \tilde{\mathcal{O}}_n(t) - \zeta_n^{(0)}(t) \mathbb{1} - \zeta_n^{(2)}(t) m^2 \mathbb{1},\tag{29}$$

which leads to the flowed OPE for the current correlator:

$$T(Q) \xrightarrow{Q^2 \rightarrow \infty} \tilde{C}^{(0)}(Q^2, t) + \tilde{C}^{(2)}(Q^2, t) m^2 + \sum_n \tilde{C}_n(Q^2, t) \langle \tilde{\mathcal{O}}_n(t) \rangle + O(t).\tag{30}$$

The flowed Wilson coefficients are related to the regular Wilson coefficients through

$$\tilde{C}_n(Q^2, t) = \sum_k C_k(Q^2) \zeta_{kn}^{-1}(t), \quad \tilde{C}^{(0,2)}(Q^2, t) = C^{(0,2)}(Q^2) - \sum_n \tilde{C}_n(Q^2, t) \zeta_n^{(0,2)}(t).\tag{31}$$

The regular Wilson coefficients $C^{(0)}$ and $C^{(2)}$ are given by the first two terms in m^2/Q^2 of the large- Q^2 expansion of the VPFs. Through the required order, they can be found in Ref. [66] for vector-, in Ref. [67] for axial-vector-, and in Ref. [68] for scalar- and pseudo-scalar currents, for example. The dimension-four coefficients can be found in Refs. [69, 70].³

In Ref. [14] we determined the mixing matrix ζ through NNLO with the help of the method of projectors [71, 72]. Most of the matrix elements can already be extracted from the calculation of the VEVs in Section 3 (or rather Ref. [13]) as well as from the calculation of the EMT in Ref. [15]. The remaining elements correspond to higher-order corrections in the bare mass to the VEVs. By combining ζ with the known results for the regular Wilson coefficients, one can determine the flowed coefficients of Eq. (31) to the same order. Together with an evaluation of the flowed operator matrix elements on the lattice, the VPFs can be extracted and used in the determination of various physical quantities.

6 Conclusion

In this contribution we discussed the perturbative gradient flow and stressed its powerful renormalization properties. Then, we outlined the calculation of some VEVs of gauge-invariant operators at finite flow time which enable the construction of a gradient flow coupling. Afterwards, we discussed the flowed OPE which can be used to replace regular operators by their better behaved flowed counterparts and applied it to VPFs which might lead to new results for quantities like anomalous magnetic moments from lattice simulations.

³Since the latter reference is only available in German, they have also been included in Ref. [14].

Acknowledgements

We thank Johannes Artz, Robert V. Harlander, Tobias Neumann, and Mario Prausa for their collaboration on the projects presented in this contribution. Furthermore, we thank Robert V. Harlander and Tobias Neumann for comments on the manuscript.

Funding information The research projects presented in this contribution were supported by the *Deutsche Forschungsgemeinschaft* (DFG, German Research Foundation) through grant [386986591](#). Some of the calculations in Section 3 were performed with computing resources granted by RWTH Aachen University under project “rwth0244”.

References

- [1] R. Narayanan and H. Neuberger, *Infinite N phase transitions in continuum Wilson loop operators*, J. High Energy Phys. **03**, 064 (2006), doi:[10.1088/1126-6708/2006/03/064](https://doi.org/10.1088/1126-6708/2006/03/064).
- [2] M. Lüscher, *Trivializing Maps, the Wilson Flow and the HMC Algorithm*, Commun. Math. Phys. **293**, 899 (2009), doi:[10.1007/s00220-009-0953-7](https://doi.org/10.1007/s00220-009-0953-7).
- [3] M. Lüscher, *Properties and uses of the Wilson flow in lattice QCD*, J. High Energy Phys. **08**, 071 (2010), doi:[10.1007/JHEP08\(2010\)071](https://doi.org/10.1007/JHEP08(2010)071).
- [4] Sz. Borsányi et al. , *High-precision scale setting in lattice QCD*, J. High Energy Phys. **09**, 010 (2012), doi:[10.1007/JHEP09\(2012\)010](https://doi.org/10.1007/JHEP09(2012)010).
- [5] R. Sommer, *Scale setting in lattice QCD*, Proc. Sci. **187**, 015 (2014), doi:[10.22323/1.187.0015](https://doi.org/10.22323/1.187.0015).
- [6] M. Lüscher and P. Weisz, *Perturbative analysis of the gradient flow in non-abelian gauge theories*, J. High Energy Phys. **02**, 051 (2011), doi:[10.1007/JHEP02\(2011\)051](https://doi.org/10.1007/JHEP02(2011)051).
- [7] M. Dalla Brida, *Past, present, and future of precision determinations of the QCD coupling from lattice QCD*, Eur. Phys. J. A **57**, 66 (2021), doi:[10.1140/epja/s10050-021-00381-3](https://doi.org/10.1140/epja/s10050-021-00381-3).
- [8] H. Suzuki, *Energy-momentum tensor from the Yang–Mills gradient flow*, Progr. Theor. Exp. Phys. 083B03 (2013), doi:[10.1093/ptep/ptt059](https://doi.org/10.1093/ptep/ptt059).
- [9] H. Makino and H. Suzuki, *Lattice energy-momentum tensor from the Yang–Mills gradient flow—incusion of fermion fields*, Progr. Theor. Exp. Phys. 063B02 (2014), doi:[10.1093/ptep/ptu070](https://doi.org/10.1093/ptep/ptu070).
- [10] C. Monahan and K. Orginos, *Locally smeared operator product expansions in scalar field theory*, Phys. Rev. D **91**, 074513 (2015), doi:[10.1103/PhysRevD.91.074513](https://doi.org/10.1103/PhysRevD.91.074513).
- [11] M. Dalla Brida and M. Lüscher, *SMD-based numerical stochastic perturbation theory*, Eur. Phys. J. C **77**, 308 (2017), doi:[10.1140/epjc/s10052-017-4839-0](https://doi.org/10.1140/epjc/s10052-017-4839-0).
- [12] M. Lüscher, *Chiral symmetry and the Yang–Mills gradient flow*, J. High Energy Phys. **04**, 123 (2013), doi:[10.1007/JHEP04\(2013\)123](https://doi.org/10.1007/JHEP04(2013)123).
- [13] J. Artz, R. V. Harlander, F. Lange, T. Neumann and M. Prausa, *Results and techniques for higher order calculations within the gradient-flow formalism*, J. High Energy Phys. **06**, 121 (2019), doi:[10.1007/JHEP06\(2019\)121](https://doi.org/10.1007/JHEP06(2019)121).

- [14] R. V. Harlander, F. Lange and T. Neumann, *Hadronic vacuum polarization using gradient flow*, J. High Energy Phys. **08**, 109 (2020), doi:[10.1007/JHEP08\(2020\)109](https://doi.org/10.1007/JHEP08(2020)109).
- [15] R. V. Harlander, Y. Kluth and F. Lange, *The two-loop energy-momentum tensor within the gradient-flow formalism*, Eur. Phys. J. C **78**, 944 (2018), doi:[10.1140/epjc/s10052-018-6415-7](https://doi.org/10.1140/epjc/s10052-018-6415-7).
- [16] J. P. Ellis, *TikZ-Feynman: Feynman diagrams with TikZ*, Comput. Phys. Commun. **210**, 103 (2017), doi:[10.1016/j.cpc.2016.08.019](https://doi.org/10.1016/j.cpc.2016.08.019).
- [17] P. Nogueira, *Automatic Feynman Graph Generation*, J. Comput. Phys. **105**, 279 (1993), doi:[10.1006/jcph.1993.1074](https://doi.org/10.1006/jcph.1993.1074).
- [18] P. Nogueira, *Abusing qgraf*, Nucl. Instrum. Meth. Phys. Res. Sec. A **559**, 220 (2006), doi:[10.1016/j.nima.2005.11.151](https://doi.org/10.1016/j.nima.2005.11.151).
- [19] R. Harlander, T. Seidensticker and M. Steinhauser, *Corrections of $\mathcal{O}(\alpha\alpha_s)$ to the decay of the Z boson into bottom quarks*, Phys. Lett. B **426**, 125 (1998), doi:[10.1016/S0370-2693\(98\)00220-2](https://doi.org/10.1016/S0370-2693(98)00220-2).
- [20] T. Seidensticker, *Automatic application of successive asymptotic expansions of Feynman diagrams*, arXiv:[hep-ph/9905298](https://arxiv.org/abs/hep-ph/9905298).
- [21] J. A. M. Vermaasen, *New features of FORM*, arXiv:[math-ph/0010025](https://arxiv.org/abs/math-ph/0010025).
- [22] J. Kuipers, T. Ueda, J. A. M. Vermaasen and J. Vollinga, *FORM version 4.0*, Comput. Phys. Commun. **184**, 1453 (2013), doi:[10.1016/j.cpc.2012.12.028](https://doi.org/10.1016/j.cpc.2012.12.028).
- [23] T. van Ritbergen, A. N. Schellekens and J. A. M. Vermaasen, *Group theory factors for Feynman diagrams*, Int. J. Mod. Phys. A **14**, 41 (1999), doi:[10.1142/S0217751X99000038](https://doi.org/10.1142/S0217751X99000038).
- [24] F. V. Tkachov, *A theorem on analytical calculability of 4-loop renormalization group functions*, Phys. Lett. B **100**, 65 (1981), doi:[10.1016/0370-2693\(81\)90288-4](https://doi.org/10.1016/0370-2693(81)90288-4).
- [25] K. G. Chetyrkin and F. V. Tkachov, *Integration by parts: The algorithm to calculate β -functions in 4 loops*, Nucl. Phys. B **192**, 159 (1981), doi:[10.1016/0550-3213\(81\)90199-1](https://doi.org/10.1016/0550-3213(81)90199-1).
- [26] S. Laporta, *High-precision calculation of multiloop Feynman integrals by difference equations*, Int. J. Mod. Phys. A **15**, 5087 (2000), doi:[10.1016/S0217-751X\(00\)00215-7](https://doi.org/10.1016/S0217-751X(00)00215-7).
- [27] P. Maierhöfer, J. Usovitsch and P. Uwer, *Kira—A Feynman integral reduction program*, Comput. Phys. Commun. **230**, 99 (2018), doi:[10.1016/j.cpc.2018.04.012](https://doi.org/10.1016/j.cpc.2018.04.012).
- [28] J. Klappert, F. Lange, P. Maierhöfer and J. Usovitsch, *Integral reduction with Kira 2.0 and finite field methods*, Comput. Phys. Commun. **266**, 108024 (2021), doi:[10.1016/j.cpc.2021.108024](https://doi.org/10.1016/j.cpc.2021.108024).
- [29] J. Klappert and F. Lange, *Reconstructing rational functions with FireFly*, Comput. Phys. Commun. **247**, 106951 (2020), doi:[10.1016/j.cpc.2019.106951](https://doi.org/10.1016/j.cpc.2019.106951).
- [30] J. Klappert, S. Y. Klein and F. Lange, *Interpolation of dense and sparse rational functions and other improvements in FireFly*, Comput. Phys. Commun. **264**, 107968 (2021), doi:[10.1016/j.cpc.2021.107968](https://doi.org/10.1016/j.cpc.2021.107968).
- [31] R. V. Harlander and T. Neumann, *The perturbative QCD gradient flow to three loops*, J. High Energy Phys. **06**, 161 (2016), doi:[10.1007/JHEP06\(2016\)161](https://doi.org/10.1007/JHEP06(2016)161).

- [32] T. Binoth and G. Heinrich, *An automatized algorithm to compute infrared divergent multi-loop integrals*, Nucl. Phys. B **585**, 741 (2000), doi:[10.1016/S0550-3213\(00\)00429-6](https://doi.org/10.1016/S0550-3213(00)00429-6).
- [33] T. Binoth and G. Heinrich, *Numerical evaluation of multi-loop integrals by sector decomposition*, Nucl. Phys. B **680**, 375 (2004), doi:[10.1016/j.nuclphysb.2003.12.023](https://doi.org/10.1016/j.nuclphysb.2003.12.023).
- [34] A. V. Smirnov and M. N. Tentyukov, *Feynman Integral Evaluation by a Sector decompositiOn Approach (FIESA)*, Comput. Phys. Commun. **180**, 735 (2009), doi:[10.1016/j.cpc.2008.11.006](https://doi.org/10.1016/j.cpc.2008.11.006).
- [35] A. V. Smirnov, V. A. Smirnov and M. Tentyukov, *FIESA 2: Parallelizeable multiloop numerical calculations*, Comput. Phys. Commun. **182**, 790 (2011), doi:[10.1016/j.cpc.2010.11.025](https://doi.org/10.1016/j.cpc.2010.11.025).
- [36] A. V. Smirnov, *FIESA 3: Cluster-parallelizable multiloop numerical calculations in physical regions*, Comput. Phys. Commun. **185**, 2090 (2014), doi:[10.1016/j.cpc.2014.03.015](https://doi.org/10.1016/j.cpc.2014.03.015).
- [37] A. C. Genz and A. A. Malik, *An Imbedded Family of Fully Symmetric Numerical Integration Rules*, SIAM J. Numer. Anal. **20**, 580 (1983), doi:[10.1137/0720038](https://doi.org/10.1137/0720038).
- [38] Wolfram Research, Inc., *Mathematica, Version 11.3*, Champaign, IL (2018).
- [39] T. Huber and D. Maître, *HypExp, a Mathematica package for expanding hypergeometric functions around integer-valued parameters*, Comput. Phys. Commun. **175**, 122 (2006), doi:[10.1016/j.cpc.2006.01.007](https://doi.org/10.1016/j.cpc.2006.01.007).
- [40] T. Huber and D. Maître, *HypExp 2, Expanding hypergeometric functions about half-integer parameters*, Comput. Phys. Commun. **178**, 755 (2008), doi:[10.1016/j.cpc.2007.12.008](https://doi.org/10.1016/j.cpc.2007.12.008).
- [41] E. Panzer, *Algorithms for the symbolic integration of hyperlogarithms with applications to Feynman integrals*, Comput. Phys. Commun. **188**, 148 (2015), doi:[10.1016/j.cpc.2014.10.019](https://doi.org/10.1016/j.cpc.2014.10.019).
- [42] M. Lüscher, *Step scaling and the Yang-Mills gradient flow*, J. High Energy Phys. **06**, 105 (2014), doi:[10.1007/JHEP06\(2014\)105](https://doi.org/10.1007/JHEP06(2014)105).
- [43] Z. Fodor, K. Holland, J. Kuti, D. Nogradi and C. Him Wong, *A new method for the beta function in the chiral symmetry broken phase*, EPJ Web Conf. **175**, 08027 (2018), doi:[10.1051/epjconf/201817508027](https://doi.org/10.1051/epjconf/201817508027).
- [44] A. Hasenfratz and O. Witzel, *Continuous renormalization group β function from lattice simulations*, Phys. Rev. D **101**, 034514 (2020), doi:[10.1103/PhysRevD.101.034514](https://doi.org/10.1103/PhysRevD.101.034514).
- [45] K. Holland, Z. Fodor, J. Kuti, D. Nogradi and C.-H. Wong, *Case studies of near-conformal beta-functions*, Proc. Sci. **363**, 121 (2019), doi:[10.22323/1.363.0121](https://doi.org/10.22323/1.363.0121).
- [46] M. Asakawa, T. Hatsuda, E. Itou, M. Kitazawa and H. Suzuki, *Thermodynamics of SU(3) gauge theory from gradient flow on the lattice*, Phys. Rev. D **90**, 011501 (2014), doi:[10.1103/PhysRevD.90.011501](https://doi.org/10.1103/PhysRevD.90.011501).
- [47] Y. Taniguchi, S. Ejiri, R. Iwami, K. Kanaya, M. Kitazawa, H. Suzuki, T. Umeda and N. Wakabayashi, *Exploring $N_f = 2+1$ QCD thermodynamics from the gradient flow*, Phys. Rev. D **96**, 014509 (2017), doi:[10.1103/PhysRevD.96.014509](https://doi.org/10.1103/PhysRevD.96.014509).
- [48] M. Kitazawa, T. Iritani, M. Asakawa, T. Hatsuda and H. Suzuki, *Equation of state for SU(3) gauge theory via the energy-momentum tensor under gradient flow*, Phys. Rev. D **94**, 114512 (2016), doi:[10.1103/PhysRevD.94.114512](https://doi.org/10.1103/PhysRevD.94.114512).

- [49] M. Kitazawa, T. Iritani, M. Asakawa and T. Hatsuda, *Correlations of the energy-momentum tensor via gradient flow in $SU(3)$ Yang-Mills theory at finite temperature*, Phys. Rev. D **96**, 111502 (2017), doi:[10.1103/PhysRevD.96.111502](https://doi.org/10.1103/PhysRevD.96.111502).
- [50] R. Yanagihara, T. Iritani, M. Kitazawa, M. Asakawa and T. Hatsuda, *Distribution of stress tensor around static quark–anti-quark from Yang–Mills gradient flow*, Phys. Lett. B **789**, 210 (2019), doi:[10.1016/j.physletb.2018.09.067](https://doi.org/10.1016/j.physletb.2018.09.067).
- [51] T. Iritani, M. Kitazawa, H. Suzuki and H. Takaura, *Thermodynamics in quenched QCD: energy–momentum tensor with two-loop order coefficients in the gradient-flow formalism*, Progr. Theor. Exp. Phys. 023B02 (2019), doi:[10.1093/ptep/ptz001](https://doi.org/10.1093/ptep/ptz001).
- [52] M. Kitazawa, S. Mogliacci, I. Kolbé and W. A. Horowitz, *Anisotropic pressure induced by finite-size effects in $SU(3)$ Yang-Mills theory*, Phys. Rev. D **99**, 094507 (2019), doi:[10.1103/PhysRevD.99.094507](https://doi.org/10.1103/PhysRevD.99.094507).
- [53] Y. Taniguchi, S. Ejiri, K. Kanaya, M. Kitazawa, H. Suzuki and T. Umeda, $N_f = 2+1$ QCD thermodynamics with gradient flow using two-loop matching coefficients, Phys. Rev. D **102**, 014510 (2020), doi:[10.1103/PhysRevD.102.014510](https://doi.org/10.1103/PhysRevD.102.014510).
- [54] R. Yanagihara, M. Kitazawa, M. Asakawa and T. Hatsuda, *Distribution of energy-momentum tensor around a static quark in the deconfined phase of $SU(3)$ Yang-Mills theory*, Phys. Rev. D **102**, 114522 (2020), doi:[10.1103/PhysRevD.102.114522](https://doi.org/10.1103/PhysRevD.102.114522).
- [55] M. D. Rizik, C. J. Monahan and A. Shindler, *Short flow-time coefficients of CP-violating operators*, Phys. Rev. D **102**, 034509 (2020), doi:[10.1103/PhysRevD.102.034509](https://doi.org/10.1103/PhysRevD.102.034509).
- [56] A. Suzuki, Y. Taniguchi, H. Suzuki and K. Kanaya, *Four quark operators for kaon bag parameter with gradient flow*, Phys. Rev. D **102**, 034508 (2020), doi:[10.1103/PhysRevD.102.034508](https://doi.org/10.1103/PhysRevD.102.034508).
- [57] F. Jegerlehner, *The Anomalous Magnetic Moment of the Muon*, Springer International Publishing, Cham, ISBN 9783319635750 (2017), doi:[10.1007/978-3-319-63577-4](https://doi.org/10.1007/978-3-319-63577-4).
- [58] T. Aoyama et al., *The anomalous magnetic moment of the muon in the Standard Model*, Phys. Rep. **887**, 1 (2020), doi:[10.1016/j.physrep.2020.07.006](https://doi.org/10.1016/j.physrep.2020.07.006).
- [59] K. G. Chetyrkin, J. H. Kühn, A. Maier, P. Maierhöfer, P. Marquard, M. Steinhauser and C. Sturm, *Precise charm- and bottom-quark masses: Theoretical and experimental uncertainties*, Theor. Math. Phys. **170**, 217 (2012), doi:[10.1007/s11232-012-0024-7](https://doi.org/10.1007/s11232-012-0024-7).
- [60] A. Crivellin, M. Hoferichter, C. Andrea Manzari and M. Montull, *Hadronic Vacuum Polarization: $(g - 2)_\mu$ versus Global Electroweak Fits*, Phys. Rev. Lett. **125**, 091801 (2020), doi:[10.1103/PhysRevLett.125.091801](https://doi.org/10.1103/PhysRevLett.125.091801).
- [61] A. Keshavarzi, W. J. Marciano, M. Passera and A. Sirlin, *Muon $g - 2$ and $\Delta\alpha$ connection*, Phys. Rev. D **102**, 033002 (2020), doi:[10.1103/PhysRevD.102.033002](https://doi.org/10.1103/PhysRevD.102.033002).
- [62] Sz. Borsanyi et al., *Leading hadronic contribution to the muon magnetic moment from lattice QCD*, Nature **593**, 51 (2021), doi:[10.1038/s41586-021-03418-1](https://doi.org/10.1038/s41586-021-03418-1).
- [63] C. A. Dominguez, *Analytical determination of the QCD quark masses*, Int. J. Mod. Phys. A **29**, 1430069 (2014), doi:[10.1142/S0217751X14300695](https://doi.org/10.1142/S0217751X14300695).
- [64] V. P. Spiridonov, *Anomalous Dimension of $G_{\mu\nu}^2$ and β -function*, IYAI-P-0378 (1984).

- [65] V. P. Spiridonov and K. G. Chetyrkin, *Nonleading mass corrections and renormalization of the operators $m\bar{\psi}\psi$ and $G_{\mu\nu}^2$* , Sov. J. Nucl. Phys. **47**, 522 (1988).
- [66] K. G. Chetyrkin, R. Harlander, J. H. Kühn and M. Steinhauser, *Mass corrections to the vector current correlator*, Nucl. Phys. B **503**, 339 (1997), doi:[10.1016/S0550-3213\(97\)00383-0](https://doi.org/10.1016/S0550-3213(97)00383-0).
- [67] R. Harlander and M. Steinhauser, $\mathcal{O}(\alpha_s^2)$ corrections to top quark production at e^+e^- colliders, Eur. Phys. J. C **2**, 151 (1998), doi:[10.1007/s100520050129](https://doi.org/10.1007/s100520050129).
- [68] R. Harlander and M. Steinhauser, *Higgs boson decay to top quarks at $O(\alpha_s^2)$* , Phys. Rev. D **56**, 3980 (1997), doi:[10.1103/PhysRevD.56.3980](https://doi.org/10.1103/PhysRevD.56.3980).
- [69] K. G. Chetyrkin, S. G. Gorishny and V. P. Spiridonov, *Wilson expansion for correlators of vector currents at the two-loop level: Dimension-four operators*, Phys. Lett. B **160**, 149 (1985), doi:[10.1016/0370-2693\(85\)91482-0](https://doi.org/10.1016/0370-2693(85)91482-0).
- [70] R. Harlander, *Quarkmasseneffekte in der Quantenchromodynamik und asymptotische Entwicklung von Feynman-Integralen*, Ph.D. thesis, Universität Karlsruhe (1998).
- [71] S. G. Gorishny, S. A. Larin and F. V. Tkachov, *The algorithm for OPE coefficient functions in the MS scheme*, Phys. Lett. B **124**, 217 (1983), doi:[10.1016/0370-2693\(83\)91439-9](https://doi.org/10.1016/0370-2693(83)91439-9).
- [72] S. G. Gorishny and S. A. Larin, *Coefficient functions of asymptotic operator expansions in the minimal subtraction scheme*, Nucl. Phys. B **283**, 452 (1987), doi:[10.1016/0550-3213\(87\)90283-5](https://doi.org/10.1016/0550-3213(87)90283-5).