

Non-zero transverse single spin asymmetry of very forward π^0 in polarized $p + p$ collisions at $\sqrt{s} = 510$ GeV

Minho H. Kim^{1,2*} for the RHICf Collaboration

¹ RIKEN BNL Research Center, Brookhaven National Laboratory,
Upton, New York 11973-5000, USA

² Korea University, Seoul 02841, Korea

* jipangie@korea.ac.kr



*Proceedings for the XXVIII International Workshop
on Deep-Inelastic Scattering and Related Subjects,
Stony Brook University, New York, USA, 12-16 April 2021*
doi:[10.21468/SciPostPhysProc.8](https://doi.org/10.21468/SciPostPhysProc.8)

Abstract

The RHICf experiment measured transverse single spin asymmetry of very forward ($\eta > 6$) π^0 from polarized $p + p$ collisions at $\sqrt{s} = 510$ GeV. In order to measure it precisely, we installed a new electromagnetic calorimeter at zero-degree area of the STAR experiment at the Relativistic Heavy Ion Collider (RHIC) and measured the π^0 s over the kinematic range of $x_F > 0.25$ and $0 < p_T < 1$ GeV/c in June 2017. A clear non-zero asymmetry was observed even in low $p_T < 1$ GeV/c showing a similar x_F dependence with that of the forward ($2 < \eta < 4$) π^0 . A possible diffractive contribution may need to be taken into account to explain the very forward π^0 asymmetry. RHICf-STAR combined analysis and follow-up experiment will give a clue to understand it qualitatively.



Copyright M. H. Kim *et al.*
This work is licensed under the Creative Commons
[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).
Published by the SciPost Foundation.

Received 02-08-2021

Accepted 06-04-2022

Published 12-07-2022

doi:[10.21468/SciPostPhysProc.8.060](https://doi.org/10.21468/SciPostPhysProc.8.060)



Check for
updates

1 Introduction

Transverse single spin asymmetry (A_N) is a powerful observable for understanding the spin-involved production mechanism in the polarized $p + p$ collision. It is defined by a left-right cross section asymmetry,

$$A_N = \frac{\sigma_L - \sigma_R}{\sigma_L + \sigma_R}, \quad (1)$$

where the $\sigma_{L(R)}$ is the cross section of a particle produced in the left (right) side with respect to the beam polarization.

Large non-zero asymmetries in the π^0 production have been discussed only in the quarks and gluons' degrees of freedom. It has been measured in a wide range of collision energies

[1–5] and these results have been explained by transverse momentum dependent (TMD) [6–8] and higher twist functions [9–11] in an initial or final state with the transverse motions of quarks and gluons.

Recently, further analysis results [12–14] showed a strange behavior in the forward π^0 asymmetry. Bigger asymmetry was observed in more isolated π^0 event which is connected to the diffractive process. In this analysis, the π^0 was considered as isolated one when its energy fraction to the detected electromagnetic particles was close to 1. In this case, since the π^0 carries larger energy fraction, the asymmetry nature could be biased. However, it also gives us a possibility of the diffractive contribution to the π^0 asymmetry because it is also expected that there is few particles around it when it is produced by the diffractive process, thereby the π^0 asymmetry induced by the diffractive process may be observed in this analysis. This means the non-zero asymmetry could be correlated with not only the partonic interaction but also the diffractive one.

In order to study the diffractive contribution to the π^0 asymmetry, the RHICf experiment measured the A_N in the very forward π^0 production. The diffractive process is expected to dominate when the produced particle is measured in the very forward region.

2 RHICf Experiment

We installed an electromagnetic calorimeter (RHICf detector) which had been originally developed for the LHCf experiment [15] in front of a STAR hadronic calorimeter, zero-degree




Figure 1: Schematic drawing of the RHICf detector installed in front of the ZDC. We moved the detector vertically to cover p_T from 0.0 to 1.0 GeV/c.

calorimeter (ZDC) [16], which was located 18 m away from the beam interaction point and took the data in June 2017. Figure 1 shows the schematic drawing of the experimental setup.

The RHICf detector consists of two sampling calorimeters, the large (TL, 40 mm dimension) and small (TS, 20 mm dimension) towers. Both towers are composed of 17 layers of tungsten absorber with a 44 radiation lengths in total, 16 layers of GSO plate for energy measurement, and 4 layers of GSO bars. Particle energy is reconstructed by the energy deposits of the GSO plates. Position is reconstructed by the GSO bar layers, which are covered by X-Y pairs of GSO bars with 1 mm dimension.

We measured the very forward π^0 from the polarized $p + p$ collisions at $\sqrt{s} = 510$ GeV. In order to measure the wide p_T coverage, we used the radial polarization which was the 90° -

rotated one than usual vertical polarization. We also moved the detector vertically so that the zero-degree direction of the beam faced the center of the TL, the center of the TS, and 24 mm below the center of the TS. We requested larger β^* value of 8 m and lower luminosity $\sim 10^{31} \text{ cm}^{-2} \text{ s}^{-1}$ than usual to make the systematic uncertainty by the angular beam divergence small. The β^* is an indicator for how much the beam is squeezed. With these detector positions and special settings, π^0 s over the longitudinal momentum fraction range of $x_F > 0.25$ and the transverse momentum range of $0 < p_T < 1 \text{ GeV}/c$ were measured. For the measured π^0 s, the RHICf detector has an energy resolution of 2.5%~3.5% and p_T resolution of 3.0%~4.5%.

π^0 s could be identified by measuring two decayed photons in each tower (Type-I) or both ones in same tower (Type-II). A Type-I π^0 trigger was used for the measurement of the Type-I events. It was operated when the energy deposits of three successive layers of both towers are larger than 45 MeV. Since the electromagnetic shower stops the development in the middle of the detector, only the upstream seven layers were used for this trigger. The Type-II events were measured by a high electromagnetic (high-EM) trigger. It was operated when the energy deposit of the fourth layer of either tower is larger than 500 MeV.

3 Data analysis

Due to the detector geometry, we used following luminosity formula to calculate the π^0 asymmetry

$$A_N = \frac{1}{PD_\phi} \left(\frac{N_L - RN_R}{N_L + RN_R} \right), \quad (2)$$

where P is the beam polarization, $N_{L(R)}$ is the number of detected π^0 s in the left (right) side of the beam polarization. R is the luminosity ratio of the spin directions resulting into the events to right and left sides. It was estimated using STAR beam beam counter (BBC) [17] and vertex




Figure 2: Reconstructed two photon invariant mass distribution of Type-I event in $x_F > 0.25$ and $0.0 < p_T < 1.0 \text{ GeV}/c$.

position detector (VPD) [18]. D_ϕ is a correction factor to compensate the diluted asymmetry by the azimuthal angle distribution of the detected π^0 s. Only the Type-I π^0 triggered events

were used for the asymmetry calculation of the Type-I events and high-EM triggered events for the Type-II events to be free from the effect of the different detection efficiencies.

π^0 s to be analyzed were identified using invariant mass distribution of two detected photons. Figure 2 shows the reconstructed invariant mass distribution. In this distribution, a clear π^0 peak (blue dashed line) was observed in the widely smeared background events (black filled histogram). The background mostly comes from two accidental photons from different π^0 s. Superposition of the Gaussian (for describing the π^0 events) and the 6th order of polynomial (for describing the background events) function was used to fit the invariant mass distribution. 3σ width of the Gaussian peak was considered as the final π^0 candidate. The effect of the background events included in the final π^0 sample was subtracted using the following equation

$$A_N^S = \left(1 + \frac{N_B}{N_S}\right)A_N^{S+B} - \left(\frac{N_B}{N_S}\right)A_N^B, \quad (3)$$

where A_N^{S+B} , A_N^S , and A_N^B are the estimated asymmetries in the signal+background, signal only and background only regions. The A_N^B was estimated using the events whose invariant mass was further than 5σ of the Gaussian peak and the signal to background ratio, $\frac{N_B}{N_S}$, was calculated from the fit result.

The π^0 asymmetry was calculated as functions of x_F and p_T to study their correlation. Systematic uncertainties by the polarization and beam center estimations, and the background subtraction were included. The effect of the smearing due to the finite x_F and p_T resolutions were studied by GEANT4 using the single π^0 beam which was artificially weighted to reproduce the asymmetry, but they were negligible. For each bin divided by selected x_F and p_T values, more than 90% of migrated events were from $\delta x_F < 0.025$ and $\delta p_T < 0.035$ GeV/c of the bin boundaries. The differences between the calculated and true $\langle x_F \rangle$, $\langle p_T \rangle$, A_N values of each bin were less than 0,008, 0.009 GeV/c, and 0.0015, respectively.

To find any missing systematic uncertainties, a "bunch shuffling" analysis was proceeded. It is done by randomizing the spin patterns and reconstructing the A_N again. Since the spin patterns are randomized, ideally, the reconstructed A_N should be fluctuated around 0 with its statistical uncertainty if there is no systematic source missed. The bunch shuffled asymmetries were consistent with zero in the comparable fluctuation with the statistical uncertainties. We conclude that there are no noticeable false asymmetries in this analysis.

4 Results

Figure 3 shows the A_N of very forward π^0 as functions of x_F and p_T . Increasing A_N as a function of p_T up to ~ 0.2 is clearly shown in Figure 3 (a) and the corresponding p_T range is where the diffractive process dominates. In Figure 3 (b), the backward asymmetries which are the ones in the opposite side of the polarized beam are consistent with zero. The forward asymmetry is also consistent with zero at very low $p_T < 0.07$ GeV/c, but it starts to increase as a function of x_F as p_T increases.

Figure 4 shows the comparison of the RHICf data with the forward π^0 asymmetries previously measured by FNAL, PHENIX, and STAR. It shows that the increasing asymmetries of the very forward π^0 get comparable with the forward π^0 ones at higher p_T though it is still smaller than 1 GeV/c. The RHICf result also shows the same x_F scaling with the forward π^0 ones. The diffractive process may contribute to the asymmetries at higher p_T than it has been expected. Since the RHICf data may be also a tail of the partonic one, more detailed analysis than inclusive one is necessary. Correlations with STAR's central detectors and Roman pots will make it possible by identifying from which process the π^0 with non-zero asymmetry




Figure 3: A_N of the very forward π^0 s as functions of (a) p_T for several x_F ranges and (b) x_F for several p_T ranges. Only forward A_N was presented in (a). Error bars represent the statistical uncertainties, and the boxes represent the systematic uncertainties.

comes. It is also desirable to investigate the same observables in the unexplored kinematic region between $0.8 \sim 2.0$ GeV/c where the fractions of both partonic and diffractive processes are comparable.

5 Conclusion

The role of the diffractive process to the π^0 asymmetry can be studied by measuring the A_N of the very forward π^0 . At low $p_T < 1$ GeV/c where the diffractive process is expected to dominate, large non-zero asymmetries increasing as functions of both x_F and p_T have been observed for the first time from the polarized $p+p$ collisions at $\sqrt{s} = 510$ GeV. The asymmetries show an approximate x_F scaling with the forward π^0 ones, which may indicate a possible diffractive contribution to the π^0 asymmetries. The RHICf result will be further investigated by RHICf-STAR combined analysis and a follow-up experiment, RHICf-II.




Figure 4: Comparison of the RHICf data with the previously measured A_N of the forward π^0 s as a function of x_F .

6 Future Prospect

In order to more deeply understand the RHICf result, RHICf-STAR combined analysis has been started. Using STAR detectors covering each η region, the type of the production mechanism is expected to be identified.

A follow-up experiment (RHICf-II) is also being prepared. To measure more various particles and for higher statistics, a larger detector of 8 cm \times 18 cm dimension is planned to be constructed. Technology of ALICE FoCal-E [19] will be transferred to the construction of the RHICf-II detector. Currently, we're optimizing the definite design of the detector and planning to take the data in 2024 at STAR.

Funding information This program is partly supported by the U.S.-Japan Science and Technology Cooperation Program in High Energy Physics, JSPS KAKENHI (No. JP26247037 and No. JP18H01227), the joint research program of the Institute for Cosmic Ray Research (ICRR), University of Tokyo, and the National Research Foundation of Korea (No. 2016R1A2B2008505 and No. 2018R1A5A1025563), and "UNICT 2020-22 Linea 2" program, University of Catania.

References

- [1] A. Adare et al. (PHENIX Collaboration), *Measurement of transverse-single-spin asymmetries for midrapidity and forward-rapidity production of hadrons in polarized $p+p$ collisions at $\sqrt{s} = 200$ and 62.4 GeV*, Phys. Rev. D **90**, 012006 (2014), doi:[10.1103/PhysRevD.90.012006](https://doi.org/10.1103/PhysRevD.90.012006).
- [2] B. I. Abelev et al. (STAR Collaboration), *Forward Neutral Pion Transverse Single Spin Asymmetries in $p+p$ Collisions at $\sqrt{s} = 200$ GeV*, Phys. Rev. Lett. **101**, 222001 (2008), doi:[10.1103/PhysRevLett.101.222001](https://doi.org/10.1103/PhysRevLett.101.222001).
- [3] D. L. Adams et al. (E581 and E704 Collaborations), *Comparison of spin asymmetries and cross-sections in π^0 production by 200-GeV polarized anti-protons and protons*, Phys. Lett. B **261**, 201 (1991), doi:[10.1016/0370-2693\(91\)91351-U](https://doi.org/10.1016/0370-2693(91)91351-U).

- [4] B. E. Bonner et al., *Analyzing Power Measurement in Inclusive π_0 Production at High x_F* , Phys. Rev. Lett. **61**, 1918 (1988), doi:[10.1103/PhysRevLett.61.1918](https://doi.org/10.1103/PhysRevLett.61.1918).
- [5] R. D. Klem, J. E. Bowers, H. W. Courant, H. Kagan, M. L. Marshak, E. A. Peterson, K. Ruddick, W. H. Dragoset and J. B. Roberts, *Measurement of Asymmetries of Inclusive Pion Production in Proton Proton Interactions at 6-GeV/c and 11.8-GeV/c*, Phys. Rev. Lett. **36**, 929 (1976), doi:[10.1103/PhysRevLett.36.929](https://doi.org/10.1103/PhysRevLett.36.929).
- [6] D. Sivers, *Single-spin production asymmetries from the hard scattering of pointlike constituents*, Phys. Rev. D **41**, 83 (1990), doi:[10.1103/PhysRevD.41.83](https://doi.org/10.1103/PhysRevD.41.83).
- [7] J. Collins, *Fragmentation of transversely polarized quarks probed in transverse momentum distributions*, Nucl. Phys. B **396**, 161 (1993), doi:[10.1016/0550-3213\(93\)90262-N](https://doi.org/10.1016/0550-3213(93)90262-N).
- [8] J. C. Collins, S. F. Heppelmann and G. A. Ladinsky, *Measuring transversity densities in singly polarized hadron-hadron and lepton-hadron collisions*, Nucl. Phys. B **420**, 565 (1994), doi:[10.1016/0550-3213\(94\)90078-7](https://doi.org/10.1016/0550-3213(94)90078-7).
- [9] J. Qiu and G. Sterman, *Single transverse spin asymmetries in direct photon production*, Nucl. Phys. B **378**, 52 (1992), doi:[10.1016/0550-3213\(92\)90003-T](https://doi.org/10.1016/0550-3213(92)90003-T).
- [10] H. Eguchi, Y. Koike and K. Tanaka, *Twist-3 formalism for single transverse spin asymmetry reexamined: Semi-inclusive deep inelastic scattering*, Nucl. Phys. B **763**, 198 (2007), doi:[10.1016/j.nuclphysb.2006.11.016](https://doi.org/10.1016/j.nuclphysb.2006.11.016).
- [11] K. Kanazawa, Y. Koike, A. Metz and D. Pitonyak, *Towards an explanation of transverse single-spin asymmetries in proton-proton collisions: The role of fragmentation in collinear factorization*, Phys. Rev. D **89**, 111501 (2014), doi:[10.1103/PhysRevD.89.111501](https://doi.org/10.1103/PhysRevD.89.111501).
- [12] S. Heppelmann et al. (STAR Collaboration), *Large p_T Forward Transverse Single Spin Asymmetries of π^0 Mesons at $\sqrt{s} = 200$ and 500 GeV from STAR*, Proc. Sci. **191**, 240 (2013), doi:[10.22323/1.191.0240](https://doi.org/10.22323/1.191.0240).
- [13] M. M. Mondal et al., (STAR Collaboration), *Measurement of the Transverse Single-Spin Asymmetries for π^0 and Jet-like Events at Forward Rapidities at STAR in $p + p$ Collisions at $\sqrt{s} = 510$ GeV*, Proc. Sci. **203**, 216 (2014), doi:[10.22323/1.203.0216](https://doi.org/10.22323/1.203.0216).
- [14] J. Adam et al. (STAR Collaboration), *Measurement of transverse single-spin asymmetries of π^0 and electromagnetic jets at forward rapidity in 200 and 500 GeV transversely polarized proton-proton collisions*, Phys. Rev. D **103**, 092009, doi:[10.1103/PhysRevD.103.092009](https://doi.org/10.1103/PhysRevD.103.092009).
- [15] O. Adriani et al. (LHCf Collaboration), *Measurements of longitudinal and transverse momentum distributions for neutral pions in the forward-rapidity region with the LHCf detector*, Phys. Rev. D **94**, 032007 (2016), doi:[10.1103/PhysRevD.94.032007](https://doi.org/10.1103/PhysRevD.94.032007).
- [16] C. Adler, A. Denisov, E. Garcia, M. Murray, H. Stroebele and S. White, *The RHIC zero degree calorimeters*, Nucl. Instrum. Meth. A **470**, 488 (2001), doi:[10.1016/S0168-9002\(01\)00627-1](https://doi.org/10.1016/S0168-9002(01)00627-1).
- [17] C. A. Whitten Jr. et al. (STAR Collaboration), *The Beam-Beam Counter: A Local Polarimeter at STAR*, <https://www.star.bnl.gov/~eca/LocalPol>.
- [18] W. J. Llope et al., *The STAR Vertex Position Detector*, Nucl. Instrum. Methods Phys. Res., Sect. A **759**, 23 (2014), doi:[10.1016/j.nima.2014.04.080](https://doi.org/10.1016/j.nima.2014.04.080).
- [19] ALICE Collaboration, *Letter of Intent: A Forward Calorimeter (FoCal) in the ALICE experiment*, CERN-LHCC-2020-009, <https://cds.cern.ch/record/2719928>.