

Non-Standard Neutrino Interactions and Neutral Gauge Bosons

Julian Heeck^{1,2*}, Manfred Lindner³, Werner Rodejohann³, Stefan Vogl³

1 Service de Physique Théorique, Université Libre de Bruxelles, Boulevard du Triomphe, CP225, 1050 Brussels, Belgium

2 Department of Physics and Astronomy, University of California, Irvine, CA 92697-4575, USA

3 Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany

* Corresponding Author: julian.heeck@uci.edu

February 26, 2019

1 Abstract

2 We investigate Non-Standard Neutrino Interactions (NSI) arising from a flavor-sensitive Z' boson of a new $U(1)'$ symmetry. **3** We compare the limits from neutrino oscillations, coherent elastic neutrino–nucleus scattering, and Z' searches **4** at different beam and collider experiments for a variety of straightforward **5** anomaly-free $U(1)'$ models generated by linear combinations of $B - L$ and **6** lepton-family-number differences $L_\alpha - L_\beta$. Depending on the flavor structure **7** of those models it is easily possible to avoid NSI signals in long-baseline neu- **8** trino oscillation experiments or change the relative importance of the various **9** experimental searches. We also point out that kinetic $Z-Z'$ mixing gives van- **10** ishing NSI in long-baseline experiments if a direct coupling between the $U(1)'$ **11** gauge boson and matter is absent. In contrast, $Z-Z'$ mass mixing generates **12** such NSI, which in turn means that there is a Higgs multiplet charged under **13** both the Standard Model and the new $U(1)'$ symmetry. **14**

15

16 Contents

| | | |
|----|---|-----------|
| 17 | 1 Introduction | 2 |
| 18 | 2 Non-Standard Neutrino Interactions: Formalism and Limits | 3 |
| 19 | 3 Calculating NSI Operators from Z' Bosons | 6 |
| 20 | 3.1 NSI without $Z-Z'$ mixing | 8 |
| 21 | 3.1.1 Electrophobic NSI | 10 |
| 22 | 3.1.2 Electrophilic NSI | 12 |
| 23 | 3.2 NSI with $Z-Z'$ mixing | 14 |
| 24 | 4 Conclusions | 17 |
| 25 | References | 18 |

26

27

28 Introduction

29 The precision era of neutrino physics implies that small effects beyond the standard
 30 paradigm of three massive neutrinos may be detected. In particular new physics with
 31 a non-trivial flavor structure deserves careful consideration since it will modify neutrino
 32 oscillation probabilities in matter and may hinder our abilities to determine the unknown
 33 neutrino parameters at upcoming neutrino oscillation facilities, as discussed in Refs. [1–7].
 34 The effects of Non-Standard neutrino Interactions (NSI) on low-energy observables are tra-
 35 ditionally parametrized by an effective Lagrangian that describes couplings of neutrinos
 36 to quarks or electrons via [8–11]

$$\mathcal{L}_{\text{eff}} \propto \epsilon_{\alpha\beta}^f (\bar{\nu}_\alpha \gamma_\mu \nu_\beta) (\bar{f} \gamma^\mu f) \quad \text{with } f = e, u, d. \quad (1)$$

37 This effective interaction is clearly not $SU(2)_L \times U(1)_Y$ gauge invariant, begging the
 38 question how this Lagrangian is generated in a complete theory and what the mass scale
 39 of that theory is. The scale is of particular relevance for phenomenological studies since
 40 only processes with a momentum transfer smaller than the mass of the new physics can be
 41 described accurately by Eq. (1). Comparing NSI limits to other experimental data that
 42 probes much higher momentum transfers then typically requires a discussion of the full
 43 UV-complete theory. Several approaches have been followed in the literature to generate
 44 and study the interactions of Eq. (1) [12–21], here we discuss the origin of non-standard
 45 interactions in flavor-sensitive $U(1)'$ models [7, 22–29]. The presence of additional Abelian
 46 symmetries is quite natural and can, for example, be motivated by Grand Unified Theories,
 47 string constructions, solutions to the hierarchy problem or extra dimensional models, see
 48 Ref. [30] for details and references.

49 We assume here the presence of a flavor-sensitive gauged $U(1)'$. In these theories the
 50 Z' belonging to the $U(1)'$ is integrated out and generates the effective NSI Lagrangian
 51 Eq. (1).¹ Limits on the strength of the interaction can be translated into limits on the Z'
 52 mass and gauge coupling. Those limits have to be compared with direct beam and collider
 53 searches, as well as neutrino–electron and elastic coherent neutrino–nucleus scattering
 54 results. In our discussion we will refer to the low-energy four-fermion operators and their
 55 impact on neutrino oscillations as NSI, while we discuss all observables with non-vanishing
 56 momentum transfer in terms of the high-energy $U(1)'$. This is the preferable notation for
 57 NSI mediated by rather light particles for which the effective NSI Lagrangian fails to
 58 describe all the relevant phenomenology.

59 The necessary ingredients for Z' -induced NSI are Z' couplings to matter, i.e. elec-
 60 trons, protons or neutrons, as well as non-universal couplings to neutrinos. Neutrino
 61 oscillations would not be affected by flavor-*universal* NSI, $\epsilon \propto \mathbb{1}$, so NSI are actually a
 62 probe of *lepton non-universality*. This is interesting in view of the accumulating hints for
 63 lepton non-universality in B meson decays (see Ref. [32] for a recent overview). While
 64 we will not attempt to make a direct connection between NSI and these tantalizing hints
 65 for new physics, it should be kept in mind as a motivation. The NSI model-building
 66 challenge is then to find realistic $U(1)'$ models with lepton non-universal Z' couplings.
 67 As is well known, the classical Standard Model (SM) Lagrangian already contains the
 68 global symmetry $U(1)_B \times U(1)_{L_e} \times U(1)_{L_\mu} \times U(1)_{L_\tau}$ associated with conserved baryon
 69 and lepton numbers. A simple extension of the SM by three right-handed neutrinos

¹The current–current structure of Eq. (1) for neutrino–quark scattering could also be induced by lep-
 toquarks. The leptoquark Yukawa couplings automatically bring the desired lepton non-universality, but
 typically also lead to lepton-flavor and even baryon-number violation, which forces them to be very weakly
 coupled. While it is possible to eliminate some of the undesired couplings by means of a (flavor) symme-
 try [31], we will not pursue this direction here.

70 – which are in any case useful to generate neutrino masses – allows one to promote
 71 $U(1)_{B-L} \times U(1)_{L_\mu-L_\tau} \times U(1)_{L_\mu-L_e}$ or any subgroup thereof to a local gauge symme-
 72 try [33]. We will focus on simple $U(1)_X$ subgroups, which are hence generated by

$$X = r_{BL}(B - L) + r_{\mu\tau}(L_\mu - L_\tau) + r_{\mu e}(L_\mu - L_e) \quad (2)$$

73 for arbitrary real coefficients r_x [33] (see also Refs. [34–38]), potentially including $Z-Z'$
 74 mixing. We stress that these $U(1)_X$ models are anomaly free and UV-complete, allowing
 75 us to reliably compare limits from NSI and other experiments. In their simplest form these
 76 models are also safe from proton decay and lepton flavor violation without the need for
 77 any fine-tuning, and can furthermore accommodate neutrino masses via a seesaw mecha-
 78 nism [33, 38]. This makes them perfect benchmark models for NSI, ideal to illustrate the
 79 importance of neutrino-oscillation limits compared to e.g. neutrino scattering constraints.

80 While Z' bosons and NSI have been considered before [7, 22, 23, 25–27, 29], our work is
 81 distinct due to the following aspects: we stress the importance of whether the Z' couples
 82 directly to matter particles (i.e. electrons, up- and down-quarks), or whether it couples to
 83 matter only via $Z-Z'$ mixing. We demonstrate that in the latter case $Z-Z'$ mass mixing
 84 is required to generate observable NSI in long-baseline oscillation experiments, implying
 85 non-trivial Higgs phenomenology. This is because mass mixing requires a Higgs multi-
 86 plet which is charged under both the $U(1)'$ and SM gauge groups. Working with simple
 87 anomaly-free $U(1)'$ symmetries we furthermore stress the importance of the flavor struc-
 88 ture of the underlying models, which strongly influences the size of the limits (via the
 89 sign of the generated ϵ), as well as the importance of other constraints on the Z' mass
 90 and gauge coupling. We also demonstrate that within simple UV-complete models it is
 91 possible to make terrestrial neutrino oscillation experiments insensitive to NSI, such that
 92 only scattering or collider limits apply.

93
 94 The paper is organized as follows: In Section 2 we introduce the formalism of NSI and
 95 summarize current limits from neutrino oscillations. The interplay of the flavor structure
 96 of the ϵ is stressed by comparing COHERENT limits in different cases. Section 3 deals
 97 with the calculation of NSI operators when Z' bosons are integrated out, with particular
 98 focus on whether kinetic or mass mixing is present. Specific examples from explicit models,
 99 which are anomaly-free when only right-handed neutrinos are introduced, are given. We
 100 conclude in Section 4.

101 Non-Standard Neutrino Interactions: Formalism and Limits

102 NSI relevant for neutrino propagation in matter are usually described by the effective
 103 Lagrangian

$$\mathcal{L}_{\text{eff}} = -2\sqrt{2}G_F \epsilon_{\alpha\beta}^{fX} (\bar{\nu}_\alpha \gamma_\mu P_L \nu_\beta) (\bar{f} \gamma^\mu P_X f), \quad (3)$$

104 where $X = L, R$ depends on the chirality of the interaction with $P_{L,R} = \frac{1}{2}(1 \mp \gamma_5)$ and
 105 $f \in \{e, u, d\}$ encodes the coupling to matter; $2\sqrt{2}G_F \simeq (174 \text{ GeV})^{-2}$ is a normalization
 106 factor that makes ϵ dimensionless. Relevant for neutrino oscillation experiments is only
 107 the vector part

$$\epsilon_{\alpha\beta}^f \equiv \epsilon_{\alpha\beta}^{fL} + \epsilon_{\alpha\beta}^{fR}, \quad (4)$$

108 because this induces coherent forward scattering of neutrinos in unpolarized matter. For
 109 non-trivial flavor structures, $\epsilon \not\propto \mathbb{1}$, this modifies neutrino propagation and oscillation
 110 in the Sun and Earth. In the following, we will denote this oscillation effect of the La-
 111 grangian in Eq. (3) as NSI, in contrast to various other places where the Lagrangian and

| f | $\epsilon_{ee}^f - \epsilon_{\mu\mu}^f$ | $\epsilon_{\tau\tau}^f - \epsilon_{\mu\mu}^f$ |
|-------|---|---|
| u | $[-0.020, +0.456]$ | $[-0.005, +0.130]$ |
| d | $[-0.027, +0.474]$ | $[-0.005, +0.095]$ |
| p | $[-0.041, +1.312]$ | $[-0.015, +0.426]$ |
| n | $[-0.114, +1.499]$ | $[-0.015, +0.222]$ |
| $p+n$ | $[-0.038, +0.707]$ | $[-0.008, +0.180]$ |

Table 1: 2σ bounds on the diagonal NSI $\epsilon_{\ell\ell}^f - \epsilon_{\mu\mu}^f$ assuming scattering on the fermions $f \in \{u, d, p, n, p+n\}$ from neutrino oscillation data assuming LMA, as derived in Ref. [40].

112 its UV-complete realization may show up. Limits on NSI parameters can be obtained by
 113 fitting neutrino oscillation data, which is modified due to the additional Hermitian matter
 114 potential in flavor space

$$H_{\text{mat}} = \sqrt{2}G_F N_e(x) \begin{pmatrix} 1 + \epsilon_{ee}(x) & \epsilon_{e\mu}(x) & \epsilon_{e\tau}(x) \\ \epsilon_{e\mu}^*(x) & \epsilon_{\mu\mu}(x) & \epsilon_{\mu\tau}(x) \\ \epsilon_{e\tau}^*(x) & \epsilon_{\mu\tau}^*(x) & \epsilon_{\tau\tau}(x) \end{pmatrix}, \quad (5)$$

115 with normalized NSI $\epsilon_{\alpha\beta} = \sum_f \frac{N_f(x)}{N_e(x)} \epsilon_{\alpha\beta}^f$ and position-dependent fermion densities $N_f(x)$.²
 116 Since neutrino oscillations are not sensitive to a matter potential $H_{\text{mat}} \propto \mathbb{1}$, one can
 117 constrain only *two* diagonal entries, usually written in the form of differences as $\epsilon_{ee} - \epsilon_{\mu\mu}$
 118 and $\epsilon_{\tau\tau} - \epsilon_{\mu\mu}$. Limits are typically obtained assuming a neutrino scattering only off one
 119 species $f \in \{e, u, d\}$. Recently, Ref. [40] has generalized this approach to allow for an
 120 arbitrary linear combination of up- and down-quark NSI, which in particular includes the
 121 case of scattering off protons ($f = p$: $\epsilon_{\alpha\beta}^p \equiv 2\epsilon_{\alpha\beta}^u + \epsilon_{\alpha\beta}^d$) or neutrons ($f = n$: $\epsilon_{\alpha\beta}^n \equiv$
 122 $\epsilon_{\alpha\beta}^u + 2\epsilon_{\alpha\beta}^d$). Limits on the diagonal NSI from oscillation data are given in Tab. 1, derived
 123 under the Large Mixing Angle (LMA) assumption for θ_{12} [40].³ Three combinations will
 124 turn out to be of particular interest for our study: (i) $p+n$, (ii) n , and (iii) p . The
 125 combination $p+n$ corresponds to NSI couplings $-2\sqrt{2}G_F \epsilon_{\alpha\beta}^{p+n} (\bar{\nu}_\alpha \gamma_\mu P_L \nu_\beta) j_B^\mu$ to the baryon
 126 current

$$j_B^\mu = \frac{1}{3} \sum_q \bar{q} \gamma^\mu q \supset \bar{p} \gamma^\mu p + \bar{n} \gamma^\mu n, \quad (6)$$

127 from which we can obtain the relation with $\epsilon^{u,d}$ via $\epsilon_{\alpha\beta}^{p+n} \equiv (\epsilon_{\alpha\beta}^p + \epsilon_{\alpha\beta}^n)/2 = (3\epsilon_{\alpha\beta}^u + 3\epsilon_{\alpha\beta}^d)/2$.
 128 Pure neutron NSI are realized if the couplings to protons and electrons cancel in matter,
 129 a situation we will encounter for instance in Sec. 3.2. Pure coupling to protons, on the
 130 other hand, can under certain assumptions be used as a proxy for electron NSI.⁴

131 NSI mediated by a new neutral vector boson Z' with coupling strength g' and mass
 132 $M_{Z'}$ are generically of the form $\epsilon \sim (2\sqrt{2}G_F)^{-1} (g'/M_{Z'})^2$, even if the Z' mass is tiny. The

²Crossing through electrically neutral matter consisting of protons, neutrons and electrons, coherent forward scattering picks up NSI effects proportional to the number densities: $\epsilon_{\alpha\beta}^{\text{Matter}} = \epsilon_{\alpha\beta}^e + \epsilon_{\alpha\beta}^p + Y_n^{\text{Matter}} \epsilon_{\alpha\beta}^n$, where $Y_n^{\text{Matter}} = n_n/n_e$ is the ratio of neutron and electron number densities. For Earth matter, $Y_n^{\text{Earth}} = 1.051$ on average [39].

³See e.g. Refs. [5, 7] for recent discussions on the LMA-Dark solution.

⁴Limits on ϵ^p are not equivalent to ϵ^e despite the same electron and proton abundance in electrically neutral matter because they modify the neutrino detection process differently [40]. However, in the models considered in the following neutrino–electron scattering provides an independent constraint on the strength of the interaction which restricts the new-physics impact on the neutrino detection process in oscillation experiments such as Super-Kamiokande substantially. We stress that this is only an estimate and encourage a dedicated analysis of the interplay of ϵ^e and ϵ^p . A summary of independent constraints on NSI from electrons $\epsilon_{\alpha\beta}^e$ which do not come from a global fit can be found in Ref. [11].

133 values of Tab. 1 then correspond to scales $M_{Z'}/g'$ from 140 GeV to 2.5 TeV, depending on
 134 α , β , f , and the sign of the coefficient. These have to be compared to limits from other
 135 processes, e.g. resonance searches for Z' at the LHC or meson decays. Among the various
 136 processes which could be used to test a Z' , neutrino scattering off electrons [41, 42] or
 137 nucleons [27] has the greatest similarity to NSI and the main difference between scattering
 138 experiments and NSI constraints is the momentum transfer: neutrino oscillations probe
 139 zero-momentum forward scattering and thus give limits on $M_{Z'}/g'$ that are independent
 140 of $M_{Z'}$ [25]. In contrast, the observations of neutrino scattering off quarks and electrons
 141 always requires a non-vanishing momentum transfer. Neutrino–electron scattering exper-
 142 iments are sensitive to $\mathcal{O}(1 \text{ MeV})$ momentum transfer while Coherent Elastic ν –Nucleus
 143 Scattering (CE ν NS), which has been measured by COHERENT [43] recently, currently
 144 allows to probe a momentum transfer q of the order of $\sim 50 \text{ MeV}$. Future data from CO-
 145 HERENT and other experiments such as CONUS [44] will further improve this probe [7].
 146 With initial neutrinos of flavor α (that is $\alpha = e$ for experiments with reactor neutrinos
 147 such as CONUS and $\alpha = e, \mu$ for experiments with pion beams such as COHERENT), the
 148 cross section for CE ν NS on a nucleus i with Z_i protons and N_i neutrons is proportional
 149 to the effective charge-squared

$$\tilde{Q}_{i,\alpha}^2 \equiv \left[N_i \left(-\frac{1}{2} + \epsilon_{\alpha\alpha}^n \right) + Z_i \left(\frac{1}{2} - 2s_W^2 + \epsilon_{\alpha\alpha}^p \right) \right]^2 + \sum_{\beta \neq \alpha} \left[N_i \epsilon_{\alpha\beta}^n + Z_i \epsilon_{\alpha\beta}^p \right]^2, \quad (7)$$

150 assuming real NSI for simplicity. Due to the short neutrino propagation length one can
 151 neglect neutrino oscillations here. The COHERENT [43] experiment uses neutrinos from
 152 pion decay at rest, scattering on cesium and iodine, which leads to an expression for the
 153 number of CE ν NS events

$$N_{\text{CE}\nu\text{NS}} \propto \sum_{i \in \{\text{Cs}, \text{I}\}} \left[f_{\nu_e} \tilde{Q}_{i,e}^2 + (f_{\nu_\mu} + f_{\bar{\nu}_\mu}) \tilde{Q}_{i,\mu}^2 \right], \quad (8)$$

154 with $f_{\nu_e} = 0.31$, $f_{\nu_\mu} = 0.19$, and $f_{\bar{\nu}_\mu} = 0.50$ as appropriate neutrino-flavor fractions for
 155 COHERENT. Note that experiments with reactor neutrinos such as CONUS are only sen-
 156 sitive to $\tilde{Q}_{i,e}^2$. CE ν NS is obviously sensitive to different NSI combinations than oscillation
 157 data and therefore perfectly complementary. To assess NSI limits from COHERENT we
 158 follow Refs. [40, 43, 45] and construct a $\chi^2(\epsilon)$ function that is marginalized over system-
 159 atic nuisance parameters.⁵ Compared to oscillation-based limits on NSI, the limits from
 160 scattering experiments always imply a non-zero momentum exchange q , which has to be
 161 taken into account in NSI realizations with light mediators. Specifically for Z' models, the
 162 above expression is only valid for $M_{Z'} \gg q \simeq 10 \text{ MeV}$, otherwise there is a suppression of
 163 the form $\epsilon \rightarrow \epsilon M_{Z'}^2/q^2$ [25]. In addition, neutrino scattering experiments are also sensitive
 164 to $\epsilon_{\alpha\beta} \propto \delta_{\alpha\beta}$ and are therefore invaluable as a probe of new flavor-*universal* interactions.

165 As examples we consider diagonal muon- and electron-neutrino NSI that come from
 166 scattering on baryons, i.e. e^{p+n} . Setting $\epsilon_{\tau\tau} = 0$ implies a strong bound from oscillation
 167 data due to the stringent constraint on $|\epsilon_{\tau\tau} - \epsilon_{\mu\mu}|$ (Tab. 1), so that COHERENT limits
 168 are weaker (Fig. 1 (left)). Setting on the other hand $\epsilon_{\tau\tau} = \epsilon_{\mu\mu}$ completely eliminates one
 169 of the two diagonal NSI constraints from oscillation data and thus renders COHERENT
 170 crucial to constrain the parameter space (Fig. 1 (right)). Although counterintuitive due to
 171 the absence of tau-neutrinos in the experiment, the COHERENT limits are particularly
 172 important for $\epsilon_{\tau\tau} \neq 0$, because this can weaken the strong oscillation constraints. As we
 173 will see in the following, COHERENT is indeed mainly relevant for simple Z' models with
 174 $\epsilon_{\tau\tau} \sim \epsilon_{\mu\mu}$.

⁵See also Refs. [46–51] for discussions of NSI at coherent scattering experiments.




Figure 1: Allowed regions for diagonal muon- and electron-neutrino NSI coupled to baryon number, assuming $\epsilon_{\tau\tau} = 0$ (left) and $\epsilon_{\tau\tau} = \epsilon_{\mu\mu}$ (right).

175 One lesson learned so far is that a possible underlying flavor structure of the $\epsilon_{\alpha\beta}$
 176 strongly influences which experiment is most sensitive to them.

177 Calculating NSI Operators from Z' Bosons

178 A particularly popular class of NSI realizations uses new neutral gauge bosons Z' as t -
 179 channel mediators in neutrino scattering. Here we will derive the general expressions for ϵ
 180 in terms of the Z' couplings and then discuss the simplest possible UV-complete scenarios.
 181 In addition to the direct coupling of the new $U(1)'$ gauge boson to SM fermions we will also
 182 allow for mixing between the Z' and the Z and start with the most general Lagrangian de-
 183 scribing the mixing. The formalism for Z - Z' mixing [52,53] has been frequently discussed
 184 in the literature, see for example Refs. [30,54].⁶ The Lagrangian contains a term with the
 185 usual SM expressions, the Z' part, and a term describing kinetic and mass mixing:

$$\begin{aligned}
 \mathcal{L}_{\text{SM}} &= -\frac{1}{4}\hat{B}_{\mu\nu}\hat{B}^{\mu\nu} - \frac{1}{4}\hat{W}_{\mu\nu}^a\hat{W}^{a\mu\nu} + \frac{1}{2}\hat{M}_Z^2\hat{Z}'_\mu\hat{Z}'^\mu - \frac{\hat{e}}{\hat{c}_W}j_Y^\mu\hat{B}_\mu - \frac{\hat{e}}{\hat{s}_W}j_W^{a\mu}\hat{W}_\mu^a, \\
 \mathcal{L}_{Z'} &= -\frac{1}{4}\hat{Z}'_{\mu\nu}\hat{Z}'^{\mu\nu} + \frac{1}{2}\hat{M}_Z'^2\hat{Z}'_\mu\hat{Z}'^\mu - \hat{g}'j'^\mu\hat{Z}'_\mu, \\
 \mathcal{L}_{\text{mix}} &= -\frac{\sin\chi}{2}\hat{Z}'^{\mu\nu}\hat{B}_{\mu\nu} + \delta\hat{M}^2\hat{Z}'_\mu\hat{Z}'^\mu.
 \end{aligned} \tag{9}$$

186 Hatted fields indicate here that those fields have neither canonical kinetic nor mass terms.
 187 The two Abelian gauge bosons \hat{B} and \hat{Z}' couple to each other via the term $\hat{Z}'^{\mu\nu}\hat{B}_{\mu\nu}$, which
 188 induces kinetic mixing of \hat{Z}' with the other gauge bosons [52]. It is allowed by the gauge
 189 symmetry and hence should be expected. Even if zero at some scale, this term is generated
 190 at loop level if there are particles charged under hypercharge and $U(1)'$ [53]. Tree-level
 191 mass mixing via the term $\delta\hat{M}^2\hat{Z}'_\mu\hat{Z}'^\mu$ requires that there is a scalar with a nonzero vacuum
 192 expectation value (VEV) charged under the SM and $U(1)'$.

⁶An analysis for Z - Z' - Z'' mixing was performed in Ref. [55].

193 The currents are defined as

$$j_Y^\mu = -\frac{1}{2} \sum_{\ell=e,\mu,\tau} [\bar{L}_\ell \gamma^\mu L_\ell + 2\bar{\ell}_R \gamma^\mu \ell_R] + \frac{1}{6} \sum_{\text{quarks}} [\bar{Q}_L \gamma^\mu Q_L + 4\bar{u}_R \gamma^\mu u_R - 2\bar{d}_R \gamma^\mu d_R],$$

$$j_W^{a\mu} = \sum_{\ell=e,\mu,\tau} \bar{L}_\ell \gamma^\mu \frac{\sigma^a}{2} L_\ell + \sum_{\text{quarks}} \bar{Q}_L \gamma^\mu \frac{\sigma^a}{2} Q_L, \quad (10)$$

194 with the left-handed $SU(2)$ -doublets Q_L and L_ℓ and the Pauli matrices σ^a . The final
195 electric current after electroweak symmetry breaking is given as $j_{\text{EM}} \equiv j_W^3 + j_Y$ and the
196 weak neutral current is $j_{\text{NC}} \equiv 2j_W^3 - 2\hat{s}_W^2 j_{\text{EM}}$. The new neutral current j' of the $U(1)'$
197 is left unspecified here, but has to contain flavor *non-universal* neutrino interactions in
198 order to generate NSI:

$$j'_\mu \supset \sum_{\alpha,\beta} q_{\alpha\beta} \bar{\nu}_\alpha \gamma_\mu P_L \nu_\beta, \quad (11)$$

199 with some flavor-dependent coupling matrix $q \neq \mathbf{1}$. Below we will consider some simple
200 models that lead to such couplings.

201 After diagonalization, the physical massive gauge bosons $Z_{1,2}$ and the massless photon
202 couple to a linear combination of j' , j_{NC} and j_{EM} :

$$\mathcal{L}_{\text{int}} = - \left(e j_{\text{EM}}, \frac{e}{2\hat{s}_W \hat{c}_W} j_{\text{NC}}, g' j' \right) \begin{pmatrix} 1 & a_1 & a_2 \\ 0 & b_1 & b_2 \\ 0 & d_1 & d_2 \end{pmatrix} \begin{pmatrix} A \\ Z_1 \\ Z_2 \end{pmatrix}. \quad (12)$$

203 Here the entries of the matrix are

$$\begin{aligned} a_1 &= -\hat{c}_W \sin \xi \tan \chi, \\ b_1 &= \cos \xi + \hat{s}_W \sin \xi \tan \chi, \\ d_1 &= \frac{\sin \xi}{\cos \chi}, \\ a_2 &= -\hat{c}_W \cos \xi \tan \chi, \\ b_2 &= \hat{s}_W \cos \xi \tan \chi - \sin \xi, \\ d_2 &= \frac{\cos \xi}{\cos \chi}. \end{aligned} \quad (13)$$

204 The angles χ and ξ in the above expressions come from diagonalizing the kinetic and the
205 mass terms of the massive gauge bosons Z and Z' , respectively. The diagonalization of
206 the mass matrix is achieved via

$$\begin{pmatrix} \cos \xi & \sin \xi \\ -\sin \xi & \cos \xi \end{pmatrix} \begin{pmatrix} a & b \\ b & c \end{pmatrix} \begin{pmatrix} \cos \xi & -\sin \xi \\ \sin \xi & \cos \xi \end{pmatrix} = \begin{pmatrix} M_1^2 & 0 \\ 0 & M_2^2 \end{pmatrix} \equiv \begin{pmatrix} M_Z^2 & 0 \\ 0 & M_{Z'}^2 \end{pmatrix}, \quad (14)$$

207 where

$$\tan 2\xi = \frac{2b}{a-c} \text{ with } \begin{cases} a = \hat{M}_Z^2, \\ b = \hat{s}_W \tan \chi \hat{M}_Z^2 + \frac{\delta \hat{M}^2}{\cos \chi}, \\ c = \frac{1}{\cos^2 \chi} \left(\hat{M}_Z^2 \hat{s}_W^2 \sin^2 \chi + 2\hat{s}_W \sin \chi \delta \hat{M}^2 + \hat{M}_{Z'}^2 \right). \end{cases} \quad (15)$$

208 At energies $E \ll M_{1,2}$, one can integrate out the Z_1 and Z_2 bosons to obtain the following
209 effective operators:

$$\mathcal{L}_{\text{eff}} = - \sum_{i=1,2} \frac{1}{2M_i^2} \left(e j_{\text{EM}} a_i + \frac{e}{2\hat{s}_W \hat{c}_W} j_{\text{NC}} b_i + g' j' d_i \right)^2. \quad (16)$$

210 If more Z' bosons are present, the sum would extend over all their mass states [55]. Note
 211 that \hat{s}_W reduces to the known weak angle $\sin\theta_W$ for small Z - Z' mixing angle ξ [54].

212 Comparing the effective Lagrangian from Eq. (16) with the NSI operators in Eqs. (3,4)
 213 gives from the mixed j' - j_{EM} and j' - j_{NC} terms the following NSI coefficients for coupling
 214 to electrons, up- and down-quarks:

$$\begin{aligned}\epsilon_{\alpha\beta}^e &= \sum_{i=1,2} q_{\alpha\beta} \frac{g'd_i}{\sqrt{2}M_i^2 G_F} \left(-ea_i + \frac{eb_i}{2s_W c_W} \left(-\frac{1}{2} + 2s_W^2 \right) + g'd_i \frac{\partial j'_\alpha}{\partial \bar{e}\gamma_\alpha e} \right), \\ \epsilon_{\alpha\beta}^u &= \sum_{i=1,2} q_{\alpha\beta} \frac{g'd_i}{\sqrt{2}M_i^2 G_F} \left(\frac{2}{3}ea_i + \frac{eb_i}{2s_W c_W} \left(\frac{1}{2} - \frac{4}{3}s_W^2 \right) + g'd_i \frac{\partial j'_\alpha}{\partial \bar{u}\gamma_\alpha u} \right), \\ \epsilon_{\alpha\beta}^d &= \sum_{i=1,2} q_{\alpha\beta} \frac{g'd_i}{\sqrt{2}M_i^2 G_F} \left(-\frac{1}{3}ea_i + \frac{eb_i}{2s_W c_W} \left(-\frac{1}{2} + \frac{2}{3}s_W^2 \right) + g'd_i \frac{\partial j'_\alpha}{\partial \bar{d}\gamma_\alpha d} \right).\end{aligned}\quad (17)$$

215 The origin of the a_i (b_i) terms from the electric and neutral currents is obvious, whereas
 216 the d_i terms take into account that the Z' might have direct couplings to matter particles
 217 (i.e. first generation charged fermions) even in the absence of Z - Z' mixing. Later we will
 218 consider cases with and without direct couplings to matter particles.

219 Forward scattering of neutrinos in matter corresponds to zero momentum exchange,
 220 so the above expressions are valid even for very light Z' masses, contrary to e.g. neutrino
 221 scattering in COHERENT. Note however that Z' masses below ~ 5 MeV are strongly
 222 disfavored by cosmology, in particular the number of relativistic degrees of freedom N_{eff} ,
 223 unless the coupling is made tiny [56–58]. One can still consider minuscule g' and Z' mass
 224 with $M_{Z'}/g' \sim 100$ GeV so as to evade N_{eff} constraints and still have testable NSI [59],
 225 but this typically requires an analysis in terms of long-range potentials [60–62] instead of
 226 the contact interactions of Eq. (3) and will not be considered here.

227 NSI without Z - Z' mixing

228 Let us first consider the case of vanishing Z - Z' mixing, $\xi = \chi = 0$, which simplifies Eq. (17)
 229 substantially. We must then find a Z' that has couplings to matter particles as well as
 230 non-universal neutrino couplings. Flavor-violating neutrino couplings $\bar{\nu}_\alpha \not{Z}' P_L \nu_{\beta \neq \alpha}$ are
 231 typically difficult to obtain and often, but not always, run into problems with constraints
 232 from charged-lepton flavor violation (LFV) [11, 27]. We will therefore focus on flavor-
 233 *diagonal* neutrino couplings in the following, which are much easier to obtain. This is also
 234 motivated by the recent hints for lepton-flavor non-universality in B -meson decays, which
 235 can be explained with models that typically give at least diagonal NSI.

236 There is a very simple class of Z' models that lead to diagonal NSI that will be the
 237 focus of this work. We use the fact that, introducing only right-handed neutrinos to the
 238 particle content of the SM, the most general anomaly-free $U(1)_X$ symmetry is generated
 239 by Eq. (2),

$$X = r_{BL}(B - L) + r_{\mu\tau}(L_\mu - L_\tau) + r_{\mu e}(L_\mu - L_e)$$

240 for arbitrary real coefficients r_x [33] (see also Refs. [34–38]). This gives the current $j'_\alpha =$
 241 $\sum_f X(f) \bar{f} \gamma_\alpha f$, which is vector-like for all charged particles. The first term in Eq. (2)
 242 can couple the Z' to matter even in the absence of Z - Z' mixing, while the last two terms
 243 induce the neutrino-flavor non-universality necessary for NSI, to be discussed below. Aside
 244 from being anomaly-free, the above symmetries can also easily accommodate the observed
 245 pattern of neutrino masses and mixing. The key point is that one can break the $U(1)_X$
 246 symmetry using only electroweak singlets which then generate a non-trivial right-handed

247 neutrino Majorana mass matrix that leads to the seesaw mechanism [33]. Despite our
 248 flavor symmetry we therefore do not have to worry about LFV, as these effects are still
 249 heavily suppressed.

250 Assuming negligible $Z-Z'$ mixing, the effective Lagrangian from Eq. (16) becomes very
 251 simple:

$$\begin{aligned} \mathcal{L}_{\text{eff}} &= -\frac{(g')^2}{2M_{Z'}^2} j'_\alpha j'^\alpha \\ &\supset -\frac{(g')^2}{M_{Z'}^2} [r_{BL}(\bar{p}\gamma^\alpha p + \bar{n}\gamma^\alpha n) - (r_{BL} + r_{\mu e})\bar{e}\gamma^\alpha e] \\ &\quad \times [-(r_{BL} + r_{\mu e})\bar{\nu}_e\gamma_\alpha P_L\nu_e - (r_{BL} - r_{\mu e} - r_{\mu\tau})\bar{\nu}_\mu\gamma_\alpha P_L\nu_\mu - (r_{BL} + r_{\mu\tau})\bar{\nu}_\tau\gamma_\alpha P_L\nu_\tau], \end{aligned} \quad (18)$$

252 where we used the new-physics current generated by Eq. (2) and only kept the terms
 253 relevant for NSI. The NSI coefficients with coupling to baryons then take the form

$$\epsilon_{ee}^{p,n} - \epsilon_{\mu\mu}^{p,n} = -\frac{(g')^2}{2\sqrt{2}G_F M_{Z'}^2} r_{BL}(2r_{\mu e} + r_{\mu\tau}), \quad (19)$$

$$\epsilon_{\tau\tau}^{p,n} - \epsilon_{\mu\mu}^{p,n} = -\frac{(g')^2}{2\sqrt{2}G_F M_{Z'}^2} r_{BL}(2r_{\mu\tau} + r_{\mu e}), \quad (20)$$

254 and similar for those with electrons

$$\epsilon_{ee}^e - \epsilon_{\mu\mu}^e = +\frac{(g')^2}{2\sqrt{2}G_F M_{Z'}^2} (r_{BL} + r_{\mu e})(2r_{\mu e} + r_{\mu\tau}), \quad (21)$$

$$\epsilon_{\tau\tau}^e - \epsilon_{\mu\mu}^e = +\frac{(g')^2}{2\sqrt{2}G_F M_{Z'}^2} (r_{BL} + r_{\mu e})(2r_{\mu\tau} + r_{\mu e}). \quad (22)$$

255 Neutral matter necessarily contains an equal number of protons and electrons, so the
 256 relevant combination is actually the sum $\epsilon^p + \epsilon^e$:

$$(\epsilon_{ee}^p + \epsilon_{ee}^e) - (\epsilon_{\mu\mu}^p + \epsilon_{\mu\mu}^e) = +\frac{(g')^2}{2\sqrt{2}G_F M_{Z'}^2} r_{\mu e}(2r_{\mu e} + r_{\mu\tau}), \quad (23)$$

$$(\epsilon_{\tau\tau}^p + \epsilon_{\tau\tau}^e) - (\epsilon_{\mu\mu}^p + \epsilon_{\mu\mu}^e) = +\frac{(g')^2}{2\sqrt{2}G_F M_{Z'}^2} r_{\mu e}(2r_{\mu\tau} + r_{\mu e}). \quad (24)$$

257 Non-vanishing NSI in neutrino oscillations without $Z-Z'$ mixing thus require either $r_{BL} \neq$
 258 0 in order to generate a coupling to neutrons or $r_{\mu e} \neq 0$ in order to couple to electrons.
 259 Naturally, the phenomenology of a Z' depends sensitively on the SM fermions it couples
 260 to. In the following we will go through the basic simple coupling structures which arise in
 261 this class of $U(1)'$ groups. We first introduce the various experimental probes and then
 262 discuss how these compare to the limits on the NSI derived from neutrino oscillations.⁷

263 Before moving on let us briefly discuss the possibility of realizing the LMA-Dark [63]
 264 solution within our $U(1)'$ framework. As is well known, neutrino oscillations in the presence
 265 of NSI contain a generalized mass-ordering degeneracy [64–67] that in principle allows for
 266 large ϵ if the neutrino mixing parameters take on different values from the non-NSI LMA
 267 scenario. This LMA-Dark region of parameter space requires a large $\epsilon_{ee} - \epsilon_{\mu\mu} = -\mathcal{O}(1)$
 268 but all other NSI much smaller in magnitude, currently compatible with zero [40]. In our
 269 $U(1)'$ models the condition $|\epsilon_{\tau\tau} - \epsilon_{\mu\mu}| \ll |\epsilon_{ee} - \epsilon_{\mu\mu}|$ essentially requires that muons and

⁷See e.g. Ref. [42] for a discussion of future limits on some of the models under study here.

270 taus carry the same $U(1)'$ charge, which translates into $r_{\mu\tau} = -r_{\mu e}/2$ above. The only
 271 non-vanishing NSI are then

$$(\epsilon_{ee}^p + \epsilon_{ee}^e) - (\epsilon_{\mu\mu}^p + \epsilon_{\mu\mu}^e) = + \frac{3(g')^2}{4\sqrt{2}G_F M_{Z'}^2} r_{\mu e}^2, \quad (25)$$

$$\epsilon_{ee}^n - \epsilon_{\mu\mu}^n = - \frac{3(g')^2}{4\sqrt{2}G_F M_{Z'}^2} r_{\mu e} r_{BL}. \quad (26)$$

272 The proton plus electron NSI are strictly positive and thus incapable of realizing the
 273 LMA-Dark solution; the neutron NSI on the other hand can be negative and even dom-
 274 inant over the proton plus electron NSI by choosing $|r_{\mu e}| \ll |r_{BL}|$. It has however been
 275 shown in Ref. [40] that neutron NSI by themselves ($\eta = \pm 90^\circ$ in their notation) do not
 276 admit the LMA-Dark solution. This can be easily understood from the highly varying
 277 neutron-to-proton density inside the Sun, which explicitly breaks the generalized mass-
 278 ordering degeneracy and thus distinguishes between LMA-Dark and LMA [65], the latter
 279 providing a significantly better fit [40]. As a result, none of our simple $U(1)'$ models can
 280 accommodate the LMA-Dark solution, and so we will not discuss it further. Note that
 281 this conclusion remains true if we allow for Z - Z' mixing, because this can at best generate
 282 neutron NSI as we will see below.

283 Electrophobic NSI

284 Coming back to the LMA scenario, an interesting special case arises for $r_{\mu e} = -r_{BL} \neq 0$.
 285 This assignment of the charges eliminates the coupling to electrons and thus leads to NSI
 286 that are generated by the baryon density (i.e. by protons plus neutrons). This simply
 287 corresponds to a $U(1)_X$ symmetry generated by $X = B - 2L_\mu - L_\tau + r_{\mu\tau}(L_\mu - L_\tau)$.

288 Irrespective of the flavor of the leptonic interactions these $U(1)'$ can be probed by
 289 purely baryonic processes. In the presence of a light new resonance with a mass below
 290 the QCD scale the scattering rates between baryons are modified. The most stringent
 291 limits come from measurements of neutron-lead scattering [68, 69]. In addition, a light
 292 Z' could play a role in meson decays. For $M_{Z'} \lesssim m_{\pi^0}$ the strongest limits come from
 293 $\pi^0 \rightarrow \gamma + \text{invisible}$, while at higher masses the production of additional hadrons via the
 294 Z' can be constrained by a close scrutiny of η , η' , Ψ or Υ decays [25]. Limits derived
 295 from these observables can be applied to all $U(1)'$ groups that include a coupling to the
 296 baryonic current, see for example Fig. 2.

297 The leptonic couplings of the Z' lead to additional observables which can be used to
 298 constrain the interaction strength. On the one hand, couplings to τ leptons are hard to
 299 constrain for Z' 's in the mass range considered here. The short lifetime and large mass of
 300 the τ prevents a detailed scrutiny of its interaction in low-energy experiments such that
 301 we need to rely on the baryonic probes mentioned previously. One of the few relevant
 302 τ constraint comes from the one-loop vertex correction to the $Z\tau\tau$ and $Z\nu_\tau\nu_\tau$ couplings,
 303 which for $M_{Z'} \ll M_Z$ are given by

$$\frac{g_{V,A}}{g_{V,A}^{\text{SM}}} \simeq 1 + \frac{(X(\tau)g')^2}{(4\pi)^2} \left[\frac{\pi^2}{3} - \frac{7}{2} - 3 \log \left(\frac{M_{Z'}^2}{M_Z^2} \right) - \log^2 \left(\frac{M_{Z'}^2}{M_Z^2} \right) - 3i\pi - 2i\pi \log \left(\frac{M_{Z'}^2}{M_Z^2} \right) \right], \quad (27)$$

304 with $X(\tau)$ the $U(1)_X$ charge of the tau. The Z' corrections suppress the Z couplings to
 305 taus, which have been precisely measured at LEP [72]. We show the naive 2σ constraint
 306 from the axial $Z\tau\tau$ coupling, $|g_A - g_A^{\text{SM}}| < 2 \times 0.00064$ in Fig. 2. While stronger than
 307 most $U(1)_B$ limits for $M_{Z'} \sim \text{GeV}$, these limits will not be relevant for $U(1)_X$ models with
 308 muon or electron couplings, which are strongly constrained by other observables.




Figure 2: Limits on $U(1)_{B-3L_\tau}$ gauge coupling and Z' mass from Refs. [27, 70] together with the strong NSI constraint (blue). For limits that include (radiative) kinetic mixing, see Ref. [71].

309 Muons, for example, allow for precision experiments. Rare neutrino-induced processes
 310 such as neutrino trident production, which has been measured by the CCFR experi-
 311 ment [73], can test the interaction between neutrinos and muons [74]. As is well known,
 312 a light Z' can alleviate the tension between the SM prediction and the measured value of
 313 the anomalous magnetic moment of the muon $(g-2)_\mu$. The parameter space in which the
 314 tension is reduced to 2σ (1σ) is indicated by the dark (light) green band in Fig. 3. In the
 315 region above the green band $(g-2)_\mu$ is dominated by the new-physics contribution while
 316 $(g-2)_\mu$ asymptotes to the SM value below the green band. Since the new physics can
 317 drive the expected anomalous magnetic moment further away from the measurement than
 318 the SM a large fraction of the upper region is disfavored compared to the lower regions.
 319 We omit this constraint in the figure since this regions is already in tension with CCFR.
 320 Additional constraints on a light mediator coupling of muons can be derived from searches
 321 for $e^+e^- \rightarrow \mu^+\mu^-Z'$ in four-muon final states at BaBar [75]. This search is sensitive down
 322 to the two-muon threshold and excludes $g' \gtrsim 10^{-3}$ for $M_{Z'} \simeq 200$ MeV. Finally, there are
 323 also constraints from cosmology which are largely insensitive to the details of the particle-
 324 physics model. A light Z' can be produced copiously in the early Universe if coupled to
 325 light SM fermions, even if just to neutrinos. Bosons with mass below $M_{Z'} \lesssim 5$ MeV then
 326 either contribute themselves to the relativistic degrees of freedom N_{eff} at the time of Big
 327 Bang nucleosynthesis [56], or heat up the decoupled neutrino bath via $Z' \rightarrow \nu\nu$ [57, 58],
 328 putting strong constraints on our models.

329 The relevant NSI limits from a global fit to neutrino oscillation data can be readily
 330 read off from Tab. 1. We give the three most extreme cases for $r_{\mu\tau}$ in Tab. 2 which also
 331 illustrates the importance of the NSI sign:

- 332 • For $B-3L_\tau$ [76–78], corresponding to $r_{\mu\tau} = 2$, we obtain negative NSI coefficients,
 333 which are much more constrained than positive NSI. As a result, NSI impose a very
 334 strong constraint $M_{Z'}/|g'| > 4.8$ TeV on this scenario, to be compared to extremely
 335 weak limits from other experiments (see Fig. 2). This is the scenario where neutrino
 336 oscillations are most important. COHERENT does not set a limit here because it
 337 does not involve tau neutrinos.




Figure 3: Constraints on $U(1)_{B-\frac{3}{2}(L_\mu+L_\tau)}$ (left) and $U(1)_{B-3L_\mu}$ (right) together with the 2σ NSI bound from neutrino oscillations (Tab. 2) and the 2σ constraint from COHERENT. Also shown is the preferred region to resolve the muon's $(g-2)$ at 1 and 2σ in green and exclusions from ΔN_{eff} , BaBar [75] and neutrino trident production in CCFR [73, 74].

| $U(1)_X$ | $\epsilon_{ee}^{p+n} - \epsilon_{\mu\mu}^{p+n}$ | $\epsilon_{\tau\tau}^{p+n} - \epsilon_{\mu\mu}^{p+n}$ | $M_{Z'}/ g' $ |
|-----------------------------------|---|---|---------------------|
| $B - 3L_\tau$ | 0 | $-\frac{3(g')^2}{\sqrt{2}G_F M_{Z'}^2}$ | $> 4.8 \text{ TeV}$ |
| $B - \frac{3}{2}(L_\mu + L_\tau)$ | $+\frac{3(g')^2}{2\sqrt{2}G_F M_{Z'}^2}$ | 0 | $> 360 \text{ GeV}$ |
| $B - 3L_\mu$ | $+\frac{3(g')^2}{\sqrt{2}G_F M_{Z'}^2}$ | $+\frac{3(g')^2}{\sqrt{2}G_F M_{Z'}^2}$ | $> 1.0 \text{ TeV}$ |

Table 2: Examples for NSI from electrophobic anomaly-free $U(1)_X$ without $Z-Z'$ mass mixing, as well as the NSI limit [40] on the Z' mass and coupling. See Figs. 2 and 3 for additional limits on the parameter space.

- $B - \frac{3}{2}(L_\mu + L_\tau)$ [79], corresponding to $r_{\mu\tau} = 1/2$, gives positive NSI and a rather weak limit of $M_{Z'}/|g'| > 360 \text{ GeV}$. Thanks to the condition $\epsilon_{\tau\tau} = \epsilon_{\mu\mu}$, COHERENT can give better constraints than oscillation data (Fig. 1) and in fact provides the best limit for $40 \text{ MeV} < M_{Z'} < 800 \text{ MeV}$, but is overpowered at higher masses by BaBar [75] and neutrino trident production as measured by CCFR [73, 74] (see Fig. 3). At no point can one resolve the longstanding $(g-2)_\mu$ anomaly [80].
- $B - 3L_\mu$ [81], corresponding to $r_{\mu\tau} = -1$, only gives $\epsilon_{\mu\mu}$ and a rather strong limit $M_{Z'}/|g'| > 1 \text{ TeV}$ from neutrino oscillations, which is however weaker than neutrino-trident limits if $M_{Z'} > 700 \text{ MeV}$ (see Fig. 3). As expected from Fig. 1, COHERENT is currently not competitive with oscillation constraints here.

As can be seen, the bounds on hadronic interactions of a Z' are weaker than those arising from interactions with muons. Consequently, we only show the hadronic limits in Fig. 2 and focus on the other constraints in Fig. 3. In all these cases neutrino oscillations provide the strongest limits for light Z' , $M_{Z'} = \mathcal{O}(1-100) \text{ MeV}$, and NSI with a strength that might impair future neutrino oscillation experiments can not be excluded.

Electrophilic NSI

Moving on from the electrophobic NSI to Z' scenarios with electron couplings, we again focus on some simple examples to illustrate the different possibilities. Prime examples for relevant $U(1)_X$ generators that lead to ϵ^e are $B - 3L_e$ [82], $L_e - L_\mu$ [83, 84], and $L_e - L_\tau$, collected in Tab. 3.

| | | | | |
|----------------|---|---|------------------------|---------------------|
| $U(1)_X$ | $\epsilon_{ee}^{e+p} - \epsilon_{\mu\mu}^{e+p}$ | $\epsilon_{ee}^n - \epsilon_{\mu\mu}^n$ | $M_{Z'}/ g' $ (TEXONO) | $M_{Z'}/ g' $ (NSI) |
| $B - 3L_e$ | $+\frac{3(g')^2}{\sqrt{2}G_F M_{Z'}^2}$ | $-\frac{3(g')^2}{2\sqrt{2}G_F M_{Z'}^2}$ | $> 2 \text{ TeV}$ | $> 0.2 \text{ TeV}$ |
| $U(1)_X$ | $\epsilon_{ee}^e - \epsilon_{\mu\mu}^e$ | $\epsilon_{\tau\tau}^e - \epsilon_{\mu\mu}^e$ | $M_{Z'}/ g' $ (TEXONO) | $M_{Z'}/ g' $ (NSI) |
| $L_e - L_\mu$ | $+\frac{(g')^2}{\sqrt{2}G_F M_{Z'}^2}$ | $+\frac{(g')^2}{2\sqrt{2}G_F M_{Z'}^2}$ | $> 0.7 \text{ TeV}$ | $> 0.3 \text{ TeV}$ |
| $L_e - L_\tau$ | $+\frac{(g')^2}{2\sqrt{2}G_F M_{Z'}^2}$ | $-\frac{(g')^2}{2\sqrt{2}G_F M_{Z'}^2}$ | $> 0.7 \text{ TeV}$ | $> 1.4 \text{ TeV}$ |

Table 3: Examples for NSI from electrophilic anomaly-free $U(1)_X$ without Z – Z' mass mixing, as well as the TEXONO e – ν -scattering limit [85] on the Z' mass and coupling and approximate NSI constraints.

358 Models with couplings between neutrinos and electrons allow for additional ways to
 359 test the $U(1)'$. First of all, this coupling directly modifies the scattering of neutrinos
 360 off electrons. The best limits on the contribution of a light Z' to ν – e scattering come
 361 from a reanalysis [41, 85] of data collected during the TEXONO-CsI run [86]. In addition,
 362 bounds on new interactions with electrons can be derived from positron–electron collisions.
 363 The best limits in the mass range of interest here come from the BaBar search for dark
 364 photons [87]. When translated into the parameters of the Z' model considered here these
 365 limits exclude $g' \gtrsim 10^{-4}$ in a wide range of masses, see e.g. Fig. 4. In addition, there are
 366 constraints on light Z' from beam-dump experiments. These bounds can be translated to
 367 a given Z' model once the couplings and Z' branching ratios are known [88]. We use the
 368 code `Darkcast` [71] to translate the relevant beam-dump limits [89–95] to the $B - 3L_e$
 369 model, see Fig. 4.

370 Since there is no recent analysis of global neutrino oscillation data for NSI that come
 371 from the electron density, we have to make some approximations. In principle, the electron
 372 matter density and the proton matter density are identical; one is therefore tempted to
 373 assume that the limits on proton NSI are the same as those on electron NSI. However,
 374 one has to keep in mind that interactions with electrons will not only affect the matter
 375 potential (i.e. neutrino propagation) but also the neutrino *detection* process and so bounds
 376 of ϵ^p are not strictly identical to bounds on ϵ^e . Nevertheless, the independent bounds on
 377 the interaction of Z' with electrons mentioned above ensure that the neutrino detection
 378 process is basically unaffected by new physics. In the following we will hence assume that
 379 the limits on proton NSI from the global fit of Ref. [40] are a good proxy for the electron
 380 NSI.

381 Now we can use the limits from Tab. 1 to constrain straightforwardly $L_e - L_{\mu,\tau}$. For
 382 $L_e - L_\mu$ the best NSI limit comes from $\epsilon_{\tau\tau}^e - \epsilon_{\mu\mu}^e$ and gives $M_{Z'}/|g'| > 0.3 \text{ TeV}$, a factor of
 383 two weaker than the TEXONO limit (Tab. 3). For $L_e - L_\tau$ the best NSI limit also comes
 384 from the $\epsilon_{\tau\tau}^e - \epsilon_{\mu\mu}^e$ entry, but is much stronger due to the opposite sign compared to $L_e - L_\mu$;
 385 the limit reads $M_{Z'}/|g'| > 1.4 \text{ TeV}$ and is thus a factor two stronger than TEXONO's.
 386 This once again illustrates the importance of the NSI sign and the complementarity of
 387 the different experiments and observables. Current and future limits in the $M_{Z'}-g'$ plane
 388 for these two scenarios (without the NSI bounds) can be found in Ref. [42]. In the last
 389 example, $B - 3L_e$, we only generate the $\epsilon_{ee} - \epsilon_{\mu\mu}$ NSI combination, but with contributions
 390 from electron, protons, and neutrons of the form $\epsilon^n/\epsilon^{e+p} = -1/2$. Overall this leads to
 391 positive $\epsilon_{ee} - \epsilon_{\mu\mu}$ which is then only weakly constrained, $M_{Z'}/|g'| > 0.2 \text{ TeV}$, so that
 392 TEXONO is more relevant. We strongly encourage a global analysis of ϵ^e NSI seeing as
 393 they give crucial limits on the parameter space of flavored gauge bosons. Of our three
 394 examples, only $B - 3L_e$ can lead to CE ν NS, but this process does not give better limits
 395 than TEXONO (Fig. 4).




Figure 4: Constraints on $U(1)_{B-3L_e}$ from beam dumps and BaBar (adapted from Refs. [71, 88]) together with COHERENT and TEXONO (2σ) neutrino scattering bounds [41, 42, 85, 88] as well as approximate NSI constraints.

396 Going back to the effective Lagrangian (18) one can find another interesting limit
 397 around $r_{\mu e} \simeq +r_{BL} \neq 0$, as this would imply a vanishing $\epsilon^p + \epsilon^e + \epsilon^n$ in matter with equal
 398 number of protons, neutrons, and electrons. This relation is approximately satisfied inside
 399 Earth, which would then be insensitive to this kind of NSI, all the while one could still
 400 have large effects in *solar* neutrino oscillations. This corresponds to the case $\eta \simeq -44^\circ$
 401 analyzed in Ref. [40], where it was shown that this scenario indeed severely weakens NSI
 402 constraints. Analogously, one can easily imagine a scenario with non-vanishing NSI inside
 403 Earth but with $\epsilon \simeq 0$ at one specific radius inside the Sun, once again covered in Ref. [40].
 404 This again weakens the NSI bounds and makes other experimental probes, such as neutrino
 405 scattering off electrons and nucleons, more important.

406 We see again, now more explicitly within UV-complete models, that the flavor structure
 407 is crucial to determine which experimental approach can provide the best limits on the
 408 model.

409 NSI with $Z-Z'$ mixing

410 In the cases discussed above, the Z' already had couplings to matter particles u, d, e ,
 411 allowing for NSI without the need for $Z-Z'$ mixing. To see the effect of $Z-Z'$ mixing, let
 412 us consider a simple $U(1)_X$ under which no matter particles are charged. As is obvious
 413 from Eq. (2), this singles out $U(1)_{L_\mu-L_\tau}$ [83, 84, 96]. Starting from Eq. (17) it is instructive
 414 to obtain the NSI coefficients for protons and neutrons instead of quarks:

$$\begin{aligned}
 \epsilon_{\alpha\beta}^n &= \sum_{i=1,2} q_{\alpha\beta} \frac{eg'd_i}{\sqrt{2}M_i^2 G_F} \frac{b_i}{2s_W c_W} \left(-\frac{1}{2}\right), \\
 \epsilon_{\alpha\beta}^p &= \sum_{i=1,2} q_{\alpha\beta} \frac{eg'd_i}{\sqrt{2}M_i^2 G_F} \left(a_i + \frac{b_i}{2s_W c_W} \left(\frac{1}{2} - 2s_W^2\right)\right), \\
 \epsilon_{\alpha\beta}^e &= \sum_{i=1,2} q_{\alpha\beta} \frac{eg'd_i}{\sqrt{2}M_i^2 G_F} \left(-a_i - \frac{b_i}{2s_W c_W} \left(\frac{1}{2} - 2s_W^2\right)\right),
 \end{aligned} \tag{28}$$

415 where now $q = \text{diag}(0, 1, -1)$ due to the $U(1)_{L_\mu - L_\tau}$ coupling. Interestingly, proton and
416 electron NSI cancel each other exactly in electrically neutral matter:

$$\epsilon_{\alpha\beta}^p + \epsilon_{\alpha\beta}^e = 0. \quad (29)$$

417 Note that this result is independent of $L_\mu - L_\tau$, and holds for any $U(1)'$ model one may
418 imagine that has $Z-Z'$ mixing but no direct coupling to electrons, up- or down-quarks.
419 Therefore, if the NSI-matter couplings come from $Z-Z'$ mixing, the only effects are from
420 coupling to *neutrons* [22], and the limits can be read off Table 1.

421 Let us take a closer look at the neutron part. An important combination of parameters
422 in the previous expressions is the sum over $b_i d_i / M_i^2$. Using Eqs. (12-14), we can rewrite
423 it as follows:

$$\begin{aligned} \sum_{i=1,2} \frac{d_i b_i}{M_i^2} &= \frac{1}{c_\chi} \left[c_\xi s_\xi \left(\frac{1}{M_1^2} - \frac{1}{M_2^2} \right) + s_W t_\chi \left(\frac{s_\xi^2}{M_1^2} + \frac{c_\xi^2}{M_2^2} \right) \right] \\ &= \frac{\delta \hat{M}^2}{(\delta \hat{M}^2)^2 - \hat{M}_{Z'}^2 \hat{M}_Z^2} \\ &= -\frac{\delta \hat{M}^2}{M_1^2 M_2^2 c_\chi^2}. \end{aligned} \quad (30)$$

424 Hence, if there is no, or sufficiently suppressed, mass mixing $\delta \hat{M}^2$, no NSI effects will
425 be generated in neutrino oscillations. In particular, *kinetic mixing* cannot by itself lead
426 to such NSI, even if the Z' has non-universal couplings to neutrinos; *mass mixing* is
427 required, which is a much bigger model-building challenge. Kinetic mixing will of course
428 still lead to effects in neutrino scattering experiments, with the best constraint coming
429 from Borexino [97, 98] rather than COHERENT [99]. Below we will focus on the opposite
430 case where kinetic mixing is absent but mass mixing is present and can thus lead to NSI.

431 Using Eq. (30), the final NSI for the $L_\mu - L_\tau$ plus mass mixing case are

$$\epsilon_{\tau\tau}^n - \epsilon_{\mu\mu}^n = 2(\epsilon_{ee}^n - \epsilon_{\mu\mu}^n) = -2 \frac{eg'}{4\sqrt{2}G_F s_W c_W} \frac{\delta \hat{M}^2}{M_Z^2 M_{Z'}^2 c_\chi^2}, \quad (31)$$

432 where we denote $M_{1,2} \rightarrow M_{Z,Z'}$. These NSI are best constrained by the $\tau\tau - \mu\mu$ NSI:
433 $\epsilon_{\tau\tau}^n - \epsilon_{\mu\mu}^n \in [-0.015, +0.222]$ (see Tab. 1). It is clear from the above expression that the
434 NSI now depend on more parameters of the new physics sector and knowledge of g' and
435 $M_{Z'}$ is no longer sufficient to predict $\epsilon_{\alpha\beta}^n$. Similarly, the neutrino-nucleus scattering cross
436 section tested by COHERENT is sensitive to the $Z-Z'$ mixing parameter. As expected
437 from Fig. 1, however, the current COHERENT limit is weaker than the NSI limit due to
438 $\epsilon_{\mu\mu} = -\epsilon_{\tau\tau}$.

439 Using the $Z-Z'$ mixing angle ξ from Eq. (15) the NSI can be expressed as

$$\epsilon_{\tau\tau}^n - \epsilon_{\mu\mu}^n = 2(\epsilon_{ee}^n - \epsilon_{\mu\mu}^n) \simeq -0.04 \left(\frac{550 \text{ GeV}}{M_{Z'}/g'} \right) \left(\frac{1 \text{ TeV}}{M_{Z'}/\xi} \right), \quad (32)$$

440 showing explicitly that NSI are the result of a cross-coupling of the $L_\mu - L_\tau$ current $g' j'$
441 and the neutral current ξj_{NC} . The former is only weakly constrained due to the absence of
442 first-generation particles in j' , illustrated in Fig. 5. For light Z' , values $M_{Z'}/g' \sim 10 \text{ GeV}$
443 are possible, whereas heavier Z' are constrained conservatively by CCFR [73] as $M_{Z'}/g' \gtrsim$
444 550 GeV [74].

445 The Z' coupling to the non-conserved neutral current ξj_{NC} on the other hand gives
446 potentially strong constraints. The most generally applicable bounds are due to additional




Figure 5: Constraints on $U(1)_{L_\mu-L_\tau}$ without any $Z-Z'$ mixing. Shown are the preferred region to resolve the muon's $(g-2)$ at 1 and 2σ in green and exclusions from ΔN_{eff} [57,58], BaBar [75], CMS [100], and neutrino trident production in CCFR [73,74].

447 parity violation and lead to $M_{Z'}/\xi \gtrsim 1$ TeV with little dependence on the details of the UV-
 448 completion of the mass-mixing [101–103]. In addition, processes that are sensitive to the
 449 emission of the longitudinal Z' are naively expected to receive a $1/M_{Z'}$ enhanced amplitude
 450 and, therefore, meson decays such as $K \rightarrow \pi Z'$ and $B \rightarrow K Z'$ promise strong constraints.
 451 However, a certain amount of care is required when dealing with these constraints. In a
 452 theory with only mass mixing added to the SM the amplitude is divergent [103]. In the full
 453 UV-theory this divergence is canceled by the new physics omitted in the low energy theory
 454 and the divergence is replaced by a term $\propto \log(\Lambda^2/M_W^2)$, where Λ is the mass scale of the
 455 additional degrees of freedom. It has been shown that this estimate reproduces the full
 456 result of an exemplary UV-completions well provided that no cancellations occur [103]. In
 457 this case $K \rightarrow \pi Z'$ gives a limit $M_{Z'}/\xi \gtrsim 10^3$ TeV for $M_{Z'} < 100$ MeV and the CHARM
 458 beam-dump gives $\xi < 10^{-8}$ for $\text{MeV} < M_{Z'} < 350$ MeV. This indicates that the induced
 459 NSI will most likely be severely suppressed for light Z' but we would like to caution that
 460 the final answer to this question cannot be given in a model-independent fashion. We note
 461 in particular that the $(g-2)_\mu$ -motivated region of parameter space cannot give large NSI.
 462 Taken together with the constraints from Fig. 5 we see that the largest NSI in this
 463 model can be achieved with a Z' with mass above ~ 100 GeV, giving NSI as large as
 464 0.04 (Eq. (32)). Depending on the sign of $g'\xi$ this can already be in violation with the
 465 global-fit constraints of Tab. 1. However, for such an electroweak-scale Z' one does not
 466 just have rare-decay constraints [103] but also direct searches at colliders, e.g. in dilepton
 467 channels. From the LHC these are typically only given for Z' masses above 150 GeV (see
 468 e.g. Ref. [104]), leaving a gap of currently weakly constrained parameter space [105]. If
 469 future neutrino data ever hints at a large $\epsilon_{\tau\tau}^n - \epsilon_{\mu\mu}^n$ then a dedicated search for ~ 100 GeV-
 470 scale Z' would be highly desirable.

471 As we have seen above, the NSI discussion does not depend on the UV-origin of the Z -
 472 Z' mass-mixing angle ξ , although some of the constraints on ξ do. Let us briefly mention
 473 other implications of the UV completion. $Z-Z'$ mass mixing unavoidably requires a new
 474 scalar that carries both $L_\mu - L_\tau$ and electroweak charge, the simplest example being an
 475 additional scalar doublet ϕ' with the same hypercharge as the lepton doublet and $L_\mu - L_\tau$

476 charge $q_{\phi'}$. This gives [30]

$$\delta\hat{M}^2 = \frac{eg'q_{\phi'}}{s_Wc_W}\langle\phi'\rangle^2, \quad (33)$$

477 and hence

$$\epsilon_{\tau\tau}^n - \epsilon_{\mu\mu}^n = 2(\epsilon_{ee}^n - \epsilon_{\mu\mu}^n) = -\frac{1}{2\sqrt{2}G_F} \left(\frac{eg'}{s_Wc_W}\right)^2 \frac{q_{\phi'}\langle\phi'\rangle^2}{M_Z^2 M_{Z'}^2 c_\chi^2}. \quad (34)$$

478 The vacuum expectation value $\langle\phi'\rangle$ cannot be the only contribution to $M_{Z'}$, so additional
 479 electroweak singlets with $L_\mu - L_\tau$ charge are required [22, 106]. The value of $q_{\phi'}$ determines
 480 additional signatures that go beyond the simple Z - Z' mass mixing relevant for NSI. For
 481 example, in models with $q_{\phi'} = \pm 1$ off-diagonal terms in the charged lepton mass matrix
 482 are allowed which induce LFV decays $\mu \rightarrow e$ or $\tau \rightarrow e$ sector [22]; in models with $q_{\phi'} = \pm 2$
 483 on the other hand the structure is such that LFV can appear in the tau-mu sector, e.g. in
 484 $\tau \rightarrow \mu\gamma$ or $h \rightarrow \mu\tau$ [106]. Other assignments of $q_{\phi'}$ will not have any impact on LFV
 485 and essentially look like a type-I 2HDM. Since these signatures depend additionally on
 486 the scalar mixing angle(s) and the scalar mass spectrum, it is difficult to make definite
 487 predictions.

488 Conclusions

489 The origin of NSI may be a flavor-sensitive $U(1)'$. Such scenarios face a number of
 490 constraints from beam, neutrino scattering and of course oscillation measurements. We
 491 demonstrated in this paper that it is quite easy to obtain large *diagonal* NSI in anomaly-
 492 free $U(1)'$ models. The models we studied are very well motivated as they are anomaly-free
 493 when only right-handed neutrinos are introduced to the particle content of the SM. Neu-
 494 trino oscillations can often place the strongest constraints on such models if the Z' is
 495 in the 10–100 MeV region. These arguably simplest realizations of NSI lead to neutrino
 496 scattering off neutrons, protons and electrons in specific combinations.

497 Some of our key messages may be formulated as follows:

- 498 • Large *diagonal* NSI coefficients are possible via a light Z' from an anomaly-free
 499 $U(1)_X$ with $X = r_{BL}(B - L) + r_{\mu\tau}(L_\mu - L_\tau) + r_{\mu e}(L_\mu - L_e)$.
- 500 • Instead of analyzing NSI for up- and down-quarks one should rather use protons and
 501 neutrons as the natural basis.
- 502 • The sign of the NSI is fixed by the $U(1)_X$, as is which linear combination of e , p , and
 503 n is relevant for the model. NSI effects in long-baseline experiments can be easily
 504 avoided.
- 505 • For light Z' one has to carefully distinguish between NSI in oscillations (i.e. for-
 506 ward scattering) and scattering off electrons or nucleons with non-zero momentum
 507 transfer.
- 508 • NSI and neutrino scattering limits (both ν - e and (coherent) ν - q) are complementary
 509 and depend strongly on X .
- 510 • *Kinetic* mixing is not relevant for NSI, but for all other probes.

- If the $U(1)_X$ does not couple to first generation charged fermions, electron and proton NSI cancel each other exactly, and Z - Z' mass mixing is required to generate effects on neutrons. This mass mixing requires a Higgs multiplet charged under the SM and $U(1)'$ symmetries, and thus in principle testable non-standard Higgs phenomenology.

NSI effects in neutrino oscillations were shown here to be connected to various experimental probes beyond long-baseline or solar neutrino experiments, and surely a broad approach to disentangle their origin will become necessary if any sign of those effects were to be found. On the other hand, well-motivated Z' models were shown to generate NSI effects in oscillations, and should be taken into account when limits on those models are discussed.

Acknowledgments

We would like to thank Michele Maltoni for providing the values of Tab. 1 and Ivan Esteban for discussions. JH is a postdoctoral researcher of the F.R.S.-FNRS and furthermore supported, in part, by the National Science Foundation under Grant No. PHY-1620638, and by a Feodor Lynen Research Fellowship of the Alexander von Humboldt Foundation. WR is supported by the DFG with grant RO 2516/7-1 in the Heisenberg program.

References

- [1] M. Masud, A. Chatterjee and P. Mehta, *Probing CP violation signal at DUNE in presence of non-standard neutrino interactions*, J. Phys. **G43**(9), 095005 (2016), doi:10.1088/0954-3899/43/9/095005, 1510.08261.
- [2] A. de Gouvêa and K. J. Kelly, *Non-standard Neutrino Interactions at DUNE*, Nucl. Phys. **B908**, 318 (2016), doi:10.1016/j.nuclphysb.2016.03.013, 1511.05562.
- [3] M. Blennow, S. Choubey, T. Ohlsson, D. Pramanik and S. K. Raut, *A combined study of source, detector and matter non-standard neutrino interactions at DUNE*, JHEP **08**, 090 (2016), doi:10.1007/JHEP08(2016)090, 1606.08851.
- [4] S. K. Agarwalla, S. S. Chatterjee and A. Palazzo, *Degeneracy between θ_{23} octant and neutrino non-standard interactions at DUNE*, Phys. Lett. **B762**, 64 (2016), doi:10.1016/j.physletb.2016.09.020, 1607.01745.
- [5] P. Coloma, P. B. Denton, M. C. Gonzalez-Garcia, M. Maltoni and T. Schwetz, *Curtailling the Dark Side in Non-Standard Neutrino Interactions*, JHEP **04**, 116 (2017), doi:10.1007/JHEP04(2017)116, 1701.04828.
- [6] K. N. Deepthi, S. Goswami and N. Nath, *Challenges posed by non-standard neutrino interactions in the determination of δ_{CP} at DUNE*, Nucl. Phys. **B936**, 91 (2018), doi:10.1016/j.nuclphysb.2018.09.004, 1711.04840.
- [7] P. B. Denton, Y. Farzan and I. M. Shoemaker, *Testing large non-standard neutrino interactions with arbitrary mediator mass after COHERENT data*, JHEP **07**, 037 (2018), doi:10.1007/JHEP07(2018)037, 1804.03660.
- [8] L. Wolfenstein, *Neutrino Oscillations in Matter*, Phys. Rev. **D17**, 2369 (1978), doi:10.1103/PhysRevD.17.2369.

- 550 [9] M. M. Guzzo, A. Masiero and S. T. Petcov, *On the MSW effect with massless neutrinos and no mixing in the vacuum*, Phys. Lett. **B260**, 154 (1991), doi:10.1016/0370-
551 2693(91)90984-X.
552
- 553 [10] T. Ohlsson, *Status of non-standard neutrino interactions*, Rept. Prog. Phys. **76**,
554 044201 (2013), doi:10.1088/0034-4885/76/4/044201, 1209.2710.
- 555 [11] Y. Farzan and M. Tortola, *Neutrino oscillations and Non-Standard Interactions*,
556 Front.in Phys. **6**, 10 (2018), doi:10.3389/fphy.2018.00010, 1710.09360.
- 557 [12] S. Antusch, J. P. Baumann and E. Fernandez-Martinez, *Non-Standard Neutrino*
558 *Interactions with Matter from Physics Beyond the Standard Model*, Nucl. Phys.
559 **B810**, 369 (2009), doi:10.1016/j.nuclphysb.2008.11.018, 0807.1003.
- 560 [13] M. Malinsky, T. Ohlsson and H. Zhang, *Non-Standard Neutrino Inter-*
561 *actions from a Triplet Seesaw Model*, Phys. Rev. **D79**, 011301 (2009),
562 doi:10.1103/PhysRevD.79.011301, 0811.3346.
- 563 [14] T. Ohlsson, T. Schwetz and H. Zhang, *Non-standard neutrino interactions in the*
564 *Zee-Babu model*, Phys. Lett. **B681**, 269 (2009), doi:10.1016/j.physletb.2009.10.025,
565 0909.0455.
- 566 [15] D. V. Forero and W.-C. Huang, *Sizable NSI from the $SU(2)_L$ scalar*
567 *doublet-singlet mixing and the implications in DUNE*, JHEP **03**, 018 (2017),
568 doi:10.1007/JHEP03(2017)018, 1608.04719.
- 569 [16] J. Herrero-García, T. Ohlsson, S. Riad and J. Wirén, *Full parameter scan of*
570 *the Zee model: exploring Higgs lepton flavor violation*, JHEP **04**, 130 (2017),
571 doi:10.1007/JHEP04(2017)130, 1701.05345.
- 572 [17] U. K. Dey, N. Nath and S. Sadhukhan, *Non-Standard Neutrino In-*
573 *teractions in a Modified $\nu 2HDM$* , Phys. Rev. **D98**, 055004 (2018),
574 doi:10.1103/PhysRevD.98.055004, 1804.05808.
- 575 [18] T. Wang and Y.-L. Zhou, *Neutrino non-standard interactions as a portal to test*
576 *flavour symmetries* (2018), 1801.05656.
- 577 [19] I. Bischer, W. Rodejohann and X.-J. Xu, *Loop-induced Neutrino Non-Standard*
578 *Interactions*, JHEP **10**, 096 (2018), doi:10.1007/JHEP10(2018)096, 1807.08102.
- 579 [20] I. Bischer and W. Rodejohann, *General Neutrino Interactions at the DUNE Near*
580 *Detector*, Phys. Rev. **D99**, 036006 (2019), doi:10.1103/PhysRevD.99.036006, 1810.
581 02220.
- 582 [21] W. Altmannshofer, M. Tammara and J. Zupan, *Non-standard neutrino interactions*
583 *and low energy experiments* (2018), 1812.02778.
- 584 [22] J. Heeck and W. Rodejohann, *Gauged $L_\mu - L_\tau$ Symmetry at the Electroweak Scale*,
585 Phys. Rev. **D84**, 075007 (2011), doi:10.1103/PhysRevD.84.075007, 1107.5238.
- 586 [23] M. B. Wise and Y. Zhang, *Effective Theory and Simple Completions for Neutrino*
587 *Interactions*, Phys. Rev. **D90**, 053005 (2014), doi:10.1103/PhysRevD.90.053005,
588 1404.4663.
- 589 [24] A. Crivellin, G. D'Ambrosio and J. Heeck, *Addressing the LHC flavor anoma-*
590 *lies with horizontal gauge symmetries*, Phys. Rev. **D91**, 075006 (2015),
591 doi:10.1103/PhysRevD.91.075006, 1503.03477.

- 592 [25] Y. Farzan, *A model for large non-standard interactions of neutri-*
593 *nos leading to the LMA-Dark solution*, Phys. Lett. **B748**, 311 (2015),
594 doi:10.1016/j.physletb.2015.07.015, 1505.06906.
- 595 [26] Y. Farzan and I. M. Shoemaker, *Lepton Flavor Violating Non-Standard Interactions*
596 *via Light Mediators*, JHEP **07**, 033 (2016), doi:10.1007/JHEP07(2016)033, 1512.
597 09147.
- 598 [27] Y. Farzan and J. Heeck, *Neutrinophilic nonstandard interactions*, Phys. Rev. **D94**,
599 053010 (2016), doi:10.1103/PhysRevD.94.053010, 1607.07616.
- 600 [28] C. Bonilla, T. Modak, R. Srivastava and J. W. F. Valle, *$U(1)_{B_3-3L_\mu}$ gauge symme-*
601 *try as a simple description of $b \rightarrow s$ anomalies*, Phys. Rev. **D98**, 095002 (2018),
602 doi:10.1103/PhysRevD.98.095002, 1705.00915.
- 603 [29] K. S. Babu, A. Friedland, P. A. N. Machado and I. Mocioiu, *Flavor Gauge Models*
604 *Below the Fermi Scale*, JHEP **12**, 096 (2017), doi:10.1007/JHEP12(2017)096, 1705.
605 01822.
- 606 [30] P. Langacker, *The Physics of Heavy Z' Gauge Bosons*, Rev. Mod. Phys. **81**, 1199
607 (2009), doi:10.1103/RevModPhys.81.1199, 0801.1345.
- 608 [31] T. Hambye and J. Heeck, *Proton decay into charged leptons*, Phys. Rev. Lett. **120**,
609 171801 (2018), doi:10.1103/PhysRevLett.120.171801, 1712.04871.
- 610 [32] S. Bifani, S. Descotes-Genon, A. Romero Vidal and M.-H. Schune, *Review of Lepton*
611 *Universality tests in B decays*, J. Phys. **G46**, 023001 (2019), doi:10.1088/1361-
612 6471/aaf5de, 1809.06229.
- 613 [33] T. Araki, J. Heeck and J. Kubo, *Vanishing Minors in the Neutrino Mass Matrix from*
614 *Abelian Gauge Symmetries*, JHEP **07**, 083 (2012), doi:10.1007/JHEP07(2012)083,
615 1203.4951.
- 616 [34] L. N. Chang, O. Lebedev, W. Loinaz and T. Takeuchi, *Constraints on*
617 *gauged $B - 3L_\tau$ and related theories*, Phys. Rev. **D63**, 074013 (2001),
618 doi:10.1103/PhysRevD.63.074013, hep-ph/0010118.
- 619 [35] M.-C. Chen, A. de Gouvêa and B. A. Dobrescu, *Gauge Trimming of Neu-*
620 *trino Masses*, Phys. Rev. **D75**, 055009 (2007), doi:10.1103/PhysRevD.75.055009,
621 hep-ph/0612017.
- 622 [36] M.-C. Chen and J. Huang, *TeV scale seesaw model and a flavorful Z' at the LHC*,
623 Phys. Rev. **D81**, 055007 (2010), doi:10.1103/PhysRevD.81.055007, 0910.5029.
- 624 [37] H.-S. Lee and E. Ma, *Gauged $B - x_i L$ origin of R Parity and its implications*, Phys.
625 Lett. **B688**, 319 (2010), doi:10.1016/j.physletb.2010.04.032, 1001.0768.
- 626 [38] E. Salvioni, A. Strumia, G. Villadoro and F. Zwirner, *Non-universal mini-*
627 *mal Z' models: present bounds and early LHC reach*, JHEP **03**, 010 (2010),
628 doi:10.1007/JHEP03(2010)010, 0911.1450.
- 629 [39] A. M. Dziewonski and D. L. Anderson, *Preliminary reference earth model*, Phys.
630 Earth Planet. Interiors **25**, 297 (1981), doi:10.1016/0031-9201(81)90046-7.
- 631 [40] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, I. Martinez-Soler and J. Salvado,
632 *Updated Constraints on Non-Standard Interactions from Global Analysis of Oscilla-*
633 *tion Data*, JHEP **08**, 180 (2018), doi:10.1007/JHEP08(2018)180, 1805.04530.

- 634 [41] M. Lindner, F. S. Queiroz, W. Rodejohann and X.-J. Xu, *Neutrino-electron scat-*
635 *tering: general constraints on Z' and dark photon models*, JHEP **05**, 098 (2018),
636 doi:10.1007/JHEP05(2018)098, 1803.00060.
- 637 [42] M. Bauer, P. Foldenauer and J. Jaeckel, *Hunting All the Hidden Photons*, JHEP
638 **07**, 094 (2018), doi:10.1007/JHEP07(2018)094, 1803.05466.
- 639 [43] D. Akimov *et al.*, *Observation of Coherent Elastic Neutrino-Nucleus Scattering*,
640 Science **357**(6356), 1123 (2017), doi:10.1126/science.aao0990, 1708.01294.
- 641 [44] W. Maneschg, *The Status of CONUS*, In *NEUTRINO 2018, Heidelberg, Germany,*
642 *June 4-9, 2018*, doi:10.5281/zenodo.1286926.
- 643 [45] P. Coloma, M. C. Gonzalez-Garcia, M. Maltoni and T. Schwetz, *COHER-*
644 *ENT enlightenment of the neutrino dark side*, Phys. Rev. **D96**, 115007 (2017),
645 doi:10.1103/PhysRevD.96.115007, 1708.02899.
- 646 [46] J. Barranco, O. G. Miranda and T. I. Rashba, *Probing new physics with co-*
647 *herent neutrino scattering off nuclei*, JHEP **12**, 021 (2005), doi:10.1088/1126-
648 6708/2005/12/021, hep-ph/0508299.
- 649 [47] B. Dutta, R. Mahapatra, L. E. Strigari and J. W. Walker, *Sensitivity to Z' -prime and*
650 *nonstandard neutrino interactions from ultralow threshold neutrino-nucleus coher-*
651 *ent scattering*, Phys. Rev. **D93**, 013015 (2016), doi:10.1103/PhysRevD.93.013015,
652 1508.07981.
- 653 [48] D. K. Papoulias and T. S. Kosmas, *Standard and Nonstandard Neutrino-Nucleus*
654 *Reactions Cross Sections and Event Rates to Neutrino Detection Experiments*, Adv.
655 High Energy Phys. **2015**, 763648 (2015), doi:10.1155/2015/763648, 1502.02928.
- 656 [49] M. Lindner, W. Rodejohann and X.-J. Xu, *Coherent Neutrino-Nucleus*
657 *Scattering and new Neutrino Interactions*, JHEP **03**, 097 (2017),
658 doi:10.1007/JHEP03(2017)097, 1612.04150.
- 659 [50] J. Liao and D. Marfatia, *COHERENT constraints on nonstandard neutrino interac-*
660 *tions*, Phys. Lett. **B775**, 54 (2017), doi:10.1016/j.physletb.2017.10.046, 1708.04255.
- 661 [51] J. Billard, J. Johnston and B. J. Kavanagh, *Prospects for exploring New*
662 *Physics in Coherent Elastic Neutrino-Nucleus Scattering*, JCAP **1811**, 016 (2018),
663 doi:10.1088/1475-7516/2018/11/016, 1805.01798.
- 664 [52] P. Galison and A. Manohar, *Two Z 's or not two Z 's?*, Phys. Lett. **136B**, 279
665 (1984), doi:10.1016/0370-2693(84)91161-4.
- 666 [53] B. Holdom, *Two $U(1)$'s and ϵ Charge Shifts*, Phys. Lett. **166B**, 196 (1986),
667 doi:10.1016/0370-2693(86)91377-8.
- 668 [54] K. S. Babu, C. F. Kolda and J. March-Russell, *Implications of generalized Z - Z'*
669 *mixing*, Phys. Rev. **D57**, 6788 (1998), doi:10.1103/PhysRevD.57.6788, hep-ph/
670 9710441.
- 671 [55] J. Heeck and W. Rodejohann, *Kinetic and mass mixing with three abelian groups*,
672 Phys. Lett. **B705**, 369 (2011), doi:10.1016/j.physletb.2011.10.050, 1109.1508.

- 673 [56] B. Ahlgren, T. Ohlsson and S. Zhou, *Comment on “Is Dark Matter with Long-Range*
674 *Interactions a Solution to All Small-Scale Problems of Λ Cold Dark Matter Cosmol-*
675 *ogy?”*, Phys. Rev. Lett. **111**, 199001 (2013), doi:10.1103/PhysRevLett.111.199001,
676 1309.0991.
- 677 [57] A. Kamada and H.-B. Yu, *Coherent Propagation of PeV Neutrinos and the*
678 *Dip in the Neutrino Spectrum at IceCube*, Phys. Rev. **D92**, 113004 (2015),
679 doi:10.1103/PhysRevD.92.113004, 1504.00711.
- 680 [58] A. Kamada, K. Kaneta, K. Yanagi and H.-B. Yu, *Self-interacting dark mat-*
681 *ter and muon $g - 2$ in a gauged $U(1)_{L_\mu - L_\tau}$ model*, JHEP **06**, 117 (2018),
682 doi:10.1007/JHEP06(2018)117, 1805.00651.
- 683 [59] P. B. Denton, Y. Farzan and I. M. Shoemaker, *Activating the fourth neutrino of the*
684 *$3 + 1$ scheme*, Phys. Rev. **D99**, 035003 (2019), doi:10.1103/PhysRevD.99.035003,
685 1811.01310.
- 686 [60] J. Heeck and W. Rodejohann, *Gauged $L_\mu - L_\tau$ and different Muon Neutrino and*
687 *Anti-Neutrino Oscillations: MINOS and beyond*, J. Phys. **G38**, 085005 (2011),
688 doi:10.1088/0954-3899/38/8/085005, 1007.2655.
- 689 [61] M. B. Wise and Y. Zhang, *Lepton Flavorful Fifth Force and Depth-dependent Neu-*
690 *trino Matter Interactions*, JHEP **06**, 053 (2018), doi:10.1007/JHEP06(2018)053,
691 1803.00591.
- 692 [62] M. Bustamante and S. K. Agarwalla, *Universe’s Worth of Electrons to Probe Long-*
693 *Range Interactions of High-Energy Astrophysical Neutrinos*, Phys. Rev. Lett. **122**,
694 061103 (2019), doi:10.1103/PhysRevLett.122.061103, 1808.02042.
- 695 [63] O. G. Miranda, M. A. Tortola and J. W. F. Valle, *Are solar neutrino oscillations ro-*
696 *bust?*, JHEP **10**, 008 (2006), doi:10.1088/1126-6708/2006/10/008, hep-ph/0406280.
- 697 [64] M. C. Gonzalez-Garcia, M. Maltoni and J. Salvado, *Testing matter effects in*
698 *propagation of atmospheric and long-baseline neutrinos*, JHEP **05**, 075 (2011),
699 doi:10.1007/JHEP05(2011)075, 1103.4365.
- 700 [65] M. C. Gonzalez-Garcia and M. Maltoni, *Determination of matter poten-*
701 *tial from global analysis of neutrino oscillation data*, JHEP **09**, 152 (2013),
702 doi:10.1007/JHEP09(2013)152, 1307.3092.
- 703 [66] P. Bakhti and Y. Farzan, *Shedding light on LMA-Dark solar neutrino solution by*
704 *medium baseline reactor experiments: JUNO and RENO-50*, JHEP **07**, 064 (2014),
705 doi:10.1007/JHEP07(2014)064, 1403.0744.
- 706 [67] P. Coloma and T. Schwetz, *Generalized mass ordering degeneracy*
707 *in neutrino oscillation experiments*, Phys. Rev. **D94**, 055005 (2016),
708 doi:10.1103/PhysRevD.94.055005, [Erratum: Phys. Rev. D95, 079903 (2017)],
709 1604.05772.
- 710 [68] R. Barbieri and T. E. O. Ericson, *Evidence Against the Existence of a Low Mass*
711 *Scalar Boson from Neutron-Nucleus Scattering*, Phys. Lett. **57B**, 270 (1975),
712 doi:10.1016/0370-2693(75)90073-8.
- 713 [69] V. Barger, C.-W. Chiang, W.-Y. Keung and D. Marfatia, *Proton size anomaly*, Phys.
714 Rev. Lett. **106**, 153001 (2011), doi:10.1103/PhysRevLett.106.153001, 1011.3519.

- 715 [70] S. Tulin, *New weakly-coupled forces hidden in low-energy QCD*, Phys. Rev. **D89**,
716 114008 (2014), doi:10.1103/PhysRevD.89.114008, 1404.4370.
- 717 [71] P. Ilten, Y. Soreq, M. Williams and W. Xue, *Serendipity in dark photon searches*,
718 JHEP **06**, 004 (2018), doi:10.1007/JHEP06(2018)004, 1801.04847.
- 719 [72] M. Tanabashi *et al.*, *Review of Particle Physics*, Phys. Rev. **D98**, 030001 (2018),
720 doi:10.1103/PhysRevD.98.030001.
- 721 [73] S. Mishra *et al.*, *Neutrino tridents and $W-Z$ interference*, Phys. Rev. Lett. **66**, 3117
722 (1991), doi:10.1103/PhysRevLett.66.3117.
- 723 [74] W. Altmannshofer, S. Gori, M. Pospelov and I. Yavin, *Neutrino Trident Production:
724 A Powerful Probe of New Physics with Neutrino Beams*, Phys. Rev. Lett. **113**,
725 091801 (2014), doi:10.1103/PhysRevLett.113.091801, 1406.2332.
- 726 [75] J. P. Lees *et al.*, *Search for a muonic dark force at BABAR*, Phys. Rev. **D94**, 011102
727 (2016), doi:10.1103/PhysRevD.94.011102, 1606.03501.
- 728 [76] E. Ma, *Gauged $B-3L_\tau$ and radiative neutrino masses*, Phys. Lett. **B433**, 74 (1998),
729 doi:10.1016/S0370-2693(98)00599-1, hep-ph/9709474.
- 730 [77] E. Ma and D. P. Roy, *Phenomenology of the $B-3L_\tau$ gauge boson*, Phys. Rev. **D58**,
731 095005 (1998), doi:10.1103/PhysRevD.58.095005, hep-ph/9806210.
- 732 [78] E. Ma and U. Sarkar, *Gauged $B-3L_\tau$ and baryogenesis*, Phys. Lett. **B439**, 95
733 (1998), doi:10.1016/S0370-2693(98)01019-3, hep-ph/9807307.
- 734 [79] E. Ma and D. P. Roy, *Minimal seesaw model for atmospheric and solar neutrino
735 oscillations*, Phys. Rev. **D59**, 097702 (1999), doi:10.1103/PhysRevD.59.097702,
736 hep-ph/9811266.
- 737 [80] M. Lindner, M. Platscher and F. S. Queiroz, *A Call for New Physics: The Muon
738 Anomalous Magnetic Moment and Lepton Flavor Violation*, Phys. Rept. **731**, 1
739 (2018), doi:10.1016/j.physrep.2017.12.001, 1610.06587.
- 740 [81] S. Davidson, S. Forte, P. Gambino, N. Rius and A. Strumia, *Old and new physics
741 interpretations of the NuTeV anomaly*, JHEP **02**, 037 (2002), doi:10.1088/1126-
742 6708/2002/02/037, hep-ph/0112302.
- 743 [82] E. Ma, D. P. Roy and U. Sarkar, *A Seesaw model for atmospheric and solar neutrino
744 oscillations*, Phys. Lett. **B444**, 391 (1998), doi:10.1016/S0370-2693(98)01395-1,
745 hep-ph/9810309.
- 746 [83] X. G. He, G. C. Joshi, H. Lew and R. R. Volkas, *New- Z' phenomenology*, Phys.
747 Rev. **D43**, 22 (1991), doi:10.1103/PhysRevD.43.R22.
- 748 [84] X.-G. He, G. C. Joshi, H. Lew and R. R. Volkas, *Simplest Z' model*, Phys. Rev.
749 **D44**, 2118 (1991), doi:10.1103/PhysRevD.44.2118.
- 750 [85] S. Bilmis, I. Turan, T. M. Aliev, M. Deniz, L. Singh and H. T. Wong, *Constraints
751 on Dark Photon from Neutrino-Electron Scattering Experiments*, Phys. Rev. **D92**,
752 033009 (2015), doi:10.1103/PhysRevD.92.033009, 1502.07763.
- 753 [86] M. Deniz *et al.*, *Measurement of $\bar{\nu}_e$ -Electron Scattering Cross Section with a CsI(Tl)
754 Scintillating Crystal Array at the Kuo-Sheng Nuclear Power Reactor*, Phys. Rev.
755 **D81**, 072001 (2010), doi:10.1103/PhysRevD.81.072001, 0911.1597.

- 756 [87] J. P. Lees *et al.*, *Search for a Dark Photon in e^+e^- Collisions at BaBar*, Phys. Rev.
757 Lett. **113**, 201801 (2014), doi:10.1103/PhysRevLett.113.201801, 1406.2980.
- 758 [88] J. Heeck, *Unbroken $B - L$ symmetry*, Phys. Lett. **B739**, 256 (2014),
759 doi:10.1016/j.physletb.2014.10.067, 1408.6845.
- 760 [89] E. M. Riordan *et al.*, *A Search for Short Lived Axions in an Electron Beam Dump*
761 *Experiment*, Phys. Rev. Lett. **59**, 755 (1987), doi:10.1103/PhysRevLett.59.755.
- 762 [90] J. D. Bjorken, S. Ecklund, W. R. Nelson, A. Abashian, C. Church, B. Lu, L. W.
763 Mo, T. A. Nunamaker and P. Rassmann, *Search for Neutral Metastable Penetrat-*
764 *ing Particles Produced in the SLAC Beam Dump*, Phys. Rev. **D38**, 3375 (1988),
765 doi:10.1103/PhysRevD.38.3375.
- 766 [91] A. Bross, M. Crisler, S. H. Pordes, J. Volk, S. Errede and J. Wrbanek, *A Search*
767 *for Shortlived Particles Produced in an Electron Beam Dump*, Phys. Rev. Lett. **67**,
768 2942 (1991), doi:10.1103/PhysRevLett.67.2942.
- 769 [92] M. Davier and H. Nguyen Ngoc, *An Unambiguous Search for a Light Higgs Boson*,
770 Phys. Lett. **B229**, 150 (1989), doi:10.1016/0370-2693(89)90174-3.
- 771 [93] J. Blümlein *et al.*, *Limits on neutral light scalar and pseudoscalar particles in a*
772 *proton beam dump experiment*, Z. Phys. **C51**, 341 (1991), doi:10.1007/BF01548556.
- 773 [94] J. Blümlein *et al.*, *Limits on the mass of light (pseudo)scalar particles from Bethe-*
774 *Heitler e^+e^- and $\mu^+\mu^-$ pair production in a proton-iron beam dump experiment*,
775 Int. J. Mod. Phys. **A7**, 3835 (1992), doi:10.1142/S0217751X9200171X.
- 776 [95] D. Banerjee *et al.*, *Search for a Hypothetical 16.7 MeV Gauge Boson and Dark*
777 *Photons in the NA64 Experiment at CERN*, Phys. Rev. Lett. **120**, 231802 (2018),
778 doi:10.1103/PhysRevLett.120.231802, 1803.07748.
- 779 [96] R. Foot, *New Physics From Electric Charge Quantization?*, Mod. Phys. Lett. **A6**,
780 527 (1991), doi:10.1142/S0217732391000543.
- 781 [97] G. Bellini *et al.*, *Precision measurement of the ${}^7\text{Be}$ solar neutrino*
782 *interaction rate in Borexino*, Phys. Rev. Lett. **107**, 141302 (2011),
783 doi:10.1103/PhysRevLett.107.141302, 1104.1816.
- 784 [98] Y. Kaneta and T. Shimomura, *On the possibility of a search for the $L_\mu - L_\tau$ gauge*
785 *boson at Belle-II and neutrino beam experiments*, PTEP **2017**(5), 053B04 (2017),
786 doi:10.1093/ptep/ptx050, 1701.00156.
- 787 [99] P. Foldenauer, *Light dark matter in a gauged $U(1)_{L_\mu-L_\tau}$ model*, Phys. Rev. **D99**,
788 035007 (2019), doi:10.1103/PhysRevD.99.035007, 1808.03647.
- 789 [100] A. M. Sirunyan *et al.*, *Search for an $L_\mu - L_\tau$ gauge boson using $Z \rightarrow 4\mu$ events in*
790 *proton-proton collisions at $\sqrt{s} = 13$ TeV* (2018), 1808.03684.
- 791 [101] H. Davoudiasl, H.-S. Lee and W. J. Marciano, *'Dark' Z implications for Parity*
792 *Violation, Rare Meson Decays, and Higgs Physics*, Phys. Rev. **D85**, 115019 (2012),
793 doi:10.1103/PhysRevD.85.115019, 1203.2947.
- 794 [102] H. Davoudiasl, H.-S. Lee and W. J. Marciano, *Muon $g - 2$, rare kaon de-*
795 *cays, and parity violation from dark bosons*, Phys. Rev. **D89**, 095006 (2014),
796 doi:10.1103/PhysRevD.89.095006, 1402.3620.

- 797 [103] J. A. Dror, R. Lasenby and M. Pospelov, *Light vectors coupled to bosonic currents*
798 (2018), 1811.00595.
- 799 [104] M. Aaboud *et al.*, *Search for new high-mass phenomena in the dilepton final state*
800 *using 36 fb⁻¹ of proton-proton collision data at $\sqrt{s} = 13$ TeV with the ATLAS*
801 *detector*, JHEP **10**, 182 (2017), doi:10.1007/JHEP10(2017)182, 1707.02424.
- 802 [105] B. Batell, M. Pospelov and B. Shuve, *Shedding Light on Neutrino Masses with Dark*
803 *Forces*, JHEP **08**, 052 (2016), doi:10.1007/JHEP08(2016)052, 1604.06099.
- 804 [106] J. Heck, M. Holthausen, W. Rodejohann and Y. Shimizu, *Higgs $\rightarrow \mu\tau$ in*
805 *Abelian and non-Abelian flavor symmetry models*, Nucl. Phys. **B896**, 281 (2015),
806 doi:10.1016/j.nuclphysb.2015.04.025, 1412.3671.