

Emergence of \mathcal{PT} -symmetry breaking in open quantum systems

J. Huber^{1*}, P. Kirton^{1,2}, S. Rotter³, P. Rabl¹,

1 Vienna Center for Quantum Science and Technology, Atominstitut, TU Wien, 1040 Vienna, Austria

2 Department of Physics and SUPA, University of Strathclyde, Glasgow G4 0NG, UK

3 Institute for Theoretical Physics, TU Wien, 1040 Vienna, Austria

* julian.huber@tuwien.ac.at

September 28, 2020

Abstract

The effect of \mathcal{PT} -symmetry breaking in coupled systems with balanced gain and loss has recently attracted considerable attention and has been demonstrated in various photonic, electrical and mechanical systems in the classical regime. However, it is still an unsolved problem how to generalize the concept of \mathcal{PT} symmetry to the quantum domain, where the conventional definition in terms of non-Hermitian Hamiltonians is not applicable. Here we introduce a symmetry relation for Liouville operators that describe the dissipative evolution of arbitrary open quantum systems. Specifically, we show that the invariance of the Liouvillian under this symmetry transformation implies the existence of stationary states with preserved and broken parity symmetry. As the dimension of the Hilbert space grows, the transition between these two limiting phases becomes increasingly sharp and the classically expected \mathcal{PT} -symmetry breaking transition is recovered. This quantum-to-classical correspondence allows us to establish a common theoretical framework to identify and accurately describe \mathcal{PT} -symmetry breaking effects in a large variety of physical systems, operated both in the classical and quantum regimes.

Contents

1	Introduction	2
2	\mathcal{PT}-symmetric quantum systems	3
3	Phenomenology	5
4	Existence of a fully symmetric steady state	6
5	Symmetry-breaking transition	7
6	Generalizations	8
7	Discussion and Conclusions	9
A	Mean field equations of motion	11

B Fully symmetric steady state	12
C Random jump operators	13
References	14

1 Introduction

The breaking of parity and time-reversal (\mathcal{PT}) symmetry has been widely studied in dissipative systems with an exact balance between gain and loss [1–5]. Owing to this symmetry, the dynamical matrix describing such systems may exhibit a purely real eigenvalue spectrum, despite a constant exchange of energy with the environment. As the dissipation rates are increased above a critical value, at least one pair of eigenvalues develops imaginary parts with opposite signs and the corresponding gain and loss eigenmodes no longer exhibit the symmetry of the underlying equations of motion. Over the past years, this effect has attracted considerable attention and has been demonstrated in various optical [6–8], electrical [9] and mechanical [10] settings. Apart from purely fundamental interest, this mechanism also has many important practical consequences, for example, for the operation of multi-mode lasers [11, 12], enhanced measurements [13–18], the band-structure of dissipative lattice systems [19–22] or energy transport at macroscopic [23] and microscopic [24, 25] scales.

In connection with \mathcal{PT} -symmetric systems it is common to use the terminology of non-Hermitian ‘Hamilton operators’. However, the effect described above is *a priori* only defined for classical systems that can be modeled in terms of a complex-valued dynamical matrix [26]. Indeed, in a full master equation (ME) formulation of open quantum system [28], there is no such transition between purely real and complex conjugated eigenvalues of the corresponding Liouville operator. Also, at a microscopic level, the time-reversal equivalence between loss and gain is broken by quantum fluctuations [29–32]. Therefore, it is still an unresolved question how to formally define \mathcal{PT} -symmetry for dissipative quantum systems [33] and if the breaking of this symmetry can exist at all at a microscopic level [32]. In several previous studies this question has been addressed by looking at coupled quantum oscillators [17, 29–32, 35–40] or bosonic atoms [41] with gain and loss, or at equivalent coherent, but unstable systems [42]. In such settings, the symmetry-breaking effect can still be observed in the dynamics of the mean amplitudes, which simply reproduce the classical equations of motion, while quantum effects lead to increased fluctuations. However, these findings cannot be generalized to non-bosonic systems with a finite dimensional Hilbert space, where mean-field approximations are neither justified nor uniquely defined. The study of purely linear models also provides no insights into the stationary states of \mathcal{PT} -symmetric systems, which do not exist unless nonlinear saturation effects are taken into account [24, 31].

In this work we introduce a symmetry transformation for Liouville operators, which extends the conventional definition of \mathcal{PT} symmetry to arbitrary open quantum systems. We show that under very generic conditions, the existence of this symmetry implies that the steady state of the system can be tuned between a fully symmetric and a symmetry-broken phase. While the change from one to the other limiting state is always continuous, it becomes more and more pronounced as the dimension of the Hilbert space is increased, and a sharp \mathcal{PT} -symmetry breaking transition emerges in the semiclassical limit. This

quantum-to-classical correspondence allows us to establish a unified theoretical framework for analyzing \mathcal{PT} -symmetry breaking effects in a wide range of physical systems and to identify characteristic properties and experimentally observable features that are common to all of them.

2 \mathcal{PT} -symmetric quantum systems

We consider a generic bipartite quantum system with a total Hamiltonian H . The two subsystems, A and B, have the same Hilbert space dimension, d , and they are subject to dissipation described by the local jump operators c_A and c_B , respectively. The ME for the system density operator ρ can then be written as ($\hbar = 1$) [28]

$$\begin{aligned} \dot{\rho} &= -i[H, \rho] + \mathcal{D}[c_A]\rho + \mathcal{D}[c_B]\rho \\ &= -i\left(H_{\text{eff}}\rho - \rho H_{\text{eff}}^\dagger\right) + 2c_A\rho c_A^\dagger + 2c_B\rho c_B^\dagger \equiv \mathcal{L}\rho. \end{aligned} \quad (1)$$

Here $\mathcal{D}[c]\rho = 2c\rho c^\dagger - c^\dagger c\rho - \rho c^\dagger c$ and $\mathcal{L} \equiv \mathcal{L}[H; c_A, c_B]$ is the Liouvillian superoperator, which is specified, according to the first line in Eq. (1), in terms of the Hermitian Hamiltonian H and the two local jump operators c_A and c_B . In the second line of Eq. (1), the ME equation has been rewritten such that the first part describes the evolution of a quantum state under the action of the effective non-Hermitian Hamiltonian

$$H_{\text{eff}} = H - ic_A^\dagger c_A - ic_B^\dagger c_B. \quad (2)$$

This term does not conserve the norm of the state and thus the recycling terms $\sim 2c\rho c^\dagger$ must be added to obtain a trace-preserving dynamics.

Given the decomposition of a ME in Eq. (1), it is tempting to define \mathcal{PT} -symmetric quantum systems in analogy to the classical case [4, 5], namely as open quantum systems where $(\mathcal{PT})H_{\text{eff}}(\mathcal{PT})^{-1} = H_{\text{eff}}$. Here \mathcal{P} is the parity operator with $\mathcal{P}(A \otimes B)\mathcal{P}^{-1} = B \otimes A$ and $\mathcal{T}i\mathcal{T}^{-1} = -i$. However, H_{eff} has only negative imaginary parts because the norm of a state evolving under H_{eff} always decreases and this symmetry relation can only be satisfied in closed systems. The same is then also true for the eigenvalues of the full Liouville operator \mathcal{L} whose real part must always be negative or zero. Therefore, while according to Eq. (2) there is a natural way to introduce non-Hermitian Hamiltonians in open quantum systems and even probe them via conditional measurements [43–47], there are no \mathcal{PT} -symmetric (super-)operators in the conventional sense. To overcome this problem, one could be less strict and only demand that the \mathcal{PT} symmetry criterion applies to the resulting equations of motion for the expectation values of system operators. However, this identification is restricted to linear bosonic systems, where the quantum and classical dynamics of mean values is the same. For fermions, spins or other finite-dimensional quantum system the same method is not applicable, as illustrated by a simple example in Appendix A. We conclude that none of approaches used in the literature so far offers meaningful way to define \mathcal{PT} symmetry for generic quantum systems.

To provide such a definition, it is important to keep in mind that the relevant physical effect of the \mathcal{T} -operator is to exchange loss and gain and not to implement a time-reversal transformation. While for classical systems both operations are equivalent and usually no distinction is made, this is no longer true for quantum systems. In the simplest example of a quantum harmonic oscillator the effect of loss with rate Γ is modeled by a jump operator $c = \sqrt{\Gamma}a$, where a is the annihilation operator. In turn, the effect of gain with the same rate can be described by modifying the jump operator to be $c = \sqrt{\Gamma}a^\dagger$. Therefore, in this case we find that the transformation between loss and gain is implemented in the ME

formalism by replacing the jump operator by its adjoint, $c \rightarrow c^\dagger$, and not by replacing i with $-i$ in any part of the ME.

Guided by this explicit example, we introduce the following anti-unitary transformation for operators O ,

$$\mathbb{P}\mathbb{T}(O) = \mathcal{P}O^\dagger\mathcal{P}^{-1}, \quad (3)$$

and define an open quantum system to be $\mathcal{P}\mathcal{T}$ -symmetric, if the corresponding Liouvillian satisfies

$$\mathcal{L}[\mathbb{P}\mathbb{T}(H); \mathbb{P}\mathbb{T}(c_A), \mathbb{P}\mathbb{T}(c_B)] = \mathcal{L}[H; c_A, c_B]. \quad (4)$$

This condition implies that the Hamiltonian H is parity-symmetric and that the local jump operators are of the form

$$c_A = \sqrt{\Gamma}O \otimes \mathbb{1}, \quad c_B = \sqrt{\Gamma}\mathbb{1} \otimes O^\dagger, \quad (5)$$

where O can be an arbitrary dimensionless operator.

Before we proceed with a discussion of the physical implications of Eq. (4), let us briefly comment on some related symmetries and classifications of open quantum systems. First of all, we remark that the definition given in Eq. (4) differs from the $\mathcal{P}\mathcal{T}$ -symmetric Liouville operators introduced originally in Ref. [33] and analyzed for different spin models in Refs. [34, 48, 49]. In those systems, the eigenvalues of the Liouvillian exhibit a particular symmetry in terms of a cross-shaped pattern in the complex plane, which is broken at larger values of Γ . However, this property is rather specific and while the systems studied in Refs. [33, 34, 48, 49] satisfy Eq. (4) with a redefinition of \mathcal{P} , none of the examples discussed below exhibits the symmetry considered in these references when $d > 2$. Therefore, based on currently known examples, it seems difficult to link this symmetry in the eigenvalue structure to the conventional effect of $\mathcal{P}\mathcal{T}$ -symmetry breaking in the appropriate semi-classical limit.

Moreover, there has been considerable interest recently in the classifications of the symmetries of non-Hermitian Hamiltonians [19–22] and Liouvillian [50, 51] lattice models for dissipative fermions. Connected to the classifications of non-Hermitian Hamiltonians, we emphasize that according to Eq. (3) the $\mathbb{P}\mathbb{T}$ map is applied to the *Hermitian* Hamiltonian H and the jump operators separately, i.e., the transformed Liouvillian is constructed as

$$\mathcal{L}[\mathbb{P}\mathbb{T}(H); \mathbb{P}\mathbb{T}(c_A), \mathbb{P}\mathbb{T}(c_B)] = -i[\mathbb{P}\mathbb{T}(H), \rho] + \mathcal{D}[\mathbb{P}\mathbb{T}(c_A)]\rho + \mathcal{D}[\mathbb{P}\mathbb{T}(c_B)]\rho. \quad (6)$$

This ensures that the whole transformation remains physically meaningful, which would not be the case when applying the same map to the non-Hermitian Hamiltonian H_{eff} , as defined in Eq. (2). Second, the Hermitian adjoint operation $O \rightarrow O^\dagger$ in Eq. (3) is not related to a state transformation [50–52]. Also, it is applied to the full operators H , c_A and c_B , rather than to individual fermionic operators. Therefore, although reminiscent of the well-known particle-hole or chiral transformations for fermions [52], the symmetry introduced here is, in general, a different one. For example, consider a jump operator $c = a^\dagger a$, where a is a fermionic annihilation operator. Then, leaving parity aside, $\mathbb{P}\mathbb{T}(a^\dagger a) = a^\dagger a \neq aa^\dagger$. While basic $\mathbb{P}\mathbb{T}$ -symmetric systems with linear jump operators, i.e., $c_A = a^\dagger$ and $c_B = a$, may also exhibit a particle-hole symmetry as well (see, e.g., the example in Appendix A) this symmetry is not, per se, relevant for the physics discussed below. Indeed, it is always true that $H^\dagger = H$ and, apart from parity, no additional symmetries of the Hamiltonian are required to satisfy Eq. (4). Therefore, the Liouvillian symmetry introduced in Eq. (3) goes beyond the scope of previous classifications of dissipative fermionic systems and is also more generally applicable. In particular, it doesn't rely on a specific operator algebra or a local Hilbert space dimension, which allows us to discuss the physical consequences




Figure 1: Two basic examples of \mathcal{PT} -symmetric quantum systems with a finite Hilbert space dimension d : (a) two coupled spin $S = (d - 1)/2$ systems and (b) two coupled harmonic oscillators with a finite number of energy levels. In (c) and (d) we plot the corresponding dependence of the symmetry parameter Δ defined in Eq. (8) on the ratio Γ/g . In (c) the line for $S = \infty$ is obtained from a Holstein-Primakoff approximation [see discussion of Eq. (14)].

of this symmetry and the quantum-to-classical correspondence of \mathcal{PT} -symmetry breaking in a much broader context.

3 Phenomenology

As a first step, let us illustrate the physical implications of Eq. (4) in terms of two simple examples: (i) Two coupled spin $S = (d - 1)/2$ systems with $O = S^+$, where $S^+ = S^x + iS^y$ is the spin raising operator, and (ii) two coupled harmonic oscillators with $O = a^\dagger$. In the second example we introduce a finite cutoff occupation number, i.e., $a^\dagger|n = d - 1\rangle = 0$. This cutoff mimics the effect of saturation in realistic systems [24] and allows us to vary the Hilbert space dimension. In both examples we consider a Hamiltonian of the form

$$H = g(O_A O_B^\dagger + O_A^\dagger O_B), \quad (7)$$

where $O_A = O \otimes \mathbb{1}$ and $O_B = \mathbb{1} \otimes O$. This Hamiltonian describes the coherent exchange of energy between the two subsystems with a strength g . The resulting Liouvillian, $\mathcal{L}[H; \sqrt{\Gamma}O_A, \sqrt{\Gamma}O_B^\dagger]$, then satisfies Eq. (4).

We calculate the steady state, ρ_0 , satisfying $\mathcal{L}\rho_0 = 0$, for different ratios Γ/g and show in Fig. 1 the symmetry parameter [31]

$$\Delta = \frac{|\langle O_A^\dagger O_A - O_B^\dagger O_B \rangle|}{\langle O_A^\dagger O_A + O_B^\dagger O_B \rangle} \leq 1. \quad (8)$$

This is an experimentally observable quantity, only requiring measurements of local operators, which provides a measure for the symmetry of the system, i.e., $\Delta = 0$ for a parity-symmetric density operator, $\mathcal{P}\rho\mathcal{P}^{-1} = \rho$. For the current examples, Δ represents the normalized population imbalance between the two subsystems. For small dimensions

d , this parameter changes gradually from 0 to 1 with increasing Γ . This smooth variation is expected since observables of finite dimensional quantum systems cannot exhibit any non-analytic behavior. However, as the system size increases, Δ vanishes for $\Gamma/g < 1$ in the limit $d \rightarrow \infty$, while it retains a finite value for $\Gamma/g > 1$. In both examples, the critical ratio is $\Gamma/g = 1$, which corresponds to the dynamical \mathcal{PT} -symmetry breaking point of an equivalent linear oscillator system with gain and loss [4, 5]. We thus conclude that \mathcal{PT} -symmetry breaking, i.e., a non-analytic transition between two steady states with different symmetries, exists even for non-harmonic and finite dimensional quantum systems, but only as an emergent phenomenon in the semiclassical limit.

To obtain better insights into the nature of the two phases, we plot in Fig. 2(a) the purity, $P = \text{Tr}\{\rho_0^2\}$, for the steady state of the spin system. This quantity again exhibits a sharp transition around $\Gamma = g$ and shows that the symmetric and symmetry-broken phases are characterized by a highly mixed and an almost pure steady state, respectively. More precisely, the scaling $P(\Gamma \rightarrow 0) \simeq d^{-2}$ implies that in the symmetric phase the steady state is close to the maximally mixed state, $\rho_0(\Gamma \ll g) \simeq \mathbb{1}/d^2$. This indicates that for $\Gamma < g$ the gain and loss processes cancel out on average while quantum fluctuations still occur with rate Γ and completely randomize the system's long-time dynamics [24, 31]. In contrast, for $\Gamma > g$, the incoherent processes dominate and pump the spins into the polarised pure state, $\rho_0(\Gamma \gg g) \simeq |\psi_0\rangle\langle\psi_0|$, which satisfies $O_A|\psi_0\rangle = O_B^\dagger|\psi_0\rangle = 0$. Closer to the transition point, the coherent coupling creates excitations $\sim O_A^\dagger O_B|\psi_0\rangle$ on top of this state, which are strongly correlated. As shown in Fig. 2(b), this results in a characteristic peak in the entanglement negativity \mathcal{N} around the transition point, which is a measure of non-classical correlations between the two subsystems [53, 54]. These correlations vanish again in the symmetric phase due to fluctuations. Consistent with similar features observed in saturable oscillator systems [24], this peak in the entanglement shows that even for $d \gg 1$ the \mathcal{PT} -symmetry breaking transition retains genuine quantum mechanical properties.

4 Existence of a fully symmetric steady state

We will now show that the properties discussed above for specific examples are indeed a general consequence of the symmetry relation in Eq. (4). Firstly, we demonstrate that, for any Liouvillian that satisfies this condition and where the spectrum of H is non-degenerate, the fully mixed state,

$$\rho_0(\Gamma \rightarrow 0^+) = \frac{\mathbb{1}}{d^2}, \quad (9)$$

is a stationary state of \mathcal{L} in the limit of a vanishingly small, but finite Γ . To do so we decompose $\mathcal{L} = \mathcal{L}_H + \mathcal{L}_\Gamma$, where $\mathcal{L}_H\rho = -i[H, \rho]$ describes the coherent evolution and $\mathcal{L}_\Gamma\rho = \sum_{\eta=A,B}(2c_\eta\rho c_\eta^\dagger - c_\eta^\dagger c_\eta\rho - \rho c_\eta^\dagger c_\eta)$. Further, we write the density operator in the eigenbasis of H as

$$\rho = \sum_{n,m} \rho_{n,m} |E_n\rangle\langle E_m|, \quad (10)$$

where $H|E_n\rangle = E_n|E_n\rangle$. For $\Gamma = 0$ any diagonal state with $\rho_{n,m} = 0$ for $n \neq m$ is a stationary solution of the ME, but the populations $p_n = \rho_{n,n}$ are not uniquely determined. To show that for small but finite Γ only the fully mixed state is dynamically stable, we write $p_n = 1/d^2 + \delta p_n$. Up to first order in Γ we then obtain

$$\delta \dot{p}_n = \frac{2}{d^2} \langle E_n | \left([c_A, c_A^\dagger] + [c_B, c_B^\dagger] \right) | E_n \rangle. \quad (11)$$

We now make use of the relation $\mathcal{P}c_B\mathcal{P}^{-1} = c_A^\dagger$, which follows from Eq. (5), and the fact that the eigenstates of H are also eigenstates of the parity operator, i.e., $\mathcal{P}|E_n\rangle = \pm|E_n\rangle$.




Figure 2: (a) Plot of the purity P of the steady state of a \mathcal{PT} -symmetric spin dimer [see Fig. 1(a)] as a function of the dissipation rate and for different values of S . The inset shows that the purity satisfies $P \simeq 1/d^2$ for $\Gamma \ll g$. (b) Plot of the entanglement negativity \mathcal{N} [53, 54] for the same model. In (c) and (d) the same quantities are plotted for \mathcal{PT} -symmetric systems with random jump operators, as described in Appendix C, and in (e) and (f) for the generalized spin model defined in Sec. 6.

This allows us to rewrite

$$\begin{aligned} \langle E_n | [c_B, c_B^\dagger] | E_n \rangle &= \langle E_n | \mathcal{P} [c_B, c_B^\dagger] \mathcal{P}^{-1} | E_n \rangle \\ &= - \langle E_n | [c_A, c_A^\dagger] | E_n \rangle, \end{aligned} \quad (12)$$

and $\delta \dot{p}_n = 0$. This result shows that for \mathcal{PT} -symmetric quantum systems the fully mixed state is stationary in the presence of a small amount of dissipation, even when each individual jump operator $c_{A,B}$ would drive the system into a polarized state. Note that the analysis presented here assumed a non-degenerate spectrum, i.e., the absence of any additional symmetries, \mathcal{S} , other than parity. In the general case the same arguments still hold as long as all eigenstates with different parity are separated by a finite energy gap, or, more formally, as long as $[\mathcal{S}, \mathcal{P}] = 0$. For a detailed proof and explanation see Appendix B.

5 Symmetry-breaking transition

While the existence of a fully symmetric steady state follows directly from Eq. (4), there are many trivial cases where this is also the only stationary state, for example, when O is Hermitian. Therefore, we are interested in systems where there is a competing asymmetric phase in the limit $\Gamma \rightarrow \infty$. To ensure that such a phase exists we now restrict ourselves to a Hamiltonian as given in Eq. (7) and a non-Hermitian jump operator of rank $d - 1$ with $\text{Tr}\{O\} = 0$. This implies that there are dark states $|D\rangle$ and $|D^*\rangle$, which satisfy $O|D\rangle = 0$ and $O^\dagger|D^*\rangle = 0$. Under these assumptions we obtain the symmetry-broken phase

$$\rho_0(\Gamma \rightarrow \infty) = |D\rangle\langle D| \otimes |D^*\rangle\langle D^*|, \quad (13)$$

which is fully asymmetric, $\Delta = 1$, and has maximal purity, $P = 1$. Note, however, that for observing symmetry-breaking effects it is not essential that $\rho_0(\Gamma \rightarrow \infty)$ is a pure state and, later in this manuscript, we discuss examples where the symmetry-broken state is mixed.

Given the two distinct limiting phases, the remaining question is, if there is a sharp phase transition between them at a critical intermediate value Γ_c . For the spin system discussed above this question can be rigorously answered in the limit $S \gg 1$ by examining the stability of linear fluctuations on top of the fully polarized state. This can be done using a Holstein-Primakoff approximation [55], where the spin operators are replaced by a pair of bosonic operators, $S_A^- \simeq \sqrt{2S} a^\dagger$, $S_A^+ \simeq \sqrt{2S} a$, $S_B^- \simeq \sqrt{2S} b$ and $S_B^+ \simeq \sqrt{2S} b^\dagger$, where $[a, a^\dagger] = [b, b^\dagger] = 1$. This approximate transformation brings the ME into a quadratic form,

$$\dot{\rho} = -i[H_{\text{lin}}, \rho] + \Gamma \mathcal{D}[a]\rho + \Gamma \mathcal{D}[b]\rho, \quad (14)$$

with Hamiltonian $H_{\text{lin}} = g(ab + a^\dagger b^\dagger)$. From the analytic solution of this linearized model we find that the fluctuations $\langle a^\dagger a \rangle$ and $\langle b^\dagger b \rangle$ diverge at the point $\Gamma_c = g$. Explicitly, in terms of the original spin expectation values we obtain

$$\langle S_{A/B}^z \rangle_0 = \pm S \mp \frac{g^2}{2(\Gamma^2 - g^2)}. \quad (15)$$

Similarly, we can use well-known results for Gaussian states [56] and derive analytic expressions for the purity and the entanglement negativity,

$$P = 1 - \frac{g^2}{\Gamma^2}, \quad \mathcal{N} = \frac{g}{2\Gamma}. \quad (16)$$

These predictions are shown as the curves labeled by $S \rightarrow \infty$ in Fig. 2(a)–(b). Within this Holstein-Primakoff approximation, the substantial amount of entanglement with a maximum of $\mathcal{N}(\Gamma = \Gamma_c) = 1/2$ at the transition point can be directly understood from the form of H_{lin} , which represents a two-mode squeezing interaction.

In general, such an analytic treatment is not possible and, in many situations, \mathcal{PT} -symmetry breaking can occur as a smooth crossover, rather than a sharp phase transition. Nevertheless, it turns out that the appearance of a sharp transition in the limit of large d does not require any specific fine tuning of the dissipation mechanism. This point is illustrated in Fig. 2(c)–(d), where we consider a set of \mathcal{PT} -symmetric quantum systems with randomly generated jump operators O . For each individual line in this plot a jump operator O has been constructed by a Cholesky decomposition of an operator $R = OO^\dagger$, which is drawn randomly from the Gaussian orthogonal ensemble (GOE) (see Appendix C for more details). This operator O is then used to obtain both the dissipative and coherent terms as in Eqs. (5) and (7). For each individual instance, we observe the characteristic transition between the fully mixed and pure states and the asymmetric entanglement peak. These features sharpen as the Hilbert space dimension is increased. Therefore, this study demonstrates that sharp \mathcal{PT} -symmetry breaking transitions are not restricted to simple systems with a direct classical counterpart and are expected to occur in a large range of systems that obey Eq. (4).

6 Generalizations

The symmetry defined in Eq. (4) and the proof of the fully mixed symmetric phase presented in Sec. 4 can be generalized in a straightforward manner to systems with multiple

jump operators. For example, we see the same symmetry-breaking effect in a spin system, with Hamiltonian as above, but considering two competing jump operators for each site,

$$c_A^{1,2} = \sqrt{\frac{1 \pm p}{2}} S_A^\pm, \quad c_B^{1,2} = \sqrt{\frac{1 \mp p}{2}} S_B^\pm. \quad (17)$$

This model, $\mathcal{L}[H; \{\sqrt{\Gamma}c_A^{1,2}\}, \{\sqrt{\Gamma}c_B^{1,2}\}]$, represents two coupled spins, where one is coupled to a positive temperature reservoir while the other is coupled to a negative temperature reservoir. Crucially, this model still obeys the symmetry relation defined in Eq. (4). In Fig. 2(e)–(f) we plot the purity and entanglement negativity for this model with $p = 0.8$. Although in this case the symmetry-broken phase in the limit $\Gamma \rightarrow \infty$ is mixed and the transition is shifted to $\Gamma/g = 1/p$, all the signatures of \mathcal{PT} -symmetry breaking described above are still clearly visible.

Even more relevant is the fact that all the arguments presented above still apply to systems where parity is complemented by another unitary symmetry, $\mathcal{P} \rightarrow \mathcal{P}U$. For example, by choosing $U = e^{i\pi(S_A^x + S_B^x)}$ and a Hamiltonian $H = g(S_A^+ S_B^+ + S_A^- S_B^-)$, we obtain a \mathcal{PT} -symmetric quantum system $\mathcal{L}[H; \sqrt{\Gamma}S_A^-, \sqrt{\Gamma}S_B^-]$. While this model contains only loss processes and the occupation numbers $\langle S_A^+ S_A^- \rangle = \langle S_B^+ S_B^- \rangle$ remain symmetric for all ratios of Γ/g , the Liouvillian respects the symmetry of Eq. (4) with a generalized anti-unitary map

$$\mathbb{P}\mathbb{T}(O) = \mathcal{P}UO^\dagger(U\mathcal{P})^{-1}. \quad (18)$$

As a consequence one observes the same transition from a fully mixed to a low-entropy state, as in the spin model discussed above. The symmetry relation in Eq. (4) is thus a powerful tool to identify \mathcal{PT} -symmetry breaking effects, even in systems where our naive intuition fails.

7 Discussion and Conclusions

In summary, we have introduced the symmetry relation, Eq. (4), for Liouville operators, which extends the notion of \mathcal{PT} symmetry to bipartite open quantum systems. This definition is consistent with previous examples of linear \mathcal{PT} -symmetric quantum systems for which the conventional definition of \mathcal{PT} symmetry is recovered in the limit of large oscillation amplitudes. At the same time the map, $\mathbb{P}\mathbb{T}$, in Eq. (3) is completely general and can be used to define \mathcal{PT} symmetry in highly nonlinear systems or for dissipation processes that have no direct classical counterpart.

In this paper we have mainly focused on the steady state ρ_0 , which is determined for all parameters by the zero eigenvector of \mathcal{L} . In classical systems, \mathcal{PT} -symmetry breaking is usually discussed in terms of a transition from purely oscillatory to exponentially damped or amplified dynamics, which is associated with the appearance of exceptional points in the eigenspectrum of the dynamical matrix. This has motivated similar studies of the appearance of exceptional points in the spectra of Liouville operators [57, 58]. While symmetries as introduced in Ref. [33] lead to remarkable patterns in the complex eigenvalues, which can be broken above a certain dissipation strength, such transitions are not necessarily associated with a qualitative change in the dynamics of physical observables. With the symmetry relations we introduce here, such dynamical changes can be well observed, such as in Fig. 3, where we consider the example of two spin $S = 4$ systems, as shown in Fig. 1(a),(c). The two plots show the full Liouville spectrum below and above the transition point in Fig. 3(a)–(b) and the associated dynamics in panels (c)–(d). For the two cases we don't observe any significant differences in the overall eigenvalue structure. Still




Figure 3: Plot of all complex eigenvalues λ_i of the Liouvillian \mathcal{L} for the \mathcal{PT} -symmetric spin system introduced in Fig. 1(a) with $S = 4$. In (a) the spectrum is shown below ($\Gamma/g = 0.5$) and in (b) above ($\Gamma/g = 1.5$) the transition point. For the same parameters, (c) and (d) show the corresponding time evolution of the observables $\langle S_{A,B}^z \rangle(t)$, starting from the initial state $\rho(t=0) = |-S\rangle\langle -S| \otimes |S\rangle\langle S|$.

the evolution of the observables $\langle S_{A,B}^z \rangle$ undergoes the classically expected change from an oscillatory to an overdamped behavior.

This final example confirms our previous conclusion, namely that \mathcal{PT} -symmetry breaking is an emergent phenomenon in the dynamics and stationary expectation values of macroscopic observables, which, in general, depend little on individual eigenvalues of H_{eff} or \mathcal{L} . Based on the symmetry in Eq. (4), this effect can now be studied more systematically and used to make physically consistent predictions for real experiments. This will be important, for example, for trapped atoms [59], optomechanics [60] or circuit QED systems [61], where gain and loss but also much more complex dissipation processes can be engineered [62, 63]. Our discussion also shows that there are still many interesting conceptual questions to address. This concerns, in particular, the existence and the nature of the \mathcal{PT} -symmetry breaking transition in extended lattices [24, 64] and interacting quantum many-body system, for which exact numerical simulations are no longer available.

Acknowledgements

Funding information. This work was supported through an ESQ fellowship (P.K.) and a DOC Fellowship (J.H.) from the Austrian Academy of Sciences (ÖAW) and by the Austrian Science Fund (FWF) through the DK CoQuS (Grant No. W 1210), Grant No. P32299 (PHONED) and Grant No. P32300 (WAVELAND).

A Mean field equations of motion

In this appendix we discuss a basic example of a bipartite quantum system, which illustrates how the same structure of the Liouville operator results in very different equations of motion for the mean values of bosonic, fermionic and spin operators. To do so we consider the following ME

$$\dot{\rho} = -ig[O_A^\dagger O_B + O_A O_B^\dagger, \rho] + \Gamma_g \mathcal{D}[O_A^\dagger] \rho + \Gamma_l \mathcal{D}[O_B] \rho = \mathcal{L} \rho, \quad (19)$$

which has the same structure as the one assumed in most other examples in this paper. In particular, for $\Gamma_l = \Gamma_g$ the Liouvillian \mathcal{L} satisfies the symmetry relation in Eq. (4) for arbitrary operators O_A and O_B .

In a first step we assume that the two subsystems are represented by two bosonic modes with annihilation operators a and b . We identify $O_A = a$ and $O_B = b$ and define the vector of expectation values $\vec{\psi}_b = (\langle a \rangle, \langle b \rangle)^T$. This vector obeys the equation of motion

$$\partial_t \vec{\psi}_b = -i \mathcal{H}_b \vec{\psi}_b, \quad \text{where} \quad \mathcal{H}_b = \begin{pmatrix} i\Gamma_g & g \\ g & -i\Gamma_l \end{pmatrix}. \quad (20)$$

We see that for $\Gamma_l = \Gamma_g$ the Liouvillian \mathcal{PT} symmetry is directly reflected in the non-Hermitian two-by-two matrix \mathcal{H}_b , which satisfies $(\mathcal{PT}) \mathcal{H}_b (\mathcal{PT})^{-1} = \mathcal{H}_b$. We now perform the same calculations with a and b representing fermionic annihilation operators. The mean values of these operators, $\vec{\psi}_f = (\langle a \rangle, \langle b \rangle)^T$, obey a very similar equation of motion

$$\partial_t \vec{\psi}_f = -i \mathcal{H}_f \vec{\psi}_f, \quad \text{where} \quad \mathcal{H}_f = \begin{pmatrix} -i\Gamma_g & g \\ g & -i\Gamma_l \end{pmatrix}, \quad (21)$$

but in this case $(\mathcal{PT}) \mathcal{H}_f (\mathcal{PT})^{-1} \neq \mathcal{H}_f$.

Non-interacting bosons and fermions are rather special, since in both cases the equations of motion for $\langle a \rangle$ and $\langle b \rangle$ are closed. In general, this is not the case and, for example, when considering two $S = 1/2$ particles with $O_A = \sigma_A^-$ and $O_B = \sigma_B^-$ we already obtain 16 coupled equations for the expectation values of all possible combinations of spin operators. While here we do not write out the resulting matrix \mathcal{H}_s explicitly, it is straight forward to show that also in this case $(\mathcal{PT}) \mathcal{H}_s (\mathcal{PT})^{-1} \neq \mathcal{H}_s$. Basically, this result can already understood by looking at a single spin with loss and gain, i.e.,

$$\dot{\rho} = \Gamma_g \mathcal{D}[\sigma^+] \rho + \Gamma_l \mathcal{D}[\sigma^-] \rho = \mathcal{L} \rho. \quad (22)$$

For this system we define the vector $\vec{\psi}_s = (\langle \sigma^- \rangle, \langle \sigma^+ \rangle, \langle \sigma^+ \sigma^- \rangle, \langle \sigma^- \sigma^+ \rangle)^T$, which obeys

$$\partial_t \vec{\psi}_s = -i \mathcal{H}_s \vec{\psi}_s, \quad \text{where} \quad \mathcal{H}_s = \begin{pmatrix} -i(\Gamma_l + \Gamma_g) & 0 & 0 & 0 \\ 0 & -i(\Gamma_l + \Gamma_g) & 0 & 0 \\ 0 & 0 & -2i\Gamma_l & +2i\Gamma_g \\ 0 & 0 & +2i\Gamma_l & -2i\Gamma_g \end{pmatrix}. \quad (23)$$

This simple example shows that for spin systems, exchanging gain and loss is not the same as replacing $i \rightarrow -i$ in the dynamical matrix for the evolution of mean values.

We conclude that whether or not the symmetry of the Liouville operator defined in Eq. (4) maps onto a conventional \mathcal{PT} symmetry condition for the equations of motion depends on the type of quantum system under consideration. For this simple example we also find that both the Liouville operator as well as the corresponding non-Hermitian effective Hamiltonian are invariant under a particle-hole transformation,

$$a \rightarrow b^\dagger, \quad b \rightarrow \pm a^\dagger, \quad (24)$$

where the plus (minus) sign applies to bosons (fermions). However, this invariance is lost when additional energy terms are added, e.g., $H \rightarrow H' = H + \omega_0(a^\dagger a + b^\dagger b)$, while it does not affect the \mathcal{PT} symmetry.

B Fully symmetric steady state

In this section we detail and extend the proof for the linear stability of the fully mixed symmetric phase in the limit $\Gamma \rightarrow 0$ discussed above. As a starting point we write the density operator as

$$\rho = \sum_{n,m} \rho_{n,m} |E_n\rangle \langle E_m|, \quad (25)$$

where $|E_n\rangle$ are the energy eigenstates of H , i.e. $H|E_n\rangle = E_n|E_n\rangle$. From the \mathcal{PT} -criterion in Eq. (4), we know that $[H, \mathcal{P}] = 0$, and hence we may simultaneously diagonalise the parity operator $\mathcal{P}|E_n\rangle = \zeta_n|E_n\rangle$, where $|\zeta_n|^2 = 1$ without loss of generality.

For $\Gamma = 0$ the fully mixed state, $\rho = \mathbb{1}/d^2$, is a stationary solution of the ME $\dot{\rho} = \mathcal{L}_H \rho = -i[H, \rho]$, but this is also true for any other diagonal state. Therefore, we make the ansatz $\rho_{n,m} = \delta_{n,m}/d^2 + \delta\rho_{n,m}$ and evaluate the evolution of $\delta\rho_{n,m}$ up to first order in Γ [noting that $c_{A,B} \sim O(\sqrt{\Gamma})$],

$$\delta\dot{\rho}_{n,m} = -\frac{i}{\hbar}(E_n - E_m)\rho_{n,m} + \frac{2}{d^2} \langle E_n | [c_A, c_A^\dagger] + [c_B, c_B^\dagger] | E_m \rangle. \quad (26)$$

We first assume that $E_n \neq E_m$. In this case the elements $\rho_{n,m}$ represent coherences between non-degenerate eigenstates and we obtain

$$\delta\rho_{n,m}(t) \simeq -i \frac{2\hbar}{d^2(E_n - E_m)} \langle E_n | [c_A, c_A^\dagger] + [c_B, c_B^\dagger] | E_m \rangle \times \left(1 - e^{-i(E_n - E_m)t/\hbar}\right). \quad (27)$$

Therefore, to lowest order in Γ all these off-diagonal elements of the density matrix remain bounded and $|\delta\rho_{n,m}| \rightarrow 0$ for $\Gamma \rightarrow 0$.

For all other matrix elements with $E_n = E_m$ the coherent evolution vanishes and

$$\delta\dot{\rho}_{n,m} = \frac{2}{d^2} \langle E_n | [c_A, c_A^\dagger] + [c_B, c_B^\dagger] | E_m \rangle. \quad (28)$$

This results in a linear growth in time, unless the matrix element on the right-hand side is zero. We now make use of the relation

$$\mathcal{P}c_B\mathcal{P}^{-1} = c_A^\dagger, \quad (29)$$

which follows from the \mathcal{PT} -symmetry relation for the Liouville operator. Based on this transformation we obtain

$$\begin{aligned} \langle E_n | [c_B, c_B^\dagger] | E_m \rangle &= \langle E_n | \mathcal{P}^{-1} \mathcal{P} [c_B, c_B^\dagger] \mathcal{P}^{-1} \mathcal{P} | E_m \rangle \\ &= \langle E_n | \mathcal{P}^{-1} [c_A^\dagger, c_A] \mathcal{P} | E_m \rangle \\ &= -\zeta_n^* \zeta_m \langle E_n | [c_A, c_A^\dagger] | E_m \rangle, \end{aligned} \quad (30)$$

and the evolution equation from above can be written as

$$\delta\dot{\rho}_{n,m} = \frac{2}{d^2} \langle E_n | [c_A, c_A^\dagger] | E_m \rangle (1 - \zeta_n^* \zeta_m). \quad (31)$$

In the case of a Hamiltonian H with a non-degenerate spectrum, Eq. (31) only applies to the populations $p_n = \rho_{n,n}$, in which case $|\zeta_n|^2 = 1$ and the right hand side vanishes. This is the result given in the main text.

A bit more care must be taken for Hamiltonians with degeneracies imposed by extra symmetries beyond that generated by \mathcal{P} . Even though the populations in a given basis still remain fixed, the build-up of coherences between degenerate levels leads to a deviation from the fully mixed state. If the Hamiltonian has a symmetry, \mathcal{S} , such that $[H, \mathcal{S}] = 0$, then the states generated by applying \mathcal{S} to $|E_n\rangle$ are degenerate. From Eq. (31) we see that this leads to a non-identity steady state when two states $|E_n\rangle$ and $|E_m\rangle$ with the same energy have a different parity, $\zeta_n \neq \zeta_m$. However, if $[\mathcal{P}, \mathcal{S}] = 0$ then it is straightforward to see that $\zeta_n = \zeta_m$. Therefore, for the existence of a fully mixed symmetric phase it is in general not enough that $[H, \mathcal{P}] = 0$. In addition, we require that all other non-trivial symmetries of the Hamiltonian also commute with the parity operator, at least within each degenerate subspace.

A simple example where such non-trivial symmetries play a role is the spin model described by the Hamiltonian

$$H = g(S_A^+ S_B^+ + S_A^- S_B^-) \quad (32)$$

and the \mathcal{PT} -symmetric ME

$$\dot{\rho} = -i[H, \rho] + \Gamma \mathcal{D}[S_A^-] \rho + \Gamma \mathcal{D}[S_B^+] \rho. \quad (33)$$

This model has a symmetry generated by $\mathcal{S} = S_A^z - S_B^z$ which does not commute with \mathcal{P} and indeed one can show that the steady state for this model has spin- A pointing down and spin- B pointing up independent of the value of Γ/g .

C Random jump operators

In Fig. 2(c)–(d) we calculate the steady state of random \mathcal{PT} -symmetric finite dimensional quantum systems. Here we describe how these random models are constructed. For simplicity we keep the relationship between jump operators and the Hamiltonian as described in the main text. We also wish to ensure that the jump operators have a single dark state, such that in the limit $\Gamma \rightarrow \infty$ the purity $P \rightarrow 1$.

The procedure we use is then as follows: We first create a random matrix R from the Gaussian orthogonal ensemble (GOE), i.e., a symmetric matrix with real entries which follow a Gaussian distribution [65]. This matrix is then shifted by its lowest eigenvalue such that $R' = R - \lambda_0 \mathbb{I}$ is positive semidefinite with a guaranteed zero eigenvalue. To obtain the jump operator O we then perform a Cholesky decomposition on the resulting matrix,

$$R' = OO^\dagger, \quad (34)$$

such that O is a lower triangular matrix. Since the Cholesky decomposition for positive semi-definite matrices is not unique, we implement this step by first diagonalizing the random matrix R' ,

$$R' = UDU^\dagger, \quad (35)$$

with U a unitary matrix and $D = \text{diag}(0, \lambda_1, \dots, \lambda_{d-1})$, a diagonal matrix where λ_n are non-zero eigenvalues. The diagonal matrix D can be decomposed as $D = LL^\dagger$, where only the first super diagonal of L^\dagger is non-zero with $(\sqrt{\lambda_1}, \sqrt{\lambda_2}, \dots, \sqrt{\lambda_{d-1}})$. As a result the jump operator is

$$O = ULU^\dagger. \quad (36)$$

This procedure of constructing a random jump operator ensures that most of the resulting decay rates are $O(1)$, due to the fact that the spacing between the eigenvalues of R will

follow a Wigner surmise distribution $P(\Delta E) \sim \Delta E \exp(-A\Delta E^2)$ [65], meaning that there are very few almost degenerate states. By enforcing L^\dagger to only have non-vanishing elements in the first upper diagonal ensures that it is possible to observe the \mathcal{PT} -symmetry breaking transition. This is not guaranteed in general. For example, by decomposing the diagonal matrix D in Eq. (35) in terms of two diagonal matrices $D = \sqrt{D}\sqrt{D}$, the resulting jump operator would be Hermitian and there would be no phase transition since the trivial identity state is always a steady state of such a model.

References

- [1] C. M. Bender and S. Boettcher, Real spectra in non-hermitian hamiltonians having PT symmetry, *Phys. Rev. Lett.* **80**, 5243 (1998).
- [2] L. Feng, R. El-Ganainy, and L. Ge, Non-Hermitian photonics based on parity–time symmetry, *Nature Photon.* **11**, 752 (2017).
- [3] S. Longhi, Parity-time symmetry meets photonics: A new twist in non-Hermitian optics, *EPL* **120**, 64001 (2017).
- [4] R. El-Ganainy, K. G. Makris, M. Khajavikhan, Z. H. Musslimani, S. Rotter, and D. N. Christodoulides, Non-Hermitian physics and PT symmetry, *Nature Phys.* **14**, 11 (2018).
- [5] S. K. Özdemir, S. Rotter, F. Nori, and L. Yang, Parity-Time Symmetry and Exceptional Points in Photonics, *Nature Mater.* **18**, 783 (2019).
- [6] C. E. Rüter, K. G. Makris, R. El-Ganainy, D. N. Christodoulides, M. Segev, and D. Kip, Observation of Parity-Time Symmetry in Optics, *Nature Phys.* **6**, 192 (2010).
- [7] B. Peng, S. K. Özdemir, F. Lei, F. Monifi, M. Gianfreda, G. L. Long, S. Fan, F. Nori, C. M. Bender, and L. Yang, Parity-Time-Symmetric Whispering-Gallery Microcavities, *Nature Phys.* **10**, 394 (2014).
- [8] L. Chang, X. Jiang, S. Hua, C. Yang, J. Wen, L. Jiang, G. Li, G. Wang, and M. Xiao, Parity–time symmetry and variable optical isolation in active–passive-coupled microresonators, *Nature Photon.* **8**, 524 (2014).
- [9] J. Schindler, A. Li, M. C. Zheng, F. M. Ellis, and T. Kottos, Experimental study of active LRC circuits with PT symmetries, *Phys. Rev. A* **84**, 040101(R) (2011).
- [10] C. M. Bender, B. K. Berntson, D. Parker, and E. Samuel, Observation of PT phase transition in a simple mechanical system, *Am. J. Phys.* **81**, 173 (2013).
- [11] A. U. Hassan, H. Hodaei, M. A. Miri, M. Khajavikhan, and D. N. Christodoulides, Nonlinear reversal of the PT-symmetric phase transition in a system of coupled semiconductor microring resonators, *Phys. Rev. A* **92**, 063807 (2015).
- [12] L. Ge and R. El-Ganainy, Nonlinear modal interactions in parity-time (PT) symmetric lasers, *Sci. Rep.* **6**, 24889 (2016).
- [13] J. Wiersig, Enhancing the sensitivity of frequency and energy splitting detection by using exceptional points: application to microcavity sensors for single-particle detection, *Phys. Rev. Lett.* **112**, 203901 (2014).

- [14] M. Am-Shallem, R. Kosloff, and N. Moiseyev, Parameter estimation in atomic spectroscopy using exceptional points, *Phys. Rev. A* **93**, 032116 (2016).
- [15] H.-K. Lau and A. A. Clerk, Fundamental limits and non-reciprocal approaches in non-Hermitian quantum sensing, *Nat. Commun.* **9**, 4320 (2018).
- [16] C. Chen, L. Jin, and R.-B. Liu, Sensitivity of parameter estimation near the exceptional point of a non-Hermitian system, *New J. Phys.* **21**, 083002 (2019).
- [17] M. Zhang, W. Sweeney, C. W. Hsu, L. Yang, A. D. Stone, and L. Jiang, Quantum Noise Theory of Exceptional Point Amplifying Sensors, *Phys. Rev. Lett.* **123**, 180501 (2019).
- [18] M. P. Hokmabadi, A. Schumer, D. N. Christodoulides, and M. Khajavikhan, Non-Hermitian Ring Laser Gyroscopes with Enhanced Sagnac Sensitivity, *Nature* **576**, 70 (2019).
- [19] E. J. Bergholtz, J. Carl Budich, and F. K. Kunst, Exceptional Topology of Non-Hermitian Systems, arXiv:1912.10048 (2019).
- [20] K. Kawabata, K. Shiozaki, M. Ueda, and M. Sato, Symmetry and Topology in Non-Hermitian Physics, *Phys. Rev. X* **9**, 041015 (2019).
- [21] H. Zhou and J. Y. Lee, Periodic table for topological bands with non-Hermitian symmetries, *Phys. Rev. B* **99**, 235112 (2019).
- [22] Y. Ashida, Z. Gong, and M. Ueda, Non-Hermitian Physics, arXiv:2006.01837 (2020).
- [23] S. Assaworrorarit, X. Yu, and S. Fan, Robust wireless power transfer using a nonlinear parity-time-symmetric circuit, *Nature* **546**, 387 (2017).
- [24] J. Huber and P. Rabl, Active energy transport and the role of symmetry breaking in microscopic power grids, *Phys. Rev. A* **100**, 012129 (2019).
- [25] M. Partanen, *et al.*, Exceptional points in tunable superconducting resonators, *Phys. Rev. B* **100**, 134505 (2019).
- [26] This excludes possible non-Hermitian quantum theories beyond the standard formulation of quantum mechanics [27].
- [27] C. M. Bender, Making sense of non-Hermitian Hamiltonians, *Rep. Prog. Phys.* **70**, 947 (2007).
- [28] H. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, UK, 2002).
- [29] H. Schomerus, Quantum Noise and Self-Sustained Radiation of PT-Symmetric Systems, *Phys. Rev. Lett.* **104**, 233601 (2010).
- [30] G. S. Agarwal and K. Qu, Spontaneous generation of photons in transmission of quantum fields in PT-symmetric optical systems, *Phys. Rev. A* **85**, 031802 (2012).
- [31] K. V. Kepesidis, T. J. Milburn, J. Huber, K. G. Makris, S. Rotter, and P. Rabl, PT-symmetry breaking in the steady state of microscopic gain-loss systems, *New J. Phys.* **18**, 095003 (2016).

- [32] S. Scheel and A. Szameit, PT-symmetric photonic quantum systems with gain and loss do not exist, *EPL* **122**, 34001 (2018).
- [33] T. Prosen, PT-Symmetric Quantum Liouvillean Dynamics, *Phys. Rev. Lett.* **109**, 090404 (2012).
- [34] T. Prosen, *Generic examples of PT-symmetric qubit (spin-1/2) Liouvillian dynamics*, *Phys. Rev. A* **86**, 044103 (2012).
- [35] B. He, L. Yang, Z. Zhang, and M. Xiao, Cyclic permutation-time symmetric structure with coupled gain-loss microcavities, *Phys. Rev. A* **91**, 033830 (2015).
- [36] J. Naikoo, K. Thapliyal, S. Banerjee, and A. Pathak, Quantum Zeno effect and non-classicality in a PT-symmetric system of coupled cavities, *Phys. Rev. A* **99**, 023820 (2019).
- [37] F. Roccati, S. Lorenzo, G. Massimo Palma, G. T. Landi, M. Brunelli, and F. Ciccarello, Quantum correlations in PT-symmetric systems, arXiv:2002.11127 (2020).
- [38] I. I. Arkhipov, A. Miranowicz, F. Minganti, and F. Nori, Quantum and semiclassical exceptional points of a linear system of coupled cavities with losses and gain within the Scully-Lamb laser theory, *Phys. Rev. A* **101**, 013812 (2020).
- [39] B. Jaramillo Avila, C. Ventura-Velazquez, R. de J. Leon-Montiel, Y. N. Joglekar, and B. M. Rodriguez-Lara, PT-symmetry from Lindblad dynamics in a linearized optomechanical system, *Scientific Rep.* **10**, 1761 (2020).
- [40] A. Purkayastha, M. Kulkarni, and Y. N. Joglekar, Emergent PT symmetry in a double-quantum-dot circuit QED set-up, arXiv:2004.07541 (2020).
- [41] D. Dast, D. Haag, H. Cartarius, and G. Wunner, Quantum master equation with balanced gain and loss, *Phys. Rev. A* **90**, 052120 (2014).
- [42] Y.-X. Wang and A. A. Clerk, Non-Hermitian dynamics without dissipation in quantum systems, *Phys. Rev. A* **99**, 063834 (2019).
- [43] M. Naghiloo, M. Abbasi, Y. N. Joglekar, and K. W. Murch, Quantum state tomography across the exceptional point in a single dissipative qubit, *Nature Phys.* **15**, 1232 (2019).
- [44] Y. Wu, W. Liu, J. Geng, X. Song, X. Ye, C.-K. Duan, X. Rong, and J. Du, Observation of parity-time symmetry breaking in a single-spin system, *Science* **364**, 878 (2019).
- [45] J. Dalibard, Y. Castin, and K. Mølmer, Wave-function approach to dissipative processes in quantum optics, *Phys. Rev. Lett.* **68**, 580 (1992).
- [46] T. E. Lee and C.-K. Chan, Heralded Magnetism in Non-Hermitian Atomic Systems, *Phys. Rev. X* **4**, 041001 (2014).
- [47] Y. Ashida, S. Furukawa, and M. Ueda, Parity-time-symmetric quantum critical phenomena, *Nature Commun.* **8**, 15791 (2017).
- [48] M. van Caspel and V. Gritsev, Symmetry-protected coherent relaxation of open quantum systems, *Phys. Rev. A* **97**, 052106 (2018).

- [49] D. Huybrechts, F. Minganti, F. Nori, M. Wouters, and N. Shammah, Validity of mean-field theory in a dissipative critical system: Liouvillian gap, PT-symmetric antigap, and permutational symmetry in the XYZ model, *Phys. Rev. B* **101**, 214302 (2020).
- [50] S. Lieu, M. McGinley, and N. R. Cooper, Tenfold Way for Quadratic Lindbladians, *Phys. Rev. Lett.* **124**, 040401 (2020).
- [51] A. Altland, M. Fleischhauer, and S. Diehl, Symmetry classes of open fermionic quantum matter, arXiv:2007.10448 (2020).
- [52] C.-K. Chiu, J. C. Y. Teo, A. P. Schnyder, and S. Ryu, Classification of topological quantum matter with symmetries, *Rev. Mod. Phys.* **88**, 035005 (2016).
- [53] K. Zyczkowski, P. Horodecki, A. Sanpera, and M. Lewenstein, Volume of the set of separable states, *Phys. Rev. A* **58**, 883 (1998).
- [54] G. Vidal and R. F. Werner, Computable measure of entanglement, *Phys. Rev. A* **65**, 032314 (2002).
- [55] T. Holstein and H. Primakoff, Field dependence of the intrinsic domain magnetization of a ferromagnet, *Phys. Rev.* **58**, 1098 (1940).
- [56] A. Serafini, F. Illuminati, M. G. Paris, and S. De Siena, Entanglement and purity of two-mode Gaussian states in noisy channels, *Phys. Rev. A*, **69**, 022318 (2014).
- [57] A. Pick, S. Silberstein, N. Moiseyev, and N. Bar-Gill, Robust mode conversion in NV centers using exceptional points, *Phys. Rev. Research* **1**, 013015 (2019).
- [58] F. Minganti, A. Miranowicz, R. W. Chhajlany, and F. Nori, Quantum exceptional points of non-Hermitian Hamiltonians and Liouvillians: The effects of quantum jumps, *Phys. Rev. A* **100**, 062131 (2019).
- [59] D. Leibfried, R. Blatt, C. Monroe, and D. Wineland, Quantum dynamics of single trapped ions, *Rev. Mod. Phys.* **75**, 281 (2003).
- [60] M. Aspelmeyer, T. J. Kippenberg, and F. Marquardt, Cavity optomechanics, *Rev. Mod. Phys.* **86**, 1391 (2014).
- [61] X. Gu, A. Frisk Kockum, A. Miranowicz, Y.-X. Liu, and F. Nori, Microwave photonics with superconducting quantum circuits, *Phys. Rep.* **718**, 1 (2017).
- [62] J. T. Barreiro, M. Müller, P. Schindler, D. Nigg, T. Monz, M. Chwalla, M. Hennrich, C. F. Roos, P. Zoller, and R. Blatt, An open-system quantum simulator with trapped ions, *Nature* **470**, 486 (2011).
- [63] M. Müller, S. Diehl, G. Pupillo, and P. Zoller, Engineered Open Systems and Quantum Simulations with Atoms and Ions, *Adv. At. Mol. Opt. Phys.* **61**, 1 (2012).
- [64] J. Huber, P. Kirton, and P. Rabl, Non-equilibrium magnetic phases in spin lattices with gain and loss, *Phys. Rev. A* **102**, 012219 (2020).
- [65] M. L. Mehta, *Random matrices* (Elsevier, 2004).