

Area law and OPE blocks in conformal field theory

Jiang Long

School of Physics, Huazhong University of Science and Technology,
Wuhan, Hubei 430074, China

★ longjiang@hust.edu.cn

*4th International Conference on Holography,
String Theory and Discrete Approach
Hanoi, Vietnam, 2020*
doi:[10.21468/SciPostPhysProc.4](https://doi.org/10.21468/SciPostPhysProc.4)

Abstract

This is an introduction to the relationship between area law and OPE blocks in conformal field theory.

Copyright J.Long

This work is licensed under the Creative Commons
[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Published by the SciPost Foundation.

Received ??-??-20??

Accepted ??-??-20??

Published ??-??-20??

doi:[10.21468/SciPostPhysProc.4](https://doi.org/10.21468/SciPostPhysProc.4)??

1

2 Contents

3	1 Introduction	2
4	2 Setup	2
5	2.1 Area law	3
6	2.2 OPE block	4
7	2.3 Modular Hamiltonian and area law	6
8	2.4 Deformed reduced density matrix and connected correlation function	6
9	3 Area law	8
10	3.1 Continuation	9
11	3.2 Kinematic information	10
12	3.3 UV/IR relation	11
13	3.4 Discussion	14
14	4 Generalizations	14
15	5 Summary and outlook	17
16	References	17

17
18

19 1 Introduction

20 This report consists a summary of our recent progress on the relationship between area law and
21 OPE blocks. Area law has been a continuous topic in physics. The prototype of area law dates
22 back to black hole physics in general relativity. The unusual property that the thermal entropy
23 of a black hole is proportional to the event horizon of the black hole [1, 2] has stimulated varies
24 modern idea of theoretical physics, including the famous holographic principle.

25 OPE block [3, 4], on the other hand, is a relatively new topic in conformal field theory, though
26 it has been noticed at the early stages of conformal field theory [5, 6]. The operator product
27 expansion of two primary operators is equivalent to a summation of OPE blocks with corre-
28 sponding three point function coefficients. It is a smeared operator which is generated from
29 (quasi-)primary operator.

30 Modular Hamiltonian, the logarithm of the reduced density matrix [7], plays a central role
31 in the context of geometric entanglement entropy [8–11]. Entanglement entropy is a von
32 Neumann entropy generated from reduced density matrix of a subregion of spacetime. An
33 intriguing fact of entanglement entropy is that it obeys area law in the leading order, though
34 one should introduce a cutoff to secure the divergent behaviour. Its connection to gravity
35 has been established by the work of Ryu and Takayanagi [12], in which they proposed that
36 the entanglement entropy of a CFT is equal to the area of a minimal surface in the bulk AdS
37 spacetime.

38 Modular Hamiltonian is a special OPE block generated by stress energy-momentum tensor
39 for a ball region. This leads to the conjecture that OPE block may be related to area law as
40 modular Hamiltonian. Indeed, in a series of papers [13–16], we have shown that the quantity
41 which satisfies area law is type- (m) connected correlation function (CCF). More explicitly, the
42 leading term of the type- (m) CCF is proportional to the area of the boundary of the ball. In the
43 subleading terms, we find a logarithmic divergence with degree q . The degree q is a natural
44 number which is no larger than 2 in general dimensions. The coefficient p_q for the logarithmic
45 term with degree q is cutoff independent. We establish a relationship between p_q and type-
46 $(m - 1, 1)$ CCF of OPE blocks for two balls which are far away to each other. The coefficient
47 p_q obeys a cyclic identity which is independent of the order of the operators.

48 This paper is organised as follows. In section 2, we will introduce basic concepts and conven-
49 tions used in this paper. Section 3 is devoted to the study of the new area law which is related
50 to OPE blocks. Varies generalizations have been given in section 4. We conclude in section 5
51 with a number of general open problems that deserve, in our opinion, more work.

52 2 Setup

53 In this section, we introduce some basic concepts and conventions used in this paper.

54 **2.1 Area law**

55 In continues quantum field theory(QFT), physical degrees exist at each point $(t, x^i), i = 1, \dots, d-1$
 56 of spacetime M . At each time slice $t = t_0$, data on the Cauchy surface Σ determines the evalu-
 57 ation of fields. One can divide the surface Σ into a spacelike subregion A and its complement
 58 $\bar{A}, \Sigma = A \cup \bar{A}$. The boundary ∂A is a codimension 2 surface whose area is \mathcal{A} . The causal devel-
 59 opment of A is denoted by $\mathcal{D}(A)$. Physical data on A can only determine the evaluation of fields
 60 in $\mathcal{D}(A)$. The causal development $\mathcal{D}(A)$ is an independent subsystem of original spacetime M .
 61 Operators in this subsystem are collected to form an algebra $\mathfrak{a}(A)$. Assume QFT in spacetime
 62 M is described by a density matrix ρ , then by integrating out the degree of freedom in the
 63 complement of \bar{A} , one achieves a reduced density matrix ρ_A

$$\rho_A = \text{tr}_{\bar{A}} \rho. \tag{2.1}$$

64 Reduced density matrix ρ_A is a special operator in $\mathfrak{a}(A)$ since it describes the subsystem $\mathcal{D}(A)$
 65 effectively. A general quantity $Q(A)$ in $\mathfrak{a}(A)$ is said to obey area law if its leading term is
 66 proportional to the area of boundary ∂A ,

$$Q(A) \propto \mathcal{A} + \dots. \tag{2.2}$$

67 One typical example is the black hole entropy in Einstein gravity. Black hole entropy is propor-
 68 tional to the area of the event horizon,

$$S_{bh} = \frac{\mathcal{A}}{4G} \tag{2.3}$$

69 where G is Newton constant. At the loop level, black hole entropy requires logarithmic correc-
 70 tions [17–22]. Usually, the logarithmic correction is in the form $C \log \mathcal{A}$ where the constant C
 71 may encode useful information of the black hole.

72 Sometimes the area law is divergent, one typical example is the geometric entanglement en-
 73 tropy

$$S_A = -\text{tr}_A \rho_A \log \rho_A. \tag{2.4}$$

74 In this case, one should insert a cutoff $\epsilon > 0$,

$$S_A = \gamma \frac{\mathcal{A}}{\epsilon^{d-2}} + \dots. \tag{2.5}$$

75 In the subleading terms, there may be a logarithmic term whose coefficient is independent of
 76 the cutoff,

$$S_A = \gamma \frac{R^{d-2}}{\epsilon^{d-2}} + \dots + p \log \frac{R}{\epsilon} + \dots \tag{2.6}$$

77 where the parameter R is the characteristic length of region A .

78 In this report, we will present a quantity $Q(A)$ which has a slightly different logarithmic be-
 79 haviour

$$Q(A) = \gamma \frac{R^{d-2}}{\epsilon^{d-2}} + \dots + p_q \log^q \frac{R}{\epsilon} + \dots. \tag{2.7}$$

80 The maximum power q of the logarithmic terms is a natural number. We will call it the degree
 81 of the quantity $Q(A)$. The coefficient p_q is cutoff independent, it encodes useful information
 82 of the theory. In the special case that the subregion A is a ball, R could be chosen as its radius.
 83 Subregion A and its causal development $\mathcal{D}(A)$ are in one-to-one correspondence, we will not
 84 distinguish them in the following.

85 In two dimensions, there is no polynomial terms of $\frac{R}{\epsilon}$, the modified “area law” is

$$Q(A) = p_q \log^q \frac{R}{\epsilon} + \dots. \tag{2.8}$$

86 **2.2 OPE block**

87 In d dimensional CFT, operators are classified into (quasi-)primary operators \mathcal{O} and their de-
 88 scendants $\partial_\mu \partial_\nu \dots \mathcal{O}$. A general primary operator is characterized by two quantum numbers,
 89 conformal weight Δ and $so(d-1)$ spin J_{ij} with magnitude J . Under a global conformal trans-
 90 formation $x \rightarrow x'$, a primary spin 0 operator transforms as

$$\mathcal{O}(x) \rightarrow \left| \frac{\partial x'}{\partial x} \right|^{-\Delta/d} \mathcal{O}(x). \tag{2.9}$$

91 where $|\partial x'/\partial x|$ is the Jacobian of the conformal transformation of the coordinates, Δ is the
 92 conformal weight of the primary operator. Operator product expansion(OPE) of two separated
 93 primary scalar operators $\mathcal{O}_i(x_1)\mathcal{O}_j(x_2)$ is to expand their product in a local orthogonal and
 94 complete basis around a suitable point

$$\mathcal{O}_i(x_1)\mathcal{O}_j(x_2) = \sum_k C_{ijk} |x_{12}|^{\Delta_k - \Delta_i - \Delta_j} (\mathcal{O}_k(x_2) + \dots), \tag{2.10}$$

95 where \dots are descendants of the primary operator \mathcal{O}_k . Its form is fixed by global conformal
 96 symmetry, therefore it just contains kinematic information of the CFT. Here we expand the
 97 product around the point x_2 . The distance of any two points x_i, x_j is written as $|x_{ij}|$. The
 98 constant C_{ijk} is called OPE coefficient which is related to the three point function of primary
 99 operators

$$\langle \mathcal{O}_i(x_1)\mathcal{O}_j(x_2)\mathcal{O}_k(x_3) \rangle = \frac{C_{ijk}}{|x_{12}|^{\Delta_{12,3}} |x_{23}|^{\Delta_{23,1}} |x_{13}|^{\Delta_{13,2}}}, \quad \Delta_{ij,k} = \Delta_i + \Delta_j - \Delta_k. \tag{2.11}$$

100 They are the only dynamical parameters in a CFT. The constants $\Delta_i, \Delta_j, \Delta_k$ are conformal
 101 weights of the corresponding primary operators. By collecting all kinematic terms in the sum-
 102 mation, we can rewrite OPE (2.10) as

$$\mathcal{O}_i(x_1)\mathcal{O}_j(x_2) = |x_{12}|^{-\Delta_i - \Delta_j} \sum_k C_{ijk} Q_k^{ij}(x_1, x_2). \tag{2.12}$$

103 The objects $Q_k^{ij}(x_1, x_2)$ are called OPE blocks [3,5,6]. They are non-local operators in the CFT
 104 and depend on the position x_1 and x_2 of external operators. The upper index i and j show
 105 that it also depends on the quantum number of the external operators \mathcal{O}_i and \mathcal{O}_j . It is easy to
 106 see that OPE block has dimension zero. Under a global conformal transformation $x \rightarrow x'$, an
 107 OPE block $Q_k^{ij}(x_1, x_2)$ will transform as

$$Q_k^{ij}(x_1, x_2) \rightarrow f(x'_1, x'_2) Q_k^{ij}(x'_1, x'_2). \tag{2.13}$$

108 The explicit form of $f(x'_1, x'_2)$ is not important in this work. When the two external operators
 109 are the same, we have $f(x'_1, x'_2) = 1$ and OPE block will be invariant under global conformal
 110 transformation. One can also show that the OPE block is independent of the external operator
 111 in this special case. Due to this reason, we relabel such kind of OPE block as

$$Q_A[\mathcal{O}_k] = Q_k^{ii}(x_1, x_2). \tag{2.14}$$

112 The subscript A denotes the region determined by the two points x_1 and x_2 where the two
 113 external operators insert into. The operator in square bracket reflects the fact that OPE block is
 114 generated by a primary operator \mathcal{O}_k . We omit the information of i since OPE block is insensitive
 115 to the external operators in this case. We will classify the primary operator \mathcal{O}_k into conserved

116 currents \mathcal{J} and non-conserved operators \mathcal{O} . A general symmetric traceless primary operator
 117 obeys the following unitary bound [23]

$$\begin{cases} \Delta \geq J + d - 2, & J \geq 1, \\ \Delta \geq \frac{d-2}{2}, & J = 0. \end{cases}$$

118 A conserved current \mathcal{J} with spin $J (J \geq 1)$ will satisfy $\Delta = J + d - 2$. All other primary operators
 119 are non-conserved operators. Correspondingly, the OPE block (2.14) generated by conserved
 120 currents \mathcal{J} will be called type-J OPE block. On the other hand, the OPE block (2.14) generated
 121 by non-conserved operators \mathcal{O} will be called type-O OPE block.

122 When two operators are time-like separated, region A is a causal diamond. The two operators
 123 are at the sharp corner of the diamond A . We can use conformal transformation to fix

$$x_1 = (1, \vec{x}_0), \quad x_2 = (-1, \vec{x}_0), \quad (2.15)$$

124 then the causal diamond A intersects $t = 0$ slice with a unit ball which we will also denote it
 125 as A

$$A = \{(0, \vec{x}) | (\vec{x} - \vec{x}_0)^2 \leq 1\}. \quad (2.16)$$

126 The center of the ball is \vec{x}_0 . The boundary of the ball A is a unit sphere ∂A . In the context of
 127 geometric entanglement entropy, the surface ∂A is an entanglement surface which separates
 128 the ball A and its complement. Leading term of entanglement entropy is proportional to the
 129 area of surface ∂A in general higher dimensions ($d > 2$). In two dimensions, the entanglement
 130 entropy is logarithmically divergent with the logarithmic degree $q = 1$. There is a conformal
 131 Killing vector K which preserves the diamond A ,

$$K^\mu = \frac{1}{2}(1 - (\vec{x} - \vec{x}_A)^2 - t^2, -2t\vec{x}). \quad (2.17)$$

132 Conformal Killing vector K is null on the boundary of the diamond A . It generates modular
 133 flow of the diamond A . Type-O OPE block corresponds to point pair (2.15) or unit ball A (2.16)
 134 is [4]

$$Q_A[\mathcal{O}_{\mu_1 \dots \mu_J}] = c_{\mathcal{O}_{\mu_1 \dots \mu_J}} \int_{\mathcal{D}(A)} d^d x K^{\mu_1} \dots K^{\mu_J} |K|^{\Delta-d-J} \mathcal{O}_{\mu_1 \dots \mu_J}, \quad (2.18)$$

135 where the primary operator $\mathcal{O}_{\mu_1 \dots \mu_J}$ is non-conserved

$$\partial^{\mu_1} \mathcal{O}_{\mu_1 \dots \mu_J} \neq 0. \quad (2.19)$$

136 It has dimension Δ and spin J . When the operator is a conserved current

$$\partial^{\mu_1} \mathcal{J}_{\mu_1 \dots \mu_J} = 0, \quad (2.20)$$

137 the corresponding type-J OPE block is

$$Q_A[\mathcal{J}_{\mu_1 \dots \mu_J}] = c_{\mathcal{J}_{\mu_1 \dots \mu_J}} \int_A d^{d-1} \vec{x} (K^0)^{J-1} \mathcal{J}_{0 \dots 0}. \quad (2.21)$$

138 It can be obtained from (2.18) by using conservation law (2.20) and reducing it to a lower
 139 $d-1$ dimensional integral. The coefficient $c_{\mathcal{J}_{\mu_1 \dots \mu_J}}$ is also redefined at the same time. In (2.18)
 140 and (2.21), the coefficients $c_{\mathcal{O}_{\mu_1 \dots \mu_J}}$ and $c_{\mathcal{J}_{\mu_1 \dots \mu_J}}$ are free parameters, we set them to be 1.

141 **2.3 Modular Hamiltonian and area law**

142 A very special type-J OPE block is modular Hamiltonian [7, 24] of the ball A ,

$$H_A = 2\pi \int_A d^{d-1} \vec{x} K^0 T_{00} = 2\pi \int_A d^{d-1} \vec{x} \frac{1 - (\vec{x} - \vec{x}_0)^2}{2} T_{00}(0, \vec{x}). \quad (2.22)$$

143 Modular Hamiltonian is the logarithm of the reduced density matrix ρ_A

$$H_A = -\log \rho_A. \quad (2.23)$$

144 It plays a central role in the context of entanglement entropy,

$$S_A = -\text{tr}_A \rho_A \log \rho_A = \text{tr}_A e^{-H_A} H_A. \quad (2.24)$$

145 More generally, Rényi entanglement entropy

$$S_A^{(n)} = \frac{1}{1-n} \log \text{tr}_A \rho_A^n \quad (2.25)$$

146 has been shown to satisfy an area law generally

$$S_A^{(n)} = \gamma \frac{\mathcal{A}}{\epsilon^{d-2}} + \dots, \quad (2.26)$$

147 where \mathcal{A} is the area of the entanglement surface ∂A and ϵ is a UV cutoff. The constant γ is
 148 cutoff dependent. The subleading terms \dots contain a logarithmic term with degree $q = 1$ in
 149 even dimensions

$$S_A^{(n)} = \gamma \frac{\mathcal{A}}{\epsilon^{d-2}} + \dots + p_1(n) \log \frac{R}{\epsilon} + \dots, \quad (2.27)$$

150 where we have inserted back the radius $R = 1$. The area \mathcal{A} is related to the radius R through
 151 the power law

$$\mathcal{A} \sim R^{d-2}. \quad (2.28)$$

152 The coefficient $p_1(n)$ encodes useful information of the CFT. The relation between modular
 153 Hamiltonian and area law motivates the conjecture that OPE block maybe related to area
 154 law in a suitable way. We will give the framework to discuss this problem in the following
 155 subsection.

156 **2.4 Deformed reduced density matrix and connected correlation function**

157 Given a primary operator \mathcal{O} in a ball A , one can always define a corresponding OPE block
 158 $Q_A[\mathcal{O}]$. We construct an exponential operator formally [14]

$$\rho_A = e^{-\mu Q_A} \quad (2.29)$$

159 which is still in subregion A . The constant μ is free. Operators of the form (2.29) is called
 160 deformed reduced density matrix. Recall that modular Hamiltonian is a special OPE block, if
 161 one replaces OPE block by modular Hamiltonian in (2.29) and set $\mu = 1$, the deformed reduced
 162 density matrix becomes reduced density matrix exactly. We can relax the definition, namely,
 163 allow Q_A in (2.29) is a linear superposition of several OPE blocks. Note we use the same symbol
 164 ρ_A to label deformed reduced density matrix. As a naive generalization of Rényi entanglement
 165 entropy, we construct logarithm of the vacuum expectation value of the deformed reduced
 166 density matrix,

$$T_A(\mu) = \log \langle \rho_A \rangle = \log \langle e^{-\mu Q_A} \rangle. \quad (2.30)$$

167 When Q_A is modular Hamiltonian, the above quantity is related to Rényi entropy for vacuum
 168 state.

169 However, a direct computation of $T_A(\mu)$ is hard in general. A much more severe problem is
 170 that OPE block has no lower bound in general, therefore the definition is not valid for general
 171 OPE blocks. To solve this problem, we observe that $T_A(\mu)$ could be expanded for small μ ,

$$T_A(\mu) = \sum_{m=1}^{\infty} \frac{(-\mu)^m}{m!} \langle Q_A^m \rangle_c. \quad (2.31)$$

172 The Taylor expansion coefficient

$$\langle Q_A^m \rangle_c = (-1)^m \frac{\partial^m}{\partial \mu^m} T_A(\mu)|_{\mu \rightarrow 0} \quad (2.32)$$

173 is called Type-(m) connected correlation function (CCF) of OPE block Q_A . For each definite m ,
 174 one can always calculate the corresponding CCF without knowing $T_A(\mu)$. The first few CCFs
 175 are

$$\begin{aligned} \langle Q_A^2 \rangle_c &= \langle Q_A^2 \rangle - \langle Q_A \rangle^2, \\ \langle Q_A^3 \rangle_c &= \langle Q_A^3 \rangle - 3\langle Q_A^2 \rangle \langle Q_A \rangle + 2\langle Q_A \rangle^3. \end{aligned} \quad (2.33)$$

176 Using CCF, there is no issue of lower bound of OPE block. As an application of the concept CCF,
 177 we set the OPE block to modular Hamiltonian, then it is easy to show that CCF of modular
 178 Hamiltonian H_A satisfies area law with logarithmic degree $q = 1$ in even dimensions,

$$\langle H_A^m \rangle_c = \tilde{\gamma} \frac{A}{\epsilon^{d-2}} + \dots + \tilde{p}_1^{(m)} \log \frac{R}{\epsilon} + \dots, \quad m \geq 1. \quad (2.34)$$

179 The coefficient $\tilde{p}_1^{(m)}$ is determined from $p_1(n)$ by

$$\tilde{p}_1^{(m)} = (-1)^m \partial_n^m (1-n)p_1(n)|_{n \rightarrow 1}. \quad (2.35)$$

180 There could be multiple spacelike-separated balls A_1, A_2, \dots , each region has associate OPE
 181 block Q_{A_i} . We insert m_i OPE blocks into region A_i , then we can define corresponding type-Y
 182 CCF

$$\langle Q_{A_1}^{m_1} Q_{A_2}^{m_2} \dots \rangle_c \quad (2.36)$$

183 where the Young diagram Y is

$$Y = (m_1, m_2, \dots), \quad m_1 \geq m_2 \geq \dots \geq 1. \quad (2.37)$$

184 The generator of all type-Y CCF is

$$T_{\cup A_i}(\mu_1, \mu_2, \dots) = \log \frac{\langle e^{-\sum_i \mu_i Q_{A_i}} \rangle}{\prod_i \langle e^{-\mu_i Q_{A_i}} \rangle}. \quad (2.38)$$

185 When there are only two balls A and B , the generator is

$$T_{A \cup B}(\mu_1, \mu_2) = \log \frac{\langle e^{-\mu_1 Q_A - \mu_2 Q_B} \rangle}{\langle e^{-\mu_1 Q_A} \rangle \langle e^{-\mu_2 Q_B} \rangle} = \sum_{m_1 \geq 1, m_2 \geq 1} \frac{(-1)^{m_1+m_2} \mu_1^{m_1} \mu_2^{m_2}}{m_1! m_2!} \langle Q_A^{m_1} Q_B^{m_2} \rangle_c. \quad (2.39)$$

186 We parameterize A and B as

$$A = \{(0, \vec{x}) | (\vec{x} - \vec{x}_0)^2 \leq 1\}, \quad B = \{(0, \vec{x}) | \vec{x} \leq R^2\}. \quad (2.40)$$

187 There is only one cross ratio

$$\xi = \frac{4R'}{x_0^2 - (1 - R')^2}. \quad (2.41)$$

188 When the two regions A and B are spacelike-separated, $|x_0| > 1 + R'$, the cross ratio is between
189 0 and 1,

$$0 < \xi < 1. \quad (2.42)$$

190 In some cases, it is more convenient to use an equivalent cross ratio

$$\eta = \frac{\xi}{1 - \xi} = \frac{4R'}{x_0^2 - (1 + R')^2}. \quad (2.43)$$

191 For spacelike-separated regions A and B , the range of the cross ratio η is

$$0 < \eta < \infty. \quad (2.44)$$

192 Since OPE block $Q_A[\mathcal{O}]$ is invariant under conformal transformation, any type- (m_1, m_2) CCF
193 should be a function of cross ratio ξ or η . Actually the OPE block is an eigenvector of the
194 conformal Casimir

$$[L^2, Q_A[\mathcal{O}]] = C_{\Delta, J} Q_A[\mathcal{O}] \quad (2.45)$$

195 where L^2 is the Casimir operator of global conformal group. The eigenvalue $C_{\Delta, J}$ is

$$C_{\Delta, J} = -\Delta(\Delta - d) - J(J + d - 2). \quad (2.46)$$

196 Therefore, any type- $(m - 1, 1)$ CCF should be a conformal block

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_{m-1}] Q_B[\mathcal{O}_m] \rangle_c = D^{(d)}[\mathcal{O}_1, \cdots, \mathcal{O}_m] G_{\Delta_m, J_m}^{(d)}(\xi). \quad (2.47)$$

197 The subscript Δ_m, J_m are the conformal weight and spin of the primary operator \mathcal{O}_m . The index
198 (d) is used to label the dimension of spacetime. The conformal block can be constructed ex-
199 plicitly in even dimensions [25, 26]. In this paper, we just need the diagonal limit of conformal
200 block [27]. Any type- (m_1, m_2) CCF with $m_1 \geq m_2 \geq 2$ is not a conformal block .

201 3 Area law

202 We conjecture that the type- (m) CCF of OPE blocks obeys the following area law

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_m] \rangle_c = \gamma \frac{R^{d-2}}{\epsilon^{d-2}} + \cdots + p_q \log^q \frac{R}{\epsilon} + \cdots. \quad (3.1)$$

203 The leading term is proportional to the area of the boundary ∂A . We inserted the radius $R = 1$
204 into the formula to balance the dimension. The small positive constant ϵ is the UV cutoff which
205 is roughly the distance from the cutoff to the boundary ∂A . The constant γ depends on the
206 choice of the cutoff and the method of regularization, we will not be interested in its explicit
207 value. The \cdots terms are subleading and cutoff dependent. Therefore we omit their forms. The
208 degree q characterizes the maximal power of the logarithmic terms. The coefficient p_q is not
209 invariant under the rescaling of the cutoff, therefore it encodes detail universal information of
210 the theory. When all the OPE blocks are equal to modular Hamiltonian, the degree $q = 1$ for
211 even dimensions according to (2.34). However, as we will see, q is not necessary equal to 1
212 in general. To distinguish different type- (m) CCFs in different dimensions, we write the area
213 law (3.1) more explicitly as

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_m] \rangle_c = \gamma[\mathcal{O}_1, \cdots, \mathcal{O}_m] \frac{R^{d-2}}{\epsilon^{d-2}} + \cdots + p_q^{(d)}[\mathcal{O}_1, \cdots, \mathcal{O}_m] \log^q \frac{R}{\epsilon} + \cdots. \quad (3.2)$$

214 **3.1 Continuation**

215 The two formulas (2.47) and (3.2) are actually related to each other through an analytic
 216 continuation. We use the example of two dimensional modular Hamiltonian to illustrate this
 217 relation. For CFT_2 , the modular Hamiltonian can be decomposed into holomorphic and anti-
 218 holomorphic part

$$H_A = - \int_{-1}^1 dz \frac{1-z^2}{2} T(z+x_0) + c. \tag{3.3}$$

219 The constant c can be fixed by the normalization condition

$$\text{tr}_A \rho_A = \text{tr}_A e^{-H_A} = 1. \tag{3.4}$$

220 Its value doesn't affect the type-Y CCF for any $\sum_i m_i \geq 2$. We also used the convention
 221 $T(z) = -2\pi T_{zz}$ where the subscript z is the holomorphic coordinate $z = t + x$. The radius
 222 of the interval A is 1, we have shifted variable z such that the dependence of the center x_0 is
 223 in the stress tensor. The modular Hamiltonian of region B can be obtained by setting $x_0 = 0$
 224 and restoring the radius R' . The type- $(m-1, 1)$ CCF of modular Hamiltonian is

$$\langle H_A^{m-1} H_B \rangle_c = D^{(2)}[T_{\mu_1 \nu_1}, \dots, T_{\mu_m \nu_m}] G_2^{(2)}(\eta). \tag{3.5}$$

225 The two dimensional conformal block for a chiral operator can be labeled by the conformal
 226 weight h of the operator

$$G_h^{(2)}(\eta) = (-\eta)^h {}_2F_1(h, h, 2h, -\eta). \tag{3.6}$$

227 We can move the interval A to B such that they coincide. In this limit, type- $(m-1, 1)$ CCF
 228 should approach type- (m) CCF. This is equivalent to set $\eta \rightarrow -1$. We can set $x_0 \rightarrow 0$ and then
 229 take the limit $R' \rightarrow 1$,

$$x_A \rightarrow 0, \quad R' = 1 - \epsilon, \quad \epsilon \rightarrow 0. \tag{3.7}$$

230 The cross ratio $\xi \rightarrow -\infty$ or $\eta \rightarrow -1$ by

$$\xi = -\frac{4(1-\epsilon)}{\epsilon^2} \approx -\frac{4}{\epsilon^2}, \quad \eta = -\frac{4(1-\epsilon)}{(2-\epsilon)^2} \approx -1 + \frac{\epsilon^2}{4}. \tag{3.8}$$

231 On the right hand side of (3.5), we find a logarithmic divergent term in this limit

$$G_2^{(2)}(\eta) = 12 \log \frac{2}{\epsilon} + \dots = 12 \log \frac{R}{\epsilon} + \dots \tag{3.9}$$

232 The left hand side of (3.5) approaches type- (m) CCF, therefore

$$\langle H_A^m \rangle_c = 12D^{(2)}[T_{\mu_1 \nu_1}, \dots, T_{\mu_m \nu_m}] \log \frac{R}{\epsilon} + \dots. \tag{3.10}$$

233 We read out the cutoff independent coefficient

$$p_1^{(2)}[T_{\mu_1 \nu_1}, \dots, T_{\mu_m \nu_m}] = 12D^{(2)}[T_{\mu_1 \nu_1}, \dots, T_{\mu_m \nu_m}]. \tag{3.11}$$

234 The relation (3.11) is a typical UV/IR relation for modular Hamiltonian. The left hand side is
 235 the universal coefficient for B and A coincides (UV). On the right hand side, the D coefficient
 236 characterizes the leading order behaviour of CCF when B and A are far away to each other
 237 (IR). They provide equivalent information of the CFT since the constant 12 is completely fixed
 238 by conformal symmetry. The continuation of conformal block can be generalized to higher

239 dimensions. For example, in four dimensions, the conformal block associated with stress tensor
 240 becomes divergent as A approaches B ,

$$G_{4,2}^{(4)} \approx \tilde{\gamma} \frac{R^2}{\epsilon^2} + \dots - 120 \log \frac{R}{\epsilon} + \dots \quad (3.12)$$

241 The leading term is exactly area law and the logarithmic divergent term also appears in the
 242 subleading terms. We can read type- (m) CCF of modular Hamiltonian in four dimensions

$$\langle H_A^m \rangle_c = \gamma \frac{R^2}{\epsilon^2} + \dots + p_1^{(4)} [T_{\mu_1 \nu_1}, \dots, T_{\mu_m \nu_m}] \log \frac{R}{\epsilon} + \dots \quad (3.13)$$

243 with

$$p_1^{(4)} [T_{\mu_1 \nu_1}, \dots, T_{\mu_m \nu_m}] = -120 D^{(4)} [T_{\mu_1 \nu_1}, \dots, T_{\mu_m \nu_m}]. \quad (3.14)$$

244 Note we obtain the area law and logarithmic behaviour of type- (m) CCF of modular Hamil-
 245 tonian without using any knowledge of Rényi entanglement entropy. The method of ana-
 246 lytic continuation can be applied for general dimensions and OPE blocks. A conformal block
 247 $G_{\Delta,J}^{(d)}(\xi)$ obeys area law in the limit $\xi \rightarrow -\infty$ in even dimensions. It has degree $q = 1$ only for
 248 $\Delta = J + d - 2$,

$$G_{\Delta,J}^{(d)}(\xi) = \tilde{\gamma} \frac{R^{d-2}}{\epsilon^{d-2}} + \dots + E^{(d)}[\Delta, J] \log \frac{R}{\epsilon} + \dots, \quad \xi \rightarrow -\infty. \quad (3.15)$$

249 This means that type- (m) CCF of type- J OPE blocks may always obey area law with degree
 250 $q = 1$, the cutoff independent coefficient is

$$p_q^{(d)}[\mathcal{O}_1, \dots, \mathcal{O}_m] = E^{(d)}[\mathcal{O}_m] \times D^{(d)}[\mathcal{O}_1, \dots, \mathcal{O}_m]. \quad (3.16)$$

251 We have replaced the quantum numbers in E function by the corresponding primary opera-
 252 tor. For non-conserved operators, the conformal block $G_{\Delta,J}^{(d)}$ also obeys area law in the limit
 253 $\xi \rightarrow -\infty$ in even dimension, though it has degree $q = 2$

$$G_{\Delta,J}^{(d)}(\xi) = \tilde{\gamma} \frac{R^{d-2}}{\epsilon^{d-2}} + \dots + E^{(d)}[\Delta, J] \log^2 \log \frac{R}{\epsilon} + \dots, \quad \xi \rightarrow -\infty. \quad (3.17)$$

254 Therefore, type- (m) CCF of type- O OPE blocks obeys area law with degree $q = 2$. We can
 255 obtain similar UV/IR relations as (3.16). In odd dimensions, the story is the same. The degree
 256 q is 0 for type- (m) CCF of type- J OPE blocks and 1 for type- O OPE blocks.

257 3.2 Kinematic information

258 The function $E^{(d)}[\mathcal{O}]$ is completely fixed by conformal symmetry. It can be obtained by reading
 259 out the coefficient of the logarithmic term with degree q . For each fixed quantum number Δ
 260 and J , there is a unique number $E^{(d)}[\mathcal{O}]$. For type- J OPE block in two dimensions, the primary
 261 operator \mathcal{O} has dimension $\Delta = J = h$. The conformal block (3.6) has degree $q = 1$ in the limit
 262 $\eta \rightarrow -1$. The function $E^{(2)}[\mathcal{O}]$ is

$$E^{(2)}[\mathcal{O}] = \frac{2\Gamma(2h)}{\Gamma(h)^2}, \quad \Delta = J = h. \quad (3.18)$$

263 For type- O OPE block, the primary operator \mathcal{O} has dimension $\Delta = h + \bar{h}$ and spin $J = h - \bar{h}$.
 264 The conformal block has degree $q = 2$ in the limit $\eta \rightarrow -1$. The function $E^{(2)}[\mathcal{O}]$ is

$$E^{(2)}[\mathcal{O}] = \begin{cases} \frac{2^{4h} \Gamma(h + \frac{1}{2})^2}{\pi \Gamma(h)^2} & J = 0, h > 0 \\ -\frac{4^{2h-1} \Gamma(h - \frac{1}{2}) \Gamma(h + \frac{1}{2})}{\pi \Gamma(h-1) \Gamma(h)} & J = 1, h > 1 \\ \frac{4^{2h-3} (h-2)(h-1)(2h-3)(2h-1) \Gamma(h - \frac{3}{2})^2}{\pi \Gamma(h)^2} & J = 2, h > 2 \\ \dots & \dots \end{cases} \quad (3.19)$$

265 In four dimensions, we also find

$$E^{(4)}[\mathcal{O}] = \begin{cases} 12 & \Delta = 3, J = 1 \\ -120 & \Delta = 4, J = 2 \\ 840 & \Delta = 5, J = 3 \\ \dots & \dots \end{cases} \quad (3.20)$$

266 for conserved currents and

$$E^{(4)}[\mathcal{O}] = \begin{cases} -\frac{2^{2\Delta-1}\Gamma(\frac{\Delta-1}{2})\Gamma(\frac{\Delta+1}{2})}{\pi\Gamma(\frac{\Delta-2}{2})^2} & \Delta > 1, J = 0, \\ \frac{2^{2\Delta-1}\Gamma(\frac{\Delta}{2})\Gamma(\frac{\Delta+2}{2})}{\pi\Gamma(\frac{\Delta-3}{2})\Gamma(\frac{\Delta+1}{2})} & \Delta > 3, J = 1, \\ -\frac{4^{\Delta-1}(\Delta-2)\Gamma(\frac{\Delta-3}{2})\Gamma(\frac{\Delta+3}{2})}{\pi\Gamma(\frac{\Delta-4}{2})\Gamma(\frac{\Delta+2}{2})} & \Delta > 4, J = 2, \\ \dots & \dots \end{cases} \quad (3.21)$$

267 for non-conserved operators. In three dimensions, we find

$$E^{(3)}[\mathcal{O}] = \begin{cases} -\frac{2^{2\Delta-1}(\Delta-1)\Gamma(\Delta-\frac{1}{2})}{\sqrt{\pi}\Gamma(\Delta-1)} & \Delta > \frac{1}{2}, J = 0, \\ \frac{2^{\Delta+1}\Delta\Gamma(\Delta-\frac{1}{2})}{\Gamma(\frac{\Delta-2}{2})\Gamma(\frac{\Delta+1}{2})} & \Delta > 2, J = 1, \\ -\frac{2^{2\Delta-1}(\Delta^2-1)\Gamma(\Delta-\frac{1}{2})}{\sqrt{\pi}(\Delta-2)^2\Delta\Gamma(\Delta-3)} & \Delta > 3, J = 2, \\ \dots & \dots \end{cases} \quad (3.22)$$

268 for non-conserved operators. Note for conserved currents in odd dimensions, the function
 269 $E^{(3)}[\mathcal{O}]$ may depend on explicit choice of cutoff. For example, a transformation $\epsilon \rightarrow \epsilon(1 + a\epsilon)$
 270 may shift its value. This is because the degree is 0, there is no logarithmic divergence at all.

271 3.3 UV/IR relation

272 The UV/IR relation (3.16) relates type-(m) CCF to type-($m - 1, 1$) CCF. This relation may
 273 simplify computation in many cases. To see this point, let's compute the following type-(2)
 274 CCF in two dimensions

$$\begin{aligned} \langle Q_A[\mathcal{O}]^2 \rangle_c &= \int_{-1}^1 dz_1 \int_{-1}^1 dz_2 \frac{(1-z_1^2)^{h-1}(1-z_2^2)^{h-1}}{(z_1-z_2)^{2h}} \\ &= \frac{(-1)^{-h}\sqrt{\pi}\Gamma(h)}{\Gamma(h+\frac{1}{2})} \int_{-1}^1 dz_1 \frac{1}{1-z_1^2} \\ &= \frac{(-1)^{-h}\sqrt{\pi}\Gamma(h)}{\Gamma(h+\frac{1}{2})} \log \frac{2}{\epsilon}. \end{aligned} \quad (3.23)$$

275 This is a double integral with poles at $z_1 = z_2$. We regularize the integral by ignoring these
 276 poles at the second step. At the last step, we insert a UV cutoff to regularize the integral.
 277 However, using UV/IR relation, one just need to fix the coefficient D which is related to the
 278 large distance behaviour of the type-(1, 1) CCF,

$$\langle Q_A[\mathcal{O}]Q_B[\mathcal{O}] \rangle_c = \int_{-1}^1 dz_1 \int_{-1}^1 dz_2 \frac{(1-z_1^2)^{h-1}(1-z_2^2)^{h-1}}{(z_1-z_2+x_0)^{2h}}. \quad (3.24)$$

279 In the large distance limit, $x_0 \rightarrow \infty$, the integral becomes simpler

$$\begin{aligned} \langle Q_A[\mathcal{O}]Q_B[\mathcal{O}] \rangle_c &\approx \int_{-1}^1 dz_1 \int_{-1}^1 dz_2 \frac{(1-z_1^2)^{h-1}(1-z_2^2)^{h-1}}{x_0^{2h}} \\ &= 4^{-h} \left(\frac{\sqrt{\pi}\Gamma(h)}{\Gamma(h+\frac{1}{2})} \right)^2 \eta^h. \end{aligned} \tag{3.25}$$

280 We have used the relation $\eta \approx \frac{4}{x_0^2}$ in the large distance limit. Then we can read out

$$D^{(2)}[\mathcal{O}, \mathcal{O}] = (-1)^{-h} 4^{-h} \left(\frac{\sqrt{\pi}\Gamma(h)}{\Gamma(h+\frac{1}{2})} \right)^2. \tag{3.26}$$

281 Combining UV/IR relation and (3.18), we find

$$p_1^{(2)}[\mathcal{O}, \mathcal{O}] = E^{(2)}[\mathcal{O}] \times D^{(2)}[\mathcal{O}, \mathcal{O}] = \frac{(-1)^{-h} \sqrt{\pi}\Gamma(h)}{\Gamma(h+\frac{1}{2})}. \tag{3.27}$$

282 The result is exactly the same as (3.23). We use UV/IR relation to obtain type-(3) CCF for
283 type-J OPE blocks in two dimensions, the cutoff independent coefficient is

$$p_1^{(2)}[\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3] = \frac{C_{123} \pi^{3/2} (-1)^{\frac{h_1+h_2+h_3}{2}} \Gamma(h_1)\Gamma(h_2)\Gamma(h_3)\kappa}{\Gamma(\frac{1+h_1+h_2-h_3}{2})\Gamma(\frac{1+h_1+h_3-h_2}{2})\Gamma(\frac{1+h_2+h_3-h_1}{2})\Gamma(\frac{h_1+h_2+h_3}{2})}, \tag{3.28}$$

284 where the constant $\kappa = \frac{1}{2}[1 + (-1)^{h_1+h_2+h_3}]$. We notice that the result is totally symmetric
285 under exchange of any two conformal weights. Since there are different ways to uplift type-(m)
286 to type-($m-1, 1$), the cutoff independent coefficient should be identical since they characterize
287 the same CCF after taking the limit $A \rightarrow B$. For $m = 3$, this is a cyclic identity

$$p_q^{(d)}[\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3] = p_q^{(d)}[\mathcal{O}_2, \mathcal{O}_3, \mathcal{O}_1] = p_q^{(d)}[\mathcal{O}_3, \mathcal{O}_1, \mathcal{O}_2]. \tag{3.29}$$

288 UV/IR relation and the cyclic identity has been checked for type-(m) CCF ($m=2,3$) in four
289 dimensions. We list the cutoff independent coefficients below [16].

290 • Type-(2). The normalization constants are set to 1.

291 – Spin 1-1 conserved currents.

$$p_1^{(4)}[\mathcal{J}_\mu, \mathcal{J}_\nu] = -\frac{\pi^2}{3}. \tag{3.30}$$

292 – Spin 2-2 conserved currents.

$$p_1^{(4)}[T_{\mu\nu}, T_{\rho\sigma}] = -\frac{\pi^2}{40}. \tag{3.31}$$

293 – Spin 0-0 non-conserved operators.

$$p_2^{(4)}[\mathcal{O}, \mathcal{O}] = -\frac{4\pi^2(\Delta-1)\Gamma(\Delta-2)^2\Gamma(\frac{\Delta}{2})^4}{\Gamma(\Delta)^2\Gamma(\Delta-1)^2}. \tag{3.32}$$

294 – Spin 1-1 non-conserved operators.

$$p_2^{(4)}[\mathcal{O}_\mu, \mathcal{O}_\nu] = -\frac{4^{1-\Delta}\pi^3\Delta\Gamma(\frac{\Delta-3}{2})\Gamma(\frac{\Delta+1}{2})}{\Gamma(\frac{\Delta}{2}+1)^2}, \quad \Delta > 3. \tag{3.33}$$

295 – Spin 2-2 non-conserved operators.

$$p_2^{(4)}[\mathcal{O}_{\mu\nu}, \mathcal{O}_{\rho\sigma}] = -\frac{3\pi^2(\Delta-2)\Delta^2\Gamma(\frac{\Delta}{2}-2)^2\Gamma(\frac{\Delta}{2}-1)^2}{64\Gamma(\Delta-4)\Gamma(\Delta+2)}, \quad \Delta > 4. \quad (3.34)$$

296 • Type-(3).

297 – Spin 1-1-2 conserved currents. The three point function of zero components are
298 fixed by conformal symmetry

$$\langle T_{00}(x_1)\mathcal{J}_0(x_2)\mathcal{J}_0(x_3) \rangle_c = \frac{C_{T\mathcal{J}\mathcal{J}}}{x_{12}^4 x_{13}^2 x_{23}^2}. \quad (3.35)$$

299 Then the coefficient

$$p_1^{(4)}[\mathcal{J}_\mu, \mathcal{J}_\nu, T_{\rho\sigma}] = -\frac{\pi^3}{2} C_{T\mathcal{J}\mathcal{J}}. \quad (3.36)$$

300 – Spin 2-2-2 conserved currents. The three point function of zero components are
301 fixed by conformal symmetry

$$\langle T_{00}(x_1)T_{00}(x_2)T_{00}(x_3) \rangle_c = \frac{C_{TTT}}{x_{12}^4 x_{13}^4 x_{23}^4}. \quad (3.37)$$

302 Then the coefficient

$$p_1^{(4)}[T_{\mu\nu}, T_{\rho\sigma}, T_{\alpha\beta}] = \frac{\pi^3}{12} C_{TTT}. \quad (3.38)$$

303 – Spin 0-0-0 non-conserved currents.

$$p_2^{(4)}[\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3] = -2^{4-\Delta_1-\Delta_2-\Delta_3} \pi^3 C_{123} \int_{\mathbb{D}^2} d\zeta d\bar{\zeta} (\zeta + \bar{\zeta})^2 \int_{\mathbb{D}^2} d\zeta' d\bar{\zeta}' (\zeta' + \bar{\zeta}')^2 \\ \times (1-\zeta^2)^{\frac{\Delta_1-4}{2}} (1-\bar{\zeta}^2)^{\frac{\Delta_1-4}{2}} (1-\zeta'^2)^{\frac{\Delta_2-4}{2}} (1-\bar{\zeta}'^2)^{\frac{\Delta_2-4}{2}} \int_0^\pi d\theta \frac{\sin \theta}{(a+b \cos \theta)^{\frac{\Delta_{12,3}}{2}}}, \quad (3.39)$$

304 Though the expression (3.39) is not symmetric superficially under exchange of any two con-
305 formal weights, we checked explicitly that it satisfies the cyclic identity for integer conformal
306 weights. For $m = 4$, the UV/IR relation and the cyclic identity are much more harder to check.
307 We considered type-(4) CCF for massless free scalar theory [13, 14]. In this theory, one can
308 construct an infinite tower of conserved currents with even spin [28]. The four point functions
309 can be calculated explicitly. Therefore we can find type-(3, 1) and type-(4) CCFs and read out
310 the corresponding coefficients. For example, for spin-2-2-2-4 conserved currents [14],

$$D[2, 2, 2, 4] = \frac{3}{70} D[2, 2, 4, 2]. \quad (3.40)$$

311 Both of them leads to the cutoff coefficients

$$p_1^{(2)}[2, 2, 2, 4] = \frac{2\Gamma(8)}{\Gamma(4)^2} D[2, 2, 2, 4] = \frac{2\Gamma(4)}{\Gamma(2)^2} D[2, 2, 4, 2] = p_1^{(2)}[2, 2, 4, 2]. \quad (3.41)$$

312 The cyclic identity is obeyed.

313 **3.4 Discussion**

314 The UV/IR relation should be slightly modified when the CCF contains both type-J and type-O
 315 OPE blocks. One simple example is the following type-(3) CCF

$$\langle Q_A[\mathcal{J}]Q_A[\mathcal{O}]Q_A[\tilde{\mathcal{O}}] \rangle_c \tag{3.42}$$

316 where $Q_A[\mathcal{J}]$ is a type-J OPE block while $Q_A[\mathcal{O}]$ and $Q_A[\tilde{\mathcal{O}}]$ are type-O OPE blocks. This CCF
 317 is related to the following two type-(2, 1) CCFs

$$\langle Q_A[\tilde{\mathcal{O}}]Q_A[\mathcal{J}]Q_B[\mathcal{O}] \rangle_c = D^{(d)}[\tilde{\mathcal{O}}, \mathcal{J}, \mathcal{O}]G_{\Delta, J}^{(d)}(\xi), \tag{3.43}$$

$$\langle Q_A[\mathcal{O}]Q_A[\tilde{\mathcal{O}}]Q_B[\mathcal{J}] \rangle_c = D^{(d)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}]G_{\Delta', J'}^{(d)}(\xi). \tag{3.44}$$

318 We choose $d = 4$. Taking the limit $A \rightarrow B$ from (3.43), we find a type-(3) CCF with degree
 319 $q = 2$, the UV/IR relation reads

$$p_2^{(4)}[\tilde{\mathcal{O}}, \mathcal{J}, \mathcal{O}] = E^{(4)}[\mathcal{O}] \times D^{(4)}[\tilde{\mathcal{O}}, \mathcal{J}, \mathcal{O}] \tag{3.45}$$

320 We can also take the limit $A \rightarrow B$ from (3.44), then we will find a type-(3) CCF with degree
 321 $q = 1$, the UV/IR relation reads

$$p_1^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}] = E^{(4)}[\mathcal{J}] \times D^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}]. \tag{3.46}$$

322 The equations (3.45) and (3.46) are not identical superficially since the subscript q are not
 323 equal to each other. However, an explicit calculation for spin 2-0-0 and spin 2-2-0 in four
 324 dimensions [16] shows that the coefficient $D^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}]$ is actually divergent logarithmically,
 325

$$D^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}] = D_{\log}^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}] \log \frac{R}{\epsilon} + \dots \tag{3.47}$$

326 The terms in \dots are finite and depends on cutoff scale. Due to the logarithmic divergence
 327 behaviour of the coefficient $D^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}]$, the degree of type-(3) CCF from (3.44) increases
 328 1, the modified UV/IR relation becomes

$$p_2^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}] = E^{(4)}[\mathcal{J}] \times D_{\log}^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}]. \tag{3.48}$$

329 We checked explicitly that the two constants (3.45) and (3.48) are equal to each other. The
 330 cyclic identity is still satisfied after counting the logarithmic divergence of the D function.

331 **4 Generalizations**

332 The area law and logarithmic behaviour in the subleading terms can be extended in different
 333 directions. In this section, we mention several extensions.

- 334 • UV/IR relation. In general, one can uplift type-(m) CCF to type-($p, m - p$) CCF

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_m] \rangle_c \xrightarrow{\text{uplift}} \langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_p] Q_B[\mathcal{O}_{p+1}] \cdots Q_B[\mathcal{O}_m] \rangle_c, \quad 1 \leq p \leq m-1. \tag{4.1}$$

335 When p is not 1 and $m - 1$, the type-($p, m - p$) CCF is not a conformal block. It is still
 336 a function of cross ratio ξ , therefore it should reproduce type-(m) CCF after taking the
 337 limit $A \rightarrow B$,

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_m] \rangle_c = \lim_{\xi \rightarrow -\infty} \langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_p] Q_B[\mathcal{O}_{p+1}] \cdots Q_B[\mathcal{O}_m] \rangle_c. \tag{4.2}$$

338 Obviously, this also defines a UV/IR relation between $p_q^{(d)}$ and several coefficients in the
 339 type- $(p, m - p)$ CCF. Since the right hand side is not proportional to conformal block,
 340 it is not easy to write out an explicit formula. Nevertheless, one may still check the
 341 relation (4.2) case by case. One example is to consider the type-(2, 2) CCF of modular
 342 Hamiltonian in CFT_2 . By making use of the universal feature of CCF of stress tensor, one
 343 can fix the generator of type- (m_1, m_2) CCFs [14]

$$T_{AUB}(\mu_1, \mu_2) = -\frac{c}{2} \text{tr} \log \left[\mathbf{1} - \begin{pmatrix} A & C \\ D & B \end{pmatrix} \right], \quad (4.3)$$

344 where the matrices A, B, C and D are

$$A_{xx'} = \frac{\eta^2}{4} \int_0^\infty dy \frac{\sqrt{xx'} y \sinh \pi \mu_1 x \sinh \pi \mu_2 y}{\sinh \pi x' \sinh \pi y \sinh \pi(1 + \mu_1)x \sinh \pi(1 + \mu_2)y} \left(\frac{x_{13}}{x_{23}}\right)^{i(x-x')} \mathcal{F}(x, x', y), \quad (4.4)$$

$$B_{xx'} = \frac{\eta^2}{4} \int_0^\infty dy \frac{\sqrt{xx'} y \sinh \pi \mu_1 x \sinh \pi \mu_2 y}{\sinh \pi x' \sinh \pi y \sinh \pi(1 + \mu_1)x \sinh \pi(1 + \mu_2)y} \left(\frac{x_{13}}{x_{23}}\right)^{-i(x-x')} \mathcal{F}(x', x, y), \quad (4.5)$$

$$C_{xx'} = \frac{\eta^2}{4} \int_0^\infty dy \frac{\sqrt{xx'} y \sinh \pi \mu_1 x \sinh \pi \mu_2 y}{\sinh \pi x' \sinh \pi y \sinh \pi(1 + \mu_1)x \sinh \pi(1 + \mu_2)y} \left(\frac{x_{13}}{x_{23}}\right)^{i(x+x')} \mathcal{F}(x, -x', y) \quad (4.6)$$

$$D_{xx'} = \frac{\eta^2}{4} \int_0^\infty dy \frac{\sqrt{xx'} y \sinh \pi \mu_1 x \sinh \pi \mu_2 y}{\sinh \pi x' \sinh \pi y \sinh \pi(1 + \mu_1)x \sinh \pi(1 + \mu_2)y} \left(\frac{x_{13}}{x_{23}}\right)^{-i(x+x')} \mathcal{F}(-x, x', y) \quad (4.7)$$

345 with

$$\begin{aligned} \mathcal{F}(x, x', y) = & {}_2F_1(1 + ix, 1 - iy, 2, -\eta) {}_2F_1(1 - ix', 1 + iy, 2, -\eta) \\ & + {}_2F_1(1 + ix, 1 + iy, 2, -\eta) {}_2F_1(1 - ix', 1 - iy, 2, -\eta). \end{aligned} \quad (4.8)$$

346 \mathcal{F} and its complex conjugate obey

$$\mathcal{F}^*(x, x', y) = \mathcal{F}(x', x, y), \quad \mathcal{F}^*(-x, -x', y) = \mathcal{F}(x, x', y). \quad (4.9)$$

347 so

$$A = B^*, \quad C = D^*. \quad (4.10)$$

348 We read out the first few CCFs

$$\begin{aligned} \langle H_A^m \rangle_c &= \frac{cm!}{12} \log \frac{2}{\epsilon}, \\ \langle H_A^{m-1} H_B \rangle_c &= \frac{cm!}{144} G_2^{(2)}(\eta), \\ \langle H_A^2 H_B^2 \rangle_c &= c \left\{ \frac{1 + \eta}{\eta^2} [4\text{Li}_3(1 + \eta) - 2 \log(1 + \eta) \text{Li}_2(1 + \eta) + \frac{2 \log(1 + \eta)}{3} \text{Li}_2(-\eta) \right. \\ &\quad \left. + \frac{1 + \eta}{3} \log^2(1 + \eta) - \frac{\pi^2}{3} \log(1 + \eta) - 4\zeta(3)] + \frac{2 + \eta}{3\eta} [2\text{Li}_2(-\eta) + 3 \log(1 + \eta)] - \frac{4}{3} \right\}, \end{aligned} \quad (4.11)$$

349 where the polylogarithm $\text{Li}_n(z)$ is

$$\text{Li}_n(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^n}. \quad (4.12)$$

350 The relation (4.2) can be checked for $p = 2, m = 4$. The right hand side is

$$\lim_{\eta \rightarrow -1} \langle H_A^2 H_B^2 \rangle_c = 2c \log \frac{2}{\epsilon} + \dots. \quad (4.13)$$

351 The cutoff independent coefficient $2c$ matches with the one in $\langle H_A^4 \rangle_c$.

- New power law. In the previous discussion, we focus on the case that B and A coincide with each other. However, there are other cases that the CCFs are still divergent. One can consider the limit that A just attaches the edge of B ,

$$R' = 1, \quad x_0 = 2 + \epsilon, \quad \epsilon \rightarrow 0. \tag{4.14}$$

The cross ratio ξ does not approach $-\infty$ but 1

$$\xi = \frac{4}{(2 + \epsilon)^2} = 1 - \epsilon + \dots \tag{4.15}$$

We can define a new CCF which is also divergent from type- $(m - 1, 1)$ CCF

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_{m-1}] \odot Q_B[\mathcal{O}_m] \rangle_c = \lim_{\xi \rightarrow 1} \langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_{m-1}] Q_B[\mathcal{O}_m] \rangle_c \tag{4.16}$$

The continuation of conformal block tells us that the new CCF obeys a new power law

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_{m-1}] \odot Q_B[\mathcal{O}_m] \rangle_c = \tilde{\gamma} \left(\frac{R}{\epsilon}\right)^{\frac{d-2}{2}} + \dots + \bar{p}_q^{(d)} \log^q \frac{R}{\epsilon} + \dots \tag{4.17}$$

The leading term is proportional to

$$\mathcal{L} = R^{\frac{d-2}{2}} = \sqrt{\mathcal{A}} \tag{4.18}$$

which is the characteristic length of the region A in four dimensions. In two dimensions, the leading term is a logarithmic term with power q . In this case, there is a new UV/IR relation between \bar{p}_q and D coefficient, we write it schematically

$$\bar{p}_q = \bar{E} \times D. \tag{4.19}$$

The function $\bar{E}^{(d)}[\mathcal{O}]$ is proportional to $E^{(d)}[\mathcal{O}]$. The proportional constant is shown below.

– d is even.

- * For conserved current \mathcal{O} with conformal weight $\Delta = J + d - 2$,

$$\bar{E}^{(d)}[\mathcal{O}] = \frac{(-1)^J}{2} E^{(d)}[\mathcal{O}]. \tag{4.20}$$

- * For non-conserved current \mathcal{O} with conformal weight Δ and spin J ,

$$\bar{E}^{(d)}[\mathcal{O}] = \frac{(-1)^J}{4} E^{(d)}[\mathcal{O}]. \tag{4.21}$$

We checked the relation for $d = 2, 4$ and spin $J \leq 2$.

– d is odd.

- * For non-conserved current \mathcal{O} with conformal weight Δ and spin J ,

$$\bar{E}^{(d)}[\mathcal{O}] = \frac{(-1)^J}{2} E^{(d)}[\mathcal{O}]. \tag{4.22}$$

- * For conserved current \mathcal{O} , there is no logarithmic divergent term in the CCF.

We checked the relation for $d = 3$ and spin $J \leq 2$.

Since D function is the same, we find a relation between two cutoff independent coefficients p and \bar{p} ,

$$\frac{p}{E} = \frac{\bar{p}}{\bar{E}}. \tag{4.23}$$

374 5 Summary and outlook

375 In this report, we have introduced the area law (3.1) of type-(m) CCF of OPE blocks. It is a
 376 generalization of the area law of entanglement entropy. We will list several open problems for
 377 future work.

378 • Higher $m \geq 4$. In most of the work, we restrict to the region $m \geq 3$. This is because
 379 the structure of m -point correlation function of primary operators in CFT is fixed up to
 380 $m = 3$. For $m \geq 4$, it is harder to extract cutoff independent coefficient.

381 • UV/IR relation. The UV/IR relation

$$p = E \times D \quad (5.1)$$

382 has been checked for several examples. A rigorous proof is still lacking.

383 • Cyclic identity. The cyclic identity of p reflects the fact that p is independent of the way
 384 to regularize the type-(m) CCF. However, we feel that a direct computation is impossible
 385 to check this identity.

386 • New power law. We generalize the type-(m_1, m_2) CCF to the case that A and B just
 387 attaches with each other. The corresponding CCF is divergent with a new power law
 388 (4.17). The corresponding new UV/IR relation

$$\bar{p} = \bar{E} \times D \quad (5.2)$$

389 also needs understanding.

390 • Deformed reduced density matrix. This exponential operator is similar to “Wilson loop”
 391 in gauge theories [29,30] despite the fact that the OPE block has no lower bound in gen-
 392 eral. When the OPE block has a lower bound, the logarithm of the vacuum expectation
 393 value of the deformed reduced density matrix

$$\log \langle e^{-\mu Q_A} \rangle \quad (5.3)$$

394 should also obey area law with logarithmic divergence. There may be a gravitational
 395 dual for this quantity as [31,32]. The similarity of the area law between this program
 396 and black hole entropy implies that the classical part contributes to the area term while
 397 quantum effects lead to logarithmic corrections.

398 • Multiple integrals. According to the method of continuation of conformal block, area law
 399 of type-(m) CCF is protected by conformal invariance. However, the method of contin-
 400 uation itself cannot guarantee that it always leads to correct result. One has to develop
 401 other methods to deal with the multiple integrals. In two dimensions, one should gen-
 402 eralize Selberg integrals [33,34] to include more parameters [15].

403 References

404 [1] J. D. Bekenstein, “Black holes and entropy,” Phys. Rev. D 7 (1973), 2333-2346
 405 doi:10.1103/PhysRevD.7.2333

- 406 [2] S. W. Hawking, “Particle Creation by Black Holes,” *Commun. Math. Phys.* **43** (1975),
407 199-220 doi:10.1007/BF02345020
- 408 [3] B. Czech, L. Lamprou, S. McCandlish, B. Mosk and J. Sully, “A Stereoscopic Look into
409 the Bulk,” *JHEP* **1607**, 129 (2016) doi:10.1007/JHEP07(2016)129 [arXiv:1604.03110
410 [hep-th]].
- 411 [4] J. de Boer, F. M. Haehl, M. P. Heller and R. C. Myers, “Entanglement, hologra-
412 phy and causal diamonds,” *JHEP* **1608**, 162 (2016) doi:10.1007/JHEP08(2016)162
413 [arXiv:1606.03307 [hep-th]].
- 414 [5] S. Ferrara, A. F. Grillo and R. Gatto, “Manifestly conformal covariant operator-product
415 expansion,” *Lett. Nuovo Cim.* **2S2**, 1363 (1971) [*Lett. Nuovo Cim.* **2**, 1363 (1971)].
416 doi:10.1007/BF02770435
- 417 [6] S. Ferrara, A. F. Grillo and R. Gatto, “Manifestly conformal-covariant expansion on the
418 light cone,” *Phys. Rev. D* **5**, 3102 (1972). doi:10.1103/PhysRevD.5.3102
- 419 [7] R. Haag, “Local quantum physics: fields, particles, algebras ,” Springer, Berlin, Germany
420 (1992).
- 421 [8] L. Bombelli, R. K. Koul, J. Lee and R. D. Sorkin, “A Quantum Source of Entropy for Black
422 Holes,” *Phys. Rev. D* **34**, 373 (1986).
- 423 [9] M. Srednicki, “Entropy and area,” *Phys. Rev. Lett.* **71**, 666 (1993)
424 doi:10.1103/PhysRevLett.71.666 [hep-th/9303048].
- 425 [10] C. G. Callan, Jr. and F. Wilczek, “On geometric entropy,” *Phys. Lett. B* **333**, 55 (1994)
426 doi:10.1016/0370-2693(94)91007-3 [hep-th/9401072].
- 427 [11] H. Araki, “Relative Entropy of States of Von Neumann Algebras,” *Publ. Res. Inst. Math.*
428 *Sci. Kyoto* **1976**, 809 (1976).
- 429 [12] S. Ryu and T. Takayanagi, “Holographic derivation of entanglement entropy from
430 AdS/CFT,” *Phys. Rev. Lett.* **96**, 181602 (2006) doi:10.1103/PhysRevLett.96.181602
431 [hep-th/0603001].
- 432 [13] J. Long, “Correlation function of modular Hamiltonians,” *JHEP* **11** (2019), 163
433 doi:10.1007/JHEP11(2019)163 [arXiv:1907.00646 [hep-th]].
- 434 [14] J. Long, “Correlation function with the insertion of zero modes of modular Hamilto-
435 nians,” *JHEP* **01** (2020), 173 doi:10.1007/JHEP01(2020)173 [arXiv:1911.11487 [hep-
436 th]].
- 437 [15] J. Long, “Logarithmic behaviour of connected correlation function in CFT,”
438 [arXiv:2001.05129 [hep-th]].
- 439 [16] J. Long, “Area law of connected correlation function in higher dimensional conformal
440 field theory,” [arXiv:2007.15380 [hep-th]].
- 441 [17] S. N. Solodukhin, “The Conical singularity and quantum corrections to entropy of
442 black hole,” *Phys. Rev. D* **51** (1995), 609-617 doi:10.1103/PhysRevD.51.609 [arXiv:hep-
443 th/9407001 [hep-th]].
- 444 [18] S. N. Solodukhin, “On ‘Nongeometric’ contribution to the entropy of black hole due to
445 quantum corrections,” *Phys. Rev. D* **51** (1995), 618-621 doi:10.1103/PhysRevD.51.618
446 [arXiv:hep-th/9408068 [hep-th]].

- 447 [19] R. K. Kaul and P. Majumdar, “Logarithmic correction to the Bekenstein-Hawking entropy,”
448 Phys. Rev. Lett. **84** (2000), 5255-5257 doi:10.1103/PhysRevLett.84.5255 [arXiv:gr-
449 qc/0002040 [gr-qc]].
- 450 [20] S. Carlip, “Logarithmic corrections to black hole entropy from the Cardy formula,” Class.
451 Quant. Grav. **17** (2000), 4175-4186 doi:10.1088/0264-9381/17/20/302 [arXiv:gr-
452 qc/0005017 [gr-qc]].
- 453 [21] T. R. Govindarajan, R. K. Kaul and V. Suneeta, “Logarithmic correction to the Bekenstein-
454 Hawking entropy of the BTZ black hole,” Class. Quant. Grav. **18** (2001), 2877-2886
455 doi:10.1088/0264-9381/18/15/303 [arXiv:gr-qc/0104010 [gr-qc]].
- 456 [22] A. Sen, “Logarithmic Corrections to Schwarzschild and Other Non-extremal Black Hole
457 Entropy in Different Dimensions,” JHEP **04** (2013), 156 doi:10.1007/JHEP04(2013)156
458 [arXiv:1205.0971 [hep-th]].
- 459 [23] S. Minwalla, “Restrictions imposed by superconformal invariance on quantum field the-
460 ories,” Adv. Theor. Math. Phys. **2**, 783 (1998) doi:10.4310/ATMP.1998.v2.n4.a4 [hep-
461 th/9712074].
- 462 [24] H. Casini, M. Huerta and R. C. Myers, “Towards a derivation of holographic entanglement
463 entropy,” JHEP **1105**, 036 (2011) doi:10.1007/JHEP05(2011)036 [arXiv:1102.0440
464 [hep-th]].
- 465 [25] F. A. Dolan and H. Osborn, “Conformal four point functions and the operator product
466 expansion,” Nucl. Phys. B **599**, 459 (2001) doi:10.1016/S0550-3213(01)00013-X [hep-
467 th/0011040].
- 468 [26] F. A. Dolan and H. Osborn, “Conformal partial waves and the operator product ex-
469 pansion,” Nucl. Phys. B **678**, 491 (2004) doi:10.1016/j.nuclphysb.2003.11.016 [hep-
470 th/0309180].
- 471 [27] M. Hogervorst, H. Osborn and S. Rychkov, “Diagonal Limit for Conformal Blocks in d Di-
472 mensions,” JHEP **1308**, 014 (2013) doi:10.1007/JHEP08(2013)014 [arXiv:1305.1321
473 [hep-th]].
- 474 [28] I. Bakas and E. Kiritsis, Nucl. Phys. B **343** (1990), 185-204 [erratum: Nucl. Phys. B **350**
475 (1991), 512-512] doi:10.1016/0550-3213(90)90600-I
- 476 [29] J. M. Maldacena, “Wilson loops in large N field theories,” Phys. Rev. Lett. **80**, 4859 (1998)
477 doi:10.1103/PhysRevLett.80.4859 [hep-th/9803002].
- 478 [30] S. J. Rey and J. T. Yee, “Macroscopic strings as heavy quarks in large N gauge theory and
479 anti-de Sitter supergravity,” Eur. Phys. J. C **22**, 379 (2001) doi:10.1007/s100520100799
480 [hep-th/9803001].
- 481 [31] D. L. Jafferis and S. J. Suh, “The Gravity Duals of Modular Hamiltonians,” JHEP **09**
482 (2016), 068 doi:10.1007/JHEP09(2016)068 [arXiv:1412.8465 [hep-th]].
- 483 [32] D. L. Jafferis, A. Lewkowycz, J. Maldacena and S. J. Suh, “Relative entropy
484 equals bulk relative entropy,” JHEP **06** (2016), 004 doi:10.1007/JHEP06(2016)004
485 [arXiv:1512.06431 [hep-th]].
- 486 [33] A. Selberg, “Bemerkninger om et multipelt integral,” Norsk. Mat. Tidsskr. **24**(1944)71-78.
- 487 [34] P. Forrester and S. Warnaar, “The importance of the Selberg integral,” Bull. Amer. Math.
488 Soc. (N.S.) **45** (2008), 489-534.