

# Area law and OPE blocks in conformal field theory

Jiang Long

School of Physics, Huazhong University of Science and Technology,  
Wuhan, Hubei 430074, China

★ [longjiang@hust.edu.cn](mailto:longjiang@hust.edu.cn)

*4th International Conference on Holography,  
String Theory and Discrete Approach  
Hanoi, Vietnam, 2020*  
doi:[10.21468/SciPostPhysProc.4](https://doi.org/10.21468/SciPostPhysProc.4)

## Abstract

This is an introduction to the relationship between area law and OPE blocks in conformal field theory.

Copyright J.Long

This work is licensed under the Creative Commons  
[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Published by the SciPost Foundation.

Received ??-??-20??

Accepted ??-??-20??

Published ??-??-20??

doi:[10.21468/SciPostPhysProc.4](https://doi.org/10.21468/SciPostPhysProc.4)??

1

## 2 Contents

3	<b>1 Introduction</b>	<b>2</b>
4	<b>2 Setup</b>	<b>2</b>
5	2.1 Area law	3
6	2.2 OPE block	4
7	2.3 Modular Hamiltonian and area law	6
8	2.4 Deformed reduced density matrix and connected correlation function	6
9	<b>3 Area law</b>	<b>8</b>
10	3.1 Continuation	9
11	3.2 Kinematic information	11
12	3.3 UV/IR relation	11
13	3.4 Discussion	14
14	<b>4 Generalizations</b>	<b>15</b>
15	<b>5 Summary and outlook</b>	<b>17</b>
16	<b>References</b>	<b>18</b>

17  
18

## 19 **1 Introduction**

20 This report consists a summary of our recent progress on the relationship between area law and  
21 OPE blocks. Area law has been a continuous topic in physics. The prototype of area law dates  
22 back to black hole physics in general relativity. The unusual property that the thermal entropy  
23 of a black hole is proportional to the event horizon of the black hole [1,2] has stimulated varies  
24 modern idea of theoretical physics, including the famous holographic principle.

25 OPE block [3,4], on the other hand, is a relatively new topic in conformal field theory, though  
26 it has been noticed at the early stages of conformal field theory [5,6]. The operator product  
27 expansion of two primary operators is equivalent to a summation of OPE blocks with corre-  
28 sponding three point function coefficients. It is a smeared operator which is generated from  
29 the so-called (quasi-)primary operator.

30 Modular Hamiltonian, the logarithm of the reduced density matrix [7], plays a central role  
31 in the context of geometric entanglement entropy [8–11]. Entanglement entropy is a von  
32 Neumann entropy generated from reduced density matrix of a subregion of spacetime. An  
33 intriguing fact of entanglement entropy is that it obeys area law in the leading order, though  
34 one should introduce a cutoff to secure the divergent behaviour. Its connection to gravity  
35 has been established by the work of Ryu and Takayanagi [12], in which they proposed that  
36 the entanglement entropy of a CFT is equal to the area of a minimal surface in the bulk AdS  
37 spacetime.

38 Modular Hamiltonian is a special OPE block generated by the stress energy-momentum tensor  
39 for a ball region. This leads to the conjecture that OPE block may be related to area law as  
40 modular Hamiltonian. Indeed, in a series of papers [13,14,16,17], we have shown that the  
41 quantity which satisfies area law is the type- $(m)$  connected correlation function (CCF). More  
42 explicitly, the leading term of the type- $(m)$  CCF is proportional to the area of the boundary of  
43 the ball. In the subleading terms, we find a logarithmic divergence with degree  $q$ . The degree  
44  $q$  is a natural number which is no larger than 2 in general dimensions. The coefficient  $p_q$  for  
45 the logarithmic term with degree  $q$  is cutoff independent. We establish a relationship between  
46  $p_q$  and the type- $(m-1, 1)$  CCF of OPE blocks for two balls which are far away to each other.  
47 The coefficient  $p_q$  obeys a cyclic identity which is independent of the order of the operators.

48 This paper is organised as follows. In section 2, we will introduce some basic concepts and  
49 conventions used in this paper. Section 3 is devoted to the study of the new area law which is  
50 related to the OPE blocks. Varies generalizations have been given in section 4. We conclude  
51 in section 5 with a number of general open problems that deserve, in our opinion, more work.

## 52 **2 Setup**

53 In this section, we introduce some basic concepts and conventions used in this paper.

54 **2.1 Area law**

55 In any continuous quantum field theory (QFT), physical degrees exist at each point  $(t, x^i), i = 1, \dots, d-1$   
 56 of spacetime  $M$ . At each time slice  $t = t_0$ , the data on the Cauchy surface  $\Sigma$  determines the  
 57 evolution of the fields. One can divide the surface  $\Sigma$  into a spacelike subregion  $A$  and its com-  
 58 plement  $\bar{A}$ ,  $\Sigma = A \cup \bar{A}$ . The boundary  $\partial A$  is a codimension 2 surface whose area is  $\mathcal{A}$ . The  
 59 causal development of  $A$  is denoted by  $\mathcal{D}(A)$ . The physical data on  $A$  can only determine the  
 60 evolution of the fields in  $\mathcal{D}(A)$ . The causal development  $\mathcal{D}(A)$  is an independent subsystem of  
 61 the original spacetime  $M$ . Operators in this subsystem are collected to form an algebra  $\mathfrak{a}(A)$ .  
 62 Assume the QFT in the spacetime  $M$  is described by a density matrix  $\rho$ , then by integrating  
 63 out the degree of freedom in the complement of  $\bar{A}$ , one achieves a reduced density matrix  $\rho_A$

$$\rho_A = \text{tr}_{\bar{A}} \rho. \tag{2.1}$$

64 The reduced density matrix  $\rho_A$  is a special operator in  $\mathfrak{a}(A)$  since it describes the subsystem  
 65  $\mathcal{D}(A)$  effectively. A general quantity  $Q(A)$  in  $\mathfrak{a}(A)$  is said to obey area law if its leading term is  
 66 proportional to the area of the boundary  $\partial A$ ,

$$Q(A) \propto \mathcal{A} + \dots. \tag{2.2}$$

67 One typical example is the black hole entropy in Einstein gravity. The black hole entropy is  
 68 proportional to the area of its event horizon,

$$S_{bh} = \frac{\mathcal{A}}{4G} \tag{2.3}$$

69 where  $G$  is the Newton constant. At the loop level, black hole entropy requires logarithmic cor-  
 70 rections [18–23]. Usually, the logarithmic correction is in the form  $C \log \mathcal{A}$  where the constant  
 71  $C$  may encode useful information of the black hole.

72 Sometimes the area law is divergent, one typical example is the geometric entanglement en-  
 73 tropy

$$S_A = -\text{tr}_A \rho_A \log \rho_A. \tag{2.4}$$

74 In this case, one should insert a cutoff  $\epsilon > 0$ ,

$$S_A = \gamma \frac{\mathcal{A}}{\epsilon^{d-2}} + \dots. \tag{2.5}$$

75 In the subleading terms, there may be a logarithmic term whose coefficient is independent of  
 76 the cutoff,

$$S_A = \gamma \frac{R^{d-2}}{\epsilon^{d-2}} + \dots + p \log \frac{R}{\epsilon} + \dots \tag{2.6}$$

77 where the parameter  $R$  is the characteristic length of the region  $A$ .

78 In this report, we will present a quantity  $Q(A)$  which has a slightly different logarithmic be-  
 79 haviour

$$Q(A) = \gamma \frac{R^{d-2}}{\epsilon^{d-2}} + \dots + p_q \log^q \frac{R}{\epsilon} + \dots. \tag{2.7}$$

80 The maximum power  $q$  of the logarithmic terms is a natural number. We will call it the degree  
 81 of the quantity  $Q(A)$ . The coefficient  $p_q$  is cutoff independent and encodes useful information  
 82 of the theory. In the special case that the subregion  $A$  is a ball,  $R$  could be chosen as its radius.  
 83 The subregion  $A$  and its causal development  $\mathcal{D}(A)$  are in one-to-one correspondence, we will  
 84 not distinguish them in the following.

85 In two dimensions, there is no polynomial terms of  $\frac{R}{\epsilon}$ , the modified “area law” is

$$Q(A) = p_q \log^q \frac{R}{\epsilon} + \dots. \tag{2.8}$$

86 **2.2 OPE block**

87 In any d dimensional CFT, operators are classified into (quasi-)primary operators  $\mathcal{O}$  and their  
 88 descendants  $\partial_\mu \partial_\nu \dots \mathcal{O}$ . A general primary operator is characterized by two quantum num-  
 89 bers, conformal weight  $\Delta$  and  $so(d-1)$  spin  $J_{ij}$  with magnitude  $J$ . Under a global conformal  
 90 transformation  $x \rightarrow x'$ , a primary spin 0 operator transforms as

$$\mathcal{O}(x) \rightarrow \left| \frac{\partial x'}{\partial x} \right|^{-\Delta/d} \mathcal{O}(x). \tag{2.9}$$

91 where  $|\partial x'/\partial x|$  is the Jacobian of the conformal transformation of the coordinates,  $\Delta$  is the  
 92 conformal weight of the primary operator. Operator product expansion(OPE) of two separated  
 93 primary scalar operators  $\mathcal{O}_i(x_1)\mathcal{O}_j(x_2)$  is to expand their product in a local orthogonal and  
 94 complete basis around a suitable point

$$\mathcal{O}_i(x_1)\mathcal{O}_j(x_2) = \sum_k C_{ijk} |x_{12}|^{\Delta_k - \Delta_i - \Delta_j} (\mathcal{O}_k(x_2) + \dots), \tag{2.10}$$

95 where  $\dots$  are descendants of the primary operator  $\mathcal{O}_k$ . Its form is fixed by global conformal  
 96 symmetry, therefore it just contains kinematic information of the CFT. The summation is over  
 97 all possible primary operators of the CFT. Here we expand the product around the point  $x_2$ .  
 98 The distance of any two points  $x_i, x_j$  is written as  $|x_{ij}|$ . The constant  $C_{ijk}$  is called the OPE  
 99 coefficient which is related to the three point function of primary operators

$$\langle \mathcal{O}_i(x_1)\mathcal{O}_j(x_2)\mathcal{O}_k(x_3) \rangle = \frac{C_{ijk}}{|x_{12}|^{\Delta_{12,3}} |x_{23}|^{\Delta_{23,1}} |x_{13}|^{\Delta_{13,2}}}, \quad \Delta_{ij,k} = \Delta_i + \Delta_j - \Delta_k. \tag{2.11}$$

100 They are the only dynamical parameters in the CFT. The constants  $\Delta_i, \Delta_j, \Delta_k$  are conformal  
 101 weights of the corresponding primary operators. By collecting all kinematic terms in the sum-  
 102 mation, we can rewrite the OPE (2.10) as

$$\mathcal{O}_i(x_1)\mathcal{O}_j(x_2) = |x_{12}|^{-\Delta_i - \Delta_j} \sum_k C_{ijk} Q_k^{ij}(x_1, x_2). \tag{2.12}$$

103 The objects  $Q_k^{ij}(x_1, x_2)$  are called OPE blocks [3,5,6]. They are non-local operators in the CFT  
 104 and depend on the position  $x_1$  and  $x_2$  of the external operators. The upper index  $i$  and  $j$  show  
 105 that it also depends on the quantum number of the external operators  $\mathcal{O}_i$  and  $\mathcal{O}_j$ . It is easy to  
 106 see that OPE block has dimension zero. Under a global conformal transformation  $x \rightarrow x'$ , an  
 107 OPE block  $Q_k^{ij}(x_1, x_2)$  will transform as

$$Q_k^{ij}(x_1, x_2) \rightarrow f(x'_1, x'_2) Q_k^{ij}(x'_1, x'_2). \tag{2.13}$$

108 The explicit form of  $f(x'_1, x'_2)$  is not important in this work. When the two external operators  
 109 are the same, we have  $f(x'_1, x'_2) = 1$  and OPE block will be invariant under the global con-  
 110 formal transformation. One can also show that the OPE block is independent of the external  
 111 operator in this special case. Due to this reason, we relabel such kind of OPE block as

$$Q_A[\mathcal{O}_k] = Q_k^{ii}(x_1, x_2). \tag{2.14}$$

112 The subscript  $A$  denotes the region determined by the two points  $x_1$  and  $x_2$  where the two  
 113 external operators insert into. The operator in the square bracket reflects the fact that OPE  
 114 block is generated by a primary operator  $\mathcal{O}_k$ . We omit the information of  $i$  since the OPE block  
 115 is insensitive to the external operators in this case. We will classify the primary operators

116  $\mathcal{O}_k$  into conserved currents  $\mathcal{J}$  and non-conserved operators  $\mathcal{O}$ . A general symmetric traceless  
 117 primary operator obeys the following unitary bound [24]

$$\begin{cases} \Delta \geq J + d - 2, & J \geq 1, \\ \Delta \geq \frac{d-2}{2}, & J = 0. \end{cases}$$

118 A conserved current  $\mathcal{J}$  with spin  $J(J \geq 1)$  will satisfy  $\Delta = J + d - 2$ . All other primary  
 119 operators are non-conserved operators. Correspondingly, the OPE block (2.14) generated by  
 120 a conserved current  $\mathcal{J}$  will be called a type-J OPE block. On the other hand, the OPE block  
 121 (2.14) generated by a non-conserved operator  $\mathcal{O}$  will be called a type-O OPE block.

122 When two operators are time-like separated, the region  $A$  is a causal diamond. The two oper-  
 123 ators are at the sharp corner of the diamond  $A$ . We can use the conformal transformation to  
 124 fix

$$x_1 = (1, \vec{x}_0), \quad x_2 = (-1, \vec{x}_0), \quad (2.15)$$

125 then the causal diamond  $A$  intersects  $t = 0$  slice with a unit ball which we will also denote it  
 126 as  $A$

$$A = \{(0, \vec{x}) | (\vec{x} - \vec{x}_0)^2 \leq 1\}. \quad (2.16)$$

127 The center of the ball is  $\vec{x}_0$ . The boundary of the ball  $A$  is a unit sphere  $\partial A$ . In the context of  
 128 geometric entanglement entropy, the surface  $\partial A$  is an entanglement surface which separates  
 129 the ball  $A$  and its complement. The leading term of entanglement entropy is proportional to  
 130 the area of the surface  $\partial A$  in general higher dimensions ( $d > 2$ ). In two dimensions, the  
 131 entanglement entropy is logarithmically divergent with the logarithmic degree  $q = 1$ . There  
 132 is a conformal Killing vector  $K$  which preserves the diamond  $A$ ,

$$K^\mu = \frac{1}{2}(1 - (\vec{x} - \vec{x}_A)^2 - t^2, -2t\vec{x}). \quad (2.17)$$

133 The conformal Killing vector  $K$  is null on the boundary of the diamond  $A$ . It generates a  
 134 modular flow of the diamond  $A$ . A type-O OPE block corresponds to point pair (2.15) or unit  
 135 ball  $A$  (2.16) is [4]

$$Q_A[\mathcal{O}_{\mu_1 \dots \mu_J}] = c_{\mathcal{O}_{\mu_1 \dots \mu_J}} \int_{\mathcal{D}(A)} d^d x K^{\mu_1} \dots K^{\mu_J} |K|^{\Delta-d-J} \mathcal{O}_{\mu_1 \dots \mu_J}, \quad (2.18)$$

136 where the primary operator  $\mathcal{O}_{\mu_1 \dots \mu_J}$  is non-conserved

$$\partial^{\mu_1} \mathcal{O}_{\mu_1 \dots \mu_J} \neq 0. \quad (2.19)$$

137 It has dimension  $\Delta$  and spin  $J$ . When the operator is a conserved current

$$\partial^{\mu_1} \mathcal{J}_{\mu_1 \dots \mu_J} = 0, \quad (2.20)$$

138 the corresponding type-J OPE block is

$$Q_A[\mathcal{J}_{\mu_1 \dots \mu_J}] = c_{\mathcal{J}_{\mu_1 \dots \mu_J}} \int_A d^{d-1} \vec{x} (K^0)^{J-1} \mathcal{J}_{0 \dots 0}. \quad (2.21)$$

139 It can be obtained from (2.18) by using the conservation law (2.20) and reducing it to a lower  
 140  $d - 1$  dimensional integral. The coefficient  $c_{\mathcal{J}_{\mu_1 \dots \mu_J}}$  is also redefined at the same time. In (2.18)  
 141 and (2.21), the coefficients  $c_{\mathcal{O}_{\mu_1 \dots \mu_J}}$  and  $c_{\mathcal{J}_{\mu_1 \dots \mu_J}}$  are free parameters, we set them to be 1.

142 **2.3 Modular Hamiltonian and area law**

143 A very special type-J OPE block is the modular Hamiltonian [7, 25] of the ball  $A$ ,

$$H_A = 2\pi \int_A d^{d-1} \vec{x} K^0 T_{00} = 2\pi \int_A d^{d-1} \vec{x} \frac{1 - (\vec{x} - \vec{x}_0)^2}{2} T_{00}(0, \vec{x}). \quad (2.22)$$

144 Modular Hamiltonian is the logarithm of the reduced density matrix  $\rho_A$

$$H_A = -\log \rho_A. \quad (2.23)$$

145 It plays a central role in the context of entanglement entropy,

$$S_A = -\text{tr}_A \rho_A \log \rho_A = \text{tr}_A e^{-H_A} H_A. \quad (2.24)$$

146 More generally, Rényi entanglement entropy

$$S_A^{(n)} = \frac{1}{1-n} \log \text{tr}_A \rho_A^n \quad (2.25)$$

147 has been shown to satisfy an area law generally

$$S_A^{(n)} = \gamma \frac{\mathcal{A}}{\epsilon^{d-2}} + \dots, \quad (2.26)$$

148 where  $\mathcal{A}$  is the area of the entanglement surface  $\partial A$  and  $\epsilon$  is a UV cutoff. The constant  $\gamma$  is  
 149 cutoff dependent. The subleading terms  $\dots$  contain a logarithmic term with degree  $q = 1$  in  
 150 even dimensions

$$S_A^{(n)} = \gamma \frac{\mathcal{A}}{\epsilon^{d-2}} + \dots + p_1(n) \log \frac{R}{\epsilon} + \dots, \quad (2.27)$$

151 where we have inserted back the radius  $R = 1$ . The area  $\mathcal{A}$  is related to the radius  $R$  through  
 152 the power law

$$\mathcal{A} \sim R^{d-2}. \quad (2.28)$$

153 The coefficient  $p_1(n)$  encodes useful information of the CFT. The relation between modular  
 154 Hamiltonian and area law motivates the conjecture that OPE block maybe related to area  
 155 law in a suitable way. We will give the framework to discuss this problem in the following  
 156 subsection.

157 **2.4 Deformed reduced density matrix and connected correlation function**

158 Given a primary operator  $\mathcal{O}$  in a ball  $A$ , one can always define a corresponding OPE block  
 159  $Q_A[\mathcal{O}]$ . We construct an exponential operator formally [14]

$$\rho_A = e^{-\mu Q_A} \quad (2.29)$$

160 which is still in the subregion  $A$ . The constant  $\mu$  is free. Operators of the form (2.29) is called  
 161 deformed reduced density matrix. Note we use the same symbol  $\rho_A$  to label deformed reduced  
 162 density matrix. Recall that the modular Hamiltonian is a special OPE block, if one replaces the  
 163 OPE block by the modular Hamiltonian (2.29) and set  $\mu = 1$ , the deformed reduced density  
 164 matrix becomes the reduced density matrix exactly. We can relax the definition, namely,  $Q_A$  in  
 165 (2.29) could be a linear superposition of several OPE blocks. Note our definition of deformed  
 166 reduced density matrix is a direction extension of the generalized reduced density matrix in  
 167 the context of the so-called charged Rényi entropy [15]. In that work,  $Q_A$  is a charge which  
 168 is generated by a  $U(1)$  current. The corresponding charged Rényi entropy is holographically

169 dual to the thermal entropy of a charged black hole with hyperbolic horizon. However, in our  
 170 definition,  $Q_A$  is just a general OPE block or their linear superposition. As a naive generalization  
 171 of Rényi entanglement entropy, we construct the logarithm of the vacuum expectation value  
 172 of the deformed reduced density matrix,

$$T_A(\mu) = \log\langle\rho_A\rangle = \log\langle e^{-\mu Q_A}\rangle. \tag{2.30}$$

173 When  $Q_A$  is modular Hamiltonian, the above quantity is related to the Rényi entropy for the  
 174 vacuum state.

175 However, a direct computation of  $T_A(\mu)$  is hard in general. A much more severe problem is  
 176 that OPE block has no lower bound in general, therefore the definition is not valid for general  
 177 OPE blocks. To solve this problem, we observe that  $T_A(\mu)$  could be expanded for small  $\mu$ ,

$$T_A(\mu) = \sum_{m=1}^{\infty} \frac{(-\mu)^m}{m!} \langle Q_A^m \rangle_c. \tag{2.31}$$

178 The Taylor expansion coefficient

$$\langle Q_A^m \rangle_c = (-1)^m \frac{\partial^m}{\partial \mu^m} T_A(\mu)|_{\mu \rightarrow 0} \tag{2.32}$$

179 is called Type-(m) connected correlation function (CCF) of the OPE block  $Q_A$ . For each definite  
 180  $m$ , one can always calculate the corresponding CCF without knowing  $T_A(\mu)$ . The first few CCFs  
 181 are

$$\begin{aligned} \langle Q_A^2 \rangle_c &= \langle Q_A^2 \rangle - \langle Q_A \rangle^2, \\ \langle Q_A^3 \rangle_c &= \langle Q_A^3 \rangle - 3\langle Q_A^2 \rangle \langle Q_A \rangle + 2\langle Q_A \rangle^3. \end{aligned} \tag{2.33}$$

182 Using CCF, there is no issue of lower bound of the OPE block. As an application of the concept  
 183 of CCF, we choose the OPE block as the modular Hamiltonian, then it is easy to show that  
 184 CCF of modular Hamiltonian  $H_A$  satisfies area law with logarithmic degree  $q = 1$  in even  
 185 dimensions,

$$\langle H_A^m \rangle_c = \tilde{\gamma} \frac{\mathcal{A}}{\epsilon^{d-2}} + \dots + \tilde{p}_1^{(m)} \log \frac{R}{\epsilon} + \dots, \quad m \geq 1. \tag{2.34}$$

186 The coefficient  $\tilde{p}_1^{(m)}$  is determined from  $p_1(n)$  by

$$\tilde{p}_1^{(m)} = (-1)^m \partial_n^m (1-n)p_1(n)|_{n \rightarrow 1}. \tag{2.35}$$

187 There could be multiple spacelike-separated balls  $A_1, A_2, \dots$ , each region has associate OPE  
 188 block  $Q_{A_i}$ . We insert  $m_i$  OPE blocks into region  $A_i$ , then we can define the corresponding  
 189 type-Y CCF

$$\langle Q_{A_1}^{m_1} Q_{A_2}^{m_2} \dots \rangle_c \tag{2.36}$$

190 where the Young diagram  $Y$  is

$$Y = (m_1, m_2, \dots), \quad m_1 \geq m_2 \geq \dots \geq 1. \tag{2.37}$$

191 The generator of all type-Y CCFs is

$$T_{\cup A_i}(\mu_1, \mu_2, \dots) = \log \frac{\langle e^{-\sum_i \mu_i Q_{A_i}} \rangle}{\prod_i \langle e^{-\mu_i Q_{A_i}} \rangle}. \tag{2.38}$$

192 When there are only two balls  $A$  and  $B$ , the generator is

$$T_{A \cup B}(\mu_1, \mu_2) = \log \frac{\langle e^{-\mu_1 Q_A - \mu_2 Q_B} \rangle}{\langle e^{-\mu_1 Q_A} \rangle \langle e^{-\mu_2 Q_B} \rangle} = \sum_{m_1 \geq 1, m_2 \geq 1} \frac{(-1)^{m_1+m_2} \mu_1^{m_1} \mu_2^{m_2}}{m_1! m_2!} \langle Q_A^{m_1} Q_B^{m_2} \rangle_c. \quad (2.39)$$

193 We parameterize  $A$  and  $B$  as

$$A = \{(0, \vec{x}) | (\vec{x} - \vec{x}_0)^2 \leq 1\}, \quad B = \{(0, \vec{x}) | \vec{x} \leq R'^2\}. \quad (2.40)$$

194 There is only one cross ratio

$$\xi = \frac{4R'}{x_0^2 - (1 - R')^2}. \quad (2.41)$$

195 When the two regions  $A$  and  $B$  are spacelike-separated,  $|x_0| > 1 + R'$ , the cross ratio is between  
196 0 and 1,

$$0 < \xi < 1. \quad (2.42)$$

197 In some cases, it is more convenient to use an equivalent cross ratio

$$\eta = \frac{\xi}{1 - \xi} = \frac{4R'}{x_0^2 - (1 + R')^2}. \quad (2.43)$$

198 For spacelike-separated regions  $A$  and  $B$ , the range of the cross ratio  $\eta$  is

$$0 < \eta < \infty. \quad (2.44)$$

199 Since the OPE block  $Q_A[\mathcal{O}]$  is invariant under conformal transformation, any type- $(m_1, m_2)$   
200 CCF should be a function of cross ratio  $\xi$  or  $\eta$ . Actually the OPE block is an eigenvector of the  
201 conformal Casimir

$$[L^2, Q_A[\mathcal{O}]] = C_{\Delta, J} Q_A[\mathcal{O}] \quad (2.45)$$

202 where  $L^2$  is the Casimir operator of the global conformal group. The eigenvalue  $C_{\Delta, J}$  is

$$C_{\Delta, J} = -\Delta(\Delta - d) - J(J + d - 2). \quad (2.46)$$

203 Therefore, any type- $(m - 1, 1)$  CCF should be a conformal block

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_{m-1}] Q_B[\mathcal{O}_m] \rangle_c = D^{(d)}[\mathcal{O}_1, \cdots, \mathcal{O}_m] G_{\Delta_m, J_m}^{(d)}(\xi). \quad (2.47)$$

204 The subscript  $\Delta_m, J_m$  are the conformal weight and spin of the primary operator  $\mathcal{O}_m$ . The index  
205  $(d)$  is used to label the dimension of spacetime. The conformal block can be constructed ex-  
206 plicitly in even dimensions [26, 27]. In this paper, we just need the diagonal limit of conformal  
207 block [28]. Any type- $(m_1, m_2)$  CCF with  $m_1 \geq m_2 \geq 2$  is not a conformal block .

### 208 3 Area law

209 We conjecture that the type- $(m)$  CCF of OPE blocks obeys the following area law

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_m] \rangle_c = \gamma \frac{R^{d-2}}{\epsilon^{d-2}} + \cdots + p_q \log^q \frac{R}{\epsilon} + \cdots. \quad (3.1)$$

210 The leading term is proportional to the area of the boundary  $\partial A$ . We inserted the radius  $R = 1$   
211 into the formula to balance the dimension. The small positive constant  $\epsilon$  is the UV cutoff which  
212 is roughly the distance from the cutoff to the boundary  $\partial A$ . The constant  $\gamma$  depends on the



213 choice of the cutoff and the method of regularization, we will not be interested in its explicit  
 214 value. The  $\dots$  terms are subleading and cutoff dependent. Therefore we omit their forms.  
 215 The degree  $q$  characterizes the maximal power of the logarithmic terms. The coefficient  $p_q$  is  
 216 invariant under the rescaling of the cutoff, therefore it encodes detail universal information of  
 217 the theory. When all the OPE blocks are equal to the modular Hamiltonian, the degree  $q = 1$   
 218 for even dimensions according to (2.34). However, as we will see,  $q$  is not necessary equal to  
 219 1 in general. To distinguish different type- $(m)$  CCFs in different dimensions, we write the area  
 220 law (3.1) more explicitly as

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_m] \rangle_c = \gamma[\mathcal{O}_1, \dots, \mathcal{O}_m] \frac{R^{d-2}}{\epsilon^{d-2}} + \cdots + p_q^{(d)}[\mathcal{O}_1, \dots, \mathcal{O}_m] \log^q \frac{R}{\epsilon} + \cdots \quad (3.2)$$

### 221 3.1 Continuation

222 The two formulas (2.47) and (3.2) are actually related to each other through an analytic  
 223 continuation. We use the example of the two dimensional modular Hamiltonian to illustrate  
 224 this relation. For any  $\text{CFT}_2$ , the modular Hamiltonian can be decomposed into the holomorphic  
 225 and anti-holomorphic part, we focus on the holomorphic part

$$H_A = - \int_{-1}^1 dz \frac{1-z^2}{2} T(z+x_0) + c. \quad (3.3)$$

226 The constant  $c$  can be fixed by the normalization condition

$$\text{tr}_A \rho_A = \text{tr}_A e^{-H_A} = 1. \quad (3.4)$$

227 Its value doesn't affect the type-Y CCF with any  $\sum_i m_i \geq 2$ . We also used the convention  
 228  $T(z) = -2\pi T_{zz}$  where the subscript  $z$  is the holomorphic coordinate  $z = t + x$ . The radius of  
 229 the interval  $A$  is 1, we have shifted the variable  $z$  such that the dependence of the center  $x_0$  is  
 230 in the stress tensor. The modular Hamiltonian of region  $B$  can be obtained by setting  $x_0 = 0$   
 231 and restoring the radius  $R'$ . The type- $(m-1, 1)$  CCF of the modular Hamiltonian is

$$\langle H_A^{m-1} H_B \rangle_c = D^{(2)}[T_{\mu_1 \nu_1}, \dots, T_{\mu_m \nu_m}] G_2^{(2)}(\eta). \quad (3.5)$$

232 The two dimensional conformal block for a chiral operator can be labeled by the conformal  
 233 weight  $h$  of the operator

$$G_h^{(2)}(\eta) = (-\eta)^h {}_2F_1(h, h, 2h, -\eta). \quad (3.6)$$

234 We can move the interval  $A$  to  $B$  such that they coincide. In this limit, any type- $(m-1, 1)$  CCF  
 235 should approach a type- $(m)$  CCF. This is equivalent to set  $\eta \rightarrow -1$ . We can set  $x_0 \rightarrow 0$  and  
 236 then take the limit  $R' \rightarrow 1$ ,

$$x_A \rightarrow 0, \quad R' = 1 - \epsilon, \quad \epsilon \rightarrow 0. \quad (3.7)$$

237 The cross ratio  $\xi \rightarrow -\infty$  or  $\eta \rightarrow -1$  by

$$\xi = -\frac{4(1-\epsilon)}{\epsilon^2} \approx -\frac{4}{\epsilon^2}, \quad \eta = -\frac{4(1-\epsilon)}{(2-\epsilon)^2} \approx -1 + \frac{\epsilon^2}{4}. \quad (3.8)$$

238 On the right hand side of (3.5), we find a logarithmic divergent term in this limit

$$G_2^{(2)}(\eta) = 12 \log \frac{2}{\epsilon} + \cdots = 12 \log \frac{R}{\epsilon} + \cdots \quad (3.9)$$

239 The left hand side of (3.5) approaches type-( $m$ ) CCF, therefore

$$\langle H_A^m \rangle_c = 12D^{(2)}[T_{\mu_1\nu_1}, \dots, T_{\mu_m\nu_m}] \log \frac{R}{\epsilon} + \dots \quad (3.10)$$

240 We read out the cutoff independent coefficient

$$p_1^{(2)}[T_{\mu_1\nu_1}, \dots, T_{\mu_m\nu_m}] = 12D^{(2)}[T_{\mu_1\nu_1}, \dots, T_{\mu_m\nu_m}]. \quad (3.11)$$

241 The relation (3.11) is a typical UV/IR relation for the modular Hamiltonian. The left hand side  
 242 is the universal coefficient for  $B$  and  $A$  coincides (UV). On the right hand side, the  $D$  coefficient  
 243 characterizes the leading order behaviour of CCF when  $B$  and  $A$  are far away to each other (IR).  
 244 They provide equivalent information of the CFT since the constant 12 is completely fixed by  
 245 conformal symmetry. The continuation of the conformal block can be generalized to higher  
 246 dimensions. For example, in four dimensions, the conformal block associated with stress tensor  
 247 becomes divergent as  $A$  approaches  $B$ ,

$$G_{4,2}^{(4)} \approx \tilde{\gamma} \frac{R^2}{\epsilon^2} + \dots - 120 \log \frac{R}{\epsilon} + \dots \quad (3.12)$$

248 The leading term is exactly proportional to the area of the boundary and the logarithmic di-  
 249 vergent term also appears in the subleading terms. We can read out the type-( $m$ ) CCF of the  
 250 modular Hamiltonian in four dimensions

$$\langle H_A^m \rangle_c = \gamma \frac{R^2}{\epsilon^2} + \dots + p_1^{(4)}[T_{\mu_1\nu_1}, \dots, T_{\mu_m\nu_m}] \log \frac{R}{\epsilon} + \dots \quad (3.13)$$

251 with

$$p_1^{(4)}[T_{\mu_1\nu_1}, \dots, T_{\mu_m\nu_m}] = -120D^{(4)}[T_{\mu_1\nu_1}, \dots, T_{\mu_m\nu_m}]. \quad (3.14)$$

252 Note we obtain the area law and the logarithmic behaviour of the type-( $m$ ) CCF of the modular  
 253 Hamiltonian without using any knowledge of Rényi entanglement entropy. The method of  
 254 analytic continuation can be applied to general dimensions and OPE blocks. A conformal  
 255 block  $G_{\Delta,J}^{(d)}(\xi)$  obeys area law in the limit  $\xi \rightarrow -\infty$  in even dimensions. It has degree  $q = 1$   
 256 only for  $\Delta = J + d - 2$ ,

$$G_{\Delta,J}^{(d)}(\xi) = \tilde{\gamma} \frac{R^{d-2}}{\epsilon^{d-2}} + \dots + E^{(d)}[\Delta, J] \log \frac{R}{\epsilon} + \dots, \quad \xi \rightarrow -\infty. \quad (3.15)$$

257 This means that type-( $m$ ) CCF of type-J OPE blocks may always obey area law with degree  
 258  $q = 1$ , the cutoff independent coefficient is

$$p_q^{(d)}[\mathcal{O}_1, \dots, \mathcal{O}_m] = E^{(d)}[\mathcal{O}_m] \times D^{(d)}[\mathcal{O}_1, \dots, \mathcal{O}_m]. \quad (3.16)$$

259 We have replaced the quantum numbers in E function by the corresponding primary opera-  
 260 tor. For non-conserved operators, the conformal block  $G_{\Delta,J}^{(d)}$  also obeys area law in the limit  
 261  $\xi \rightarrow -\infty$  in even dimension, though it has degree  $q = 2$

$$G_{\Delta,J}^{(d)}(\xi) = \tilde{\gamma} \frac{R^{d-2}}{\epsilon^{d-2}} + \dots + E^{(d)}[\Delta, J] \log^2 \log \frac{R}{\epsilon} + \dots, \quad \xi \rightarrow -\infty. \quad (3.17)$$

262 Therefore, type-( $m$ ) CCF of type-O OPE blocks obeys area law with degree  $q = 2$ . We can  
 263 obtain similar UV/IR relations as (3.16). In odd dimensions, the story is the same. The degree  
 264  $q$  is 0 for type-( $m$ ) CCF of type-J OPE blocks and 1 for type-O OPE blocks.

265 **3.2 Kinematic information**

266 The function  $E^{(d)}[\mathcal{O}]$  is completely fixed by conformal symmetry. It can be obtained by reading  
 267 out the coefficient of the logarithmic term with degree  $q$ . For each fixed quantum number  $\Delta$   
 268 and  $J$ , there is a unique number  $E^{(d)}[\mathcal{O}]$ . For any type-J OPE block in two dimensions, the  
 269 primary operator  $\mathcal{O}$  has dimension  $\Delta = J = h$ . The conformal block (3.6) has degree  $q = 1$  in  
 270 the limit  $\eta \rightarrow -1$ . The function  $E^{(2)}[\mathcal{O}]$  is

$$E^{(2)}[\mathcal{O}] = \frac{2\Gamma(2h)}{\Gamma(h)^2}, \quad \Delta = J = h. \tag{3.18}$$

271 For type-O OPE block, the primary operator  $\mathcal{O}$  has dimension  $\Delta = h + \bar{h}$  and spin  $J = h - \bar{h}$ .  
 272 The conformal block has degree  $q = 2$  in the limit  $\eta \rightarrow -1$ . The function  $E^{(2)}[\mathcal{O}]$  is

$$E^{(2)}[\mathcal{O}] = \begin{cases} \frac{2^{4h}\Gamma(h+\frac{1}{2})^2}{\pi\Gamma(h)^2} & J = 0, h > 0 \\ -\frac{4^{2h-1}\Gamma(h-\frac{1}{2})\Gamma(h+\frac{1}{2})}{\pi\Gamma(h-1)\Gamma(h)} & J = 1, h > 1 \\ \frac{4^{2h-3}(h-2)(h-1)(2h-3)(2h-1)\Gamma(h-\frac{3}{2})^2}{\pi\Gamma(h)^2} & J = 2, h > 2 \\ \dots & \dots \end{cases} \tag{3.19}$$

273 In four dimensions, we also find

$$E^{(4)}[\mathcal{O}] = \begin{cases} 12 & \Delta = 3, J = 1 \\ -120 & \Delta = 4, J = 2 \\ 840 & \Delta = 5, J = 3 \\ \dots & \dots \end{cases} \tag{3.20}$$

274 for conserved currents and

$$E^{(4)}[\mathcal{O}] = \begin{cases} -\frac{2^{2\Delta-1}\Gamma(\frac{\Delta-1}{2})\Gamma(\frac{\Delta+1}{2})}{\pi\Gamma(\frac{\Delta-2}{2})^2} & \Delta > 1, J = 0, \\ \frac{2^{2\Delta-1}\Gamma(\frac{\Delta}{2})\Gamma(\frac{\Delta+2}{2})}{\pi\Gamma(\frac{\Delta-3}{2})\Gamma(\frac{\Delta+1}{2})} & \Delta > 3, J = 1, \\ -\frac{4^{\Delta-1}(\Delta-2)\Gamma(\frac{\Delta-3}{2})\Gamma(\frac{\Delta+3}{2})}{\pi\Gamma(\frac{\Delta-4}{2})\Gamma(\frac{\Delta+2}{2})} & \Delta > 4, J = 2, \\ \dots & \dots \end{cases} \tag{3.21}$$

275 for non-conserved operators. In three dimensions, we find

$$E^{(3)}[\mathcal{O}] = \begin{cases} -\frac{2^{2\Delta-1}(\Delta-1)\Gamma(\Delta-\frac{1}{2})}{\sqrt{\pi}\Gamma(\Delta-1)} & \Delta > \frac{1}{2}, J = 0. \\ \frac{2^{\Delta+1}\Delta\Gamma(\Delta-\frac{1}{2})}{\Gamma(\frac{\Delta-2}{2})\Gamma(\frac{\Delta+1}{2})} & \Delta > 2, J = 1, \\ -\frac{2^{2\Delta-1}(\Delta^2-1)\Gamma(\Delta-\frac{1}{2})}{\sqrt{\pi}(\Delta-2)^2\Delta\Gamma(\Delta-3)} & \Delta > 3, J = 2, \\ \dots & \dots \end{cases} \tag{3.22}$$

276 for non-conserved operators. Note for conserved currents in odd dimensions, the function  
 277  $E^{(3)}[\mathcal{O}]$  may depend on explicit choice of the cutoff. For example, a transformation  $\epsilon \rightarrow \epsilon(1+a\epsilon)$   
 278 may shift its value. This is because the degree is 0, there is no logarithmic divergence at all.

279 **3.3 UV/IR relation**

280 The UV/IR relation (3.16) relates type- $(m)$  CCF to type- $(m - 1, 1)$  CCF. This relation may  
 281 simplify computation in many cases. To see this point, let's compute the following type-(2)

282 CCF in two dimensions

$$\begin{aligned}
 \langle Q_A[\mathcal{O}]^2 \rangle_c &= \int_{-1}^1 dz_1 \int_{-1}^1 dz_2 \frac{(1-z_1^2)^{h-1}(1-z_2^2)^{h-1}}{(z_1-z_2)^{2h}} \\
 &= \frac{(-1)^{-h} \sqrt{\pi} \Gamma(h)}{\Gamma(h+\frac{1}{2})} \int_{-1}^1 dz_1 \frac{1}{1-z_1^2} \\
 &= \frac{(-1)^{-h} \sqrt{\pi} \Gamma(h)}{\Gamma(h+\frac{1}{2})} \log \frac{2}{\epsilon}.
 \end{aligned} \tag{3.23}$$

283 This is a double integral with poles at  $z_1 = z_2$ . We regularize the integral by ignoring these  
 284 poles at the second step. At the last step, we insert a UV cutoff to regularize the integral.  
 285 However, using UV/IR relation, one just need to fix the coefficient  $D$  which is related to the  
 286 large distance behaviour of the type-(1, 1) CCF,

$$\langle Q_A[\mathcal{O}]Q_B[\mathcal{O}] \rangle_c = \int_{-1}^1 dz_1 \int_{-1}^1 dz_2 \frac{(1-z_1^2)^{h-1}(1-z_2^2)^{h-1}}{(z_1-z_2+x_0)^{2h}}. \tag{3.24}$$

287 In the large distance limit,  $x_0 \rightarrow \infty$ , the integral becomes simpler

$$\begin{aligned}
 \langle Q_A[\mathcal{O}]Q_B[\mathcal{O}] \rangle_c &\approx \int_{-1}^1 dz_1 \int_{-1}^1 dz_2 \frac{(1-z_1^2)^{h-1}(1-z_2^2)^{h-1}}{x_0^{2h}} \\
 &= 4^{-h} \left( \frac{\sqrt{\pi} \Gamma(h)}{\Gamma(h+\frac{1}{2})} \right)^2 \eta^h.
 \end{aligned} \tag{3.25}$$

288 We have used the relation  $\eta \approx \frac{4}{x_0^2}$  in the large distance limit. Then we can read out

$$D^{(2)}[\mathcal{O}, \mathcal{O}] = (-1)^{-h} 4^{-h} \left( \frac{\sqrt{\pi} \Gamma(h)}{\Gamma(h+\frac{1}{2})} \right)^2. \tag{3.26}$$

289 Combining UV/IR relation and (3.18), we find

$$p_1^{(2)}[\mathcal{O}, \mathcal{O}] = E^{(2)}[\mathcal{O}] \times D^{(2)}[\mathcal{O}, \mathcal{O}] = \frac{(-1)^{-h} \sqrt{\pi} \Gamma(h)}{\Gamma(h+\frac{1}{2})}. \tag{3.27}$$

290 The result is exactly the same as (3.23). We use the UV/IR relation to obtain type-(3) CCF for  
 291 type-J OPE blocks in two dimensions, the cutoff independent coefficient is

$$p_1^{(2)}[\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3] = \frac{C_{123} \pi^{3/2} (-1)^{\frac{h_1+h_2+h_3}{2}} \Gamma(h_1) \Gamma(h_2) \Gamma(h_3) \kappa}{\Gamma(\frac{1+h_1+h_2-h_3}{2}) \Gamma(\frac{1+h_1+h_3-h_2}{2}) \Gamma(\frac{1+h_2+h_3-h_1}{2}) \Gamma(\frac{h_1+h_2+h_3}{2})}, \tag{3.28}$$

292 where the constant  $\kappa = \frac{1}{2} [1 + (-1)^{h_1+h_2+h_3}]$ . We notice that the result is totally symmetric  
 293 under the exchange of any two conformal weights. Since there are different ways to uplift  
 294 type-( $m$ ) to type-( $m-1, 1$ ), the cutoff independent coefficient should be identical since they  
 295 characterize the same CCF after taking the limit  $A \rightarrow B$ . For  $m = 3$ , this is a cyclic identity

$$p_q^{(d)}[\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3] = p_q^{(d)}[\mathcal{O}_2, \mathcal{O}_3, \mathcal{O}_1] = p_q^{(d)}[\mathcal{O}_3, \mathcal{O}_1, \mathcal{O}_2]. \tag{3.29}$$

296 The UV/IR relation and the cyclic identity have been checked for type-( $m$ ) CCF ( $m=2,3$ ) in  
 297 four dimensions. We list the cutoff independent coefficients below [17].

- 298 • Type-(2). The normalization constants are set to 1.

299 – Spin 1-1 conserved currents.

$$p_1^{(4)}[\mathcal{J}_\mu, \mathcal{J}_\nu] = -\frac{\pi^2}{3}. \quad (3.30)$$

300 – Spin 2-2 conserved currents.

$$p_1^{(4)}[T_{\mu\nu}, T_{\rho\sigma}] = -\frac{\pi^2}{40}. \quad (3.31)$$

301 – Spin 0-0 non-conserved operators.

$$p_2^{(4)}[\mathcal{O}, \mathcal{O}] = -\frac{4\pi^2(\Delta-1)\Gamma(\Delta-2)^2\Gamma(\frac{\Delta}{2})^4}{\Gamma(\Delta)^2\Gamma(\Delta-1)^2}. \quad (3.32)$$

302 – Spin 1-1 non-conserved operators.

$$p_2^{(4)}[\mathcal{O}_\mu, \mathcal{O}_\nu] = -\frac{4^{1-\Delta}\pi^3\Delta\Gamma(\frac{\Delta-3}{2})\Gamma(\frac{\Delta+1}{2})}{\Gamma(\frac{\Delta}{2}+1)^2}, \quad \Delta > 3. \quad (3.33)$$

303 – Spin 2-2 non-conserved operators.

$$p_2^{(4)}[\mathcal{O}_{\mu\nu}, \mathcal{O}_{\rho\sigma}] = -\frac{3\pi^2(\Delta-2)\Delta^2\Gamma(\frac{\Delta}{2}-2)^2\Gamma(\frac{\Delta}{2}-1)^2}{64\Gamma(\Delta-4)\Gamma(\Delta+2)}, \quad \Delta > 4. \quad (3.34)$$

304 • Type-(3).

305 – Spin 1-1-2 conserved currents. The three point function of zero components are  
306 fixed by conformal symmetry

$$\langle T_{00}(x_1)\mathcal{J}_0(x_2)\mathcal{J}_0(x_3) \rangle_c = \frac{C_{T\mathcal{J}\mathcal{J}}}{x_{12}^4 x_{13}^2 x_{23}^2}. \quad (3.35)$$

307 Then the coefficient

$$p_1^{(4)}[\mathcal{J}_\mu, \mathcal{J}_\nu, T_{\rho\sigma}] = -\frac{\pi^3}{2} C_{T\mathcal{J}\mathcal{J}}. \quad (3.36)$$

308 – Spin 2-2-2 conserved currents. The three point function of zero components are  
309 fixed by conformal symmetry

$$\langle T_{00}(x_1)T_{00}(x_2)T_{00}(x_3) \rangle_c = \frac{C_{TTT}}{x_{12}^4 x_{13}^4 x_{23}^4}. \quad (3.37)$$

310 Then the coefficient

$$p_1^{(4)}[T_{\mu\nu}, T_{\rho\sigma}, T_{\alpha\beta}] = \frac{\pi^3}{12} C_{TTT}. \quad (3.38)$$

311 – Spin 0-0-0 non-conserved currents.

$$p_2^{(4)}[\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3] = -2^{4-\Delta_1-\Delta_2-\Delta_3}\pi^3 C_{123} \int_{\mathbb{D}^2} d\zeta d\bar{\zeta} (\zeta + \bar{\zeta})^2 \int_{\mathbb{D}^2} d\zeta' d\bar{\zeta}' (\zeta' + \bar{\zeta}')^2 \\ \times (1-\zeta^2)^{\frac{\Delta_1-4}{2}} (1-\bar{\zeta}^2)^{\frac{\Delta_1-4}{2}} (1-\zeta'^2)^{\frac{\Delta_2-4}{2}} (1-\bar{\zeta}'^2)^{\frac{\Delta_2-4}{2}} \int_0^\pi d\theta \frac{\sin \theta}{(a+b \cos \theta)^{\frac{\Delta_{12,3}}{2}}}, \quad (3.39)$$

312 Though the expression (3.39) is not symmetric superficially under the exchange of any two  
 313 conformal weights, we checked explicitly that it satisfies the cyclic identity for integer conformal  
 314 weights.

315 For  $m = 4$ , the UV/IR relation and the cyclic identity are much more harder to check. We  
 316 considered type-(4) CCF for massless free scalar theory [13, 14]. In this theory, one can construct  
 317 an infinite tower of conserved currents with even spin [29]. The four point functions  
 318 can be calculated explicitly. Therefore we can find type-(3, 1) and type-(4) CCFs and read out  
 319 the corresponding coefficients. For example, for spin-2-2-2-4 conserved currents [14],

$$D[2, 2, 2, 4] = \frac{3}{70}D[2, 2, 4, 2]. \quad (3.40)$$

320 Both of them leads to the cutoff coefficients

$$p_1^{(2)}[2, 2, 2, 4] = \frac{2\Gamma(8)}{\Gamma(4)^2}D[2, 2, 2, 4] = \frac{2\Gamma(4)}{\Gamma(2)^2}D[2, 2, 4, 2] = p_1^{(2)}[2, 2, 4, 2]. \quad (3.41)$$

321 The cyclic identity is obeyed.

### 322 3.4 Discussion

323 The UV/IR relation should be slightly modified when the CCF contains both type-J and type-O  
 324 OPE blocks. One simple example is the following type-(3) CCF

$$\langle Q_A[\mathcal{J}]Q_A[\mathcal{O}]Q_A[\tilde{\mathcal{O}}] \rangle_c \quad (3.42)$$

325 where  $Q_A[\mathcal{J}]$  is a type-J OPE block while  $Q_A[\mathcal{O}]$  and  $Q_A[\tilde{\mathcal{O}}]$  are type-O OPE blocks. This CCF  
 326 is related to the following two type-(2, 1) CCFs

$$\langle Q_A[\tilde{\mathcal{O}}]Q_A[\mathcal{J}]Q_B[\mathcal{O}] \rangle_c = D^{(d)}[\tilde{\mathcal{O}}, \mathcal{J}, \mathcal{O}]G_{\Delta, J}^{(d)}(\xi), \quad (3.43)$$

$$\langle Q_A[\mathcal{O}]Q_A[\tilde{\mathcal{O}}]Q_B[\mathcal{J}] \rangle_c = D^{(d)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}]G_{\Delta', J'}^{(d)}(\xi). \quad (3.44)$$

327 We choose  $d = 4$ . Taking the limit  $A \rightarrow B$  from (3.43), we find a type-(3) CCF with degree  
 328  $q = 2$ , the UV/IR relation reads

$$p_2^{(4)}[\tilde{\mathcal{O}}, \mathcal{J}, \mathcal{O}] = E^{(4)}[\mathcal{O}] \times D^{(4)}[\tilde{\mathcal{O}}, \mathcal{J}, \mathcal{O}] \quad (3.45)$$

329 We can also take the limit  $A \rightarrow B$  from (3.44), then we will find a type-(3) CCF with degree  
 330  $q = 1$ , the UV/IR relation reads

$$p_1^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}] = E^{(4)}[\mathcal{J}] \times D^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}]. \quad (3.46)$$

331 The equations (3.45) and (3.46) are not identical superficially since the subscript  $q$  are not  
 332 equal to each other. However, an explicit calculation for spin 2-0-0 and spin 2-2-0 in four  
 333 dimensions [17] shows that the coefficient  $D^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}]$  is actually divergent logarithmically,  
 334

$$D^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}] = D_{\log}^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}] \log \frac{R}{\epsilon} + \dots \quad (3.47)$$

335 The terms in  $\dots$  are finite and depends on cutoff scale. Due to the logarithmic divergence  
 336 behaviour of the coefficient  $D^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}]$ , the degree of type-(3) CCF from (3.44) increases  
 337 1, the modified UV/IR relation becomes

$$p_2^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}] = E^{(4)}[\mathcal{J}] \times D_{\log}^{(4)}[\mathcal{O}, \tilde{\mathcal{O}}, \mathcal{J}]. \quad (3.48)$$

338 We checked explicitly that the two constants (3.45) and (3.48) are equal to each other. The  
 339 cyclic identity is still satisfied after counting the logarithmic divergence of the  $D$  function.

340 **4 Generalizations**

341 The area law and logarithmic behaviour in the subleading terms can be extended in different  
 342 directions. In this section, we mention several extensions.

- 343 • UV/IR relation. In general, one can uplift any type- $(m)$  CCF to a type- $(p, m - p)$  CCF

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_m] \rangle_c \xrightarrow{\text{uplift}} \langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_p] Q_B[\mathcal{O}_{p+1}] \cdots Q_B[\mathcal{O}_m] \rangle_c, \quad 1 \leq p \leq m-1. \quad (4.1)$$

344 When  $p$  is not 1 and  $m - 1$ , the type- $(p, m - p)$  CCF is not a conformal block. It is still  
 345 a function of cross ratio  $\xi$ , therefore it should reproduce the type- $(m)$  CCF after taking  
 346 the limit  $A \rightarrow B$ ,

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_m] \rangle_c = \lim_{\xi \rightarrow -\infty} \langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_p] Q_B[\mathcal{O}_{p+1}] \cdots Q_B[\mathcal{O}_m] \rangle_c. \quad (4.2)$$

347 Obviously, this also defines a UV/IR relation between  $p_q^{(d)}$  and several coefficients in  
 348 the type- $(p, m - p)$  CCF. Since the right hand side is not proportional to any conformal  
 349 block, it is not easy to write out an explicit formula. Nevertheless, one may still check  
 350 the relation (4.2) case by case. One example is to consider the type- $(2, 2)$  CCF of the  
 351 modular Hamiltonian in  $\text{CFT}_2$ . By making use of the universal feature of the CCF of the  
 352 stress tensor, one can fix the generator of type- $(m_1, m_2)$  CCFs [14]

$$T_{AUB}(\mu_1, \mu_2) = -\frac{c}{2} \text{tr} \log \left[ 1 - \begin{pmatrix} A & C \\ D & B \end{pmatrix} \right], \quad (4.3)$$

353 where the matrices  $A, B, C$  and  $D$  are

$$A_{xx'} = \frac{\eta^2}{4} \int_0^\infty dy \frac{\sqrt{xx'} y \sinh \pi \mu_1 x \sinh \pi \mu_2 y}{\sinh \pi x' \sinh \pi y \sinh \pi(1 + \mu_1)x \sinh \pi(1 + \mu_2)y} \left(\frac{x_{13}}{x_{23}}\right)^{i(x-x')} \mathcal{F}(x, x', y), \quad (4.4)$$

$$B_{xx'} = \frac{\eta^2}{4} \int_0^\infty dy \frac{\sqrt{xx'} y \sinh \pi \mu_1 x \sinh \pi \mu_2 y}{\sinh \pi x' \sinh \pi y \sinh \pi(1 + \mu_1)x \sinh \pi(1 + \mu_2)y} \left(\frac{x_{13}}{x_{23}}\right)^{-i(x-x')} \mathcal{F}(x', x, y), \quad (4.5)$$

$$C_{xx'} = \frac{\eta^2}{4} \int_0^\infty dy \frac{\sqrt{xx'} y \sinh \pi \mu_1 x \sinh \pi \mu_2 y}{\sinh \pi x' \sinh \pi y \sinh \pi(1 + \mu_1)x \sinh \pi(1 + \mu_2)y} \left(\frac{x_{13}}{x_{23}}\right)^{i(x+x')} \mathcal{F}(x, -x', y) \quad (4.6)$$

$$D_{xx'} = \frac{\eta^2}{4} \int_0^\infty dy \frac{\sqrt{xx'} y \sinh \pi \mu_1 x \sinh \pi \mu_2 y}{\sinh \pi x' \sinh \pi y \sinh \pi(1 + \mu_1)x \sinh \pi(1 + \mu_2)y} \left(\frac{x_{13}}{x_{23}}\right)^{-i(x+x')} \mathcal{F}(-x, x', y) \quad (4.7)$$

354 with

$$\begin{aligned} \mathcal{F}(x, x', y) = & {}_2F_1(1 + ix, 1 - iy, 2, -\eta) {}_2F_1(1 - ix', 1 + iy, 2, -\eta) \\ & + {}_2F_1(1 + ix, 1 + iy, 2, -\eta) {}_2F_1(1 - ix', 1 - iy, 2, -\eta). \end{aligned} \quad (4.8)$$

355  $\mathcal{F}$  and its complex conjugate obey

$$\mathcal{F}^*(x, x', y) = \mathcal{F}(x', x, y), \quad \mathcal{F}^*(-x, -x', y) = \mathcal{F}(x, x', y). \quad (4.9)$$

356 so

$$A = B^*, \quad C = D^*. \quad (4.10)$$

357 We read out the first few CCFs

$$\begin{aligned}
 \langle H_A^m \rangle_c &= \frac{cm!}{12} \log \frac{2}{\epsilon}, \\
 \langle H_A^{m-1} H_B \rangle_c &= \frac{cm!}{144} G_2^{(2)}(\eta), \\
 \langle H_A^2 H_B^2 \rangle_c &= c \left\{ \frac{1+\eta}{\eta^2} [4\text{Li}_3(1+\eta) - 2\log(1+\eta)\text{Li}_2(1+\eta) + \frac{2\log(1+\eta)}{3}\text{Li}_2(-\eta) \right. \\
 &\quad \left. + \frac{1+\eta}{3} \log^2(1+\eta) - \frac{\pi^2}{3} \log(1+\eta) - 4\zeta(3)] + \frac{2+\eta}{3\eta} [2\text{Li}_2(-\eta) + 3\log(1+\eta)] - \frac{4}{3} \right\},
 \end{aligned} \tag{4.11}$$

358 where the polylogarithm  $\text{Li}_n(z)$  is

$$\text{Li}_n(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^n}. \tag{4.12}$$

359 The relation (4.2) can be checked for  $p = 2, m = 4$ . The right hand side is

$$\lim_{\eta \rightarrow -1} \langle H_A^2 H_B^2 \rangle_c = 2c \log \frac{2}{\epsilon} + \dots. \tag{4.13}$$

360 The cutoff independent coefficient  $2c$  matches with the one in  $\langle H_A^4 \rangle_c$ .

361 • New power law. In the previous discussion, we focus on the case that  $B$  and  $A$  coincide  
 362 with each other. However, there are other cases that the CCFs are still divergent. One  
 363 can consider the limit that  $A$  just attaches the edge of  $B$ ,

$$R' = 1, \quad x_0 = 2 + \epsilon, \quad \epsilon \rightarrow 0. \tag{4.14}$$

364 The cross ratio  $\xi$  does not approach  $-\infty$  but 1

$$\xi = \frac{4}{(2 + \epsilon)^2} = 1 - \epsilon + \dots. \tag{4.15}$$

365 We can define a new CCF which is also divergent from type- $(m - 1, 1)$  CCF

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_{m-1}] \odot Q_B[\mathcal{O}_m] \rangle_c = \lim_{\xi \rightarrow 1} \langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_{m-1}] Q_B[\mathcal{O}_m] \rangle_c \tag{4.16}$$

366 The continuation of conformal block tells us that the new CCF obeys a new power law

$$\langle Q_A[\mathcal{O}_1] \cdots Q_A[\mathcal{O}_{m-1}] \odot Q_B[\mathcal{O}_m] \rangle_c = \tilde{\gamma} \left(\frac{R}{\epsilon}\right)^{\frac{d-2}{2}} + \dots + \tilde{p}_q^{(d)} \log^q \frac{R}{\epsilon} + \dots. \tag{4.17}$$

367 The leading term is proportional to

$$\mathcal{L} = R^{\frac{d-2}{2}} = \sqrt{\mathcal{A}} \tag{4.18}$$

368 which is the characteristic length of the region  $A$  in four dimensions. In two dimensions,  
 369 the leading term is a logarithmic term with power  $q$ . In this case, there is a new UV/IR  
 370 relation between  $\tilde{p}_q$  and  $D$  coefficient, we write it schematically

$$\tilde{p}_q = \tilde{E} \times D. \tag{4.19}$$

371 The function  $\tilde{E}^{(d)}[\mathcal{O}]$  is proportional to  $E^{(d)}[\mathcal{O}]$ . The proportional constant is shown  
 372 below.



373 –  $d$  is even.

374 For conserved current  $\mathcal{O}$  with conformal weight  $\Delta = J + d - 2$ ,

$$\bar{E}^{(d)}[\mathcal{O}] = \frac{(-1)^J}{2} E^{(d)}[\mathcal{O}]. \tag{4.20}$$

375 For non-conserved current  $\mathcal{O}$  with conformal weight  $\Delta$  and spin  $J$ ,

$$\bar{E}^{(d)}[\mathcal{O}] = \frac{(-1)^J}{4} E^{(d)}[\mathcal{O}]. \tag{4.21}$$

376 We checked the relation for  $d = 2, 4$  and spin  $J \leq 2$ .

377 –  $d$  is odd.

378 For non-conserved current  $\mathcal{O}$  with conformal weight  $\Delta$  and spin  $J$ ,

$$\bar{E}^{(d)}[\mathcal{O}] = \frac{(-1)^J}{2} E^{(d)}[\mathcal{O}]. \tag{4.22}$$

379 For conserved current  $\mathcal{O}$ , there is no logarithmic divergent term in the CCF.

380 We checked the relation for  $d = 3$  and spin  $J \leq 2$ .

381 Since  $D$  function is the same, we find a relation between two cutoff independent coefficients  $p$  and  $\bar{p}$ ,

$$\frac{p}{E} = \frac{\bar{p}}{\bar{E}}. \tag{4.23}$$

## 383 5 Summary and outlook

384 In this report, we have introduced the area law (3.1) of type- $(m)$  CCFs of OPE blocks. It is a  
 385 generalization of the area law of entanglement entropy. We will list several open problems for  
 386 future work.

387 • Higher  $m \geq 4$ . In most of the work, we restrict to the region  $m \geq 3$ . This is because  
 388 the structure of  $m$ -point correlation function of primary operators in CFT is fixed up to  
 389  $m = 3$ . For  $m \geq 4$ , it is harder to extract cutoff independent coefficient.

390 • UV/IR relation. The UV/IR relation

$$p = E \times D \tag{5.1}$$

391 has been checked for several examples. A rigorous proof is still lacking.

392 • Cyclic identity. The cyclic identity of  $p$  reflects the fact that  $p$  is independent of the way  
 393 to regularize the type- $(m)$  CCF. However, we feel that a direct computation is impossible  
 394 to check this identity.

395 • New power law. We generalize the type- $(m_1, m_2)$  CCF to the case that  $A$  and  $B$  just  
 396 attaches with each other. The corresponding CCF is divergent with a new power law  
 397 (4.17). The corresponding new UV/IR relation

$$\bar{p} = \bar{E} \times D \tag{5.2}$$

398 also needs understanding.

- 399 • Deformed reduced density matrix. This exponential operator is similar to the “Wilson  
400 loop” in gauge theories [30, 31] despite the fact that the OPE block has no lower bound  
401 in general. When the OPE block has a lower bound, the logarithm of the vacuum expecta-  
402 tion value of the deformed reduced density matrix

$$\log\langle e^{-\mu Q_A}\rangle \quad (5.3)$$

403 should also obey area law with logarithmic divergence. There may be a gravitational  
404 dual for this quantity as [32, 33]. The similarity of the area law between this program  
405 and black hole entropy implies that the classical part contributes to the area term while  
406 quantum effects lead to logarithmic corrections.

- 407 • Multiple integrals. According to the method of continuation of conformal block, area  
408 law of type- $(m)$  CCF is protected by conformal invariance. However, the method of  
409 continuation itself cannot guarantee that it always leads to the correct result. One has  
410 to develop other methods to deal with the multiple integrals. In two dimensions, one  
411 should generalize Selberg integrals [34, 35] to include more parameters [16].

## 412 Acknowledgements

413 This work was supported by NSFC Grant No. 12005069.

## 414 References

- 415 [1] J. D. Bekenstein, “Black holes and entropy,” *Phys. Rev. D* **7** (1973), 2333-2346  
416 doi:10.1103/PhysRevD.7.2333
- 417 [2] S. W. Hawking, “Particle Creation by Black Holes,” *Commun. Math. Phys.* **43** (1975),  
418 199-220 doi:10.1007/BF02345020
- 419 [3] B. Czech, L. Lamprou, S. McCandlish, B. Mosk and J. Sully, “A Stereoscopic Look into  
420 the Bulk,” *JHEP* **1607**, 129 (2016) doi:10.1007/JHEP07(2016)129 [arXiv:1604.03110  
421 [hep-th]].
- 422 [4] J. de Boer, F. M. Haehl, M. P. Heller and R. C. Myers, “Entanglement, hologra-  
423 phy and causal diamonds,” *JHEP* **1608**, 162 (2016) doi:10.1007/JHEP08(2016)162  
424 [arXiv:1606.03307 [hep-th]].
- 425 [5] S. Ferrara, A. F. Grillo and R. Gatto, “Manifestly conformal covariant operator-product  
426 expansion,” *Lett. Nuovo Cim.* **2S2**, 1363 (1971) [*Lett. Nuovo Cim.* **2**, 1363 (1971)].  
427 doi:10.1007/BF02770435
- 428 [6] S. Ferrara, A. F. Grillo and R. Gatto, “Manifestly conformal-covariant expansion on the  
429 light cone,” *Phys. Rev. D* **5**, 3102 (1972). doi:10.1103/PhysRevD.5.3102
- 430 [7] R. Haag, “Local quantum physics: fields, particles, algebras ,” Springer, Berlin, Germany  
431 (1992).
- 432 [8] L. Bombelli, R. K. Koul, J. Lee and R. D. Sorkin, “A Quantum Source of Entropy for Black  
433 Holes,” *Phys. Rev. D* **34**, 373 (1986).

- 434 [9] M. Srednicki, “Entropy and area,” *Phys. Rev. Lett.* **71**, 666 (1993)  
435 doi:10.1103/PhysRevLett.71.666 [hep-th/9303048].
- 436 [10] C. G. Callan, Jr. and F. Wilczek, “On geometric entropy,” *Phys. Lett. B* **333**, 55 (1994)  
437 doi:10.1016/0370-2693(94)91007-3 [hep-th/9401072].
- 438 [11] H. Araki, “Relative Entropy of States of Von Neumann Algebras,” *Publ. Res. Inst. Math.*  
439 *Sci. Kyoto* **1976**, 809 (1976).
- 440 [12] S. Ryu and T. Takayanagi, “Holographic derivation of entanglement entropy from  
441 AdS/CFT,” *Phys. Rev. Lett.* **96**, 181602 (2006) doi:10.1103/PhysRevLett.96.181602  
442 [hep-th/0603001].
- 443 [13] J. Long, “Correlation function of modular Hamiltonians,” *JHEP* **11** (2019), 163  
444 doi:10.1007/JHEP11(2019)163 [arXiv:1907.00646 [hep-th]].
- 445 [14] J. Long, “Correlation function with the insertion of zero modes of modular Hamilto-  
446 nians,” *JHEP* **01** (2020), 173 doi:10.1007/JHEP01(2020)173 [arXiv:1911.11487 [hep-  
447 th]].
- 448 [15] A. Belin, L. Y. Hung, A. Maloney, S. Matsuura, R. C. Myers and T. Sierens, “Holographic  
449 Charged Renyi Entropies,” *JHEP* **12** (2013), 059 doi:10.1007/JHEP12(2013)059  
450 [arXiv:1310.4180 [hep-th]].
- 451 [16] J. Long, “Logarithmic behaviour of connected correlation function in CFT,”  
452 [arXiv:2001.05129 [hep-th]].
- 453 [17] J. Long, “Area law of connected correlation function in higher dimensional conformal  
454 field theory,” [arXiv:2007.15380 [hep-th]].
- 455 [18] S. N. Solodukhin, “The Conical singularity and quantum corrections to entropy of  
456 black hole,” *Phys. Rev. D* **51** (1995), 609-617 doi:10.1103/PhysRevD.51.609 [arXiv:hep-  
457 th/9407001 [hep-th]].
- 458 [19] S. N. Solodukhin, “On ‘Nongeometric’ contribution to the entropy of black hole due to  
459 quantum corrections,” *Phys. Rev. D* **51** (1995), 618-621 doi:10.1103/PhysRevD.51.618  
460 [arXiv:hep-th/9408068 [hep-th]].
- 461 [20] R. K. Kaul and P. Majumdar, “Logarithmic correction to the Bekenstein-Hawking entropy,”  
462 *Phys. Rev. Lett.* **84** (2000), 5255-5257 doi:10.1103/PhysRevLett.84.5255 [arXiv:gr-  
463 qc/0002040 [gr-qc]].
- 464 [21] S. Carlip, “Logarithmic corrections to black hole entropy from the Cardy formula,” *Class.*  
465 *Quant. Grav.* **17** (2000), 4175-4186 doi:10.1088/0264-9381/17/20/302 [arXiv:gr-  
466 qc/0005017 [gr-qc]].
- 467 [22] T. R. Govindarajan, R. K. Kaul and V. Suneeta, “Logarithmic correction to the Bekenstein-  
468 Hawking entropy of the BTZ black hole,” *Class. Quant. Grav.* **18** (2001), 2877-2886  
469 doi:10.1088/0264-9381/18/15/303 [arXiv:gr-qc/0104010 [gr-qc]].
- 470 [23] A. Sen, “Logarithmic Corrections to Schwarzschild and Other Non-extremal Black Hole  
471 Entropy in Different Dimensions,” *JHEP* **04** (2013), 156 doi:10.1007/JHEP04(2013)156  
472 [arXiv:1205.0971 [hep-th]].
- 473 [24] S. Minwalla, “Restrictions imposed by superconformal invariance on quantum field theo-  
474 ries,” *Adv. Theor. Math. Phys.* **2**, 783 (1998) doi:10.4310/ATMP.1998.v2.n4.a4 [hep-  
475 th/9712074].

- 476 [25] H. Casini, M. Huerta and R. C. Myers, “Towards a derivation of holographic entanglement  
477 entropy,” JHEP **1105**, 036 (2011) doi:10.1007/JHEP05(2011)036 [arXiv:1102.0440  
478 [hep-th]].
- 479 [26] F. A. Dolan and H. Osborn, “Conformal four point functions and the operator product  
480 expansion,” Nucl. Phys. B **599**, 459 (2001) doi:10.1016/S0550-3213(01)00013-X [hep-  
481 th/0011040].
- 482 [27] F. A. Dolan and H. Osborn, “Conformal partial waves and the operator product ex-  
483 pansion,” Nucl. Phys. B **678**, 491 (2004) doi:10.1016/j.nuclphysb.2003.11.016 [hep-  
484 th/0309180].
- 485 [28] M. Hogervorst, H. Osborn and S. Rychkov, “Diagonal Limit for Conformal Blocks in  $d$  Di-  
486 mensions,” JHEP **1308**, 014 (2013) doi:10.1007/JHEP08(2013)014 [arXiv:1305.1321  
487 [hep-th]].
- 488 [29] I. Bakas and E. Kiritsis, Nucl. Phys. B **343** (1990), 185-204 [erratum: Nucl. Phys. B **350**  
489 (1991), 512-512] doi:10.1016/0550-3213(90)90600-I
- 490 [30] J. M. Maldacena, “Wilson loops in large N field theories,” Phys. Rev. Lett. **80**, 4859 (1998)  
491 doi:10.1103/PhysRevLett.80.4859 [hep-th/9803002].
- 492 [31] S. J. Rey and J. T. Yee, “Macroscopic strings as heavy quarks in large N gauge theory and  
493 anti-de Sitter supergravity,” Eur. Phys. J. C **22**, 379 (2001) doi:10.1007/s100520100799  
494 [hep-th/9803001].
- 495 [32] D. L. Jafferis and S. J. Suh, “The Gravity Duals of Modular Hamiltonians,” JHEP **09**  
496 (2016), 068 doi:10.1007/JHEP09(2016)068 [arXiv:1412.8465 [hep-th]].
- 497 [33] D. L. Jafferis, A. Lewkowycz, J. Maldacena and S. J. Suh, “Relative entropy  
498 equals bulk relative entropy,” JHEP **06** (2016), 004 doi:10.1007/JHEP06(2016)004  
499 [arXiv:1512.06431 [hep-th]].
- 500 [34] A. Selberg, “Bemerkninger om et multipelt integral,” Norsk. Mat. Tidsskr. 24(1944)71-78.
- 501 [35] P. Forrester and S. Warnaar, “The importance of the Selberg integral,” Bull. Amer. Math.  
502 Soc. (N.S.) 45 (2008), 489-534.