

MuCap: Muon Capture on the Proton

Malte Hildebrandt¹ and Claude Petitjean^{1*}

1 Paul Scherrer Institut, 5232 Villigen PSI, Switzerland

* claude.petitjean@psi.ch

January 6, 2021



Review of Particle Physics at PSI
doi:[10.21468/SciPostPhysProc.2](https://doi.org/10.21468/SciPostPhysProc.2)

Abstract

The singlet muon capture rate Λ_S on the proton $\mu^- p \rightarrow \nu_\mu n$ is determined in a high precision lifetime measurement. The main apparatus consists of a new hydrogen time projection chamber as muon detector, developed by PSI, surrounded by cylindrical wire chambers and a plastic scintillator hodoscope as electron detectors. Λ_S is evaluated as the difference between the inverse μp lifetime and that of the free μ^+ . The result $\Lambda_S^{\text{MuCap}} = (715.6 \pm 5.4^{\text{stat}} \pm 5.1^{\text{sys}}) \text{s}^{-1}$ is in excellent agreement with the prediction of chiral perturbation theory $\Lambda_S^{\chi\text{PT}} = (715.4 \pm 6.9) \text{s}^{-1}$. From Λ_S^{MuCap} a recent analysis derives for the induced pseudoscalar coupling $g_p^{\text{MuCap}} = 8.23 \pm 0.83$ whereas $\bar{g}_p^{\chi\text{PT}} = 8.25 \pm 0.25$.

17.1 Introduction

Muon capture on the proton

$$\mu^- p \rightarrow \nu_\mu n \quad (17.1)$$

is a very important elementary process in weak interactions [1]. A measurement of the singlet capture rate Λ_S is directly related to fundamental electroweak coupling constants g_A and g_P (??). While g_A is accurately known from measurements of the neutron lifetime, the induced pseudoscalar coupling g_P , can only be precisely determined from the muon capture rate. In low-energy chiral perturbation theory (χPT), g_P can be expressed as¹

$$g_P^{\chi\text{PT}}(q^2) = \frac{2m_\mu g_{\pi NN} f_\pi}{m_\pi^2 - q^2} - \frac{1}{3} g_A(0) m_\mu m_N r_A^2. \quad (17.2)$$

This leads to a theoretical prediction [3, 4] of

$$\bar{g}_p^{\chi\text{PT}} \equiv g_p^{\chi\text{PT}}(q_0^2) = 8.26 \pm 0.23, \quad (17.3)$$

where $q_0^2 = -0.88m_\mu^2$. A precise measurement of Λ_S represents therefore an important test of low-energy χPT .

Historically, many experimental attempts to determine Λ_S were already made in the 1960's at the leading accelerator labs to determine the μp capture rate. These experiments resulted however only in a precision of $\sim 15\%$, suffering mainly from two major challenges:

1) The output channel $\nu_\mu n$ consists only of neutral particles, where the ν_μ escapes detection and the neutron is very difficult to be determined with high absolute precision. Modern


¹The function $g_P(q^2) \equiv m_\mu/m_N F_P^{cc}(q^2)$ and \bar{g}_p are defined in Section 5 [2].

31 experiments avoid this problem by using the lifetime method: instead of measuring absolute
 32 neutron rates, the disappearance rate of the muon, λ_μ , is measured, i.e.

$$\frac{dN_\mu}{dt} = N_\mu e^{-\lambda_\mu t}, \quad \lambda_\mu = \lambda_0 + \Lambda_S. \quad (17.4)$$

33 Here, $\lambda_0 = 0.455 \times 10^6 \text{ s}^{-1}$ is the decay constant² of the free muon and $\Lambda_S \simeq 700 \text{ s}^{-1}$ is just
 34 a small (1.5×10^{-3}) additional component of λ_μ . Lifetime measurements therefore require
 35 high precision, i.e. large statistics. A first successful lifetime experiment was performed 1981
 36 in Saclay [5] in a target with liquid hydrogen.

37 2) Negative muons in hydrogen quickly combine to neutral (μp) atoms which behave like
 38 neutrons; they diffuse around and scatter with the surrounding nuclei. In collisions they can
 39 get easily transferred to heavier nuclei (d,N,O) contained in the hydrogen. Moreover they can
 40 form the mesic molecule ($p\mu p$)




41 with a rate of about $\lambda_{pp\mu} \simeq 2 \times 10^6 \text{ s}^{-1}$. Two species of ($p\mu p$) molecules exist, ortho-($p\mu p$)
 42 and para-($p\mu p$). In the formation process, predominantly ortho-molecules are created, which
 43 eventually convert to the energetically lower para-molecule with rate λ_{op} . Unfortunately, λ_{op}
 44 is not well known (theoretical value $\lambda_{op}^{\text{Th}} = (7.1 \pm 1.2) \times 10^4 \text{ s}^{-1}$ [6]). The capture rates differ
 45 strongly for the two states (for ortho-($p\mu p$) $\sim 545 \text{ s}^{-1}$, for para-($p\mu p$) $\sim 215 \text{ s}^{-1}$). This makes
 46 the interpretation of capture measurements in ($p\mu p$) molecules difficult. This problem can be
 47 strongly reduced in hydrogen at low density, where the ($p\mu p$) formation rate is small.

48 In addition to these two major issues, isotope and chemical purities play an important role
 49 in the experiment. Natural hydrogen contains ~ 150 ppm deuterium nuclei. Muons in such a
 50 medium get quickly transferred to the heavier isotope



51 The (μd) atoms are created at initial kinetic energy of ~ 45 eV, and have a very large diffusion
 52 rate due to a (μd)- p scattering minimum around 10 eV (Ramsauer-Townsend effect). In col-
 53 lisions with deuterium nuclei they can form ($p\mu d$) molecules leading to the muon catalyzed
 54 fusion



55 These processes would strongly interfere in a μp capture measurement. Therefore, hydrogen
 56 depleted from deuterium (so called protium) has to be used. Furthermore, the protium must
 57 be kept at highest purity to avoid transfers to higher- Z nuclei.

58 17.2 The MuCap experiment

59 The MuCap experiment was proposed in 1997 with the goal to measure the singlet μp capture
 60 rate Λ_S to 1% precision which would then determine $g_p(q_0^2)$ to $\sim 6\%$. This goal can be reached
 61 by a high precision measurement of the muon lifetime to the level 10^{-5} which requires a
 62 statistics of $\sim 10^{10}$ muon decay events.

63 Figure 17.1 shows a cross section of the MuCap experiment. The muon detector in the
 64 center consists of three components, a thin scintillator μSC providing the fast timing signal of
 65 the incoming muon, a wire chamber μPC and a time projection chamber TPC [7, 8] tracking
 66 the muon to the stopping point. The TPC is mounted inside an aluminium pressure vessel

²denoted by Γ_μ in (??)




Figure 17.1: Cross section of the full MuCap apparatus with illustration of a typical event. Every muon was tracked individually to its stopping point. The electrons were tracked back to the muon stop location. Thanks to fiducial cuts, background from accidental electrons was suppressed to the 10^{-4} level.

67 filled with 10 bar of ultra-pure protium gas. It acts simultaneously as muon stopping target
 68 and detector. The density of the protium gas is $\sim 1\%$ of liquid hydrogen, thus avoiding the
 69 problems involved with meso-molecular processes. A special isotope separation column was
 70 constructed for MuCap [9] which removed deuterium to a negligible level. A special gas cir-
 71 culation system [10] was constructed using thermo-dynamical cycles and cryo-absorption by
 72 Zeolite filters for continuous cleaning of the protium gas. The system reduced impurity levels
 73 to values below 20 ppb.

74 The TPC was operated with a 2 kV/cm vertical electrical field. The electrons from the
 75 ionizing muon tracks – after drifting downwards to a multi-wire proportional chamber at the
 76 bottom – were collected in x and z coordinates. Combined with the drift time information (y
 77 coordinate) every muon track was reconstructed in three dimensions. After suitable fiducial
 78 cuts false muon stops were suppressed below the 10^{-5} level, necessary to keep the slope of
 79 the muon decay curve free from distortions. The electron detector consists of two cylindrical
 80 wire chambers ePC1, ePC2, and a plastic scintillation hodoscope eSC. The wire chambers
 81 – originally developed by PSI for the SINDRUM rare decay experiments, Section 7 [11] –
 82 provide directional information for each electron track, while the scintillators yield the fast
 83 timing signal of the muon decay.

84 The anticipated precision was reached by collection of more than 10^{10} single good muon
 85 decay events. A significant boost of the statistics was achieved with help of the muon kicker [12]
 86 from the MuLan experiment [13] ('muons on request' method). The system transmitted single
 87 muons into the TPC without pile-up from second particles. This method increased the data
 88 collection rate by a factor 2 to 3.

89 17.3 Results

90 During three independent production runs [14, 15] 1.2×10^{10} fully reconstructed μ^- decays
 91 plus 0.6×10^{10} μ^+ decays for systematic controls were collected. The systematic corrections
 92 include distortion effects due to impurities, removal of μp scatter events, μp and μd diffu-
 93 sion, uncertainties of fiducial volume cuts, inefficiencies and electron track definitions. Aver-
 94 aging these data and using the μ^+ decay constant measured by the MuLan experiment [13],




Figure 17.2: Extracted values for g_p as a function of the poorly known molecular transition rate λ_{op} . OMC = Saclay experiment [5], RMC = TRIUMF experiment [17]. Also shown are results of two inconsistent λ_{op} measurements ($\lambda_{\text{op}}^{\text{Ex1}}$ from Saclay [18], $\lambda_{\text{op}}^{\text{Ex2}}$ from TRIUMF [19]), and the theoretical calculation $\lambda_{\text{op}}^{\text{Th}}$ [6].

95 $\lambda_{\mu^+} = (455'170.05 \pm 0.46) \text{s}^{-1}$, the final result of the singlet muon capture rate on proton is
 96 obtained as [15]

$$\Lambda_S^{\text{MuCap}} = (714.9 \pm 5.4^{\text{stat}} \pm 5.1^{\text{sys}}) \text{s}^{-1} \quad (17.8)$$

97 in excellent agreement with χ PT theory $\Lambda_S^{\chi\text{PT}} = (715.4 \pm 6.9) \text{s}^{-1}$ [16]. From this result

$$g_p^{\text{MuCap}}(q_0^2) = 8.06 \pm 0.48^{\text{exp}} \pm 0.28^{\text{th}} \quad (17.9)$$

98 is deduced [15]. This value is in agreement with χ PT (17.3).

99 Figure 17.2 shows \bar{g}_p from recent experiments as function of the poorly known transition
 100 rate λ_{op} . In contrast to previous experiments which were mostly carried out in liquid hydrogen,
 101 the MuCap experiment is virtually not sensitive to λ_{op} and, thus, avoided this longstanding
 102 problem.

103 In a refined analysis [20] a new value for $\lambda_{pp\mu}$ was derived from the MuCap data and this
 104 led to an updated value of

$$\Lambda_S^{\text{MuCap}} = (715.6 \pm 5.4^{\text{stat}} \pm 5.1^{\text{sys}}) \text{s}^{-1} \quad (17.10)$$

105 and a change of -0.045 in $g_p^{\text{MuCap}}(q_0^2)$. The change of the latter by only 8% of its uncertainty
 106 has no (visible) influence on Figure 17.2.

107 17.4 Outlook

108 The determination of \bar{g}_p from both theory and experiment requires the input of the axial vector
 109 charge radius squared r_a^2 . In a recent review [16] discussing the values and uncertainties of
 110 r_a^2 obtained by different methods, the MuCap result was re-analysed. Based on the value $r_a^2 =$
 111 $(0.46 \pm 0.22) \text{fm}^2$ evaluated from neutrino-nucleon scattering data the updated MuCap result
 112 changes to $g_p^{\text{MuCap}}(q_0^2) = 8.23 \pm 0.83$. This is in very good agreement with the updated value
 113 $\bar{g}_p^{\chi\text{PT}} = 8.25 \pm 0.25$ which is still very close to the value of the Meissner group [3, 4].

114 Following this path the value of r_a^2 is now considered to contain the largest theoretical
115 uncertainty. Fixing \bar{g}_p to the χ PT value, the MuCap result can be interpreted as an independent
116 measurement of r_a^2 : it results in the same value $r_a^2(\mu\text{H}) = (0.46 \pm 0.24) \text{ fm}^2$ as from neutrino
117 scattering.

118 Consequently, a new MuCap experiment with greatly increased statistics would allow a
119 testing of the nucleon axial radius at the correspondingly increased sensitivity. Such an effort
120 would require a newly constructed apparatus using improved detector techniques and muon
121 beam handling.

122 References

- 123 [1] P. Kammel and K. Kubodera, *Precision muon capture*, *Ann. Rev. Nucl. Part. Sci.* **60**, 327
124 (2010), doi:[10.1146/annurev-nucl-100809-131946](https://doi.org/10.1146/annurev-nucl-100809-131946).
- 125 [2] G. Colangelo and A. Signer, *A theory guide for PSI experiments*, *SciPost Phys. Proc.* **2**, ppp
126 (2021), doi:[10.21468/SciPostPhysProc.2.XXX](https://doi.org/10.21468/SciPostPhysProc.2.XXX).
- 127 [3] V. Bernard, N. Kaiser and U. G. Meissner, *QCD accurately predicts the induced pseudoscalar*
128 *coupling constant*, *Phys. Rev.* **D50**, 6899 (1994), doi:[10.1103/PhysRevD.50.6899](https://doi.org/10.1103/PhysRevD.50.6899),
129 [hep-ph/9403351](https://arxiv.org/abs/hep-ph/9403351).
- 130 [4] V. Bernard, L. Elouadrhiri and U.-G. Meissner, *Axial structure of the nucleon: Topical*
131 *Review*, *J. Phys.* **G28**, R1 (2002), doi:[10.1088/0954-3899/28/1/201](https://doi.org/10.1088/0954-3899/28/1/201), [hep-ph/0107088](https://arxiv.org/abs/hep-ph/0107088).
- 132 [5] G. Bardin, J. Duclos, A. Magnon, J. Martino, A. Richter, E. Zavattini, A. Bertin,
133 M. Piccinini, A. Vitale and D. F. Measday, *A Novel Measurement of the Muon Cap-*
134 *ture Rate in Liquid Hydrogen by the Lifetime Technique*, *Nucl. Phys.* **A352**, 365 (1981),
135 doi:[10.1016/0375-9474\(81\)90416-4](https://doi.org/10.1016/0375-9474(81)90416-4).
- 136 [6] D. D. Bakalov, M. P. Faifman, L. I. Ponomarev and S. I. Vinitsky, *μ Capture and Ortho - Para*
137 *Transitions in Mesic Molecule $pp\mu$* , *Nucl. Phys.* **A384**, 302 (1982), doi:[10.1016/0375-](https://doi.org/10.1016/0375-9474(82)90337-2)
138 [9474\(82\)90337-2](https://doi.org/10.1016/0375-9474(82)90337-2).
- 139 [7] J. Egger, M. Hildebrandt and C. Petitjean, *The 10bar hydrogen time projec-*
140 *tion chamber of the MuCap experiment*, *Nucl. Instrum. Meth.* **A628**, 199 (2011),
141 doi:[10.1016/j.nima.2010.06.316](https://doi.org/10.1016/j.nima.2010.06.316).
- 142 [8] J. Egger *et al.*, *A high-pressure hydrogen time projection chamber for the MuCap experiment*,
143 *Eur. Phys. J.* **A50**(10), 163 (2014), doi:[10.1140/epja/i2014-14163-1](https://doi.org/10.1140/epja/i2014-14163-1), [1405.2853](https://arxiv.org/abs/1405.2853).
- 144 [9] I. Alekseev, E. Arkhipov, S. Bondarenko, O. Fedorchenko, V. Ganzha, K. Ivshin, P. Kammel,
145 P. Kravtsov, C. Petitjean, V. Trofimov, A. Vasilyev, T. Vasyanina *et al.*, *Cryogenic distillation*
146 *facility for isotopic purification of protium and deuterium*, *Rev. Sci. Instr.* **86**, 125102
147 (2015), doi:[10.1063/1.4936413](https://doi.org/10.1063/1.4936413).
- 148 [10] V. A. Ganzha *et al.*, *A Circulating hydrogen ultra-high purification system for the MuCap*
149 *experiment*, *Nucl. Instrum. Meth.* **A578**, 485 (2007), doi:[10.1016/j.nima.2007.06.010](https://doi.org/10.1016/j.nima.2007.06.010),
150 [0705.1473](https://arxiv.org/abs/0705.1473).
- 151 [11] R. Eichler and C. Grab, *The SINDRUM-I Experiment*, *SciPost Phys. Proc.* **2**, ppp (2021),
152 doi:[10.21468/SciPostPhysProc.2.XXX](https://doi.org/10.21468/SciPostPhysProc.2.XXX).
- 153 [12] M. J. Barnes and G. D. Wait, *A 25-kV 75-kHz Kicker for Measurement of Muon Lifetime*,
154 *IEEE Trans. Plasma Sci.* **32**, 1932 (2004), doi:[10.1109/TPS.2004.835455](https://doi.org/10.1109/TPS.2004.835455).

- 155 [13] D. M. Webber *et al.*, *Measurement of the Positive Muon Lifetime and Determination of*
156 *the Fermi Constant to Part-per-Million Precision*, Phys. Rev. Lett. **106**, 041803 (2011),
157 doi:[10.1103/PhysRevLett.106.041803](https://doi.org/10.1103/PhysRevLett.106.041803), [10.1103/PhysRevLett.106.079901](https://doi.org/10.1103/PhysRevLett.106.079901), [Phys. Rev.
158 Lett.106,079901(2011)], [1010.0991](https://arxiv.org/abs/1010.0991).
- 159 [14] V. A. Andreev *et al.*, *Measurement of the rate of muon capture in hydrogen gas and deter-*
160 *mination of the proton's pseudoscalar coupling $g(P)$* , Phys. Rev. Lett. **99**, 032002 (2007),
161 doi:[10.1103/PhysRevLett.99.032002](https://doi.org/10.1103/PhysRevLett.99.032002), [0704.2072](https://arxiv.org/abs/0704.2072).
- 162 [15] V. A. Andreev *et al.*, *Measurement of Muon Capture on the Proton to 1% Precision and*
163 *Determination of the Pseudoscalar Coupling g_p* , Phys. Rev. Lett. **110**(1), 012504 (2013),
164 doi:[10.1103/PhysRevLett.110.012504](https://doi.org/10.1103/PhysRevLett.110.012504), [1210.6545](https://arxiv.org/abs/1210.6545).
- 165 [16] R. J. Hill, P. Kammel, W. J. Marciano and A. Sirlin, *Nucleon Axial Radius and Muonic*
166 *Hydrogen — A New Analysis and Review*, Rept. Prog. Phys. **81**(9), 096301 (2018),
167 doi:[10.1088/1361-6633/aac190](https://doi.org/10.1088/1361-6633/aac190), [1708.08462](https://arxiv.org/abs/1708.08462).
- 168 [17] D. H. Wright *et al.*, *Measurement of the induced pseudoscalar coupling using radiative*
169 *muon capture on hydrogen*, Phys. Rev. **C57**, 373 (1998), doi:[10.1103/PhysRevC.57.373](https://doi.org/10.1103/PhysRevC.57.373).
- 170 [18] G. Bardin, J. Duclos, A. Magnon, J. Martino, A. Richter, E. Zavattini, A. Bertin, M. Pic-
171 cinini and A. Vitale, *Measurement of the Ortho - Para Transition Rate in the $p\mu p$ Molecule*
172 *and Deduction of the Pseudoscalar Coupling Constant*, Phys. Lett. **104B**, 320 (1981),
173 doi:[10.1016/0370-2693\(81\)90134-9](https://doi.org/10.1016/0370-2693(81)90134-9).
- 174 [19] J. H. D. Clark, D. S. Armstrong, T. P. Gorringer, M. D. Hasinoff, P. M. King, T. J. Stocki,
175 S. Tripathi, D. H. Wright and P. A. Zolnierczuk, *Ortho-para transition rate in mu-molecular*
176 *hydrogen and the proton's induced pseudoscalar coupling $g(p)$* , Phys. Rev. Lett. **96**, 073401
177 (2006), doi:[10.1103/PhysRevLett.96.073401](https://doi.org/10.1103/PhysRevLett.96.073401), [nucl-ex/0509025](https://arxiv.org/abs/nucl-ex/0509025).
- 178 [20] V. A. Andreev *et al.*, *Measurement of the Formation Rate of Muonic Hydrogen Molecules*,
179 Phys. Rev. **C91**(5), 055502 (2015), doi:[10.1103/PhysRevC.91.055502](https://doi.org/10.1103/PhysRevC.91.055502), [1502.00913](https://arxiv.org/abs/1502.00913).