

MEG: Muon to Electron and Gamma

A. M. Baldini^{1*} and T. Mori²

¹ INFN Pisa, Largo B. Pontecorvo 2, 56127 Pisa, Italy

² ICEPP, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

* CorrespondingAuthor@email.address

May 20, 2021



Review of Particle Physics at PSI
doi:[10.21468/SciPostPhysProc.2](https://doi.org/10.21468/SciPostPhysProc.2)

Abstract

The possible existence of the $\mu \rightarrow e\gamma$ decay predicted by many new physics scenarios is investigated by stopping positive muons in a very thin target and measuring emitted photons and positrons with the best possible resolutions. Photons are measured by a 2.7 ton ultra pure liquid xenon detector while positron trajectories are measured in a specially designed gradient magnetic field by low-mass drift chambers and precisely timed by scintillation counters. A first phase of the experiment (MEG) ended in 2016, and excluded the existence of the decay with branching ratios larger than 4.2×10^{-13} (90% C.L.). This provides approximately 30 times stronger constraints on a variety of new physics models than previous experiments. In the second phase (MEG II), most of the detectors have been upgraded by adopting up-to-date technologies to improve the search sensitivity by another order of magnitude down to $\mathcal{O}(10^{-14})$. MEG II will perform a search for physics beyond the Standard Model complementary to high energy collider experiments with compatible or even higher sensitivity.

19.1 Introduction: the $\mu \rightarrow e\gamma$ decay

In early 1990's, the precision electroweak measurements at the LEP Collider, CERN, suggested that the electromagnetic, weak, and strong interactions can be unified at $\mathcal{O}(10^{16})$ GeV if TeV-scale supersymmetry exists, hinting strongly at supersymmetric grand unification (SUSY GUT) [1, 2]. It was then shown that sizable lepton flavor violating (LFV) couplings arise naturally in SUSY GUT through the renormalization group evolution, thanks to the heavy top quark, even under the assumption of no flavor mixing at the SUSY breaking scale [3, 4]. Such LFV couplings would lead to an experimentally observable rate of the muon decay, $\mu \rightarrow e\gamma$, which the Standard Model strictly prohibits.

Physicists, fascinated by the possibility of exploring SUSY GUT in muon decays, held a series of workshops in 1997 to design a possible $\mu \rightarrow e\gamma$ search experiment at PSI. PSI was considered to be the best place for such an experiment. This workshop series evolved into a Letter of Intent in 1998 [5], which was strongly supported by PSI. A research proposal for a $\mu \rightarrow e\gamma$ experiment [6] was submitted in 1999 and approved by PSI. The experimental collaboration subsequently expanded and named itself MEG (Mu-E-Gamma) with an updated sensitivity goal of $\mathcal{O}(10^{-13})$ [7]. At the time the best upper limit was 1.2×10^{-11} [8].

Discovery of neutrino oscillations in 1998 reinforced the importance of the $\mu \rightarrow e\gamma$ search. Tiny neutrino masses indicated by the oscillations are explained naturally by the see-saw mechanism. It was shown that the heavy right-handed Majorana neutrinos predicted by the see-saw

41 mechanism could induce sizable LFV couplings, contributing significantly to the $\mu \rightarrow e\gamma$ decay
42 rate in supersymmetric models [9].

43 Lepton flavor conservation is violated in the neutrino oscillations. It is the smallness of
44 the neutrino masses, not lepton flavor conservation, that suppresses the branching ratio of the
45 $\mu \rightarrow e\gamma$ decay much below 10^{-50} in the Standard Model extended for finite neutrino masses
46 [10]. Therefore, new physics scenarios that involve heavier particles coupled to leptons, such
47 as supersymmetry or extra dimensions, can naturally produce a measurable rate of the $\mu \rightarrow e\gamma$
48 decay, making this channel one of the most powerful tools to search for new physics.

49 In the final state of a $\mu \rightarrow e\gamma$ event, an electron and a photon are traveling back-to-back
50 in the rest frame of the decaying muon. Both the electron and the photon have energies of
51 half the muon mass (52.8 MeV). To take advantage of this simple 2-body kinematics in the
52 experiment, muons are stopped in a target. Positive muons are necessarily used to avoid
53 formation of muonic atoms where the nucleus gets recoiled in the muon decay.

54 The major background for the $\mu \rightarrow e\gamma$ search is the accidental coincidence of a positron
55 from normal muon decay, $\mu \rightarrow e\nu\bar{\nu}$, and a photon either from a radiative muon decay or from
56 the annihilation of a positron in material. Good energy and momentum measurements can
57 strongly suppress the physics background from radiative muon decays, $\mu \rightarrow e\nu\bar{\nu}\gamma$, to levels
58 about an order of magnitude lower than the accidental background.

59 A continuous muon beam rather than a pulsed beam is better suited as the accidental back-
60 ground increases quadratically with the muon rate. To achieve a sensitivity to the branching
61 ratio of $10^{-13} - 10^{-14}$ with a detection efficiency $\epsilon \approx \mathcal{O}(1 - 10\%)$ in a few years of data taking
62 ($T \approx \mathcal{O}(10^7)$ sec), a continuous muon rate of $(10^{13} - 10^{14})/\epsilon/T \approx (10^7 - 10^8)/\text{sec}$ proves
63 necessary. Such a high rate continuous muon beam is available only at PSI.

64 Both the MEG and the MEG II experiments were designed to satisfy the following three
65 experimental requirements to achieve a sensitivity level of $10^{-13} - 10^{-14}$:

- 66 • *A high intensity continuous muon beam of $10^7 - 10^8$ muons/sec.*
- 67 • *A photon detector with excellent energy resolution.* The energy spectrum of the back-
68 ground photons from radiative muon decays falls off towards the high energy end of
69 52.8 MeV. A photon detector with excellent energy resolution can significantly suppress
70 these backgrounds. An innovative liquid xenon scintillation photon detector, which has
71 a very good intrinsic energy resolution, not limited by impurities, was developed for
72 MEG. It also provides good resolutions in position and timing of photons to discriminate
73 the accidental background.
- 74 • *A low mass positron spectrometer that can operate at high rates.* Precision positron spec-
75 trometry must be carried out in the environment of $10^7 - 10^8$ positrons/sec. A positron
76 spectrometer with a gradient magnetic field, called COBRA (COnstant Bending RAdius),
77 was designed to avoid positrons piling up in the central part of the tracker as well as to
78 discriminate absolute momenta of positrons, together with low mass drift chambers to
79 minimize multiple scattering, and scintillation counters with excellent timing resolution.

80 19.2 The first phase of the experiment: MEG

81 In the first phase of the experiment 3×10^7 positive muons per second coming from the $\pi E5$
82 beam line were stopped in a thin (205 μm) polyethylene target placed at the center of the
83 detector. The positron spectrometer consisted of a set of drift chambers (built at PSI) and scin-
84 tillation timing counters placed inside the volume of the COBRA magnet. The photon detector,
85 located outside of the magnet, contained 900 liters of liquid xenon seen by 846 photomulti-
86 pliers (PMTs) sensitive to ultraviolet light directly immersed in xenon. The detectors covered

107 10% of the solid angle. A PSI-designed waveform digitizer based on the multi-GHz domino
 108 ring sampler chip was used to digitize each channel of the detectors.

109 One of the core activities of MEG consisted in continuously monitoring and calibrating the
 110 different detectors in order to guarantee their response stability. In particular, several tools
 111 were used for the liquid xenon detector:

- 112 • α -sources and LEDs immersed in the liquid for daily calibration of PMTs;
- 113 • a radioactive Am/Be source, the (p, γ) -reaction obtained by a dedicated Cockcroft-Walton
 114 accelerator and π^-p charge exchange and radiative capture reactions to measure the en-
 115 ergy scale and resolutions over the 4.43 MeV to 129.4 MeV energy range;
- 116 • a neutron generator to obtain a 9 MeV- γ line from the capture in nickel to check the
 117 stability of the detector even during data-taking;

118 Radiative muon decays and the two γ -rays from the $^{11}\text{B}(p, 2\gamma)^{12}\text{C}$ reaction were further used
 119 to monitor the relative timing of the positron spectrometer and xenon detectors.

120 The kinematic variables used to identify the $\mu \rightarrow e\gamma$ decays are the γ -ray and positron
 121 energies (E_γ , E_e), their opening angle ($\Theta_{e\gamma}$) and difference of their emission times ($t_{e\gamma}$).

122 The background was largely (a factor ten roughly) dominated by the accidental superpo-
 123 sition of energetic positrons from standard Michel muon decay with photons from radiative
 124 muon decay.




Figure 19.1: MEG final results: (a) E_γ vs E_e (b) $\cos(\Theta_{e\gamma})$ vs $t_{e\gamma}$. 68%, 90% and 95% C.L. signal contour lines are shown.

125 A total of 7.5×10^{14} muons were stopped in the target during the MEG experiment [11].
 126 Figure 19.1 shows the event distributions in the (E_γ, E_e) and $(\cos(\Theta_{e\gamma}), t_{e\gamma})$ planes for the full
 127 data set together with the 68%, 90% and 95% contours of the signal probability distribution
 128 function.

129 A blind analysis method was used by hiding events near (in time and energy) the signal
 130 region and using events outside this (blind) region to build probability density functions for a
 131 maximum likelihood analysis. Once analysis procedures were completed events belonging to
 132 the blind region were analyzed.

133 The analysis shows no evidence for a signal: the final branching ratio upper limit for the
 134 $\mu \rightarrow e\gamma$ decay is 4.2×10^{-13} [11] (90% Confidence Level).

135 While the signal region in Figure 19.1 does not show any significant excess of events, it
 136 does contain background events. Since, in these conditions, limits would have improved only

117 with square root of time, it was decided to end MEG data taking and proceed to a second phase
118 of the experiment with an upgraded detector able to reduce the background further.

119 The MEG dataset was also used to search for other muon decay modes such as $\mu \rightarrow eX$,
120 $X \rightarrow \gamma\gamma$, recently suggested by models (see for instance [12]) in which new physics is predicted
121 at low, rather than high, energy scales. No significant excess was found in the mass range of
122 the axion-like particle X , $m_X = 20\text{--}45 \text{ MeV}/c^2$ and $\tau_X < 40 \text{ ps}$: the upper limits established
123 [13] were lowered to $O(10^{-11})$ for $m_X = 20\text{--}30 \text{ MeV}/c^2$, up to 60 times more stringent than
124 previous results [14].

125 19.3 Towards the discovery: MEG II

126 The basic concept of the upgraded MEG experiment – MEG II – is to improve the detector
127 resolutions everywhere so that it can run at the highest muon intensity available at PSI without
128 suffering a high rate of the accidental background: MEG had to reduce the muon intensity for
129 stable detector operation, and to keep the accidental background at a sufficiently low level.
130 A significant improvement of the detector resolutions enables the higher muon stopping rate
131 with a similar level of the background as MEG, and, together with the improved detector
132 efficiency, can achieve an order of magnitude higher sensitivity than MEG.

133 The main improvements of MEG II over MEG are [15]:

134 • *Higher stopping muon rate on target:*

135 – A new single-volume drift chamber with stereo geometry instead of cathode pads
136 for stable long-term operation at the full muon intensity.

137 • *Larger detector acceptance:*

138 – Material mass and distance are minimised between the drift chamber and the tim-
139 ing counter, where nearly half of the positrons were lost in the MEG experiment.
140 – Better photon efficiency with lower material mass at the photon entrance face by
141 replacing photomultiplier tubes with silicon photosensors (SiPM).

142 • *Improved detector resolutions:*

143 – Better position resolution and more hits per track with the new drift chamber.
144 – A new pixelated timing counter system with straightforward extrapolation of positron
145 trajectory from the drift chamber for improved timing resolution.
146 – Better photon resolutions with more uniform calorimeter response by using SiPMs
147 instead of PMTs.
148 – A better energy resolution for photons entering near the lateral faces by realigning
149 photomultiplier tubes.

150 • *Background suppression:*

151 – A thinner muon stopping target.
152 – A lower-mass drift chamber.
153 – A new device to actively tag the radiative background events.

154 Furthermore, a unified trigger/digitiser data-acquisition (DAQ) board has been developed
155 to manage an increased number of read-out channels and a higher bandwidth of the analog
156 front-end.

157 A sketch of the MEG II experiment is shown in Figure 19.2.




Figure 19.2: A drawing of the MEG II experiment showing its different components

158 Re-tuning the beam line with the full intensity beam to improve the sensitivity, results in
 159 a muon stopping rate of $\sim (5 - 7) \times 10^7 \mu/\text{sec}$. Assuming 120 DAQ days per year, a final
 160 sensitivity of $(5 - 6) \times 10^{-14}$ can be reached in 3-4 years of running, an order of magnitude
 161 better than the final MEG sensitivity. The MEG II proposal was approved by PSI in January,
 162 2013.

163 19.4 Outlook

164 The MEG II experiment is currently in the detector commissioning phase at the $\pi E5$ beam line.
 165 A small fraction (about 10%) of the read-out electronics channels is available to establish the
 166 optimal data taking conditions. This includes the best beam intensity, the frequency of the
 167 detector maintenance and calibration, the gas mixture for the optimal chamber operation, etc,
 168 so that all the detectors can be operated in their best stable conditions for the entire data-taking
 169 period.

170 After all of the readout electronics is installed in 2021, the experiment will start a full
 171 engineering run. If things go well, physics data taking can then begin. It will be necessary to
 172 accumulate data for three to four years at the $\pi E5$ beam line to achieve the intended sensitivity.

173 Other experiments at PSI, J-PARC in Japan, and FNAL in the U.S., plan to start searches
 174 for other LFV muon processes, $\mu \rightarrow 3e$ [16] (see Section 20 [17]) and $\mu \rightarrow e$ conversion in
 175 the presence of a nucleus [18, 19], in this decade. MEG II, together with these experiments,
 176 will scrutinize an unexplored territory of physics beyond the Standard Model, which may not
 177 be accessible to the LHC experiments, and could even identify the dynamics of new physics
 178 from a careful comparison of these measurements. It is hoped that MEG II will lead in making
 179 important steps towards our understanding of the fundamental laws of nature.

180 The High Intensity Muon Beam (HIMB) project at PSI [20], aiming at developing new
 181 muon beam lines with intensities up to $10^{10} \mu^+/s$, could have a crucial role in the future for
 182 further studies of $\mu \rightarrow e\gamma$ to clarify new physics. It will require novel experimental technologies

183 beyond MEG II to keep the backgrounds arising from such high muon intensities under control.

184 References

- 185 [1] U. Amaldi, W. de Boer and H. Furstenau, *Comparison of grand unified theories with*
186 *electroweak and strong coupling constants measured at LEP*, Phys. Lett. B **260**, 447 (1991),
187 doi:[10.1016/0370-2693\(91\)91641-8](https://doi.org/10.1016/0370-2693(91)91641-8).
- 188 [2] U. Amaldi, W. de Boer, P. H. Frampton, H. Furstenau and J. T. Liu, *Consistency*
189 *checks of grand unified theories*, Phys. Lett. B **281**, 374 (1992), doi:[10.1016/0370-](https://doi.org/10.1016/0370-2693(92)91158-6)
190 [2693\(92\)91158-6](https://doi.org/10.1016/0370-2693(92)91158-6).
- 191 [3] R. Barbieri and L. Hall, *Signals for supersymmetric unification*, Phys. Lett. B **338**, 212
192 (1994), doi:[10.1016/0370-2693\(94\)91368-4](https://doi.org/10.1016/0370-2693(94)91368-4), [hep-ph/9408406](https://arxiv.org/abs/hep-ph/9408406).
- 193 [4] R. Barbieri, L. J. Hall and A. Strumia, *Violations of lepton flavor and CP in supersymmetric*
194 *unified theories*, Nucl. Phys. B **445**, 219 (1995), doi:[10.1016/0550-3213\(95\)00208-A](https://doi.org/10.1016/0550-3213(95)00208-A),
195 [hep-ph/9501334](https://arxiv.org/abs/hep-ph/9501334).
- 196 [5] A. Van der Schaaf *et al.*, *Letter of Intent to Paul Scherrer Institute, R-98-05.0*, available at
197 <https://meg.web.psi.ch/docs/loi/loi.ps> (1998).
- 198 [6] T. Mori *et al.*, *Research Proposal to Paul Scherrer Institute, R-99-05*, available at
199 https://meg.web.psi.ch/docs/prop_psi/proposal.pdf (1999).
- 200 [7] A. Baldini *et al.*, (MEG collaboration). *Research Proposal to INFN*, available at
201 https://meg.web.psi.ch/docs/prop_infn/nproposal.pdf (2002).
- 202 [8] M. Brooks *et al.*, Phys. Rev. Lett. **83**, 1521 (1999).
- 203 [9] J. Hisano and D. Nomura, *Solar and atmospheric neutrino oscillations and lepton flavor*
204 *violation in supersymmetric models with the right-handed neutrinos*, Phys. Rev. D **59**,
205 116005 (1999), doi:[10.1103/PhysRevD.59.116005](https://doi.org/10.1103/PhysRevD.59.116005), [hep-ph/9810479](https://arxiv.org/abs/hep-ph/9810479).
- 206 [10] W. Marciano, T. Mori and J. Roney, Annu. Rev. Nucl. Part. Sci. **58**, 315 (2008).
- 207 [11] A. Baldini *et al.*, *Search for the lepton flavour violating decay $\mu^+ \rightarrow e^+ \gamma$ with the full dataset*
208 *of the MEG experiment*, Eur. Phys. J. C **76**(8), 434 (2016), doi:[10.1140/epjc/s10052-](https://doi.org/10.1140/epjc/s10052-016-4271-x)
209 [016-4271-x](https://doi.org/10.1140/epjc/s10052-016-4271-x), [1605.05081](https://arxiv.org/abs/1605.05081).
- 210 [12] C. Cornella, P. Paradisi and O. Sumensari, *Hunting for ALPs with Lepton Flavor Violation*,
211 JHEP **01**, 158 (2020), doi:[10.1007/JHEP01\(2020\)158](https://doi.org/10.1007/JHEP01(2020)158), [1911.06279](https://arxiv.org/abs/1911.06279).
- 212 [13] A. Baldini *et al.*, *Search for lepton flavour violating muon decay mediated by a new light par-*
213 *ticle in the MEG experiment*, Eur. Phys. J. C **80**(9), 858 (2020), doi:[10.1140/epjc/s10052-](https://doi.org/10.1140/epjc/s10052-020-8364-1)
214 [020-8364-1](https://doi.org/10.1140/epjc/s10052-020-8364-1), [2005.00339](https://arxiv.org/abs/2005.00339).
- 215 [14] R. Bolton *et al.*, *Search for Rare Muon Decays with the Crystal Box Detector*, Phys. Rev. D
216 **38**, 2077 (1988), doi:[10.1103/PhysRevD.38.2077](https://doi.org/10.1103/PhysRevD.38.2077).
- 217 [15] A. Baldini *et al.*, (MEG collaboration). *The design of the MEG II experiment*, Eur. Phys. J.
218 C **78**, 380 (2018), doi:[10.1140/epjc/s10052-018-5845-6](https://doi.org/10.1140/epjc/s10052-018-5845-6).
- 219 [16] K. Arndt *et al.*, *Technical design of the phase I Mu3e experiment*, [arxiv:2009.11690v1](https://arxiv.org/abs/2009.11690v1)
220 (2020).

- 221 [17] F. Wauters, *The Mu3e experiment*, SciPost Phys. Proc. **2**, ppp (2021),
222 doi:[10.21468/SciPostPhysProc.2.XXX](https://doi.org/10.21468/SciPostPhysProc.2.XXX).
- 223 [18] R. Abramishvili *et al.*, *COMET Phase-1 Technical Design Report*, Tech. Rep. 2015-1, KEK
224 (2015).
- 225 [19] L. Bartoszek *et al.*, *Mu2e Technical Design Report*, Tech. Rep. 2014-01, Fermilab,
226 doi:[10.2172/1172555](https://doi.org/10.2172/1172555) (2014).
- 227 [20] F. Berg *et al.*, *Target Studies for Surface Muon Production*, Phys. Rev. Accel. Beams **19**(2),
228 024701 (2016), doi:[10.1103/PhysRevAccelBeams.19.024701](https://doi.org/10.1103/PhysRevAccelBeams.19.024701), [1511.01288](https://arxiv.org/abs/1511.01288).