

Thermodynamic limit and boundary energy of the spin-1 Heisenberg chain with non-diagonal boundary fields

Zhihan Zheng^{1,2}, Pei Sun^{1,2}, Xiaotian Xu^{1,2*}, Tao Yang^{1,2,3,4}, Junpeng Cao^{4,5,6,7}, Wen-Li Yang^{1,2,3,4}

1 Institute of Modern Physics, Northwest University, Xi'an 710127, China

2 Shaanxi Key Laboratory for Theoretical Physics Frontiers, Xi'an 710127, China

3 School of Physics, Northwest University, Xi'an 710127, China

4 Peng Huanwu Center for Fundamental Theory, Xi'an 710127, China

5 Beijing National Laboratory for Condensed Matter Physics, Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China

6 Songshan Lake Materials Laboratory, Dongguan, Guangdong 523808, China

7 School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

*xtxu@nwu.edu.cn

1 Abstract

The thermodynamic limit and boundary energy of the isotropic spin-1 Heisenberg chain with non-diagonal boundary fields are studied. The finite size scaling properties of the inhomogeneous term in the $T - Q$ relation at the ground state are analyzed. Based on the reduced Bethe ansatz equations (BAEs), we obtain the boundary energy of the system. These results can be generalized to the $SU(2)$ symmetric high spin Heisenberg model directly.

9 Contents

10	1 Introduction	1
11	2 Non-diagonal boundary Spin-1 Heisenberg model	3
12	3 Finite size scaling behavior	6
13	4 Boundary energy	10
14	5 Conclusions	15
15	References	15

18 1 Introduction

The study of quantum integrable models is an interesting subject in the fields of cold atoms, quantum field theory, condensed matter physics and statistic mechanics [1–5]. The spin-1/2 Heisenberg model can effectively quantify the spin-exchanging interaction and

22 plays an important role in the quantum magnetism and many-body theory. By using the
 23 Bethe ansatz method, the one-dimensional (1D) spin-1/2 Heisenberg model can be solved
 24 exactly [6]. The typical spin-exchanging couplings in the 1D spin-1 system is characterized
 25 by the bilinear biquadratic model, where the Hamiltonian reads

$$H = \sum_{k=1}^N \left[J_1 \vec{S}_k \cdot \vec{S}_{k+1} + J_2 (\vec{S}_k \cdot \vec{S}_{k+1})^2 \right]. \quad (1)$$

26 Here $\vec{S}_k(S_k^x, S_k^y, S_k^z)$ is the spin-1 operator at site k , N is the number of sites and the
 27 periodic boundary condition gives $\vec{S}_{N+1} = \vec{S}_1$. If $J_2/J_1 = 1$, the system (1) has the
 28 $SU(3)$ symmetry and is integrable. If $J_2/J_1 = -1$, the $SU(2)$ symmetry exists and
 29 the system is known as the Zamalodchikov-Fateev (ZF) model [7]. The Bethe ansatz
 30 solution and thermodynamic properties of the ZF model are studied by Takhtajan [8] and
 31 Babujian [9,10]. If $J_2 = 0$, the system is no longer integrable. Starting from the nonlinear
 32 sigma model, Haldane conjectures that the excitation of the system has a gap [11,12]. If
 33 $J_2/J_1 = 1/3$, the Hamiltonian (1) degenerates into a projector operator that is in fact
 34 the projection onto the sum of the spin-0 and spin-1 subspaces (up to a constant) and
 35 the ground state is the famous valence bond solid state [13,14]. If $J_1 = 0$, by using the
 36 Temperley-Lieb algebra, the system can be mapped into the XXZ spin chain and is also
 37 integrable [15–17].

38 Besides the periodic boundary condition, the integrable open one is also an interesting
 39 subject, which means that the system has magnetic impurity or the boundary magnetic
 40 fields [18,19]. In the past few decades, the exact results of high spin models with periodic
 41 [7,9,10,20–25] and parallel boundary fields [26–28] have been extensively studied. It is
 42 emphasized that the integrable boundary reflection matrix can have non-diagonal elements,
 43 which means that the boundary fields are unparallel. Then the $U(1)$ symmetry is broken
 44 and it is very hard to study the exact solution of the system. It is known that the integrable
 45 systems without $U(1)$ symmetry actually have many applications in the open string theory
 46 and the stochastic process of nonequilibrium statistics. Therefore, many interesting works
 47 of high spin models with non-diagonal boundary reflections have been done [29–35].

48 Recently, a systematic method, i.e., the off-diagonal Bethe ansatz (ODBA) is proposed
 49 to solve the models with or without $U(1)$ symmetry [36]. The eigenvalues and eigenstates
 50 of several typical integrable models are obtained. The next task is to derive the physical
 51 quantities in the thermodynamic limit, which is very involved in because the related Bethe
 52 ansatz equations (BAEs) are inhomogeneous and the traditional thermodynamic Bethe
 53 ansatz can not be employed. In order to overcome this difficulty, an effective method is to
 54 study the finite size scaling effects of the inhomogeneous term in the $T - Q$ relation. With
 55 the help of this idea, the thermodynamic limit, surface energy and elementary excitations
 56 of spin-1/2 XXZ spin chain with arbitrary boundary fields are studied [37]. The boundary
 57 energy of the $SU(3)$ symmetric spin-1 chain with generic integrable open boundaries is
 58 also obtained [38]. However, the corresponding thermodynamic properties of the $SU(2)$
 59 symmetric spin-1 Heisenberg model are still missing.

60 In this paper, we study the thermodynamic limit and boundary energy of the spin-1
 61 isotropic Heisenberg spin chain with non-diagonal boundary reflections. The finite size
 62 scaling analysis of the contribution of the inhomogeneous term in the $T - Q$ relation to
 63 the ground state energy is studied in detail. In the thermodynamic limit, we find that the
 64 most Bethe roots of the reduced BAEs at the ground state form 2-strings, associated with
 65 certain boundary strings and the rearrangement of Fermi sea. The different structures of
 66 Bethe roots in different regimes of model parameters are given explicitly. Based on them,
 67 we obtain the boundary energy induced by the unparallel boundary magnetic fields. We

68 also check the analytic results by the numerical extrapolation, and find that the analytic
69 results and the numerical ones coincide with each other very well. The results given in
70 this paper can be generalized to the $SU(2)$ symmetric spin- s Heisenberg model directly.

71 This paper is organized as follows. Section 2 serves as an introduction to the notations
72 for the spin-1 Heisenberg model with non-diagonal boundary fields. The ODBA exact
73 solution is also briefly reviewed. In Section 3, we focus on the contribution of the inho-
74 mogeneous term in the $T - Q$ relation to the ground state energy. In Section 4, by using
75 the patterns of Bethe roots of the reduced BAEs, we study the boundary energy of the
76 model in the thermodynamic limit. We summarize the results and give some discussions
77 in Section 5.

78 2 Non-diagonal boundary Spin-1 Heisenberg model

79 The spin-1 Heisenberg model with non-diagonal boundary fields is related with the 19-
80 vertex R -matrix

$$R_{12}(u) = \left(\begin{array}{c|cc|c} c(u) & & & \\ & b(u) & & \\ \hline & & d(u) & \\ & e(u) & & \\ \hline & & & g(u) & f(u) \\ & & g(u) & & \\ \hline & & & & b(u) & e(u) \\ & & f(u) & & & \\ \hline & & & & & g(u) & d(u) \\ & & & & & & b(u) & c(u) \end{array} \right), \quad (2)$$

81 where the non-vanishing elements are

$$\begin{aligned} a(u) &= u(u + \eta) + 2\eta^2, & b(u) &= u(u + \eta), & c(u) &= (u + \eta)(u + 2\eta), \\ d(u) &= u(u - \eta), & e(u) &= 2\eta(u + \eta), & f(u) &= 2\eta^2, & g(u) &= 2u\eta, \end{aligned} \quad (3)$$

82 u is the spectral parameter and η is the crossing parameter. Here we are dealing with the
83 isotropic model and η can be scaled out. Throughout this paper, we adopt the standard
84 notations. For any matrix $A \in \text{End}(\mathbb{V})$, A_j is an embedding operator in the tensor space
85 $\mathbb{V} \otimes \mathbb{V} \otimes \dots$, which acts as A on the j -th space and as identity on the other factor spaces.
86 For any matrix $B \in \text{End}(\mathbb{V} \otimes \mathbb{V})$, $B_{i,j}$ is an embedding operator in the tensor space, which
87 acts as identity on the factor spaces except for the i -th and j -th ones. The R -matrix
88 $R_{12}(u)$ satisfies the quantum Yang-Baxter equation (QYBE) [39, 40]

$$R_{12}(u - v)R_{13}(u)R_{23}(v) = R_{23}(v)R_{13}(u)R_{12}(u - v). \quad (4)$$

89 Besides, the R -matrix (2) also enjoys the properties

$$\text{Initial condition : } R_{12}(0) = 2\eta^2 P_{12}, \quad (5)$$

$$\text{Fusion condition : } R_{12}(-\eta) = 6\eta^2 \mathbf{P}_{12}^{(0)}, \quad (6)$$

90 where P_{12} is the permutation operator and $\mathbf{P}_{12}^{(0)}$ is the projector in the total spin-0 channel.
91 The most general off-diagonal boundary reflection at one side of the chain is quantified by
92 the reflection matrix

$$K^-(u) = (2u + \eta) \begin{pmatrix} x_1(u) & y'_4(u) & y'_6(u) \\ y_4(u) & x_2(u) & y'_5(u) \\ y_6(u) & y_5(u) & x_3(u) \end{pmatrix}, \quad (7)$$

93 where the matrix elements are

$$\begin{aligned}
x_1(u) &= (p_- + u + \frac{\eta}{2})(p_- + u - \frac{\eta}{2}) + \frac{\alpha_-^2}{2} \eta (u - \frac{\eta}{2}), \\
x_2(u) &= (p_- + u - \frac{\eta}{2})(p_- - u + \frac{\eta}{2}) + \alpha_-^2 (u + \frac{\eta}{2})(u - \frac{\eta}{2}), \\
x_3(u) &= (p_- - u - \frac{\eta}{2})(p_- - u + \frac{\eta}{2}) + \frac{\alpha_-^2}{2} \eta (u - \frac{\eta}{2}), \\
y_4(u) &= \sqrt{2} \alpha_- e^{-i\phi_-} u (p_- + u - \frac{\eta}{2}), \quad y'_4(u) = \sqrt{2} \alpha_- e^{i\phi_-} u (p_- + u - \frac{\eta}{2}), \\
y_5(u) &= \sqrt{2} \alpha_- e^{-i\phi_-} u (p_- - u + \frac{\eta}{2}), \quad y'_5(u) = \sqrt{2} \alpha_- e^{i\phi_-} u (p_- - u + \frac{\eta}{2}), \\
y_6(u) &= \alpha_-^2 e^{-2i\phi_-} u (u - \frac{\eta}{2}), \quad y'_6(u) = \alpha_-^2 e^{2i\phi_-} u (u - \frac{\eta}{2}),
\end{aligned} \tag{8}$$

94 p_- , α_- and ϕ_- are the boundary parameters which measure the strength and direction of
95 the boundary field. The reflection matrix $K^-(u)$ satisfies the reflection equation (RE)

$$R_{12}(u-v)K_1^-(u)R_{21}(u+v)K_2^-(v) = K_2^-(v)R_{21}(u+v)K_1^-(u)R_{12}(u-v). \tag{9}$$

96 The most general off-diagonal boundary reflection at the other side is quantified by the
97 dual reflection matrix

$$K^+(u) = K^-(-u - \eta) \Big|_{(p_-, \alpha_-, \phi_-) \rightarrow (p_+, -\alpha_+, \phi_+)}, \tag{10}$$

98 where p_+ , α_+ and ϕ_+ are the boundary parameters characterizing the strength and direc-
99 tion of the corresponding boundary field. The dual reflection matrix $K^+(u)$ satisfies the
100 dual RE

$$\begin{aligned}
R_{12}(v-u)K_1^+(u)R_{21}(-u-v-2\eta)K_2^+(v) \\
= K_2^+(v)R_{21}(-u-v-2\eta)K_1^+(u)R_{12}(v-u).
\end{aligned} \tag{11}$$

101 From the R -matrix (2), we construct the single row monodromy matrices $T_0(u)$ and
102 $\hat{T}_0(u)$ as

$$\begin{aligned}
T_0(u) &= R_{0N}(u - \theta_N)R_{0N-1}(u - \theta_{N-1}) \cdots R_{01}(u - \theta_1), \\
\hat{T}_0(u) &= R_{10}(u + \theta_1)R_{20}(u + \theta_2) \cdots R_{N0}(u + \theta_N),
\end{aligned} \tag{12}$$

103 where $\{\theta_k, k = 1, \dots, N\}$ are the inhomogeneous parameters, the subscript 0 means the
104 auxiliary space and $1, \dots, N$ denote the quantum spaces. The single row monodromy
105 matrices $T_0(u)$ and $\hat{T}_0(u)$ are the 3×3 matrices in the auxiliary space \mathbf{V}_0 and their elements
106 act on the quantum space $\mathbf{V}^{\otimes N}$. The transfer matrix of the system reads

$$t(u) = \text{tr}_0 \{ K_0^+(u) T_0(u) K_0^-(u) \hat{T}_0(u) \}. \tag{13}$$

107 From the QYBE (4), RE (9) and dual RE (11), one can prove that the transfer matrices
108 with different spectral parameters commute with each other, i.e.,

$$[t(u), t(v)] = 0. \tag{14}$$

109 Therefore, $t(u)$ serves as the generating functional of all the conserved quantities, which
110 ensures the integrability of the system. The model Hamiltonian is generated from the

111 transfer matrix $t(u)$ as

$$\begin{aligned}
H &= \partial_u \{ \ln[t(u)] \} \Big|_{u=0, \{\theta_k=0\}} \\
&= \frac{1}{\eta} \sum_{k=1}^{N-1} \left[\vec{S}_k \cdot \vec{S}_{k+1} - (\vec{S}_k \cdot \vec{S}_{k+1})^2 \right] \\
&\quad + \frac{1}{p_-^2 - \frac{1}{4}(1 + \alpha_-^2)\eta^2} \left[2p_- \alpha_- \cos \phi_- S_1^x + 2p_- S_1^z + \frac{1}{2}(\alpha_-^2 \eta - 2\eta)(S_1^z)^2 \right. \\
&\quad \quad - \frac{1}{2} \alpha_-^2 \eta \cos(2\phi_-) [(S_1^x)^2 - (S_1^y)^2] - \alpha_- \eta \cos \phi_- [S_1^z S_1^x + S_1^x S_1^z] \\
&\quad \quad + \frac{1}{2} \alpha_-^2 \eta \sin(2\phi_-) [S_1^x S_1^y + S_1^y S_1^x] - 2p_- \alpha_- \sin \phi_- S_1^y \\
&\quad \quad \left. + \eta \alpha_- \sin \phi_- [S_1^y S_1^z + S_1^z S_1^y] \right] \\
&\quad + \frac{1}{(3p_+^2 - \frac{3}{4}(1 + \alpha_+^2)\eta^2)\eta} \left[6p_+ \alpha_+ \eta \cos \phi_+ S_N^x - 6p_+ \eta S_N^z \right. \\
&\quad \quad + 3\alpha_+ \eta^2 \cos \phi_+ [S_N^x S_N^z + S_N^z S_N^x] - (2p_+^2 - \frac{3}{2}(1 - \alpha_+^2 \cos(2\phi_+))\eta^2)(S_N^x)^2 \\
&\quad \quad - (2p_+^2 - \frac{3}{2}(1 + \alpha_+^2 \cos(2\phi_+))\eta^2)(S_N^y)^2 - (2p_+^2 + \frac{3}{2}(1 - \alpha_+^2)\eta^2)(S_N^z)^2 \\
&\quad \quad - 3\eta^2 \alpha_+ \sin \phi_+ [S_N^y S_N^z + S_N^z S_N^y] - 6p_+ \alpha_+ \eta \sin \phi_+ S_N^y \\
&\quad \quad \left. + \frac{3}{2} \eta^2 \alpha_+^2 \sin(2\phi_+) [S_N^x S_N^y + S_N^y S_N^x] \right] \\
&\quad + \frac{\eta(1 + \alpha_+^2)}{3p_+^2 - \frac{3}{4}(1 + \alpha_+^2)\eta^2} + \frac{\eta}{p_-^2 - \frac{1}{4}(1 + \alpha_-^2)\eta^2} + \frac{1}{\eta} 3N + \frac{4}{\eta}. \tag{15}
\end{aligned}$$

112 Now, we seek the exact solution of the system (15). Let $|\Psi\rangle$ be an arbitrary eigenstate
113 of $t(u)$ with the eigenvalue $\Lambda(u)$, i.e.,

$$t(u)|\Psi\rangle = \Lambda(u)|\Psi\rangle. \tag{16}$$

114 Using the ODBA method [36] and fusion hierarchy, in the homogeneous limit $\{\theta_k = 0\}$,
115 the eigenvalue $\Lambda(u)$ can be expressed as the inhomogeneous $T - Q$ relation,

$$\Lambda(u) = -4u(u + \eta)\Lambda^{(\frac{1}{2},1)}(u + \frac{\eta}{2})\Lambda^{(\frac{1}{2},1)}(u - \frac{\eta}{2}) + 4u(u + \eta)\delta^{(1)}(u + \frac{\eta}{2}), \tag{17}$$

$$\Lambda^{(\frac{1}{2},1)}(u) = a^{(1)}(u)\frac{Q(u - \eta)}{Q(u)} + d^{(1)}(u)\frac{Q(u + \eta)}{Q(u)} + cu(u + \eta)\frac{F^{(1)}(u)}{Q(u)}, \tag{18}$$

116 where

$$\begin{aligned}
a^{(1)}(u) &= d^{(1)}(-u - \eta) \\
&= -\frac{2u + 2\eta}{2u + \eta} (\sqrt{1 + \alpha_+^2}u + p_+) (\sqrt{1 + \alpha_-^2}u - p_-) \left(u + \frac{3\eta}{2}\right)^{2N}, \tag{19}
\end{aligned}$$

$$F^{(1)}(u) = (u - \frac{\eta}{2})^{2N} (u + \frac{\eta}{2})^{2N} (u + \frac{3\eta}{2})^{2N}, \tag{20}$$

$$\delta^{(1)}(u) = a^{(1)}(u) d^{(1)}(u - \eta), \tag{21}$$

$$c = 2[\alpha_- \alpha_+ \cos(\phi_+ - \phi_-) - 1 + \sqrt{(1 + \alpha_-^2)(1 + \alpha_+^2)}], \tag{22}$$

$$Q(u) = \prod_{k=1}^{2N} (u - u_k)(u + u_k + \eta) = Q(-u - \eta), \tag{23}$$

117 and the $2N$ parameters $\{u_k|k = 1, \dots, 2N\}$ in Q -function (23) are the Bethe roots. The
118 singularity of eigenvalue $\Lambda(u)$ requires that the Bethe roots should satisfy the BAEs

$$a^{(1)}(u_k)Q(u_k - \eta) + d^{(1)}(u_k)Q(u_k + \eta) + c u_k(u_k + \eta) F^{(1)}(u_k) = 0, \quad k = 1, \dots, 2N. \quad (24)$$

119 The eigenvalue of Hamiltonian (15) reads

$$E = \sum_{k=1}^{2N} \frac{4\eta}{(u_k + \frac{3\eta}{2})(u_k - \frac{\eta}{2})} + \frac{1}{\eta} 3N + \frac{1}{\eta} E_0, \quad (25)$$

120 where $\{u_k\}$ should satisfy the BAEs (24) and

$$E_0 = \frac{8}{3} + \frac{2\sqrt{1 + \alpha_+^2} p_+ \eta}{p_+^2 - \frac{\eta^2}{4}(1 + \alpha_+^2)} - \frac{2\sqrt{1 + \alpha_-^2} p_- \eta}{p_-^2 - \frac{\eta^2}{4}(1 + \alpha_-^2)}. \quad (26)$$

121 Some remarks are in order. If the non-diagonal boundary parameters are $\alpha_+ = \alpha_- = 0$, or
122 $\alpha_+ = -\alpha_- \neq 0$ and $\phi_- = \phi_+$ (which corresponds to the parallel boundary fields case), the
123 parameter c in Eq.(22) becomes zero and the corresponding $T - Q$ relation (18) is naturally
124 reduced to the conventional diagonal one [29] obtained by the algebraic Bethe Ansatz.¹
125 For the other case with unparallel boundary fields, the parameter c does not vanish and
126 thus the corresponding $T - Q$ relation has to include a non-vanishing inhomogeneous term
127 for any finite N .

128 3 Finite size scaling behavior

129 The present BAEs (24) are inhomogeneous, thus it is very hard to investigate the thermo-
130 dynamic properties of the system by using the traditional thermodynamic Bethe ansatz.
131 In order to overcome this difficulty, we first analyze the contribution of inhomogeneous
132 term in the $T - Q$ relation (18).

133 Define the reduced $T - Q$ relation as

$$\Lambda_{hom}(u) = -4u(u + \eta)\Lambda_{hom}^{(\frac{1}{2},1)}(u + \frac{\eta}{2})\Lambda_{hom}^{(\frac{1}{2},1)}(u - \frac{\eta}{2}) + 4u(u + \eta)\delta^{(1)}(u + \frac{\eta}{2}), \quad (27)$$

$$\Lambda_{hom}^{(\frac{1}{2},1)}(u) = a^{(1)}(u)\frac{Q(u - \eta)}{Q(u)} + d^{(1)}(u)\frac{Q(u + \eta)}{Q(u)}. \quad (28)$$

134 It should be emphasized that although the non-diagonal boundary parameters $\{p_{\pm}, \alpha_{\pm}\}$
135 except ϕ_{\pm} are included in the above reduced $T - Q$ relation (28), the $\Lambda_{hom}(u)$ is not
136 the eigenvalue $\Lambda(u)$ for any finite N but in the limit of $N \rightarrow \infty$ it will give the correct
137 boundary energy (see the following parts of the paper). From the singularity analysis of
138 the reduced $T - Q$ relation (28), we obtain following reduced BAEs

$$\frac{\frac{i}{2} - \mu_k p i - \mu_k q i - \mu_k}{\frac{i}{2} + \mu_k p i + \mu_k q i + \mu_k} \left(\frac{i - \mu_k}{i + \mu_k} \right)^{2N} = \prod_{l=1}^M \frac{i - (\mu_k - \mu_l)}{i + (\mu_k - \mu_l)} \frac{i - (\mu_k + \mu_l)}{i + (\mu_k + \mu_l)}, \quad k = 1, \dots, M, \quad (29)$$

139 where $M = 1, \dots, 2N$ and we have put $\eta = 1$, $\mu_k = -iu_k - \frac{i}{2}$, $p = \frac{p_+}{\sqrt{1 + \alpha_+^2}} - \frac{1}{2}$ and
140 $q = -\frac{p_-}{\sqrt{1 + \alpha_-^2}} - \frac{1}{2}$ for convenience. From the $\Lambda_{hom}(u)$ given by Eq.(27), we obtain the

¹If the non-diagonal boundary parameters satisfy the condition $\alpha_+ = \alpha_- \neq 0$, $|\phi_- - \phi_+| = \pi$ (which corresponds to the antiparallel boundary fields case), the parameter c in Eq.(22) also becomes zero and the corresponding $T - Q$ relation naturally degenerates into the conventional diagonal one.

141 reduced energy which is defined as

$$E_{hom} = \partial_u \{ \ln \Lambda_{hom}(u) \} \Big|_{u=0} = - \sum_{k=1}^M \frac{4}{\mu_k^2 + 1} + 3N + E_0. \quad (30)$$

142 Solving the reduced BAEs (29), we could obtain the values of reduced Bethe roots $\{\mu_k\}$.
 143 Substituting the Bethe roots into Eq.(30), we obtain the values of E_{hom} .

144 Let us focus on the ground state. The reduced ground state energy can be calculated
 145 by the reduced BAEs (29). It is well-known that the even N and odd N give the same
 146 physical properties in the thermodynamic limit. Thus we set N is even. At the ground
 147 state, the number of Bethe roots in the reduced BAEs (29) is $M = N$. For simplicity,
 148 we choose the boundary parameters as $p > 0$ and $q \neq 0, -1$. We should note that at the
 149 points of $q = 0, -1$, the boundary field is divergent due to the present parameterization of
 150 the Hamiltonian (15). The distribution of reduced Bethe roots at the ground state in the
 151 thermodynamic limit is shown in Figure 1. We see that the Bethe roots can be divided
 152 into six different regimes in the $p - q$ plane.

153 1) In the regime I, where $p \geq 1/2$, $q < -1$, $-1/2 \leq q < 0$ or $q \geq 1/2$, all the Bethe
 154 roots form 2-strings, i.e., $\mu_k = \lambda_k \pm \frac{i}{2} + \mathcal{O}(e^{-\delta N})$, where λ_k denotes the position of 2-string
 155 in the real axis, δ is a small positive number and $\mathcal{O}(e^{-\delta N})$ means the finite size correction.

156 2) In the regime II, where $p < 1/2$, $q < -1$, $-1/2 \leq q < 0$ or $q \geq 1/2$, besides $N - 2$
 157 2-strings, there are two boundary strings, i.e., pi and $(p-1)i$. The boundary strings mean
 158 the pure imaginary Bethe roots which are related with the boundary parameters p and
 159 q [41].

160 3) In the regime III, where $p \geq 1/2$ and $0 < q < 1/2$, besides $N - 2$ 2-strings, there
 161 are two boundary strings, qi and $(q-1)i$.

162 4) In the regime IV, where $0 < p < 1/2$ and $0 < q < 1/2$, besides $N - 4$ 2-strings,
 163 there are four boundary strings, pi , $(p-1)i$, qi and $(q-1)i$.

164 5) In the regime V, where $p \geq 1/2$ and $-1 < q < -1/2$, besides $N - 2$ 2-strings, only
 165 the boundary string qi survives and one real Bethe root λ_0 appears which is caused by
 166 the rearrangement of Fermi sea.

167 6) In the regime VI, where $0 < p < 1/2$ and $-1 < q < -1/2$, besides $N - 4$ 2-strings,
 168 there are three boundary strings qi , $(q-1)i$, pi and one real root λ_0 .

169 Because the Bethe roots are different in the different regimes of boundary parameters,
 170 we shall discuss them separately. In the regime I, where all the Bethe roots are the
 171 2-strings. Substituting the 2-string solutions into the reduced BAEs (29), omitting the
 172 exponentially small corrections and taking the product of all the string solutions, we
 173 readily obtain

$$\begin{aligned} & \frac{i - \lambda_j}{i + \lambda_j} \frac{(p - \frac{1}{2})i - \lambda_j}{(p - \frac{1}{2})i + \lambda_j} \frac{(p + \frac{1}{2})i - \lambda_j}{(p + \frac{1}{2})i + \lambda_j} \frac{(q - \frac{1}{2})i - \lambda_j}{(q - \frac{1}{2})i + \lambda_j} \frac{(q + \frac{1}{2})i - \lambda_j}{(q + \frac{1}{2})i + \lambda_j} \\ & \times \left(\frac{\frac{1}{2}i - \lambda_j}{\frac{1}{2}i + \lambda_j} \frac{\frac{3}{2}i - \lambda_j}{\frac{3}{2}i + \lambda_j} \right)^{2N} = \prod_{l=1}^{M_1} \left[\frac{i - (\lambda_j - \lambda_l)}{i + (\lambda_j - \lambda_l)} \right]^2 \left[\frac{i - (\lambda_j + \lambda_l)}{i + (\lambda_j + \lambda_l)} \right]^2 \\ & \times \frac{2i - (\lambda_j - \lambda_l)}{2i + (\lambda_j - \lambda_l)} \frac{2i - (\lambda_j + \lambda_l)}{2i + (\lambda_j + \lambda_l)}, \quad j = 1, \dots, M_1. \end{aligned} \quad (31)$$

174 Taking the logarithm of above Eq.(31), we obtain

$$2\pi I_j = W(\lambda_j; M_1) + \theta_{2p-1}(\lambda_j) + \theta_{2p+1}(\lambda_j) + \theta_{2q-1}(\lambda_j) + \theta_{2q+1}(\lambda_j), \quad j = 1, \dots, M_1, \quad (32)$$




Figure 1: The distribution of reduced Bethe roots at the ground states with different boundary parameters p and q .

175 where

$$W(\lambda_j; M_1) = \theta_2(\lambda_j) + 2N [\theta_1(\lambda_j) + \theta_3(\lambda_j)] - \sum_{l=1}^{M_1} [2\theta_2(\lambda_j - \lambda_l) + 2\theta_2(\lambda_j + \lambda_l) + \theta_4(\lambda_j - \lambda_l) + \theta_4(\lambda_j + \lambda_l)], \quad (33)$$

176 I_j is the quantum number, $\theta_n(x) = 2 \arctan(2x/n)$ and $M_1 = N/2$. The ground state is
177 characterized by the set of quantum numbers

$$\{I_j\} = \{1, 2, \dots, M_1\}. \quad (34)$$

178 Solving the reduced BAEs (32) and substituting the values of Bethe roots into Eq.(30),
179 we obtain the reduced ground state energy as

$$E_{hom} = -2 \sum_{j=1}^{M_1} \frac{1}{\lambda_j^2 + \frac{1}{4}} + \frac{3}{\lambda_j^2 + \frac{9}{4}} + 3N + E_0 \equiv G(\lambda_j; M_1). \quad (35)$$

180 Now, we are ready to characterize the contribution of inhomogeneous term in the $T-Q$
181 relation (18) at the ground state by the quantity

$$E_{inh} = E_{hom} - E_g, \quad (36)$$

182 where E_{hom} is the reduced ground state energy given by (35) and E_g is the actual ground
183 state energy (25) of the Hamiltonian (15). The ground state energy E_g can be obtained
184 by two methods. One is solving the inhomogeneous BAEs (24) directly and the other is
185 density matrix renormalization group (DMRG) [42–44]. We have checked that the ground
186 state energy E obtained by these two methods are the same.

187 In Figure 2(a), we give the values of E_{inh} versus the system size N in the regime I. The
188 red circles are the data calculated from Eq.(36) and the blue solid line is the fitted curve.
189 From the fitted curve, we find that E_{inh} and N satisfy the power law relation $E_{inh} = \gamma N^\beta$.
190 Due to the fact that $\beta < 0$, the value of E_{inh} tends to zero when the system size N tends




Figure 2: The values of E_{inh} versus the system size N . The data can be fitted as $E_{inh} = \gamma N^\beta$. Due to the fact $\beta < 0$, when the size of system $N \rightarrow \infty$, the contribution of the inhomogeneous term tends to zero. Here (a) $p = 1.1370, q = -1.0821, \gamma = 0.06203$ and $\beta = -0.9407$ in regime I; (b) $p = 0.3263, q = -1.8931, \gamma = 0.2371$ and $\beta = -1.052$ in regime II; (c) $p = 0.2428, q = 2.3735, \gamma = 0.6236$ and $\beta = -0.8384$ in regime III; (d) $p = 0.4453, q = 0.3789, \gamma = 2.234$ and $\beta = -1.087$ in regime IV; (e) $p = 0.8410, q = -0.6990, \gamma = 0.715$ and $\beta = -1.219$ in regime V; (f) $p = 0.3971, q = -0.7985, \gamma = 4.912$ and $\beta = -1.429$ in regime VI. The insets show the distribution of Bethe roots with $N = 10$.

191 to infinity. Therefore, in the thermodynamic limit, the inhomogeneous term in the $T - Q$
 192 relation (18) can be neglected at the ground state and $E_{hom} = E_g$. The inset shows the
 193 distribution of Bethe roots with $N = 10$.

194 In the regime II, substituting the $N - 2$ 2-strings, two boundary strings $\mu_{M-1} = pi$
 195 and $\mu_M = (p - 1)i$ into the reduced BAEs (29) and taking the logarithm, we have

$$2\pi I_j = W(\lambda_j; M_2) + \theta_{2q-1}(\lambda_j) + \theta_{2q+1}(\lambda_j) - \theta_{1-2p}(\lambda_j) - \theta_{2p+1}(\lambda_j) - \theta_{3+2p}(\lambda_j) - \theta_{5-2p}(\lambda_j) - 2\theta_{3-2p}(\lambda_j), \quad j = 1, 2, \dots, M_2, \quad (37)$$

196 where $W(\lambda_j; M_2)$ is given by Eq.(33) with the replacing of M_1 by M_2 , $M_2 = N/2 - 1$ and
 197 the quantum numbers are

$$\{I_j\} = \{1, 2, \dots, M_2\}. \quad (38)$$

198 The corresponding reduced ground state energy reads

$$E_{hom} = G(\lambda_j; M_2) + \frac{4}{p^2 - 1} + \frac{4}{(p - 1)^2 - 1}, \quad (39)$$

199 where $G(\lambda_j; M_2)$ is given by Eq.(35) with the replacing of M_1 by M_2 .

200 The procedure in the regime III is similar and reduced ground state energy is

$$E_{hom} = G(\lambda_j; M_2) + \frac{4}{q^2 - 1} + \frac{4}{(q - 1)^2 - 1}. \quad (40)$$

201 In the regime IV, substituting the string solutions including four boundary strings into
 202 Eq.(29) and taking the logarithm, we have

$$2\pi I_j = W(\lambda_j; M_3) - \theta_{1-2p}(\lambda_j) - \theta_{2p+1}(\lambda_j) - \theta_{3+2p}(\lambda_j) - \theta_{5-2p}(\lambda_j) - 2\theta_{3-2p}(\lambda_j) - \theta_{1-2q}(\lambda_j) - \theta_{2q+1}(\lambda_j) - \theta_{3+2q}(\lambda_j) - \theta_{5-2q}(\lambda_j) - 2\theta_{3-2q}(\lambda_j), \quad j = 1, 2, \dots, M_3, \quad (41)$$

203 where $M_3 = N/2 - 2$ and the quantum numbers are

$$\{I_j\} = \{1, 2, \dots, M_3\}. \quad (42)$$

204 The reduced ground state energy is

$$E_{hom} = G(\lambda_j; M_3) + \frac{4}{p^2 - 1} + \frac{4}{(p-1)^2 - 1} + \frac{4}{q^2 - 1} + \frac{4}{(q-1)^2 - 1}. \quad (43)$$

205 In the regime V, the logarithm form of the BAEs are

$$2\pi I_j = W(\lambda_j; M_4) + \theta_{2p-1}(\lambda_j) + \theta_{2p+1}(\lambda_j) - \theta_{3+2q}(\lambda_j) - \theta_{3-2q}(\lambda_j) - 2\theta_{1-2q}(\lambda_j) \\ - \theta_1(\lambda_j - \lambda_0) - \theta_1(\lambda_j + \lambda_0) - \theta_3(\lambda_j - \lambda_0) - \theta_3(\lambda_j + \lambda_0), \quad j = 1, 2, \dots, M_4, \quad (44)$$

206 where $M_4 = N/2 - 1$ and the quantum numbers are $\{I_j\} = \{1, 2, \dots, M_4\}$. We shall
207 note that the quantum number corresponding to the real Bethe root λ_0 is 0. The reduced
208 ground state energy reads

$$E_{hom} = G(\lambda_j; M_4) + \frac{4}{q^2 - 1} - \frac{4}{\lambda_0^2 + 1}. \quad (45)$$

209 Similarly, the reduced ground state energy in the regime VI is

$$E_{hom} = G(\lambda_j; M_5) + \frac{4}{p^2 - 1} + \frac{4}{(p-1)^2 - 1} + \frac{4}{q^2 - 1} - \frac{4}{\lambda_0^2 + 1}, \quad (46)$$

210 where $M_5 = N/2 - 2$.

211 Substituting the reduced ground state energies in different regimes into Eq.(36), we
212 obtain the values of E_{inh} , which are shown in Figures 2(b)-(f). According to the finite
213 size scaling analysis, we see that the inhomogeneous term indeed can be neglected at the
214 ground state in the thermodynamic limit.

215 4 Boundary energy

216 In this section, we study the physical effects induced by the unparallel boundary magnetic
217 fields and compute the boundary energy [18, 35, 45–47]. The values of Bethe roots at the
218 ground state are determined by the quantum numbers $\{I_j\}$. Thus we define the counting
219 function as $Z(\lambda_j) = \frac{I_j}{2N}$. In the thermodynamic limit, the Bethe roots can take the
220 continuous values and we have $Z(\lambda_j) \rightarrow Z(u)$. Taking the derivative of $Z(u)$ with respect
221 to u , we obtain

$$\frac{dZ(u)}{du} = \rho(u) + \rho^h(u), \quad (47)$$

222 where $\rho(u)$ is the density of Bethe roots and $\rho^h(u)$ means the density of holes in the real
223 axis. Again, the distribution of Bethe roots in different regimes are different. We should
224 consider them separately. In regime I, from the BAEs (32) with the constraint $N \rightarrow \infty$
225 and using Eq.(47), we obtain the density of states as

$$\rho(u) = \frac{dZ(u)}{du} - \frac{1}{2N}[\rho^h(u) + \delta(u)] \\ = a_1(u) + a_3(u) + \frac{1}{2N} [a_2(u) + a_{2p-1}(u) + a_{2p+1}(u) + a_{2q-1}(u) + a_{2q+1}(u)] \\ - \frac{1}{2N}[\rho^h(u) + \delta(u)] - \int_{-\infty}^{\infty} [2a_2(u-v) + a_4(u+v)] \rho(v) dv, \quad (48)$$

226 where

$$a_n(u) = \frac{1}{2\pi} \frac{n}{u^2 + \frac{n^2}{4}},$$

$$\rho^h(u) = \frac{1}{2N} \left[\delta(u - \lambda_1^h) + \delta(u + \lambda_1^h) + \delta(u - \lambda_2^h) + \delta(u + \lambda_2^h) \right]. \quad (49)$$

227 We should note that the presence of delta-function in Eq.(48) is due to that $\lambda_j = 0$ is the
228 solution of BAEs (32), which should be excluded because it makes the wavefunction vanish
229 identically [48]. Note that two holes λ_1^h and λ_2^h are introduced to ensure the magnetization
230 satisfying

$$\frac{M}{N} = 2 \int_{-\infty}^{\infty} \rho(u) du = 1. \quad (50)$$

231 Thus the holes are located at the infinities in the real axis.

232 With the help of Fourier transformation

$$\tilde{F}(\omega) = \int_{-\infty}^{\infty} e^{i\omega u} F(u) du, \quad F(u) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-i\omega u} \tilde{F}(\omega) d\omega, \quad (51)$$

233 from Eq.(48), we obtain

$$\tilde{\rho}(\omega) = \tilde{\rho}_g(\omega) + \tilde{\rho}_0(\omega) + \tilde{\rho}_1(\omega) + \tilde{\rho}_2(\omega), \quad (52)$$

234 where

$$\begin{aligned} \tilde{a}_n(\omega) &= e^{-\frac{n|\omega|}{2}}, \quad \tilde{\rho}_g(\omega) = \frac{\tilde{a}_1(\omega) + \tilde{a}_3(\omega)}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)}, \quad \tilde{\rho}_0(\omega) = \frac{1}{2N} \frac{\tilde{a}_2(\omega) - 1}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)}, \\ \tilde{\rho}_1(\omega) &= \begin{cases} \frac{1}{2N} \frac{\tilde{a}_{2p+1}(\omega) - \tilde{a}_{1-2p}(\omega)}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)}, & 0 < p < \frac{1}{2}, \\ \frac{1}{2N} \frac{\tilde{a}_{2p-1}(\omega) + \tilde{a}_{2p+1}(\omega)}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)}, & p > \frac{1}{2}, \end{cases} \\ \tilde{\rho}_2(\omega) &= \begin{cases} -\frac{1}{2N} \frac{\tilde{a}_{1-2q}(\omega) + \tilde{a}_{-2q-1}(\omega)}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)}, & q < -\frac{1}{2}, \\ \frac{1}{2N} \frac{\tilde{a}_{2q+1}(\omega) - \tilde{a}_{1-2q}(\omega)}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)}, & -\frac{1}{2} < q < \frac{1}{2}, \\ \frac{1}{2N} \frac{\tilde{a}_{2q-1}(\omega) + \tilde{a}_{2q+1}(\omega)}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)}, & q > \frac{1}{2}. \end{cases} \end{aligned} \quad (53)$$

235 Then the ground state energy (35) can be expressed as

$$E_g = -2N \int_{-\infty}^{\infty} [\tilde{a}_1(\omega) + \tilde{a}_3(\omega)] \tilde{\rho}(\omega) d\omega + 3N + E_0 = N e_g + e_s, \quad (54)$$

236 where e_g is the ground state energy density which is the same as that for the periodic
237 boundary condition [9],

$$e_g = -2 \int_{-\infty}^{\infty} \frac{[\tilde{a}_1(\omega) + \tilde{a}_3(\omega)]^2}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)} d\omega + 3 = -1, \quad (55)$$

238 and e_s is boundary energy

$$e_s = 2\pi - 4 + E_0 + e_1 + e_2, \quad (56)$$

$$e_1 = \begin{cases} -\int_{-\infty}^{\infty} [\tilde{a}_1(\omega) + \tilde{a}_3(\omega)] \frac{\tilde{a}_{2p-1}(\omega) + \tilde{a}_{2p+1}(\omega)}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)} d\omega, & p > \frac{1}{2}, \\ -\int_{-\infty}^{\infty} [\tilde{a}_1(\omega) + \tilde{a}_3(\omega)] \frac{\tilde{a}_{2p+1}(\omega) - \tilde{a}_{1-2p}(\omega)}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)} d\omega, & 0 < p < \frac{1}{2}, \end{cases} \quad (57)$$

$$e_2 = \begin{cases} \int_{-\infty}^{\infty} [\tilde{a}_1(\omega) + \tilde{a}_3(\omega)] \frac{\tilde{a}_{-2q-1}(\omega) + \tilde{a}_{1-2q}(\omega)}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)} d\omega, & q < -\frac{1}{2}, \\ -\int_{-\infty}^{\infty} [\tilde{a}_1(\omega) + \tilde{a}_3(\omega)] \frac{\tilde{a}_{2q+1}(\omega) - \tilde{a}_{1-2q}(\omega)}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)} d\omega, & -\frac{1}{2} < q < \frac{1}{2}, \\ -\int_{-\infty}^{\infty} [\tilde{a}_1(\omega) + \tilde{a}_3(\omega)] \frac{\tilde{a}_{2q-1}(\omega) + \tilde{a}_{2q+1}(\omega)}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)} d\omega, & q > \frac{1}{2}. \end{cases} \quad (58)$$

239 Now, we consider the regime II. The boundary strings pi and $(p-1)i$ can give rise to
240 the rearrangement of Bethe roots in Fermi sea. From BAEs (37), the density of states
241 $\rho_p(u)$ is obtained as

$$\begin{aligned} \rho_p(u) &= a_1(u) + a_3(u) - \int_{-\infty}^{\infty} [2a_2(u-v) + a_4(u-v)] \rho_p(v) dv \\ &\quad + \frac{1}{2N} [a_2(u) - a_{1-2p}(u) + a_{2p+1}(u) + a_{2q-1}(u) + a_{2q+1}(u) - \delta(u)] \\ &\quad - \frac{1}{2N} [2a_{2p+1}(u) + 2a_{3-2p}(u) + a_{3+2p}(u) + a_{5-2p}(u)]. \end{aligned} \quad (59)$$

242 In order to show that there exist the stable boundary bound states, we denote the deviation
243 between $\rho_p(u)$ and $\rho(u)$ as $\delta\rho_p(u) = \rho_p(u) - \rho(u)$. From Eqs.(48) and (59), we obtain

$$\begin{aligned} \delta\rho_p(u) &= -\frac{1}{2N} [2a_{2p+1}(u) + 2a_{3-2p}(u) + a_{3+2p}(u) + a_{5-2p}(u)] \\ &\quad - \int_{-\infty}^{\infty} [2a_2(u-v) + a_4(u-v)] \delta\rho_p(v) dv. \end{aligned} \quad (60)$$

244 Taking the Fourier transformation of Eq.(60), we have

$$\delta\tilde{\rho}_p(\omega) = -\frac{1}{2N} \frac{2\tilde{a}_{2p+1}(\omega) + 2\tilde{a}_{3-2p}(\omega) + \tilde{a}_{3+2p}(\omega) + \tilde{a}_{5-2p}(\omega)}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)}. \quad (61)$$

245 The energy deviation δe_p induced by the density deviation $\delta\tilde{\rho}_p(\omega)$ can be expressed as

$$\begin{aligned} \delta e_p &= -2N \int_{-\infty}^{\infty} [\tilde{a}_1(\omega) + \tilde{a}_3(\omega)] \delta\tilde{\rho}_p(\omega) d\omega + \frac{4}{p^2 - 1} + \frac{4}{(p-1)^2 - 1} \\ &= 2 \int_0^{\infty} \frac{e^{-(p+1)\omega}}{1 + e^{-\omega}} d\omega + 2 \int_0^{\infty} \frac{e^{-(2-p)\omega}}{1 + e^{-\omega}} d\omega + \frac{2}{p(p-1)} < 0. \end{aligned} \quad (62)$$

246 Because of $\delta e_p < 0$, the boundary strings are stable. Then we conclude that in this regime,
247 the ground state energy of the system is $E_g = Ne_g + e_s + \delta e_p$. The total spin along the
248 z -direction is $S_z = -\int_{-\infty}^{\infty} \delta\rho_p(u) = 3/4$.

249 Next, we consider the regime III where boundary strings are qi and $(q-1)i$. Similarly,
250 the energy deviation δe_q between this case and that without boundary strings is

$$\begin{aligned} \delta e_q &= -2N \int_{-\infty}^{\infty} [\tilde{a}_1(\omega) + \tilde{a}_3(\omega)] \delta\tilde{\rho}_q(\omega) d\omega + \frac{4}{p^2 - 1} + \frac{4}{(p-1)^2 - 1} \\ &= 2 \int_0^{\infty} \frac{e^{-(q+1)\omega}}{1 + e^{-\omega}} d\omega + 2 \int_0^{\infty} \frac{e^{-(2-q)\omega}}{1 + e^{-\omega}} d\omega + \frac{2}{q(q-1)} < 0. \end{aligned} \quad (63)$$

251 Due to the fact $\delta e_q < 0$, we know that the ground state energy is $E_g = Ne_g + e_s + \delta e_q$
 252 and the total spin along the z -direction is $S_z = 3/4$.

253 In the regime IV, we combine the results (62) and (63), and conclude that the ground
 254 state energy with boundary strings pi , $(p-1)i$, qi and $(q-1)i$ equals to $E_g = Ne_g + e_s +$
 255 $\delta e_p + \delta e_q$.

256 Then, we consider the regime V where besides the $N-2$ 2-string, there also exist one
 257 real Bethe root λ_0 and a single boundary string qi . Taking the thermodynamic limit of
 258 BAEs (44), we obtain the density of states $\rho_{\lambda q}(u)$ as

$$\begin{aligned} \rho_{\lambda q}(u) &= a_1(u) + a_3(u) - \frac{1}{2N} [a_1(u - \lambda_0) + a_1(u + \lambda_0) + a_3(u - \lambda_0) + a_3(u + \lambda_0)] \\ &\quad + \frac{1}{2N} [a_2(u) + a_{2p-1}(u) + a_{2p+1}(u) - 2a_{1-2q}(u) - a_{3+2q}(u) - a_{3-2q}(u) - \delta(u)] \\ &\quad - \int_{-\infty}^{\infty} [2a_2(u-v) + a_4(u-v)] \rho_{\lambda q}(v) dv. \end{aligned} \quad (64)$$

259 Denote the deviation between $\rho_{\lambda q}(u)$ and $\rho(u)$ as $\delta\rho_{\lambda q}(u) = \rho_{\lambda q}(u) - \rho(u)$. From Eqs.(48)
 260 and (64), the value of $\delta\rho_{\lambda q}(u)$ reads

$$\begin{aligned} \delta\rho_{\lambda q}(u) &= -\frac{1}{2N} [a_1(u - \lambda_0) + a_1(u + \lambda_0) + a_3(u - \lambda_0) + a_3(u + \lambda_0)] \\ &\quad - \frac{1}{2N} [a_{1-2q}(u) - a_{-1-2q}(u) + a_{3-2q}(u) + a_{3+2q}(u)] \\ &\quad - \int_{-\infty}^{\infty} [2a_2(u) + a_4(u)] \delta\rho_{\lambda q}(v) dv. \end{aligned} \quad (65)$$

261 Taking the Fourier transformation of Eq.(65), we obtain

$$\delta\tilde{\rho}_{\lambda q}(\omega) = -\frac{1}{2N} \frac{\tilde{a}_{1-2q}(\omega) - \tilde{a}_{-1-2q}(\omega) + \tilde{a}_{3-2q}(\omega) + \tilde{a}_{3+2q}(\omega)}{1 + 2\tilde{a}_2(\omega) + \tilde{a}_4(\omega)} - \frac{1}{N} \frac{\cos(\omega\lambda_0)e^{-\frac{|\omega|}{2}}}{1 + e^{-|\omega|}}. \quad (66)$$

262 Then the deviation of energy $\delta e_{\lambda q}$ induced by $\delta\tilde{\rho}_{\lambda q}(\omega)$ is given by

$$\begin{aligned} \delta e_{\lambda q} &= -2N \int_{-\infty}^{\infty} [\tilde{a}_1(\omega) + \tilde{a}_3(\omega)] \delta\tilde{\rho}_{\lambda q}(\omega) d\omega + \frac{4}{q^2 - 1} - \frac{4}{\lambda_0^2 + 1} \\ &= 2 \int_0^{\infty} \frac{e^{-(2+q)\omega}}{1 + e^{-\omega}} d\omega - 2 \int_0^{\infty} \frac{e^{q\omega}}{1 + e^{-\omega}} d\omega - \frac{2}{1+q} < 0. \end{aligned} \quad (67)$$

263 Due to $\delta e_{\lambda q} < 0$, the ground state energy in this regime is $E_g = Ne_g + e_s + \delta e_{\lambda q}$ and the
 264 total spin along the z -direction is $S_z = 3/4$.

265 In the regime VI, there are $N-4$ 2-string, one real Bethe root λ_0 and three boundary
 266 strings qi , pi and $(p-1)i$. Combining the results (62) and (67), we obtain the ground
 267 state energy as $E_g = Ne_g + e_s + \delta e_p + \delta e_{\lambda q}$.

268 After tedious calculation, we find that the boundary energy e_b for all the regimes in
 269 Figure 1 can be expressed as

$$e_b = \begin{cases} -\frac{2}{p} - \frac{2}{q} + 2\pi - 4 + E_0, & p > 0, q > 0 \text{ or } q < -1, \\ -\frac{2}{p} - \frac{2}{q} + 2\pi \csc(q\pi) + 2\pi - 4 + E_0, & p > 0, -1 < q < 0. \end{cases} \quad (68)$$

270 The boundary energies with different boundary parameters p and q calculated by the
 271 analytical expression (68) are shown in Figure 3 as the coloured solid lines. Now we check




Figure 3: Boundary energies versus the boundary parameters p and q . The coloured curves are those calculated from the analytical expression (68) and the red points are those obtained from the DMRG. The values of q at the red points are $q = -2.6, -2.1, -1.7, -1.3, -0.7, -0.5, -0.25, 0.35, 0.7, 1.15, 1.5$ and 1.8 .




Figure 4: The values of $e_b(N)$ versus the system size N . The red points are the DMRG results with $N = 4, 14, 24, \dots, 194$. The data can be fitted as $e_b(N) = aN^b + c$, where $a = 6.7308$, $b = -1.0046$ and $c = 1.5460$. Due to the fact $b < 0$, when the system size $N \rightarrow \infty$, the values of $e_b(N)$ tend to the asymptotic value c , which gives the boundary energy. Here the boundary parameters are chosen as $p = 0.3$ and $q = 0.7$.

272 the correction of expression (68) by the numerical simulation with DMRG algorithm, and
 273 the results are shown in Figure 3 as the red points. Specifically, for each red point that
 274 is for the given boundary parameters p and q , we first calculate the ground state energy
 275 $E_g(N)$ of the model (15) with the system size $N = 10(n - 1) + 4$ and $n = 1, 2, \dots, 20$ by
 276 using the DMRG method. Then we consider the physical quantity

$$e_b(N) = E_g(N) - Ne_g, \quad (69)$$

277 where $e_g = -1$ is the ground state energy density of the system with periodic boundary
 278 condition. Obviously, in the thermodynamic limit, the value of $e_b(N \rightarrow \infty)$ gives the
 279 boundary energy. In Figure 4, we show how to extrapolate the boundary energy, where
 280 the red points are the numerical values of $e_b(N)$, the blue solid line is the fitting curve,
 281 and the red solid line is the extrapolated boundary energy. From the fitting curve, we
 282 find that the $e_b(N)$ and N satisfy the power law relation, i.e., $e_b(N) = aN^b + c$. Due to
 283 the fact that $b < 0$, the values of $e_b(N)$ tend to the asymptotic value c when the system
 284 size N tends to infinity. Therefore, in the thermodynamic limit, the asymptotic value c

285 determines the boundary energy. Repeating this process, we obtain the boundary energies
286 with other values of boundary parameters. As shown in Figure 3, the analytical and
287 numerical results agree with each other very well.

288 5 Conclusions

289 In this paper, we have studied the thermodynamic limit and boundary energy of the
290 isotropic spin-1 Heisenberg chain with generic integrable non-diagonal boundary reflec-
291 tions. It is shown that the contribution of inhomogeneous term in the associated $T - Q$
292 relation (18) (due to the unparallel boundary fields) can be neglected only in the ther-
293 modynamic limit. This fact allows us to calculate the boundary energy (68) induced by
294 the unparallel boundary magnetic fields. For the case of $\alpha_{\pm} = 0$, our result gives rise to
295 boundary energy corresponding to two parallel boundary fields which might be obtained
296 by the algebraic Bethe ansatz and thermodynamic Bethe ansatz. The method provided in
297 this paper can be used to study the thermodynamic properties of other quantum integrable
298 models associated with rational R -matrix.

299 Acknowledgments

300 Financial support from the National Natural Science Foundation of China (Grant Nos.
301 12074410, 12047502, 11934015, 11975183, 11947301, 11775177, 11775178 and 11774397),
302 the Strategic Priority Research Program of the Chinese Academy of Sciences (Grant No.
303 XDB33000000), the Major Basic Research Program of Natural Science of Shaanxi Province
304 (Grant Nos. 2017KCT-12 and 2017ZDJC-32), and the Double First-Class University Con-
305 struction Project of Northwest University is gratefully acknowledged. One of the authors,
306 Zhihan Zheng would like to thank Dr. Yangyang Chen, Dr. Fakai Wen and Dr. Yi Qiao
307 for their helpful discussions.

References

- [1] X.-W. Guan, M. T. Batchelor and C. Lee, *Fermi gases in one dimension: From Bethe ansatz to experiments*, Rev. Mod. Phys. **85**, 1633 (2013), doi:10.1103/RevModPhys.85.1633.
- [2] M. Takahashi, *Thermodynamics of one-dimensional solvable models*, Cambridge University Press, (1999), doi:10.1017/CBO9780511524332.
- [3] L. Dolan, C. R. Nappi and E. Witten, *A relation between approaches to integrability in superconformal Yang-Mills theory*, J. High Energ. Phys. **10**, 017 (2003), doi:10.1088/1126-6708/2003/10/017.
- [4] J. Sirker, R. G. Pereira, and I. Affleck, *Diffusion and ballistic transport in one-dimensional quantum systems*, Phys. Rev. Lett. **103**, 216602 (2009), doi:10.1103/PhysRevLett.103.216602.
- [5] J. de Gier and F. H. L. Essler, *Bethe ansatz solution of the asymmetric exclusion process with open boundaries*, Phys. Rev. Lett. **95**, 240601 (2005), doi:10.1103/PhysRevLett.95.240601.

- [6] H. A. Bethe, *Zur Theorie der Metalle. i. Eigenwerte und eigenfunktionen der linearen atomkette*, Zeit. für Phys. **71**, 205 (1931), doi:10.1007%2F01341708.
- [7] A. B. Zamolodchikov and V. A. Fateev, *Model factorized s matrix and an integrable heisenberg chain with spin 1., in russian*, Sov. J. Nucl. Phys. **32**, 298 (1980).
- [8] L. A. Takhtajan, *The picture of low-lying excitations in the isotropic Heisenberg chain of arbitrary spins*, Phys. Lett. A **87**, 479 (1982), doi:10.1016/0375-9601(82)90764-2.
- [9] H. M. Babujian, *Exact solution of the isotropic Heisenberg chain with arbitrary spins: Thermodynamics of the model*, Nucl. Phys. B **215**, 217 (1983), doi:10.1016/0550-3213(83)90668-5.
- [10] H. M. Babujian, *Exact solution of the one-dimensional isotropic Heisenberg chain with arbitrary spins s*, Phys. Lett. A **90**, 479 (1982), doi:10.1016/0375-9601(82)90403-0.
- [11] F. D. M. Haldane, *Continuum dynamics of the 1-D Heisenberg antiferromagnet: Identification with the O(3) nonlinear sigma model*, Phys. Lett. **93A**, 464 (1983), doi:10.1016/0375-9601(83)90631-X.
- [12] F. D. M. Haldane, *Nonlinear field theory of large-spin Heisenberg antiferromagnets: Semiclassically quantized solitons of the one-dimensional easy-axis Neel state*, Phys. Rev. Lett. **50**, 1153 (1983), doi:10.1103/PhysRevLett.50.1153.
- [13] I. Affleck, T. Kennedy, E. H. Lieb and H. Tasaki, *Rigorous results on valence-bond ground states in antiferromagnets*, Phys. Rev. Lett. **59**, 799 (1987), doi:10.1103/PhysRevLett.59.799.
- [14] I. Affleck, T. Kennedy, E. H. Lieb and H. Tasaki, *Valence bond ground states in isotropic quantum antiferromagnets*, Commun. Math. Phys. **115**, 477 (1988), doi:10.1007/BF01218021.
- [15] M. N. Barber and M. T. Batchelor, *Spectrum of the biquadratic spin-1 antiferromagnetic chain*, Phys. Rev. B **40**, 4621 (1989), doi:10.1103/PhysRevB.40.4621.
- [16] A. Klümper, *The spectra of q-state vertex models and related antiferromagnetic quantum spin chains*, J. Phys. A: Math. Gen. **23**, 809 (1990), doi:10.1088/0305-4470/23/5/023.
- [17] K.-J.-B. Lee and P. Schlottmann, *Integrable spin-1 Heisenberg chain with impurity*, Phys. Rev. B **37**, 379 (1988), doi:10.1103/PhysRevB.37.379.
- [18] F. C. Alcaraz, M. N. Barber, M. T. Batchelor, R. J. Baxter and G. R. W. Quispel, *Surface exponents of the quantum XXZ, Ashkin-Teller and Potts models*, J. Phys. A **20**, 6397 (1987), doi:10.1088/0305-4470/20/18/038.
- [19] E. K. Sklyanin, *Boundary conditions for integrable quantum systems*, J. Phys. A **21**, 2375 (1988), doi:10.1088/0305-4470/21/10/015.
- [20] P. P. Kulish and E. K. Sklyanin, *Quantum spectral transform method recent developments*, Lect. Notes in Phys. **151**, 61 (1982), doi:10.1007/3-540-11190-5_8.
- [21] P. P. Kulish, N. Yu. Reshetikhin and E. K. Sklyanin, *Yang-Baxter equation and representation theory: 1*, Lett. Math. Phys. **5**, 393 (1981), doi:10.1142/9789812798336_0027.

- [22] P. P. Kulish and N. Yu. Reshetikhin, *Quantum linear problem for the sine-Gordon equation and higher representations*, J. Sov. Math. **23**, 2435 (1983), doi:10.1007/BF01084171.
- [23] A. N. Kirillov and N. Yu. Reshetikhin, *Exact solution of the Heisenberg XXZ model of spin s* , J. Sov. Math. **35**, 2627 (1986), doi:10.1007/BF01083768.
- [24] A. N. Kirillov and N. Yu. Reshetikhin, *Exact solution of the integrable XXZ Heisenberg model with arbitrary spin. I. The ground state and the excitation spectrum*, J. Phys. A **20**, 1565 (1987), doi:10.1088/0305-4470/20/6/038.
- [25] L. A. Takhtajan, *The picture of low-lying excitations in the isotropic Heisenberg chain of arbitrary spins*, Phys. Lett. A **87**, 479 (1982), doi:10.1016/0375-9601(82)90764-2.
- [26] L. Mezincescu, R.I. Nepomechie and V. Rittenberg, *Bethe ansatz solution of the Fateev-Zamolodchikov quantum spin chain with boundary terms*, Phys. Lett. A **147**, 70 (1990), doi:10.1016/0375-9601(90)90016-H.
- [27] E. C. Fireman, A. Lima-Santos and W. Utiel, *Bethe ansatz solution for quantum spin-1 chains with boundary terms*, Nucl. Phys. B **626**, 435 (2002), doi:10.1016/S0550-3213(02)00027-5.
- [28] A. Doikou, *Fused integrable lattice models with quantum impurities and open boundaries*, Nucl. Phys. B **668**, 447 (2003), doi:10.1016/j.nuclphysb.2003.07.001.
- [29] C.S. Melo, G.A.P. Ribeiro and M.J. Martins, *Bethe ansatz for the XXX-S chain with non-diagonal open boundaries*, Nucl. Phys. B **711**, 565 (2005), doi:10.1016/j.nuclphysb.2004.12.008.
- [30] L. Frappat, R.I. Nepomechie and É. Ragoucy, *Complete Bethe ansatz solution of the open spin- s XXZ chain with general integrable boundary terms*, J. Stat. Mech. **0709**, P09009 (2007), doi:10.1088/1742-5468/2007/09/P09009.
- [31] R. Murgan, *Bethe ansatz of the open spin- s XXZ chain with nondiagonal boundary terms*, J. High Energ. Phys. **04**, 076 (2009), doi:10.1088/1126-6708/2009/04/076.
- [32] R. Baiyasi and R. Murgan, *Generalized $T - Q$ relations and the open spin- s XXZ chain with nondiagonal boundary terms*, J. Stat. Mech. **1210**, P10003 (2012), doi:10.1088/1742-5468/2012/10/P10003.
- [33] J. Cao, W.-L. Yang, K. Shi and Y. Wang, *Off-diagonal Bethe ansatz solution of the XXX spin-chain with arbitrary boundary conditions*, Nucl. Phys. B **875**, 152 (2013), doi:10.1016/j.nuclphysb.2013.06.022.
- [34] R.I. Nepomechie, *An inhomogeneous $T - Q$ equation for the open XXX chain with general boundary terms: completeness and arbitrary spin*, J. Phys. A: Math. Theor. **46**, 442002 (2013), doi:10.1088/1751-8113/46/44/442002.
- [35] R.I. Nepomechie and C. Wang, *Boundary energy of the open XXX chain with a non-diagonal boundary term*, J. Phys. A: Math. Theor. **47**, 032001 (2014), doi:10.1088/1751-8113/47/3/032001.
- [36] Y. Wang, W.-L. Yang, J. Cao and K. Shi, *Off-diagonal Bethe ansatz for exactly solvable models*, Springer Press, (2015), doi:10.1007/978-3-662-46756-5.

- [37] Y.-Y. Li, J. Cao, W.-L. Yang, K. Shi and Y. Wang, *Thermodynamic limit and surface energy of the XXZ spin chain with arbitrary boundary fields*, Nucl. Phys. B **884**, 17 (2014), doi:10.1016/j.nuclphysb.2014.04.010.
- [38] F. Wen, T. Yang, Z.-Y. Yang, J. Cao, K. Hao and W.-L. Yang, *Thermodynamic limit and boundary energy of the $su(3)$ spin chain with non-diagonal boundary fields*, Nucl. Phys. B **915**, 119 (2017), doi:10.1016/j.nuclphysb.2016.12.003.
- [39] C.-N. Yang, *Some exact results for the many-body problem in one dimension with repulsive delta-function interaction*, Phys. Rev. Lett. **19**, 1312 (1967), doi:10.1103/PhysRevLett.19.1312.
- [40] R. J. Baxter, *Exactly solved models in statistical mechanics*, Academic Press, London, (1982), doi:10.1142/9789814415255_0002.
- [41] A. Kapustin, S. Skorik, *Surface excitations and surface energy of the antiferromagnetic XXZ chain by the Bethe ansatz approach*, J. Phys. A **29**, 1629 (1996), doi:10.1088/0305-4470/29/8/011.
- [42] S. R. White, *Density matrix formulation for quantum renormalization groups*, Phys. Rev. Lett. **69**, 2863 (1992), doi:10.1103/PhysRevLett.69.2863.
- [43] S. R. White, *Density-matrix algorithms for quantum renormalization groups*, Phys. Rev. B **48**, 10345 (1993), doi:10.1103/PhysRevB.48.10345.
- [44] U. Schollwöck, *The density-matrix renormalization group*, Rev. Mod. Phys. **77**, 259 (2005), doi:10.1103/RevModPhys.77.259.
- [45] M. Gaudin, *Boundary energy of a Bose gas in one dimension*, Phys. Rev. A **4**, 386 (1971), doi:10.1103/PhysRevA.4.386.
- [46] C. J. Hamer, G.R.W. Quispel and M. T. Batchelor, *Conformal anomaly and surface energy for Potts and Ashkin-Teller quantum chains*, J. Phys. A **20**, 5677 (1987), doi:10.1088/0305-4470/20/16/040.
- [47] M. T. Batchelor and C. J. Hamer, *Surface energy of integrable quantum spin chains*, J. Phys. A **23**, 761 (1990), doi:10.1088/0305-4470/23/5/019.
- [48] P. Fendley and H. Saleur, *Deriving boundary S matrices*, Nucl. Phys. B **428**, 681 (1994), doi:10.1016/0550-3213(94)90369-7.