

Longitudinal Z-boson polarization and the Higgs boson production cross-section

S. Amoroso^{1*}, for the xFitter Collaboration

¹ Deutsches Elektronen-Synchrotron DESY, D 22607, Hamburg

* simone.amoroso@desy.de

July 31, 2021



*Proceedings for the XXVIII International Workshop
on Deep-Inelastic Scattering and Related Subjects,
Stony Brook University, New York, USA, 12-16 April 2021*
doi:[10.21468/SciPostPhysProc.?](https://doi.org/10.21468/SciPostPhysProc.)

Abstract

Lepton pairs are produced copiously in high-energy hadron collisions via electroweak gauge boson exchange, and are one of the most precisely measured final states in proton-proton collisions at the Large Hadron Collider (LHC). We propose that measurements of lepton angular distributions can be used to improve the accuracy of theoretical predictions for Higgs boson production cross sections at the LHC. To this end, we exploit the sensitivity of the lepton angular coefficient associated with the longitudinal Z-boson polarization to the parton density function (PDF) for gluons resolved from the incoming protons, in order to constrain the Higgs boson cross section from gluon fusion processes. By a detailed numerical analysis using the open-source platform xFitter, we find that high-statistics determinations of the longitudinally polarized angular coefficient at the LHC Run III and high-luminosity HL-LHC improve the PDF systematics of the Higgs boson cross section predictions by 50 % over a broad range of Higgs boson rapidities.

1 Introduction

The characterisation of the Higgs sector and the investigation of the electroweak symmetry breaking mechanism are among the main physics goals of the LHC and future High Luminosity (HL) LHC. The uncertainties from the knowledge of the parton distribution functions of the proton constitute one of the main limitations to fully exploit its physics potential [1]. In particular, the Higgs cross-section in its dominant gluon-gluon fusion production channel is extremely sensitive to the gluon PDF at intermediate Bjorken- x ranges. Currently, the gluon PDF is determined indirectly from scaling violations in deep-inelastic scattering data, with additional sensitivity coming from LHC measurements such as jet and top quark production. These latter two processes are however challenging to measure, and suffer from theoretical ambiguities making them difficult to predict to high accuracy [2, 3]. In this work [4], we propose a novel determination of the gluon PDF at intermediate Bjorken- x values using neutral current Drell-Yan data and an experimentally clean and easy to measure observable, which can be predicted theoretically to very high accuracy and exhibits a good perturbative stability. The analysis has been performed using xFitter [5, 7], an open-source package allowing to extract PDFs and other QCD parameters. It implements a large number of different datasets, and has been used in a variety of QCD studies; most recently an extraction of the pion PDF [8] and a study of the impact of forward-backward asymmetries in Drell-Yan on PDFs [9].

2 The A_0 angular coefficient

We consider Drell-Yan production [10] via Z/γ^* boson exchange at the LHC. The cross section summed over the electroweak boson polarizations has the angular distribution $1 + \cos^2 \theta$ and is sensitive to the gluon PDF for finite p_T . However, in the small- p_T region where the cross-section is the largest it is sensitive to large logarithmic corrections, while at high- p_T the missing higher order uncertainties become large. In this study we focus on the leptonic angular distributions through the angular coefficient A_0 , the ratio of the longitudinal electroweak boson cross section to the unpolarized cross section:

$$A_0(s, M, Y, p_T) = \frac{2d\sigma^{(L)}/dMdYdp_T}{d\sigma/dMdYdp_T}. \quad (1)$$

It associated with the $(3 \cos^2 \theta - 1)/2$ angular dependence of the cross-section on the polar angle θ^* [11], with θ^* measured in the Collins-Soper frame [12]. The longitudinally polarized coefficient A_0 in Eq. (1) vanishes in the parton model and receives leading-order (LO) perturbative QCD contributions at $\mathcal{O}(\alpha_s)$. It has been computed up to $\mathcal{O}(\alpha_s^3)$ [13], showing a very good perturbative stability. The A_0 coefficient is parity-conserving and sensitive to flavor singlet PDFs. The main sensitivity to the gluon distribution arises from the region where the change of A_0 with p_T is largest, i.e., around the turn-over point $\partial^2 A_0 / \partial p_T^2 = 0$. For dilepton masses near the Z -boson peak the turn-over occurs for boson p_T of the order of several ten to 100 GeVs.

The angular coefficients in Drell-Yan, and A_0 , have been measured by many fixed-target and collider experiments. We consider the A_0 distribution from the ATLAS measurement of the angular coefficients in Z boson production at $\sqrt{s} = 8$ TeV [14]. The measurement is performed in 23 p_T bins (we remove the first bins where our fixed-order predictions would not be adequate), going from 11.4 GeV to 600 GeV, and in three different rapidity bins: $0 < |Y| < 1$, $1 < |Y| < 2$, and $2 < |Y| < 3.5$. Predictions at NLO in QCD for finite boson p_T (at order $\mathcal{O}(\alpha^2)$) have been produced with mg5_amc@NLO and aMCfast [15]. A χ^2 is constructed, incorporating all experimental uncertainties and their correlations, as well as PDF uncertainties. The impact of the A_0 data on PDFs is then evaluated through an Hessian profiling technique [16,17]. We find a good description of the data, with χ^2/ndf of order one, for all modern PDFs: CT18NNLO [18], NNPDF3.1nnlo [19], ABMP16nnlo [20], and MSHT20nnlo [21]. The reduction in the PDF uncertainties after the inclusion of the A_0 data is however found to be negligible.

3 PDF constraints and the Higgs cross-section

3.1 Sensitivity to the gluon PDF

We then generate pseudodata for the A_0 distribution at $\sqrt{s} = 13$ TeV for two projected luminosity scenarios of 300 fb^{-1} (the expected integrated luminosity at the end of the LHC Run III) and 3 ab^{-1} (the design integrated luminosity of the HL-LHC stage). To do this we extrapolate the statistical uncertainties, and add a conservative estimate of 0.1% for the dominant experimental systematic on the lepton momentum scale. We perform a PDF profiling exercise using the CT18NNLO PDF set, and find that the increased statistics allows for a significant reduction in the PDF uncertainties. This reduction is particularly large for the gluon density and for the u and d sea-quark densities coupled to gluons through QCD evolution, as shown in Fig. 1. The largest sensitivity is found to come from transverse momenta in the range $p_T \sim 50$ GeV, dying out for $p_T > 100$ GeV. The gain from a 300 fb^{-1} measurement is found to dominate over the

3 ab^{-1} gain, although a small improvement is nonetheless present.




Figure 1: The gluon (xg) and gluon/Sea (xg/Σ) PDFs in the original CT18NNLO analysis (red) and profiled using A_0 pseudodata corresponding to integrated luminosities of 300 fb^{-1} (blue) and 3 ab^{-1} (green) in the region $80 \text{ GeV} < M < 100 \text{ GeV}$ and $|Y| < 3.5$.

3.2 Impact on Higgs phenomenology

The effect of the longitudinally polarized coefficient on the $Q^2 = 10^4 \text{ GeV}^2$ gluon PDF near $x \sim 10^{-2}$ will influence the Higgs boson cross section. In Fig. 2 we show the gluon-gluon luminosity for the CT18NNLO PDF, before and after the profiling using A_0 pseudodata. Most of the reduction in the uncertainties is found in the Higgs mass range. In the right plot we show the N^3LO gluon-fusion Higgs total cross section and its uncertainty obtained with `ggHiggs` [22, 23]. for the CT18NNLO, NNPDF3.1nnlo and MSHT20nnlo global PDF sets. Notwithstanding the numerical differences, a similar reduction in the PDF uncertainties is seen for the different sets. We also show The PDF4LHC15scen1 and 2 projected sets, which include pseudodata based on complete LHC data sample [24], which show a smaller, but not negligible, reduction in uncertainties.

4 Conclusion

We have studied the impact of precise measurements of the A_0 angular coefficient in neutral current Drell-Yan on the uncertainties of the gluon PDF and on the Higgs boson production cross-section in gluon-gluon fusion. We found that future measurements of A_0 have the possibility to reduce by about a factor of two the uncertainty of the gluon PDFs at intermediate Bjorken- x values. This translates in a large reduction in the PDF uncertainty on the gluon-fusion Higgs production cross-section. Our results open a new area of phenomenological studies on connections between the gauge and Higgs sectors of the SM. Further aspects may be investigated by extending the analysis to the full structure of lepton angular distributions, including polarization interferences, and to mass regions far away from the Z -boson peak.




Figure 2: Ratio of PDF uncertainties for the gluon-gluon luminosity evaluated at $\sqrt{s} = 13$ TeV. The red band shows the uncertainties of the CT18NNLO PDF set [18], reduced to 68% CL coverage. The blue and green bands show the uncertainties of the CT18NNLO including constraints from the A_0 measurement and assuming 300 fb^{-1} and 3 ab^{-1} , respectively.

References

- [1] M. Cepeda *et al.*, *Report from Working Group 2: Higgs Physics at the HL-LHC and HE-LHC*, CERN Yellow Rep. Monogr. **7**, 221 (2019), doi:[10.23731/CYRM-2019-007.221](https://doi.org/10.23731/CYRM-2019-007.221), [1902.00134](https://arxiv.org/abs/1902.00134).
- [2] R. Abdul Khalek *et al.*, *Phenomenology of NNLO jet production at the LHC and its impact on parton distributions*, Eur. Phys. J. C **80**(8), 797 (2020), doi:[10.1140/epjc/s10052-020-8328-5](https://doi.org/10.1140/epjc/s10052-020-8328-5), [2005.11327](https://arxiv.org/abs/2005.11327).
- [3] S. Bailey and L. Harland-Lang, *Differential Top Quark Pair Production at the LHC: Challenges for PDF Fits*, Eur. Phys. J. C **80**(1), 60 (2020), doi:[10.1140/epjc/s10052-020-7633-3](https://doi.org/10.1140/epjc/s10052-020-7633-3), [1909.10541](https://arxiv.org/abs/1909.10541).
- [4] S. Amoroso, J. Fiaschi, F. Giuli, A. Glazov, F. Hautmann and O. Zenaiev, *Longitudinal Z-Boson Polarization and the Higgs Boson Production Cross Section at the Large Hadron Collider* (2020), [2012.10298](https://arxiv.org/abs/2012.10298).
- [5] S. Alekhin *et al.*, *HERAFitter*, Eur. Phys. J. C **75**(7), 304 (2015), doi:[10.1140/epjc/s10052-015-3480-z](https://doi.org/10.1140/epjc/s10052-015-3480-z), [1410.4412](https://arxiv.org/abs/1410.4412).
- [6] xFitter Developers.
- [7] I. Novikov *et al.*, *Parton Distribution Functions of the Charged Pion Within The xFitter Framework*, Phys. Rev. D **102**(1), 014040 (2020), doi:[10.1103/PhysRevD.102.014040](https://doi.org/10.1103/PhysRevD.102.014040), [2002.02902](https://arxiv.org/abs/2002.02902).
- [8] E. Accomando *et al.*, *PDF Profiling Using the Forward-Backward Asymmetry in Neutral Current Drell-Yan Production*, JHEP **10**, 176 (2019), doi:[10.1007/JHEP10\(2019\)176](https://doi.org/10.1007/JHEP10(2019)176), [1907.07727](https://arxiv.org/abs/1907.07727).
- [9] S. D. Drell and T.-M. Yan, *Massive Lepton Pair Production in Hadron-Hadron Collisions at High-Energies*, Phys. Rev. Lett. **25**, 316 (1970), doi:[10.1103/PhysRevLett.25.316](https://doi.org/10.1103/PhysRevLett.25.316), [Erratum: Phys.Rev.Lett. 25, 902 (1970)].

- [10] E. Mirkes and J. Ohnemus, *Angular distributions of Drell-Yan lepton pairs at the Tevatron: Order $\alpha - s^2$ corrections and Monte Carlo studies*, Phys. Rev. D **51**, 4891 (1995), doi:[10.1103/PhysRevD.51.4891](https://doi.org/10.1103/PhysRevD.51.4891), [hep-ph/9412289](https://arxiv.org/abs/hep-ph/9412289).
- [11] J. C. Collins and D. E. Soper, *Angular Distribution of Dileptons in High-Energy Hadron Collisions*, Phys. Rev. D **16**, 2219 (1977), doi:[10.1103/PhysRevD.16.2219](https://doi.org/10.1103/PhysRevD.16.2219).
- [12] R. Gauld, A. Gehrmann-De Ridder, T. Gehrmann, E. W. N. Glover and A. Huss, *Precise predictions for the angular coefficients in Z-boson production at the LHC*, JHEP **11**, 003 (2017), doi:[10.1007/JHEP11\(2017\)003](https://doi.org/10.1007/JHEP11(2017)003), [1708.00008](https://arxiv.org/abs/1708.00008).
- [13] G. Aad *et al.*, *Measurement of the angular coefficients in Z-boson events using electron and muon pairs from data taken at $\sqrt{s} = 8$ TeV with the ATLAS detector*, JHEP **08**, 159 (2016), doi:[10.1007/JHEP08\(2016\)159](https://doi.org/10.1007/JHEP08(2016)159), [1606.00689](https://arxiv.org/abs/1606.00689).
- [14] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli and M. Zaro, *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, JHEP **07**, 079 (2014), doi:[10.1007/JHEP07\(2014\)079](https://doi.org/10.1007/JHEP07(2014)079), [1405.0301](https://arxiv.org/abs/1405.0301).
- [15] H. Paukkunen and P. Zurita, *PDF reweighting in the Hessian matrix approach*, JHEP **12**, 100 (2014), doi:[10.1007/JHEP12\(2014\)100](https://doi.org/10.1007/JHEP12(2014)100), [1402.6623](https://arxiv.org/abs/1402.6623).
- [16] S. Camarda *et al.*, *QCD analysis of W- and Z-boson production at Tevatron*, Eur. Phys. J. C **75**(9), 458 (2015), doi:[10.1140/epjc/s10052-015-3655-7](https://doi.org/10.1140/epjc/s10052-015-3655-7), [1503.05221](https://arxiv.org/abs/1503.05221).
- [17] T.-J. Hou *et al.*, *New CTEQ global analysis of quantum chromodynamics with high-precision data from the LHC*, Phys. Rev. D **103**(1), 014013 (2021), doi:[10.1103/PhysRevD.103.014013](https://doi.org/10.1103/PhysRevD.103.014013), [1912.10053](https://arxiv.org/abs/1912.10053).
- [18] R. D. Ball *et al.*, *Parton distributions from high-precision collider data*, Eur. Phys. J. C **77**(10), 663 (2017), doi:[10.1140/epjc/s10052-017-5199-5](https://doi.org/10.1140/epjc/s10052-017-5199-5), [1706.00428](https://arxiv.org/abs/1706.00428).
- [19] S. Alekhin, J. Blümlein, S. Moch and R. Placakyte, *Parton distribution functions, α_s , and heavy-quark masses for LHC Run II*, Phys. Rev. D **96**(1), 014011 (2017), doi:[10.1103/PhysRevD.96.014011](https://doi.org/10.1103/PhysRevD.96.014011), [1701.05838](https://arxiv.org/abs/1701.05838).
- [20] S. Bailey, T. Cridge, L. A. Harland-Lang, A. D. Martin and R. S. Thorne, *Parton distributions from LHC, HERA, Tevatron and fixed target data: MSHT20 PDFs*, Eur. Phys. J. C **81**(4), 341 (2021), doi:[10.1140/epjc/s10052-021-09057-0](https://doi.org/10.1140/epjc/s10052-021-09057-0), [2012.04684](https://arxiv.org/abs/2012.04684).
- [21] M. Bonvini, R. D. Ball, S. Forte, S. Marzani and G. Ridolfi, *Updated Higgs cross section at approximate N^3LO* , J. Phys. G **41**, 095002 (2014), doi:[10.1088/0954-3899/41/9/095002](https://doi.org/10.1088/0954-3899/41/9/095002), [1404.3204](https://arxiv.org/abs/1404.3204).
- [22] M. Bonvini and S. Marzani, *Double resummation for Higgs production*, Phys. Rev. Lett. **120**(20), 202003 (2018), doi:[10.1103/PhysRevLett.120.202003](https://doi.org/10.1103/PhysRevLett.120.202003), [1802.07758](https://arxiv.org/abs/1802.07758).
- [23] R. Abdul Khalek, S. Bailey, J. Gao, L. Harland-Lang and J. Rojo, *Towards Ultimate Parton Distributions at the High-Luminosity LHC*, Eur. Phys. J. C **78**(11), 962 (2018), doi:[10.1140/epjc/s10052-018-6448-y](https://doi.org/10.1140/epjc/s10052-018-6448-y), [1810.03639](https://arxiv.org/abs/1810.03639).