

Multiple relaxation times in perturbed XXZ chain

M. Mierzejewski¹, J. Pawłowski¹, P. Prelovšek^{2,3}, and J. Herbrych^{1*}

¹ Department of Theoretical Physics, Faculty of Fundamental Problems of Technology,
Wrocław University of Science and Technology, 50-370 Wrocław, Poland

² Jožef Stefan Institute, SI-1000 Ljubljana, Slovenia

³ Faculty of Mathematics and Physics, University of Ljubljana, SI-1000 Ljubljana, Slovenia

* jacek.herbrych@pwr.edu.pl

December 15, 2021

Abstract

We study numerically the relaxation of correlation functions in weakly perturbed integrable XXZ chain. While the decay of spin-current and energy-current correlations at zero magnetization are well described by single, but quite distinct, relaxation rates governed by the square of the perturbation strength g , the correlations at finite magnetization reveal multiple relaxation rates. The result can be understood in terms of decays of several quantities, conserved in the reference integrable system. On the other hand, the correlations of non-commuting quantities, being conserved at particular anisotropies Δ , decay non-exponentially with characteristic time scale linear in g .

Contents

1	Introduction	1
2	Relaxation of spin and energy currents	3
3	Memory-function analysis of the energy-current correlations	6
4	Dependence on the form of perturbation	7
5	Non-commuting conserved quantities and their non-exponential relaxation	8
6	Conclusions	10
	References	11

1 Introduction

Integrable quantum many-body systems attract a lot of attention due to their unique properties, as well as due to development of new analytical and numerical tools to deal with them (for a recent review see Ref. [1]). The crucial role in the behavior of such systems is played by the presence of extensive number of local and/or quasilocal conserved quantities (CQ), which

have important consequences for the (lack of) relaxation of observables and for the transport properties. The latter consequences are formally expressed via the Mazur bounds which relate the long-time correlations (stiffness) of observables with their projections on the local CQ (charges) [2, 3]. However, the microscopic models are integrable only for fine-tuned sets of parameters, while more realistic systems might be described in terms of nearly integrable (NI) models which contain small but non-vanishing perturbation that breaks the integrability [4, 5]. An important question concerns the details of the integrability breaking, in particular whether the asymptotic dynamics becomes consistent with the generic dissipative diffusive-type transport [6–8]. So far, the properties of NI models are better understood at intermediate time-scales, when the dynamics resembles that of integrable models, the phenomenon known as prethermalization [9–11].

The problem of asymptotic dynamics of NI models at long time scales [12, 13] appears to be more complex. It has been argued that the time evolution can be accurately described as the generalized Gibbs ensemble [14, 15] with time-dependent Lagrange parameters [16]. Several analytical and numerical studies have demonstrated that breaking of integrability leads to exponential relaxation of typical observables and that the corresponding relaxation rates scale quadratically with the strength of the perturbation [12, 17–20]. Description of NI models within the framework of generalized hydrodynamics (GHD) [21–25] seems also demanding, since the generic integrability-breaking processes involve large momenta transfers [26, 27]. Nevertheless, recent results suggest that arbitrarily weak perturbation applied to a macroscopic integrable system restores the generic chaotic (dissipative) dynamics [28]. Still, more detailed understanding or even theoretical analysis of the relaxation of different quantities in NI systems is mostly lacking so far.

In this work we numerically study relaxation of several operators in a NI XXZ model, employing the microcanonical Lanczos method (MCLM) [29–31] which allows to reach long-enough times even for NI model. In particular, we confirm that both spin-current and energy-current decay exponentially in time with relaxation rates that scale quadratically with the strength of perturbation. However, the energy-current relaxation rate turns out to be much smaller than the relaxation rate for the spin current. The presence of distinct relaxation times is consistent with the predictions of GHD [26, 27]. This result can be simply explained by single (but different for both quantities) local/quasiloal CQ (charges) involved in the relaxation process [19]. Still even in this case, a more detailed analysis using the memory-function indicates a weak contribution of other CQ from the same symmetry sector. Moreover, we can construct a single observable that clearly reveals multiple relaxation rates or, in other words, that its relaxation cannot be described by a single exponential function. We confirm the latter possibility studying relaxation of current correlations in sectors with small magnetization, i.e., with non-zero total spin $S_{\text{tot}}^z \neq 0$. Results in this case can be explained with a projection on the decay of several CQ with different relaxation times. Furthermore, studying other perturbation terms, we find that the relaxation explicitly depends on the form and the symmetry of the perturbation.

Finally, we study the weakly perturbed XXZ model for the specific anisotropy parameters $\Delta = 1$ as well as $\Delta = 0.5$, where the many-body spectra are macroscopically degenerate allowing for additional local CQ in the integrable model. Such CQ do not commute with S_{tot}^z or with other local CQ [32]. In this case, upon introducing the perturbation, correlation functions of such quantities show instead of exponential decay an approximately Gaussian decay with the characteristic relaxation time that scales linearly with the perturbation strength.

2 Relaxation of spin and energy currents

We consider the one-dimensional XXZ chain with L sites assuming periodic boundary conditions, with a specific perturbation involving the next-nearest-neighbor interaction

$$H = H_0 + gH', \quad H_0 = \sum_i h_i, \quad h_i = \frac{J}{2} (S_i^+ S_{i+1}^- + S_i^- S_{i+1}^+) + J\Delta S_i^z S_{i+1}^z, \quad H' = J \sum_i S_i^z S_{i+2}^z, \quad (1)$$

where $S^{\pm,z}$ are spin- $\frac{1}{2}$ operators. The model is integrable for $g = 0$ and the integrability is broken for $g \neq 0$. In this section we study the relaxation of the spin current, j_σ , as well as of the energy current, j_κ , obtained for the unperturbed system

$$j_\sigma = i \sum_{l,l'} l [h_{l'}, S_l^z], \quad j_\kappa = i \sum_{l,l'} l [h_{l'}, h_l], \quad (2)$$

for which we calculate the corresponding normalized current-current correlation functions

$$C_\sigma(t) = \frac{\langle j_\sigma(t) j_\sigma \rangle}{\langle j_\sigma j_\sigma \rangle}, \quad C_\kappa(t) = \frac{\langle j_\kappa(t) j_\kappa \rangle}{\langle j_\kappa j_\kappa \rangle}. \quad (3)$$

We focus on the result for large (infinite) temperature $T = 1/\beta \gg J$. Here, $\langle \dots \rangle = \text{Tr}(\dots)$ denotes averaging either over the canonical ensemble with fixed S_{tot}^z (in Secs. 2-4) or grand-canonical ensemble (in Sec. 5), and $j_\alpha(t) = e^{iHt} j_\alpha e^{-iHt}$. Note that $C_\sigma(t)$ and $C_\kappa(t)$ are related via the Fourier transform to the dynamical spin conductivity, $\sigma(\omega) = \beta \langle j_\sigma j_\sigma \rangle \tilde{C}_\sigma(\omega)$, and the thermal conductivity, $\kappa(\omega) = \beta^2 \langle j_\kappa j_\kappa \rangle \tilde{C}_\kappa(\omega)$, respectively.

For the numerical evaluation of the spectral functions $\tilde{C}_\alpha(\omega)$, on systems up to $L = 28$ sites, we use the MCLM which allows for very high frequency resolution, $\delta\omega \lesssim 10^{-3}J$, and consequently enables evaluation of $C_\alpha(t)$ up to $t \lesssim t_{\text{max}} \sim 10^3/J$. Reaching large $t_{\text{max}}J \gg 1$ is crucial to follow the slow relaxation at weak perturbations, $g \ll 1$, and to resolve possible distinct relaxation times. This is achieved within MCLM by using large number of Lanczos steps $M_L \sim 5 \cdot 10^4$. In the following, we numerically calculate canonical $C_\alpha(t)$, i.e., in sectors with fixed total spin projections S_{tot}^z . Large but finite M_L also sets the smallest reliable $C_\alpha(t) \gtrsim 2 \cdot 10^{-3}$.




Figure 1: Normalized correlation functions $C_\alpha(t)$ for (a) spin current j_σ and (b) energy current j_κ , respectively, in $S_{\text{tot}}^z = 4$ sector, calculated for $L = 28$ and $\Delta = 0.5$. Insets: the same as in main panels but with the time t rescaled by the square of the perturbation strength g^2 .

As a first step, we establish the range of the perturbation strength, g , which is relevant for NI systems. On the one hand, g should be small to remain a perturbation but, on the other

hand, g should be sufficiently large so that the decay is fully detectable within $t < t_{\max}$ for the system sizes accessible with MCLM ($L \leq 28$). To this end, we use parameters for which the decay of the correlation functions reveals clear quadratic dependence on g [12, 19, 20]. Main panels in Fig. 1(a,b) show, respectively, $C_\sigma(t)$ and $C_\kappa(t)$ within the sector $S_{\text{tot}}^z = 4$ for various $g = 0.1 - 0.4$, whereas the time in the insets is rescaled by g^2 . For $0.15 \leq g \leq 0.3$ we observe a perfect collapse of all curves, whereas the latter collapse is slightly worse for weaker g as well also for too strong perturbations. Therefore from now on, we set $g = 0.15$.




Figure 2: Correlation functions $C_\alpha(t)$ in sectors with various S_{tot}^z . Dashed curves in (a) depict exponential fits for $\alpha = j_\sigma, j_\kappa$ at $s = S_{\text{tot}}^z = 0$, while in (c,d) for $s \neq 0$ only long-time fits are presented. Calculated for $L = 28$, $\Delta = 0.5$, and $g = 0.15$.

In Fig. 2(a) we directly compare the relaxation of spin and energy currents $C_\alpha(t)$ at zero-magnetization, $S_{\text{tot}}^z = 0$. Here, H is invariant under the Z_2 spin-flip transformation, generated by the parity operator, $P = \prod_j (S_j^+ + S_j^-)$. Since the spin/energy current is odd/even under this transformation, $P j_\sigma P = -j_\sigma$ and $P j_\kappa P = j_\kappa$, in this sector both currents are mutually orthogonal, i.e., $\langle j_\sigma j_\kappa \rangle = 0$. Consequently, relaxation of both correlations are evidently different. At longer times, $tJ > 100$, the decays do not show any clear deviations from simple exponential, $C_\alpha(t) \propto \exp(-t/\tau_\alpha^0)$, $\alpha = \sigma, \kappa$, where the upper index in the relaxation times, τ_α^s , marks the value of $s = S_{\text{tot}}^z$. Nevertheless, it is clear that the relaxation of j_σ is much faster than that of j_κ , $\tau_\sigma^0 \simeq 3\tau_\kappa^0$, confirming that the breaking of integrability leads to multiple distinct relaxation times, here due to different symmetry sectors involved. The evident differences are also at short times t . Since j_σ is not CQ in the reference H_0 , there is an incoherent drop to $C_\sigma^0 \sim 0.5$ at $tJ \sim O(1)$, reflecting the nontrivial spin stiffness [3] and finite overlap with quasilocal CQ [33, 34]. On the other hand, j_κ is CQ at $g = 0$, so one might expect a single exponential decay in the whole range of t . To good approximation this is indeed the case, but there is still some visible deviation at $tJ < 100$ about which we comment in more detail in Sec. 3.

More challenging question is whether distinct relaxation rates can be observed in the dynamics of a single observable. This might happen when we consider $S_{\text{tot}}^z \neq 0$ sectors where the above symmetry arguments do not apply. In Figs. 2(b-d) we show that a departure from a simple exponential relaxation becomes increasingly visible for $S_{\text{tot}}^z \neq 0$, when also $\langle j_\sigma j_\kappa \rangle \neq 0$. Figs. 2(c,d) for $S_{\text{tot}}^z = 4, 6$ confirm that for longest $tJ > 500$ the relaxation is asymptotically

determined by the same τ_κ^s for both currents, while $S_{\text{tot}}^z = 2$ case on Fig. 2(b) is marginal due to fast decay of $C_\sigma(t)$ (and the limitation $C_\sigma(t) > 2 \cdot 10^{-3}$). Nevertheless, both correlation functions, and in particular $C_\sigma(t)$, reveal a clear deviation from a single-exponential decay.

The modest dependence of τ_κ^s on s , as extracted from Fig. 2(a-d), is shown in Fig. 3(a). More importantly, our analysis indicate that one can fit the decay of correlations $C_\alpha(t)$ for $S_{\text{tot}}^z > 0$ by a sum of two exponential functions with distinct relaxation rates $\tau_\kappa^s, \tau_\sigma^s$. This is shown in Fig. 3(b,c) for $C_\sigma(t)$, but also for $C_\kappa(t)$ in Fig. 3(d). It is indicative that the fit is consistent with only two relaxation times for both correlations, i.e., with longer relaxation rate τ_κ^s still weakly dependent on s as given in Fig. 3(a), and much faster $\tau_\sigma^s \sim \tau_\sigma^0$ which we can approximate just with the $s = 0$ result for $C_\sigma(t)$ in Fig. 2(a).




Figure 3: (a): Relaxation time for the energy current obtained from long-time fits marked by dashed lines in Fig. 2. (b,c): Continuous line shows spin-current $C_\sigma(t)$ for $S_{\text{tot}}^z = 2, 4$, respectively, while (d) energy-current $C_\kappa(t)$ for $S_{\text{tot}}^z = 4$. Results in (b,d) are fitted by a sum of two exponential functions (dashed, green curves) with the relaxation times τ_κ^s as in panel (a), and $\tau_\sigma^s = \tau_\sigma^0$. Calculated for $L = 28$, $\Delta = 0.5$, and $g = 0.15$.

Above results may serve as a motivation for a simple phenomenological description. The value of the correlation functions for $t \rightarrow \infty$ in the integrable model, $g = 0$, is determined via the Mazur bound [2, 3] by the projections of the studied operators on local/quasilocal CQ [33–39] Q_n , which should be chosen as mutually orthogonal $\langle Q_n Q_{n'} \rangle \propto \delta_{n,n'}$ in the canonical ensemble with fixed S_{tot}^z . Assuming the completeness (saturation) of the bound, this means for considered correlations,

$$C_\alpha(t \rightarrow \infty) = C_\alpha^0 = \frac{1}{\langle j_\alpha j_\alpha \rangle} \sum_n \frac{\langle j_\alpha Q_n \rangle^2}{\langle Q_n Q_n \rangle}, \quad (4)$$

where the term $\langle j_\alpha j_\alpha \rangle^{-1}$ arises from normalization in Eq. (3). Eq. (4) is invariant under the orthogonal transformation of normalized CQ

$$\frac{Q_n}{\sqrt{\langle Q_n Q_n \rangle}} = \sum_s (\hat{o})_{ns} \frac{Q_s}{\sqrt{\langle Q_s Q_s \rangle}}, \quad (5)$$

where \hat{O} is arbitrary orthogonal matrix. In NI system, Q_n are not any more CQ and decay with the characteristic times $\propto g^2$. Based on results in Fig. 3(b-d) and previous numerical studies in Ref. [19], we conjecture that projections on Q_n are essential also for the asymptotic dynamics in NI models,

$$C_\alpha(t \gg J^{-1}) \simeq \frac{1}{\langle j_\alpha j_\alpha \rangle} \sum_n \frac{\langle j_\alpha Q_n \rangle^2}{\langle Q_n Q_n \rangle} \exp\left(-\frac{t}{\tau_n}\right), \quad (6)$$

where $\tau_n \propto 1/g^2$. It should be, however, stressed that in contrast to Eq. (4) the appropriate set of Q_n in Eq. (6) is not arbitrary, but determined by the perturbation [5, 17, 27]. A numerical algorithm for identifying such modes has been discussed in Ref. [19], while here we do not attempt to determine relevant Q_n in more detail. However, for $S_{\text{tot}}^z = 0$ one can assume that one of Q_n in Eq. (6) can be well approximated by j_κ (being one of CQ in the reference system), which explains nearly perfect exponential decay of $C_\kappa(t)$ in Fig. 2(a), even though a small correction might be needed, as tested in more detail in Sec. 3. It is well known [3] that for $S_{\text{tot}}^z = 0$ and $g = 0$ the spin-current stiffness $C_\sigma^0 < 1$. Since $\langle j_\sigma j_\kappa \rangle = 0$, Q_n that are relevant for the decay time τ_σ^0 have to be related to the quasilocal CQ [19, 33, 34, 40].

On the other hand, for nonzero S_{tot}^z , we clearly need at least two distinct relaxation times $\tau_\kappa^s, \tau_\sigma^s$ to fit both decays, $C_\alpha(t)$. The necessity of multiple relaxation times is most evident for $C_\sigma(t)$ at $S_{\text{tot}}^z = 2, 4$ presented on Fig. 3(b-c). Upon changing S_{tot}^z , one has also to modify τ_κ^s , as suggested by results in Fig. 3(a), while the dependence of τ_σ^s on S_{tot}^z is less evident, so we can approximate $\tau_\sigma^s \sim \tau_\sigma^0$. In Fig. 3(d) we show that $C_\kappa(t)$ can also be well fitted by a sum of two exponential functions, with the very same relaxation times which are used for fitting $C_\sigma(t)$ in Fig. 3(c). This indicates that the same pair of Q_n is relevant for the decay of the spin and the energy currents for $S_{\text{tot}}^z \neq 0$.

3 Memory-function analysis of the energy-current correlations

It is desirable to have more explicit way to evaluate the of long-time decay of correlations, $C_\alpha(t)$, for chosen perturbations H' , and in particular a direct expression for relevant relaxation rates $1/\tau_n$. Here, it is convenient to follow the Mori formalism [41, 42], where one can generally express the current-correlation relaxation function $\phi_\alpha(\omega)$ in terms of the corresponding memory functions (MF), $M_\alpha(\omega)$,

$$\begin{aligned} \phi_\alpha(\omega) &= \frac{1}{L} \langle j_\alpha | [\mathcal{L} - \omega]^{-1} | j_\alpha \rangle = \frac{\chi_\alpha(\omega) - \chi_\alpha^0}{\omega} = \frac{-\chi_\alpha^0}{\omega + M_\alpha(\omega)}, \\ \chi_\alpha(\omega) &= \frac{i}{L} \int_0^\infty e^{i\omega t} \langle [j_\alpha^\dagger(t), j_\alpha] \rangle dt, \quad \chi_\alpha^0 = \chi_\alpha(0) > 0, \end{aligned} \quad (7)$$

where $\langle A|B \rangle = 1/L \int_0^\beta d\tau \langle AB(i\tau) \rangle$. At high temperatures, $\beta \rightarrow 0$, $\beta \tilde{C}_\alpha(\omega) = \text{Im} \phi_\alpha(\omega)$ and $\chi_\alpha^0 = \beta \langle j_\alpha j_\alpha \rangle$. Knowing numerical result for $\tilde{C}_\alpha(\omega)$ at finite g , one can evaluate (via Kramers-Kronig relation) the complex $\phi_\alpha(\omega)$, extract directly the corresponding complex MF, $M_\alpha(\omega)$, and in particular the dynamical relaxation rate $\Gamma_\alpha(\omega) = \text{Im} M_\alpha(\omega)$.

On the other hand, for a NI system one can find an explicit expression in the case where the current is a CQ in the reference system [17, 19]. This is particularly the case for the energy current, $[H_0, Q_3] = [H_0, j_\kappa] = 0$. Then, within the lowest order in the perturbation $g \ll 1$ one can approximate MF as

$$M_\kappa(\omega) = \frac{1}{\chi_\alpha^0} N_\kappa(\omega), \quad N_\kappa(\omega) = g^2 (\mathcal{F} | [\mathcal{L} - \omega]^{-1} | \mathcal{F}), \quad \mathcal{F} = [H', j_\kappa], \quad (8)$$

i.e., as the correlation function of the the force \mathcal{F} in *the unperturbed - integrable system*. Eq. (8) is derived under the assumption that only a single charge $Q_3 = j_k$ is relevant for the decay, as well as that \mathcal{F} has no overlap with any of Q_n . This can be true for the sector with $S_{\text{tot}}^z = 0$ and can be directly tested for operator \mathcal{F} for the particular H' in Eq. (1),

$$\mathcal{F} = i \sum_i \left[T_{i-1}^{i+1} S_i^z (S_{i+3}^z - S_{i-3}^z) - \Delta T_i^{i+1} (S_{i+2}^z + S_{i-1}^z) (S_{i+3}^z - S_{i+2}^z) \right], \quad (9)$$

with $T_i^l = (S_l^+ S_i^- + \text{H.c.})/2$. Due to the parity symmetry at $S_{\text{tot}}^z = 0$, the above \mathcal{F} appears orthogonal to known local/quasilocal CQ. We can then evaluate numerically $M_\kappa(\omega)$ as the correlation of \mathcal{F} in the reference H_0 system. The MCLM result for $\Gamma_\kappa(\omega)/g^2$ at $S_{\text{tot}}^z = 0$ is presented in Fig. 4, both as extracted directly via Eq. (7) from $\tilde{C}_\kappa(\omega)$ for various finite $g = 0.1 - 0.4$, as well as by calculating the result from Eq. (8) for the force given in Eq. (9). The overall agreement of perturbation result with the numerically extracted MF $\Gamma_\kappa(\omega)/g^2$, (being essentially g -independent) is very satisfactory in the whole ω regime [see the inset of Fig. 4(a)]. Still, at low $\omega/J < 0.4$ there appears a visible difference. Partly responsible is the (apparently) singular contribution in perturbative $\Gamma_\kappa^0 = \Gamma_\kappa(\omega \rightarrow 0)$, which seems to indicate a small overlap with some CQ (possibly being a finite-size effect). Apart from that, also visible is a quantitative mismatch at $\omega \rightarrow 0$ which indicates that the relevant Q_n in Eq. (6) is not just j_k , and consequently also Eq. (8) is not a full description of relaxation. In other words, even for $S_{\text{tot}}^z = 0$ more than a single mode is needed in Eq. (6) to properly describe $C_\kappa(t)$.




Figure 4: Energy-current memory function (relaxation-rate), $\Gamma_\kappa(\omega)$, (a) extracted directly from $\tilde{C}_\kappa(\omega)$ for $S_{\text{tot}}^z = 0$ with different perturbations $g = 0.1 - 0.4$, compared with the perturbation result, Eqs. (8),(9), and (b) extracted for various S_{tot}^z from $\tilde{C}_\kappa(\omega)$ for fixed $g = 0.15$. Calculated for $L = 28$ and $\Delta = 0.5$.

In Fig. 4(b) we present $\Gamma_\kappa(\omega)/g^2$ for fixed $g = 0.15$ but various S_{tot}^z . It is indicative that for $\omega/J > 0.15$ the MF is essentially independent of S_{tot}^z . On the other hand, the decrease of $\Gamma_\kappa^0 = 1/\tau_\kappa^s$ with S_{tot}^z reflects the observed increase of τ_κ^s in Fig. 3(a). Still, it is not straightforward to capture this in a perturbative approach, Eq. (8).

4 Dependence on the form of perturbation

In Sec. 2 we analysed a particular form of perturbation, i.e., the next nearest-neighbor interaction, Eq. (1), which does not break the translational symmetry or the spin parity P . This is also essential for the phenomenological explanation, Eq. (6), in terms of decaying CQ, as well as for the MF analysis in Sec. 3. However, using numerical MCLM we can check also other

perturbations. Let us consider as a perturbation

$$H'' = J \sum_i S_i^z S_{i+1}^z S_{i+2}^z, \quad (10)$$

which breaks the parity symmetry, P , of the total Hamiltonian, $H = H_0 + gH''$. Fig. 5 shows current $\alpha = \sigma, \kappa$ correlation functions $C_\alpha(t)$ for $S_{\text{tot}}^z = 0$. We observe exponential decay with a single relaxation time $\tau_\alpha \propto 1/g^2$, which is different from two distinct relaxation times obtained for the P -preserving H' , Eq. (1), i.e., the relaxation evidently depends on the form of perturbation. One may interpret this behavior in terms of conjecture, Eq. (6). All observables which have non-vanishing projection on Q_n with the longest relaxation time should asymptotically decay with the same decay rate. In order to obtain different rate, one needs to build an operator that is strictly orthogonal to the latter Q_n . It is highly nontrivial task, since approximate Q_n in Eq. (6) can be obtained numerically only for small system. For the parity-preserving perturbations, the total Hamiltonian is even and all Q_n in Eq. (6) have well defined parity being either odd or even. Then operators from one parity sectors are strictly orthogonal to Q_n from the other parity sector. Due to this orthogonality we observe different relaxation times in odd and even sectors without any fine-tuning of the studied observables. However for odd perturbations, the total Hamiltonian contains even and odd terms, hence both parity sectors are mixed during the time evolution. As a consequence, Q_n in Eq. (6) may not have well defined parity.




Figure 5: Current correlation function $C_\alpha(t)$ for the perturbation, Eq. (10), breaking the parity symmetry P , as compared with P -preserving perturbation Eq. (1). Calculated for $L = 28$, $\Delta = 0.5$, $g = 0.15$, and $S_{\text{tot}}^z = 0$.

5 Non-commuting conserved quantities and their non-exponential relaxation

In this section, we focus on specific values of the anisotropy parameter, Δ , where the many-body spectra exhibit additional massive degeneracies [31]. The latter originate from eigenstates corresponding to different S_{tot}^z with equal energies. This property allows for the presence of non-commuting local/quasilocal CQ [32, 43–45]. As a first example we take $\Delta = 1$ in which case H_0 is the $SU(2)$ -symmetric Heisenberg chain for which one can study the total S_{tot}^x spin operator

$$O_1 = \sum_j (S_j^+ + S_j^-). \quad (11)$$

It is clear that O_1 is a local operator which commutes with H_0 at $\Delta = 1$, but does not commute with other local CQ, e.g., not with the S_{tot}^z .

The second nontrivial case is related with the commensurate $\Delta = \cos(\pi/3) = 1/2$, for which the non-commuting local CQ have been derived (for $g = 0$) in Ref. [32]. Here, we study the relaxation of the operator,

$$O_3 = \sum_j (-1)^j (S_{j-1}^+ S_j^+ S_{j+1}^+ + \text{H.c.}), \quad (12)$$

which does not commute with S_{tot}^z and is not invariant under translations by odd number of sites. It should be mentioned that analogous local operators exist for other commensurate cases, in particular for $\Delta = \cos(\pi/2) = 0$, where $O_2 = \sum_j (-1)^j (S_j^+ S_{j+1}^+ + \text{H.c.})$ is local CQ for H_0 [45].

Utilizing Eq. (3), we calculate normalized correlations functions $C_1(t)$ and $C_3(t)$ for the operators O_1 and O_3 , respectively. In contrast to the preceding section, now the averaging $\langle \dots \rangle$ is carried out over grand canonical ensemble. We note that O_3 is similar to the previously studied spin-current j_σ in the sense, that it does not commute with H_0 , but has large projection on the corresponding quasilocal CQ [32]. In order to confirm this, in the inset in Fig. 6(c) we show the finite-size scaling of the relevant stiffness $\tilde{C}_3(\omega \rightarrow 0^+, g = 0)$ (with value $\simeq 0.4$ thermodynamic limit $L \rightarrow \infty$). Without imposing the translational symmetry, we studied all local non-commuting operators $\sum_i (A_i + A_i^\dagger)$, where A_i are supported on up to 3 sites and do not commute with S_{tot}^z . Utilizing the algorithm from Ref. [40], we have found for $\Delta = 1/2$, $g = 0$ and $L \leq 14$ that O_3 has the largest stiffness out of all these operators (not shown).

Next, we focus on the asymptotic decay of $C_\alpha(t)$, $\alpha = 1, 3$, and their dependence on $g \neq 0$. To this end, we first calculate the Fourier transform, $C_\alpha(\omega)$, and then its cumulative spectral function

$$\tilde{C}_\alpha(\omega, g) = \int_{-\omega}^{\omega} d\omega' C_\alpha(\omega') = \frac{\sum_{mn} \theta(\omega - |E_m - E_n|) \langle m | O_\alpha | n \rangle^2}{\sum_{mn} \langle m | O_\alpha | n \rangle^2}, \quad (13)$$

expressed in terms of eigenstates, $H|n\rangle = E_n|n\rangle$, obtained using the exact diagonalization of H on up to $L = 16$ sites. We note that $\tilde{C}_1(\omega \rightarrow 0^+, g = 0) = 1$ (because O_1 commutes with H_0 at $\Delta = 1$), whereas $\tilde{C}_3(\omega \rightarrow 0, g = 0) \simeq 0.4$. Since we are interested in the low- ω part of $C_\alpha(\omega)$, it is convenient to normalize the spectral function,

$$R_\alpha(\omega) = \frac{\tilde{C}_\alpha(\omega, g)}{\tilde{C}_\alpha(\omega \rightarrow 0^+, g = 0)}, \quad (14)$$

so that one may directly compare relaxations of both considered observables. Numerical results are shown in figures Figs. 6(a,b) and 6(d,e) for $R_3(\omega)$ at $\Delta = 1/2$ and $R_1(\omega)$ at $\Delta = 1$, respectively. In contrast to the spin- and energy-currents discussed in the preceding sections, curves for various g do not collapse when one rescales the frequency by g^2 , as it is shown in the insets in Fig. 6(b,e). However, a convincing collapse may be obtained for the scaling ω/g . Moreover, the rescaled curves can be quite accurately fitted by the error function, shown as dashed curves in the latter plots, implying $\tilde{C}_\alpha(\omega) \propto \exp[-a (\omega/g)^2]$ where the coefficient a does not depend on g . Consequently, the decay is not exponential but Gaussian, $C_\alpha(t) \propto \exp[-(t/\tau_\alpha)^2]$, where the characteristic relaxation rate, $1/\tau_\alpha \propto g$.

The Gaussian relaxation occurs in the vicinity of Δ characterized by additional degeneracies [31, 45], originating from that eigenstate with different S_{tot}^z having the same energy. The perturbation breaks integrability but also lifts the latter degeneracy. In order to disentangle these two mechanisms, we also study the correlation functions, $\tilde{C}_\alpha(\omega, g)$ for $g = 0$ but with shifted $\Delta = 1/2 + \delta$ (for $\alpha = 3$) and $\Delta = 1 + \delta$ (for $\alpha = 1$). Then, the degeneracy is lifted without destroying the integrability. Results are shown in Fig. 6(c,f) for both $\alpha = 1, 3$. One




Figure 6: Cumulative spectral functions $R_\alpha(\omega)$, Eq. (14), as calculated for $L = 16$ system. We show results for (a) O_3 with $\Delta = 1/2$ and (d) for O_1 with $\Delta = 1$. Panels (b) and (e) show respectively the same data but with rescaled frequency ω/g (main panels) and ω/g^2 (insets). Panels (c) and (f) show results for integrable system ($g = 0$) but for shifted anisotropy parameters $\Delta = 0.5 + \delta$ and $\Delta = 1 + \delta$, respectively. Dashed curves show fittings with the error function. Inset in (c) shows finite-size scaling of the stiffness $C_3(t \rightarrow \infty)$ for $\Delta = 0.5$ and $g = 0$.

observes the same behavior as for the NI system above, i.e., $C_\alpha(t) \propto \exp[-(t/\tau_\alpha)^2]$, with $1/\tau_\alpha \propto \delta$. This result explains additionally the origin of the anomalous scaling of characteristic τ_α . The nonzero stiffnesses, $\tilde{C}_\alpha(\omega \rightarrow 0^+, g = 0)$, emerges from states $|m\rangle$ and $|n\rangle$ in Eq. (13) with different S_{tot}^z , but with equal energies $E_m = E_n$. The latter degeneracy is lifted either by $g \neq 0$ or $\delta \neq 0$ already in the first order perturbation theory, $|E_m = E_n| \propto g, \delta$ [45], leading to the scaling $\omega/(g, \delta)$, shown in Fig. 6.

6 Conclusions

We numerically analyzed the decay of normalized correlation function $C_\alpha(t)$ of different local quantities in the nearly integrable XXZ model. Correlations generally reveal a fast drop at short times $tJ \sim O(1)$, consistent with finite stiffnesses of studied quantities in the integrable model H_0 . A weak integrability breaking $g \ll 1$ then leads to further slow - exponential-like - decay which is characterized with a single or multiple relaxation times, all scaling with the perturbation strength $\tau_n \propto 1/g^2$. The simplest cases appear to be the energy-current j_κ and spin-current j_σ correlations at zero magnetization $S_{\text{tot}}^z = 0$, where - due to symmetry - different CQ are involved in the relaxation of both quantities, and consequently relevant τ_n are quite distinct. Since j_κ is by itself CQ within H_0 , one can go a step further and give an explicit memory-function analysis and perturbative expression for the relaxation-rate spectral function $\Gamma_\kappa(\omega)$. In this case the extracted and perturbative $\Gamma_\kappa(\omega)$ match quite well in the whole ω range. Still, some deviations in $\Gamma_\kappa(\omega \sim 0) = 1/\tau_\kappa^0$ as well as in $C_\kappa(t)$ at shorter t/g^2 indicate on possible multiple relaxation times and more than one relevant Q_n even in this case.

The existence of multiple relaxation times is becoming evident for $S_{\text{tot}}^z \neq 0$. The conjecture Eq. (6) concerning the presence of multiple relaxation times τ_n , which are linked with (appropriately rotated) various CQ of the parent integrable model, is not in conflict with previous results [20] which report a simple exponential relaxation. In order to demonstrate at least two relaxation times, we have carefully selected observables so that they have a large projection on quickly decaying CQ for $S_{\text{tot}}^z \neq 0$ (Q_n relevant for j_σ at $S_{\text{tot}}^z = 0$), as well as much smaller projection on slowly decaying one (j_κ). Typically, the opposite holds true: one studies observables which have largest projection on simplest CQ, which are supported only on the few sites and the related τ_n are the longest relaxation times [19]. As a consequence, small and fast decaying projections on more complicated CQ in Eq. (6) may not be visible in the numerical results. Furthermore, our analysis indicates that the form, in particular the symmetry, of the perturbation is relevant for the decay of $C_\alpha(t)$.

In contrast to quantities considered above at general anisotropy Δ and their decay, the correlations $C_\alpha(t)$ of particular quantities O_l , being conserved by H_0 only at commensurate $\Delta_0 = \cos(\pi/m)$, but not commuting with S_{tot}^z , behave qualitatively different. Under finite perturbation, which here can be introduced either by $g \neq 0$ or by deviation $\Delta = \Delta_0 + \delta$, we observe effectively a (non-exponential) Gaussian-like decay of $C_\alpha(t)$ with different scaling of characteristic decay time, i.e. $\tau_\alpha \propto 1/g$. The origin here is the lifting of macroscopic degeneracy of many-body states at these particular Δ_0 .

Funding information J.H. acknowledges the support by the Polish National Agency of Academic Exchange (NAWA) under contract PPN/PPO/2018/1/00035. M.M. acknowledges the support by the National Science Centre, Poland via projects 2020/37/B/ST3/00020. P.P. acknowledges the support by the project N1-0088 of the Slovenian Research Agency. The numerical calculation were partly carried out at the facilities of the Wroclaw Centre for Networking and Supercomputing.

References

- [1] B. Bertini, F. Heidrich-Meisner, C. Karrasch, T. Prosen, R. Steinigeweg and M. Žnidarič, Finite-temperature transport in one-dimensional quantum lattice models, Rev. Mod. Phys. **93**, 025003 (2021), doi:[10.1103/RevModPhys.93.025003](https://doi.org/10.1103/RevModPhys.93.025003).
- [2] P. Mazur, Non-ergodicity of phase functions in certain systems, Physica **43**, 533 (1969), doi:[http://dx.doi.org/10.1016/0031-8914\(69\)90185-2](http://dx.doi.org/10.1016/0031-8914(69)90185-2).
- [3] X. Zotos, F. Naef and P. Prelovšek, Transport and conservation laws, Phys. Rev. B **55**, 11029 (1997), doi:[10.1103/PhysRevB.55.11029](https://doi.org/10.1103/PhysRevB.55.11029).
- [4] X. Zotos, High temperature thermal conductivity of two-leg spin-1/2 ladders, Phys. Rev. Lett. **92**, 067202 (2004), doi:[10.1103/PhysRevLett.92.067202](https://doi.org/10.1103/PhysRevLett.92.067202).
- [5] G. P. Brandino, J.-S. Caux and R. M. Konik, Glimmers of a quantum kam theorem: Insights from quantum quenches in one-dimensional bose gases, Phys. Rev. X **5**, 041043 (2015), doi:[10.1103/PhysRevX.5.041043](https://doi.org/10.1103/PhysRevX.5.041043).
- [6] T. Prosen, Time evolution of a quantum many-body system: Transition from integrability to ergodicity in the thermodynamic limit, Phys. Rev. Lett. **80**, 1808 (1998), doi:[10.1103/PhysRevLett.80.1808](https://doi.org/10.1103/PhysRevLett.80.1808).

- [7] Y. Huang, C. Karrasch and J. E. Moore, Scaling of electrical and thermal conductivities in an almost integrable chain, Phys. Rev. B **88**, 115126 (2013), doi:[10.1103/PhysRevB.88.115126](https://doi.org/10.1103/PhysRevB.88.115126).
- [8] F. H. L. Essler, S. Kehrein, S. R. Manmana and N. J. Robinson, Quench dynamics in a model with tuneable integrability breaking, Phys. Rev. B **89**, 165104 (2014), doi:[10.1103/PhysRevB.89.165104](https://doi.org/10.1103/PhysRevB.89.165104).
- [9] M. Kollar, F. A. Wolf and M. Eckstein, Generalized gibbs ensemble prediction of prethermalization plateaus and their relation to nonthermal steady states in integrable systems, Phys. Rev. B **84**, 054304 (2011), doi:[10.1103/PhysRevB.84.054304](https://doi.org/10.1103/PhysRevB.84.054304).
- [10] B. Bertini, F. H. L. Essler, S. Groha and N. J. Robinson, Prethermalization and thermalization in models with weak integrability breaking, Phys. Rev. Lett. **115**, 180601 (2015), doi:[10.1103/PhysRevLett.115.180601](https://doi.org/10.1103/PhysRevLett.115.180601).
- [11] K. Mallayya, M. Rigol and W. De Roeck, Prethermalization and thermalization in isolated quantum systems, Phys. Rev. X **9**, 021027 (2019), doi:[10.1103/PhysRevX.9.021027](https://doi.org/10.1103/PhysRevX.9.021027).
- [12] P. Jung, R. W. Helmes and A. Rosch, Transport in almost integrable models: Perturbed heisenberg chains, Phys. Rev. Lett. **96**, 067202 (2006), doi:[10.1103/PhysRevLett.96.067202](https://doi.org/10.1103/PhysRevLett.96.067202).
- [13] B. Bertini, F. H. L. Essler, S. Groha and N. J. Robinson, Thermalization and light cones in a model with weak integrability breaking, Phys. Rev. B **94**, 245117 (2016), doi:[10.1103/PhysRevB.94.245117](https://doi.org/10.1103/PhysRevB.94.245117).
- [14] M. Rigol, V. Dunjko, V. Yurovsky and M. Olshanii, Relaxation in a completely integrable many-body quantum system: An ab initio study of the dynamics of the highly excited states of 1d lattice hard-core bosons, Phys. Rev. Lett. **98**, 050405 (2007), doi:[10.1103/PhysRevLett.98.050405](https://doi.org/10.1103/PhysRevLett.98.050405).
- [15] A. C. Cassidy, C. W. Clark and M. Rigol, Generalized thermalization in an integrable lattice system, Phys. Rev. Lett. **106**, 140405 (2011), doi:[10.1103/PhysRevLett.106.140405](https://doi.org/10.1103/PhysRevLett.106.140405).
- [16] F. Lange, Z. Lenarčič and A. Rosch, Time-dependent generalized gibbs ensembles in open quantum systems, Phys. Rev. B **97**, 165138 (2018), doi:[10.1103/PhysRevB.97.165138](https://doi.org/10.1103/PhysRevB.97.165138).
- [17] P. Jung and A. Rosch, Lower bounds for the conductivities of correlated quantum systems, Phys. Rev. B **75**, 245104 (2007), doi:[10.1103/PhysRevB.75.245104](https://doi.org/10.1103/PhysRevB.75.245104).
- [18] P. Jung and A. Rosch, Spin conductivity in almost integrable spin chains, Phys. Rev. B **76**, 245108 (2007), doi:[10.1103/PhysRevB.76.245108](https://doi.org/10.1103/PhysRevB.76.245108).
- [19] M. Mierzejewski, T. Prosen and P. Prelovšek, Approximate conservation laws in perturbed integrable lattice models, Phys. Rev. B **92**, 195121 (2015), doi:[10.1103/PhysRevB.92.195121](https://doi.org/10.1103/PhysRevB.92.195121).
- [20] K. Mallayya and M. Rigol, Quantum quenches and relaxation dynamics in the thermodynamic limit, Phys. Rev. Lett. **120**, 070603 (2018), doi:[10.1103/PhysRevLett.120.070603](https://doi.org/10.1103/PhysRevLett.120.070603).
- [21] J. De Nardis, D. Bernard and B. Doyon, Hydrodynamic diffusion in integrable systems, Phys. Rev. Lett. **121**, 160603 (2018), doi:[10.1103/PhysRevLett.121.160603](https://doi.org/10.1103/PhysRevLett.121.160603).

- [22] E. Ilievski, J. De Nardis, M. Medenjak and T. Prosen, Superdiffusion in one-dimensional quantum lattice models, Phys. Rev. Lett. **121**, 230602 (2018), doi:[10.1103/PhysRevLett.121.230602](https://doi.org/10.1103/PhysRevLett.121.230602).
- [23] S. Gopalakrishnan, D. A. Huse, V. Khemani and R. Vasseur, Hydrodynamics of operator spreading and quasiparticle diffusion in interacting integrable systems, Phys. Rev. B **98**, 220303(R) (2018), doi:[10.1103/PhysRevB.98.220303](https://doi.org/10.1103/PhysRevB.98.220303).
- [24] U. Agrawal, S. Gopalakrishnan, R. Vasseur and B. Ware, Anomalous low-frequency conductivity in easy-plane XXZ spin chains, Phys. Rev. B **101**, 224415 (2020), doi:[10.1103/PhysRevB.101.224415](https://doi.org/10.1103/PhysRevB.101.224415).
- [25] V. B. Bulchandani, S. Gopalakrishnan and E. Ilievski, Superdiffusion in spin chains, J. Stat. Mech. **2021**, 084001 (2021), doi:[10.1088/1742-5468/ac12c7](https://doi.org/10.1088/1742-5468/ac12c7).
- [26] A. J. Friedman, S. Gopalakrishnan and R. Vasseur, Diffusive hydrodynamics from integrability breaking, Phys. Rev. B **101**, 180302 (2020), doi:[10.1103/PhysRevB.101.180302](https://doi.org/10.1103/PhysRevB.101.180302).
- [27] A. Bastianello, A. D. Luca and R. Vasseur, Hydrodynamics of weak integrability breaking, J. Stat. Mech. **2021**, 114003 (2021), doi:[10.1088/1742-5468/ac26b2](https://doi.org/10.1088/1742-5468/ac26b2).
- [28] T. LeBlond, D. Sels, A. Polkovnikov and M. Rigol, Universality in the Onset of Quantum Chaos in Many-Body Systems, arXiv e-prints 2012.07849 (2020), [2012.07849](https://arxiv.org/abs/2012.07849).
- [29] M. W. Long, P. Prelovšek, S. El Shawish, J. Karadamoglou and X. Zotos, Finite-temperature dynamical correlations using the microcanonical ensemble and the lanczos algorithm, Phys. Rev. B **68**, 235106 (2003), doi:[10.1103/PhysRevB.68.235106](https://doi.org/10.1103/PhysRevB.68.235106).
- [30] P. Prelovšek and J. Bonča, Ground state and finite temperature lanczos methods, In A. Avella and F. Mancini, eds., Strongly Correlated Systems - Numerical Methods. Springer, Berlin, doi:[10.1007/978-3-642-35106-8](https://doi.org/10.1007/978-3-642-35106-8) (2013).
- [31] P. Prelovšek, M. Mierzejewski and J. Herbrych, Coexistence of diffusive and ballistic transport in integrable quantum lattice models, Phys. Rev. B **104**, 115163 (2021), doi:[10.1103/PhysRevB.104.115163](https://doi.org/10.1103/PhysRevB.104.115163).
- [32] L. Zadnik, M. Medenjak and T. Prosen, Quasilocal conservation laws from semicyclic irreducible representations of $u_q(sl_2)$ in xxz spin-1/2 chains, Nucl. Phys. B. **902**, 339 (2016), doi:[10.1016/J.NuclPhysB.2015.11.023](https://doi.org/10.1016/J.NuclPhysB.2015.11.023).
- [33] T. Prosen, Open XXZ spin chain: Nonequilibrium steady state and a strict bound on ballistic transport, Phys. Rev. Lett. **106**, 217206 (2011), doi:[10.1103/PhysRevLett.106.217206](https://doi.org/10.1103/PhysRevLett.106.217206).
- [34] T. Prosen and E. Ilievski, Families of quasilocal conservation laws and quantum spin transport, Phys. Rev. Lett. **111**, 057203 (2013), doi:[10.1103/PhysRevLett.111.057203](https://doi.org/10.1103/PhysRevLett.111.057203).
- [35] M. Mierzejewski, P. Prelovšek and T. Prosen, Breakdown of the generalized Gibbs ensemble for current-generating quenches, Phys. Rev. Lett. **113**, 020602 (2014), doi:[10.1103/PhysRevLett.113.020602](https://doi.org/10.1103/PhysRevLett.113.020602).
- [36] B. Pozsgay, M. Mestyán, M. A. Werner, M. Kormos, G. Zaránd and G. Takács, Correlations after quantum quenches in the XXZ spin chain: Failure of the generalized Gibbs ensemble, Phys. Rev. Lett. **113**, 117203 (2014), doi:[10.1103/PhysRevLett.113.117203](https://doi.org/10.1103/PhysRevLett.113.117203).

- [37] G. Goldstein and N. Andrei, Failure of the local generalized Gibbs ensemble for integrable models with bound states, Phys. Rev. A **90**, 043625 (2014), doi:[10.1103/PhysRevA.90.043625](https://doi.org/10.1103/PhysRevA.90.043625).
- [38] R. G. Pereira, V. Pasquier, J. Sirker and I. Affleck, Exactly conserved quasilocal operators for the XXZ spin chain, J. Stat. Mech. **2014**, P09037 (2014), doi:[10.1088/1742-5468/2014/09/P09037](https://doi.org/10.1088/1742-5468/2014/09/P09037).
- [39] E. Ilievski, J. De Nardis, B. Wouters, J.-S. Caux, F. H. L. Essler and T. Prosen, Complete generalized gibbs ensembles in an interacting theory, Phys. Rev. Lett. **115**, 157201 (2015), doi:[10.1103/PhysRevLett.115.157201](https://doi.org/10.1103/PhysRevLett.115.157201).
- [40] M. Mierzejewski, P. Prelovšek and T. Prosen, Identifying local and quasilocal conserved quantities in integrable systems, Phys. Rev. Lett. **114**, 140601 (2015), doi:[10.1103/PhysRevLett.114.140601](https://doi.org/10.1103/PhysRevLett.114.140601).
- [41] H. Mori, Transport, collective motion and Brownian motion, Prog. Theor. Phys. **33**, 423 (1965), doi:[10.1143/PTP33.423](https://doi.org/10.1143/PTP33.423).
- [42] P. Prelovšek and J. Herbrych, Self-consistent approach to many-body localization and subdiffusion, Phys. Rev. B **96**, 035130 (2017), doi:[10.1103/PhysRevB.96.035130](https://doi.org/10.1103/PhysRevB.96.035130).
- [43] M. Fagotti, On conservation laws, relaxation and pre-relaxation after a quantum quench, J. Stat. Mech. **2014**, P03016 (2014), doi:[10.1088/1742-5468/2014/03/p03016](https://doi.org/10.1088/1742-5468/2014/03/p03016).
- [44] F. H. L. Essler and M. Fagotti, Quench dynamics and relaxation in isolated integrable quantum spin chains, J. Stat. Mech. **2016**, 064002 (2016), doi:[10.1088/1742-5468/2016/06/064002](https://doi.org/10.1088/1742-5468/2016/06/064002).
- [45] M. Mierzejewski, J. Herbrych and P. Prelovšek, Ballistic transport in integrable quantum lattice models with degenerate spectra, Phys. Rev. B **103**, 235115 (2021), doi:[10.1103/PhysRevB.103.235115](https://doi.org/10.1103/PhysRevB.103.235115).