

Searching for Wave-like Dark Matter with QSHS

I. Bailey², B. Chakraborty⁴, G. Chapman⁵, E.J. Daw⁸, J. Gallop⁵, G. Gregori⁶, E. Hardy³, L. Hao⁵, E. Laird², P. Leek⁶, S.Ó. Peatáin², Y. Pashkin², M.G. Perry^{8*}, M. Piscitelli⁶, E. Romans⁴, J. March-Russell⁶, P. Meeson⁷, S. Sarkar⁶, P.J. Smith⁸, N. Song³, M. Soni², B. K. Tan⁶, S. West⁷, S. Withington¹

1 University of Cambridge, Cambridge, UK.

2 Lancaster University, Lancaster, UK.

3 University of Liverpool, Liverpool, UK.

4 University College London, London, UK.

5 National Physics Laboratory, London, UK.


6 University of Oxford, Oxford, UK.

7 Royal Holloway, University of London, UK.

8 University of Sheffield, Sheffield, UK.

* mitchell.perry@sheffield.ac.uk

September 30, 2022



*14th International Conference on Identification of Dark Matter
Vienna, Austria, 18-22 July 2022*

Abstract

In 2021 the Quantum Sensors for the Hidden Sector (QSHS) collaboration was founded in the UK and received funding to develop and demonstrate quantum devices with the potential to detect hidden sector particles in the μeV to $100 \mu\text{eV}$ mass window. The collaboration has been developing a range of devices and has started to develop a high-field, low-temperature facility at the University of Sheffield to characterise and test the devices in a haloscope geometry. This paper introduces the motivation behind QSHS collaboration, its aims and its current progress.

1 Introduction

The nature of dark matter is an outstanding problem in physics. There is currently much interest in exploring the possibility of light dark matter candidates, in particular the QCD axion. Assuming a local dark matter halo density of approximately 0.45 GeV cm^{-3} , it can be shown that cold low mass particles ($\leq 1 \text{ eV}$) will have a high number density and a long de-Broglie wavelength. Consequently they can be treated as a classical pseudo-scalar field.

QCD axions couple electromagnetically, augmenting Maxwell's equations to give additional terms [1]. The axion field would interact with a static magnetic field to produce an effective current leading to a propagating electromagnetic wave. Achieving the necessary sensitivity for detection requires the use of resonant methods. Ultra-low cryogenic temperatures are required to permit the expected signal, having an expected power of order yocto-watts even in resonant detectors, to be discernable above the noise inherent to the detector. The QSHS collaboration

is building a test facility to integrate and test several ultra low noise detection technologies and understand their operation in a microwave cavity haloscope operating at a nominal frequency of approximately 5 GHz.

2 The Strong CP Problem

Measurements of the neutron electric dipole moment (nEDM) have failed to detect a significant EDM. Recent results from PSI [2] report an upper limit on this value of approximately 10^{-26} e cm. This is surprising as there is no reason why the neutron should not have an EDM. It would arise as a consequence of CP violation which is not forbidden by the QCD Lagrangian; this is known as the strong CP problem. Consequently, the evidence suggests that CP violation in the QCD sector is either coincidentally small or a physical principle is driving it towards zero.

One favoured mechanism for explaining the non-existence of CP violation is the Peccei-Quinn (PQ) mechanism [3]. Not only does it provide an elegant solution to the strong CP problem but it also produces a massive pseudo-scalar boson called the QCD axion, a potential dark matter candidate. However the mass of this axion remains unknown. Details of the PQ mechanism are discussed extensively in the literature [4, 5].

As the existence of QCD axions would alter Maxwell's equations they would interact with both electric and magnetic fields. Direct decay of these axions to photons is possible, but, as the coupling $g_{a\gamma\gamma}$ is expected to be very small due to the high symmetry-breaking energy scale of the axion field the QCD axion can be considered a stable particle. The rate of axion 'decay' can be increased via the inverse Primakoff effect [6, 7]. Here, QCD axions couple to virtual photons sourced from a strong magnetic field to produce real photons. The strength of the coupling is model-dependent. The effective axion to two photon coupling is given by

$$g_{a\gamma\gamma} = \frac{\alpha_{em}}{2\pi f_{PQ}} \left(\frac{E}{N} - \frac{2(4+z)}{3(1+z)} \right) \quad (1)$$

where z is the ratio of up and down quark masses and is assumed to be approximately 0.56. E is the axion electromagnetic anomaly, and N is the axion colour anomaly. f_{PQ} is a scaling constant, and is inversely proportional to the axion mass. Finally, α is the fine structure constant. Note the far right term, containing z approximately equals 1.95.

Axions must couple to colour-charge to solve the strong CP problem, but the tree-level coupling of axions to leptons is model dependent. One implementation of the PQ mechanism is the Kim-Shifman-Vainshtein-Zakharov (KSVZ) model [8] which assumes the coupling to electrons is strongly suppressed; here $E/N = 0$. A second model, the Dine-Fischler-Srednicki Zhitnitsky (DFSZ) model [9] assumes that there is coupling to both hadrons and leptons, and here $E/N = 8/3$. The bounded parameter space taken up by the axion mass to photon coupling rate as predicted by these QCD axion models is shown in figure 1.

3 QCD Axions and Resonant Detection

Microwave resonant cavity detection is suitable for searching for QCD axions in the mass range of μeV to $100 \mu\text{eV}$. In natural units, the mass of the axion is equal to the frequency of the photon produced via the inverse Primakoff effect; the axions are cold so the kinetic energy associated with their Maxwellian velocity distribution represents a small correction. Sensitivity to small values of $g_{a\gamma\gamma}$ required to detect QCD axions can be achieved using a tuneable cavity resonant in a TM_{010} mode at the frequency of interest, where the cavity is placed in a strong magnetic

field of several Tesla at cryogenic temperatures. The cavity size is constrained by the bore achievable with an affordable magnet.

Several experiments are currently operating and taking data in this mass range, see figure 1. Broadband searches, notably the CERN axion solar telescope (CAST), have already pushed the coupling rate down to $10^{-10} \text{ GeV}^{-1}$ over the mass range of interest [10]. As previously noted the signal output power from a microwave cavity experiment is expected to be very small, typically on the order of yocto-watts. It can be shown that the power output from a haloscope is given by

$$P = \left(\frac{g_{a\gamma\gamma}^2 \rho_H \hbar^2}{m_a^2 c} \right) c^2 \epsilon_0 B_0^2 V \omega_0 Q f_{nlm} \quad (2)$$

where ρ_H is the density of the halo, Q is the quality factor of the cavity and f_{nlm} represents the frequency of the cylindrical cavity mode. For cavity searches only the transverse magnetic modes are considered and primarily the TM_{010} mode as this provides the best form factor, ultimately maximising the power extracted by the cavity. Usually the terms are normalised to the experimental geometry and this gives a pre-factor of approximately a yocto-watt.

The signal to noise ratio of an antenna is given by the Dicke Radiometer equation. By suitable substitution of the power output from a microwave cavity into this equation a figure of merit for the cavity can be determined:

$$\text{Figure of Merit} \propto \frac{B^2 V}{T_S} \quad (3)$$

The cavity geometry is dictated by the search frequency. Multiple cavities may be operated in-phase to increase volume, at the cost of increased complexity. Magnets of bore of order 1 m utilising NbTi windings and achieving fields of around 8 T have not changed much in several decades, though higher field magnets using NbSn or high T_c superconductor windings can achieve higher fields, albeit at far lower volume and far higher cost. The system temperature T_S is comprised of the physical temperature of the cavity plus the noise temperature of the electronics. On the same timescale, significant improvements in the system temperature have been made due to improvements in the reliability and capabilities of dilution refrigerators, and the development of new quantum electronic amplification and photon counting techniques.

4 The QSHS facility

QSHS is a UK-based, STFC-funded, test stand that will couple various quantum sensors to a nominal 5 GHz cavity. A dilution fridge with a target temperature of 10 mK ($\geq 12 \mu\text{W}$ of cooling power at 20 mK) coupled with an 8 T magnet, has already been ordered from Oxford Instruments¹. The magnet has a nominal bore diameter of 20 cm and depth 20 cm, and can contain up to three small cavities. As part of the magnet design, field cancellation coils will give a low field region (≤ 100 gauss) a short distance away from the magnet core; this is crucial to the operation of some of the instrumentation. Refurbishment of lab space at Sheffield University has commenced. Installation of the dilution fridge is anticipated later in 2023.

Collaboration members are developing qubit arrays, bolometers, travelling wave parametric amplifiers and a superconducting low-inductance undulatory galvanometer, aka a SLUG, to provide higher frequency coupling to a cavity. Characterising the behaviour of these sensors, particularly for those sensors that may be sensitive to single photon counting in the low system temperature limit, is the principal aim of the QSHS test stand. A quantum mechanical

¹<https://www.oxinst.com/>




Figure 1: The red regions show the regions of axion parameter space explored by current and past microwave cavity experiments compared with the expected bounds of the QCD axion models shown by the diagonal yellow band [11].

understanding of the signal to noise ratio will help to improve the detector’s sensitivity to QCD axions and other hidden sector particles. It may also have implications for improving the scan rate. The facility will inform the UK expertise in haloscope development and pave the way for a large scale facility in hidden sector particle searches.

5 Conclusion

QSHS provides the UK with a technology test bench for collecting and analysing data from quantum technologies coupled to traditional microwave cavities at a target test temperature of 10 mK. QSHS has a memorandum of understanding with the Axion Dark Matter Experiment (ADMX) for cavity design and data analysis. The experience and expertise that ADMX [12] has will make a useful collaborative partner with this project.

Our longer term aim is to build up a core of expertise in the UK, so that a larger facility can be built to complement the existing searches for QCD axions and other hidden sector particles in the μeV to $100\mu\text{eV}$ mass window.

Funding information The authors would like to thank the UK Science and Technology Facilities Council (STFC) for funding this work through support for the Quantum Sensors for the Hidden Sector (QSHS) collaboration under grants ST/T006102/1, ST/T006242/1, ST/T006145/1, ST/T006277/1, ST/T006625/1, ST/T006811/1, ST/T006102/1 and ST/T006099/1.

References

- [1] L. Visinelli, *Axion electromagnetic waves*, Modern Physics Letters A **28**(35), 584 (2013), doi:[10.1142/S0217732313501629](https://doi.org/10.1142/S0217732313501629).
- [2] C. Abel, S. Afach, N. J. Ayres, C. A. Baker, G. Ban, G. Bison, K. Bodek, V. Bondar, M. Burghoff, E. Chanel, Z. Chowdhuri, P.-J. Chiu *et al.*, *Measurement of the permanent electric dipole moment of the neutron*, Phys. Rev. Lett. **124**, 081803 (2020), doi:[10.1103/PhysRevLett.124.081803](https://doi.org/10.1103/PhysRevLett.124.081803).
- [3] R. D. Peccei and H. R. Quinn, *CP conservation in the presence of pseudoparticles*, Phys. Rev. Lett. **38**, 1440 (1977), doi:[10.1103/PhysRevLett.38.1440](https://doi.org/10.1103/PhysRevLett.38.1440).
- [4] I. G. Irastorza, *An introduction to axions and their detection*, SciPost Phys. Lect. Notes p. 45 (2022), doi:[10.21468/SciPostPhysLectNotes.45](https://doi.org/10.21468/SciPostPhysLectNotes.45).
- [5] F. Chadha-Day, J. Ellis and D. J. E. Marsh, *Axion dark matter: What is it and why now?*, Science Advances **8**(8), eabj3618 (2022), doi:[10.1126/sciadv.abj3618](https://doi.org/10.1126/sciadv.abj3618).
- [6] P. Sikivie, *Experimental tests of the "invisible" axion*, Phys. Rev. Lett. **51**, 1415 (1983), doi:[10.1103/PhysRevLett.51.1415](https://doi.org/10.1103/PhysRevLett.51.1415).
- [7] P. Sikivie, *Experimental tests of the "invisible" axion - Errata*, Phys. Rev. Lett. **52**, 695 (1984), doi:[10.1103/PhysRevLett.52.695.2](https://doi.org/10.1103/PhysRevLett.52.695.2).
- [8] M. Shifman, A. Vainshtein and V. Zakharov, *Can confinement ensure natural CP invariance of strong interactions?*, Nuclear physics. B **166**(3), 493 (1980), doi:[10.1016/0550-3213\(80\)90209-6](https://doi.org/10.1016/0550-3213(80)90209-6).
- [9] M. Dine, W. Fischler and M. Srednicki, *A simple solution to the strong CP problem with a harmless axion*, Physics letters. B **104**(3), 199 (1981), doi:[10.1016/0370-2693\(81\)90590-6](https://doi.org/10.1016/0370-2693(81)90590-6).
- [10] V. Anastassopoulos, S. Aune, K. Barth, A. Belov, H. Brauning, G. Cantatore, J. M. Carmona, J. F. Castel, S. A. Cetin, F. Christensen, J. I. Collar, T. Dafni *et al.*, *New CAST limit on the axion–photon interaction*, Nature Physics **13**(6), 584 (2017), doi:[10.1038/nphys4109](https://doi.org/10.1038/nphys4109).
- [11] C. O'Hare, *cajohare/axionlimits: Axionlimits*, doi:[10.5281/zenodo.3932430](https://doi.org/10.5281/zenodo.3932430) (2020).
- [12] C. Bartram, T. Braine, E. Burns, R. Cervantes, N. Crisosto, N. Du, H. Korandla, G. Leum, P. Mohapatra, T. Nitta, L. J. Rosenberg, G. Rybka *et al.*, *Search for invisible axion dark matter in the 3.3 – 4.2 μeV mass range*, Phys. Rev. Lett. **127**, 261803 (2021), doi:[10.1103/PhysRevLett.127.261803](https://doi.org/10.1103/PhysRevLett.127.261803).