

The role of the underlying event in the charm-baryon enhancement observed in pp collisions at LHC energies

Zoltán Varga^{1,2,*}, Anett Misák^{1,3} and Róbert Vértesi¹

¹ Wigner RCP, Budapest, Hungary

² Budapest University of Technology and Economics, Budapest, Hungary

³ Eötvös University, Budapest, Hungary

* varga.zoltan@wigner.hu

October 3, 2022



51st International Symposium on Multiparticle Dynamics (ISMD2022)

Pitlochry, Scottish Highlands, 1-5 August 2022

doi:[10.21468/SciPostPhysProc.7](https://doi.org/10.21468/SciPostPhysProc.7)

Abstract

We study the enhanced production of Λ_c charmed baryons relative to that of charmed D^0 mesons in proton–proton collisions at LHC energies. We simulated collision events with the enhanced color-reconnection model in PYTHIA 8 MC generator and propose measurements based on the comparative use of different event-activity classifiers to identify the source of the charmed-baryon enhancement. We demonstrate that in this enhanced color-reconnection scenario the excess production is primarily linked to the underlying event and not to the production of high-momentum jets.

Contents

1	Introduction	1
2	Results	2
3	Conclusion	3
	References	4

1 Introduction

The factorization hypothesis states that the production cross-section of heavy-flavor hadrons can be calculated as the convolution of three independent terms: the parton distribution function of the colliding hadrons, the production cross sections of the heavy-quarks in the hard partonic process, and finally the fragmentation functions of the heavy-flavor quarks into the given heavy-flavor hadron species. The fragmentation function has been traditionally treated as universal, i.e. independent of the collision system.

Recent charmed-baryon measurements show a low-momentum enhancement over model predictions which are based on e^+e^- collisions, which challenges this traditional assumption of universality [1,2]. One of the latest measurements also shows that the charm-baryon enhancement depends on the final-state multiplicity of the collision event [3]. Several scenarios have been proposed to explain the emerging pattern, including string formation beyond leading color, the so-called enhanced color re-connection [4], which provides a qualitatively correct description of these findings for pp collisions.

In this proceedings, we briefly recapitulate our results on investigating the charm-baryon enhancement with PYTHIA 8 Monte-Carlo generator and enhanced color-reconnection models [5] and compare the calculations to new results from ALICE [3]. We propose measurements based on several event-activity classifiers to identify the source of the charm-baryon enhancement.

2 Results

We focus on two event classifiers in this proceedings, both of them are used when we have a $p_T^{\text{trig}} > 5$ GeV/c trigger particle in the selected event. One of them is the transverse event activity classifier $R_T \equiv N_{\text{ch}}^{\text{trans}} / \langle N_{\text{ch}}^{\text{trans}} \rangle$ [6], where $N_{\text{ch}}^{\text{trans}}$ is the charged-hadron multiplicity in the transverse region. It is defined with the azimuth angle relative to the trigger hadron being $\frac{\pi}{3} < |\Delta\phi| < \frac{2\pi}{3}$ within $|\eta| < 1$.

The other one is the near-side cone activity classifier $R_{\text{NC}} \equiv N_{\text{ch}}^{\text{near-side cone}} / \langle N_{\text{ch}}^{\text{near-side cone}} \rangle$ [5], where $N_{\text{ch}}^{\text{near-side cone}}$ is the charged-hadron multiplicity in a narrow cone around the trigger particle, where $\sqrt{\Delta\phi^2 + \Delta\eta^2} < 0.5$. Due to the constraints of space in this proceedings, we refer to the description of the remaining details of the analysis method in [5].

In Fig. 1 we show the Λ_c to D^0 ratios for different charged event multiplicity bins using CR Mode 2 in PYTHIA 8. The minimum-bias results, i.e. where no event-activity selection was applied, are compared to ALICE measurements [3] for reference. We note that the trends are different for the Λ_c produced directly and those coming from Σ_c and this needs further experimental investigation.




Figure 1: The Λ_c to D^0 ratios in function of p_T for different event multiplicity bins. The ratios for only the directly produced Λ_c are also shown with a dotted line.

In Fig. 2 we plot the Λ_c to D^0 ratios in function of p_T for the R_T and R_{NC} event activity

classifiers. We observe that the Λ_c enhancement significantly depends on the underlying event activity, however there is no difference among the jet region activity classes. We can draw the conclusion that the mechanism beyond this enhancement is connected to the underlying event, rather than the jet region. The exact values of the activity classifier bins can be found in [5].




Figure 2: The Λ_c to D^0 ratios are plotted in function of p_T for different R_T bins (left) and R_{NC} bins (right).

3 Conclusion

We simulated the Λ_c to D^0 yield ratios with PYTHIA 8 for the enhanced color re-connection model. Our results were compared to ALICE experimental data for different event multiplicity classes. A difference can be observed between the Λ_c yields coming from Σ_c and those produced directly, which needs to be investigated further experimentally.

By investigating the enhancement with different event activity classifiers, we conclude that within the enhanced color re-connection model, the excess Λ_c production is connected to the underlying event and not the jet production [5]. These observables provide a unique opportunity in the upcoming measurements from the high-luminosity LHC Run3 phase to understand charm fragmentation mechanisms, and will serve as valuable means for further model development.

Acknowledgements

Funding information This work was supported by the Hungarian National Research, Development and Innovation Office (NKFIH) under the contract numbers OTKA FK131979 and K135515, and the NKFIH grant 2019-2.1.11-TÉT-2019-00078. The authors acknowledge the computational resources provided by the Wigner GPU Laboratory and research infrastructure provided by the Eötvös Loránd Research Network (ELKH).

References

- [1] A. M. Sirunyan *et al.*, *Production of Λ_c^+ baryons in proton-proton and lead-lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV*, Phys. Lett. B **803**, 135328 (2020), doi:[10.1016/j.physletb.2020.135328](https://doi.org/10.1016/j.physletb.2020.135328), [1906.03322](https://arxiv.org/abs/1906.03322).
- [2] S. Acharya *et al.*, *Λ_c^+ Production and Baryon-to-Meson Ratios in pp and p-Pb Collisions at $\sqrt{s_{NN}}=5.02$ TeV at the LHC*, Phys. Rev. Lett. **127**(20), 202301 (2021), doi:[10.1103/PhysRevLett.127.202301](https://doi.org/10.1103/PhysRevLett.127.202301), [2011.06078](https://arxiv.org/abs/2011.06078).
- [3] S. Acharya *et al.*, *Observation of a multiplicity dependence in the p_T -differential charm baryon-to-meson ratios in proton-proton collisions at $\sqrt{s} = 13$ TeV*, Phys. Lett. B **829**, 137065 (2022), doi:[10.1016/j.physletb.2022.137065](https://doi.org/10.1016/j.physletb.2022.137065), [2111.11948](https://arxiv.org/abs/2111.11948).
- [4] J. R. Christiansen and P. Z. Skands, *String Formation Beyond Leading Colour*, JHEP **08**, 003 (2015), doi:[10.1007/JHEP08\(2015\)003](https://doi.org/10.1007/JHEP08(2015)003), [1505.01681](https://arxiv.org/abs/1505.01681).
- [5] Z. Varga and R. Vertesi, *The role of the underlying event in the enhancement in high-energy pp collisions*, J. Phys. G **49**(7), 075005 (2022), doi:[10.1088/1361-6471/ac7248](https://doi.org/10.1088/1361-6471/ac7248), [2111.00060](https://arxiv.org/abs/2111.00060).
- [6] T. Martin, P. Skands and S. Farrington, *Probing Collective Effects in Hadronisation with the Extremes of the Underlying Event*, Eur. Phys. J. C **76**(5), 299 (2016), doi:[10.1140/epjc/s10052-016-4135-4](https://doi.org/10.1140/epjc/s10052-016-4135-4), [1603.05298](https://arxiv.org/abs/1603.05298).