

# Muon-induced background in a next-generation dark matter experiment based on liquid xenon

V. Pěč<sup>1\*</sup> and V. A. Kudryavtsev<sup>2</sup>

<sup>1</sup> FZU – Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic<sup>i)</sup>

<sup>2</sup> Department of Physics and Astronomy, University of Sheffield, Sheffield, United Kingdom

\* viktor.pec@fzu.cz



*14th International Conference on Identification of Dark Matter*  
Vienna, Austria, 18-22 July 2022  
doi:[10.21468/SciPostPhysProc.7](https://doi.org/10.21468/SciPostPhysProc.7)

Make abstract only 8 lines long

## Abstract

We have investigated the implication of laboratory depth on the muon induced background in a future dark matter experiment capable of reaching the so-called neutrino floor. Our simulation study focuses on a xenon-based detector with 71 tonnes of active mass, surrounded by additional veto systems including an instrumented water shield. Two locations at the Boulby Underground Laboratory (UK) at depths of 2850 m w. e. and 3575 m w. e. served as a case study. Our results show that less than one event of cosmogenic background is likely to survive standard analysis cuts for 10 years of operation at either location. The largest background component that we identified comes from delayed neutron emission from  $^{17}\text{N}$  which is produced from  $^{19}\text{F}$  in the fluoropolymer components of the experiment. Our results confirm that a dark matter search with sensitivity to the neutrino floor is viable (from the point of view of cosmogenic backgrounds) in underground laboratories at these levels of rock overburden.

## 1 Introduction

The choice of underground laboratory for a future high-sensitivity experiment for rare-event searches will be heavily dependent on the depth of the site, as events triggered by cosmic-ray muons can constitute significant background to the signal events. For example, in searches for dark matter (DM) Weakly Interacting Massive Particles (WIMP), isolated neutrons originating in the muon activity can mimic dark matter interaction if scattering only once in the active region producing a single nuclear recoil (NR). We focused our attention on a potential next-generation DM experiment based on Liquid Xenon Time Projection Chamber (LXe-TPC) technology, and we investigated whether the depth of about 3 km w. e. was sufficient for such experiment.

The work was performed as a part of feasibility study for Boulby Mine (UK) to host the next-generation LXe-TPC experiment [1] and we wanted to learn whether the current underground

<sup>i)</sup>Previously at the University of Sheffield

laboratory in Boulby has sufficient depth with respect to the cosmic-ray induced backgrounds. A new larger site would have to be excavated nearby the existing laboratory which is at 1100 m level underground, or 2850 m w. e., and it is located within a layer of salt (NaCl). We also considered a new site at a deeper location at a 1400 m level, or 3575 m w. e. underground. The Boulby mine served as a case model, however, the results are relevant to sites of similar depth.

We have also compared our Geant4 results on neutron production in simplified-geometry simulations for various materials and muon energies with other simulations and experimental data. Some results and distributions of interest are included in Ref. [2].

## 2 Simulation of cosmic-ray muons

### 2.1 Geometry

We carried out simulations to determine the rate of potential background events caused by cosmic-ray muons in a next generation dark matter experiment with design based on the design of LUX-ZEPLIN (LZ) detector [3]. The main detector is a dual-phase xenon time projection chamber (hereafter LXe-TPC) containing 70 tonnes of active liquid xenon (LXe), corresponding to a  $\sim 10$ -fold upscale of existing experiments LZ [4] and XENONnT [5].

We implemented a simplified detector geometry model with the main elements included: a vacuum cryostat approximately 4 m in diameter and 5 m in height containing the xenon detector, with an anti-coincidence veto system made of gadolinium-loaded liquid scintillator (GdLS), 50 cm thick, surrounding the cryostat, and all submerged in a water tank with 12 m in diameter and 11 m in height for shielding from local radioactivity backgrounds. The water tank was placed into a cylindrical cavern of diameter and height of 30 m. The cavern was surrounded by rock material with at least 5 m on the sides and 7 m and 3 m at the top and the bottom, respectively. We tested two different rock compositions, one with salt (NaCl,  $2.17 \text{ g/cm}^3$ ) corresponding to the current laboratory location in Boulby mine, and one with polyhalite ( $\text{K}_2\text{Ca}_2\text{Mg}(\text{SO}_4)_4 \cdot 2\text{H}_2\text{O}$ ,  $2.78 \text{ g/cm}^3$ ) corresponding to the potential new site. Overview of the geometry can be seen in the cross-sectional view of the cavern in the illustration in Fig. 1a.

The central part of the cryostat was the active LXe-TPC, cylindrical, 3.5 m and 2.5 m in diameter and height. There was an 8 cm thick layer of LXe around the active volume separated by a PTFE field cage, which will be called LXe skin hereafter. Arrays of photo-multiplier tubes (PMT) meant to read the fast scintillation light from LXe and delayed electroluminescence signal from Xe gas (GXe) of the TPC were approximated with uniform cylinders made of steel with reduced density of  $0.4 \text{ g/cm}^3$ , or about 5% of the standard density of steel, simulating metal components of the structure of the arrays and matching its mass. Illustration of a cross-sectional view of the cryostat is shown in Fig. 1b.

The water tank, GdLS and LXe skin volumes were assumed to be instrumented with photo-sensitive detectors and each of them would serve as a veto system by registering either Cherenkov light in the case of the water tank, or scintillation light in the case of GdLS and LXe skin.

### 2.2 Underground muons

Cosmic-ray muons were sampled on the top and side surfaces of a 40 m cube that surrounded. Distributions of primary energies and directions were calculated using the MUSIC and MUSUN codes [6,7] (Ref. [7] describes the procedure and muon transport through rock down to experimental site). The mean muon energy and zenith angle were calculated as 261 GeV (134 GeV median) and  $30.6^\circ$  ( $30.1^\circ$  median), respectively. The rate of simulated muons was  $0.8759 \text{ s}^{-1}$  for the existing Boulby site within salt at 2850 m w. e. vertical overburden. This rate was determined based on the measured flux in the existing laboratory,  $(3.75 \pm 0.09) \times 10^{-8} \text{ cm}^{-2}\text{s}^{-1}$  [8]. The surface profile




Figure 1: Illustrations of the geometry model used in the simulations. (a) Cross-sectional view of the cavern (bluish grey) within the rock (red) with the water shield (light blue) holding the scintillator (green) and the detector cryostat. (b) Cross-sectional view of the cryostat with the main LXe-TPC (blue).

80 was assumed to be flat in these simulations (in reality, variations in elevation up to 30 m exist on  
 81 the surface over areas of a few km<sup>2</sup>). For the proposed deeper site in polyhalite at 3575 m w. e.  
 82 vertical overburden, the same muon distributions were used, but the equivalent sampling rate was  
 83 recalculated to be 0.2625 s<sup>-1</sup>.

84 We used Geant4 version 10.5 simulation toolkit to simulate the transport of muons through the  
 85 modelled experimental site. Physical processes were modelled according to the toolkit's modular  
 86 physics list Shielding. In total, 800 million muons were simulated for each rock material, salt and  
 87 polyhalite. These numbers correspond to approximately 29 years and 97 years of live time of the  
 88 experiment, respectively.

### 89 3 Neutron background analysis

90 The expected WIMP signature in a typical in a xenon-based dark matter experiment consists of a  
 91 single scatter event at low energy, usually  $\lesssim 50$  keV, which is classified as a nuclear recoil (NR)  
 92 using specific discrimination techniques, and which is in anti-coincidence with veto systems. We  
 93 adopted a simple background counting technique with the potential (irreducible) background iden-  
 94 tified as a single NR of energy above 1 keV.

95 We analysed energy depositions in active regions of the experimental setup. The detector  
 96 response to the depositions was not simulated (i.e. the digitised PMT waveforms resulting from  
 97 the fast scintillation light from LXe and delayed electroluminescence signal from GXe from each  
 98 energy deposition in the active volume) and it was considered only in terms of the characteristic  
 99 times over which signals were collected and the equivalent energy thresholds in the respective  
 100 active volumes – LXe-TPC, LXe skin, liquid scintillator, water tank. We selected events with  
 101 depositions above 1 keV from a single NR at least 5 cm from the boundary of the active region  
 102 in 1 ms readout time window of the LXe-TPC. We required there was no other NR recoil above  
 103 0.5 keV, which we deemed resolvable in the delayed signal from GXe. We also required there  
 104 were no non-NR depositions above 10 keV in total, including a quenching factor of 4 for protons  
 105 and heavier ionising particles.

106 Energy depositions in the skin, liquid scintillator and water tank were summed over 1  $\mu$ s,

irrespective of their origin. We chose this time window to emulate the shaping time of the photo-sensitive instrumentation of the systems. We chose thresholds of 100 keV, 200 keV and 200 MeV in the skin, liquid scintillator and water tank, respectively, to trigger veto signals. The selected events in the LXe-TPC were then tested for anti-coincidence with the veto signals by requiring no veto signal to be present within 0.5 ms before or after any TPC signal.

Spectra of depositions in the LXe-TPC at various stages of the selection are shown in Fig. 2a. Depositions in all events with some NR are compared to events where only NR happened. One can see the low-energy part ( $<100$  keV) of all the events is dominated by depositions from NR. The same distributions are also shown after application of the veto condition. Recognized activity near the active region of the TPC (in the LXe skin, GdLS, or the water tank) together with the requirement on only NR to be observed reduces the number of events by a factor  $10^6$ .

We also wanted to answer the question whether the scintillator veto was necessary for reduction of the cosmogenic background. We reused the simulated samples and emulated the geometry without the GdLS by assuming all depositions within its volume were part of the water tank, and applied the corresponding 200 MeV threshold on the depositions from the combined volume. Fig. 2b shows comparison of deposition spectra the same way as in Fig. 2a but this time the GdLS was not included in the veto. The number of events in the LXe-TPC passing the veto is larger than in the case with the GdLS. We must note here, that these are events before the requirement on multiplicity of NR was applied.




Figure 2: Energy depositions in the active region of the LXe-TPC for NaCl as the rock material. Events with any NR activity were considered. (a) Full configuration containing the GdLS. (b) Configuration where GdLS was considered to be part of the water shield.

After application of the full selection process described above, including the rejection of multiple scatters and exclusion of 5 cm of LXe from all sides of the TPC (64 t fiducial volume), no events passed the selection in the sample with NaCl as the material of the surrounding rock. A single event passed the selection in the sample with polyhalite. As a result, the estimated background event rate in the fiducial volume of the LXe-TPC (64 t) is  $<0.84$  events (90% CL) in 10 years for the 2850 m w. e. site in salt. For the 3578 m w. e. site in polyhalite the rate is estimated to be 0.10 (0.01–0.45 at 90% CL) events in 10 years of running.

In the case where no liquid scintillator is used as an additional veto system, no events were observed for the site in salt, and the total of 2 events were observed for the site in polyhalite. The respective estimated background rates correspond to  $<0.84$  events (90% CL) in 10 years (2850 m w. e. site, salt), and 0.31 (0.11–0.77 at 90% CL) events in 10 years (3575 m w. e. site, polyhalite). The limits and confidence intervals are statistical only. The estimated rates in both cases are well below the expected background of tens of events in 10 years extrapolated from the estimates in Table III of Ref. [9].

140 The observed single nuclear recoils are located at the boundary of the fiducial volume. Most  
141 events with isolated neutrons causing NRs in the LXe-TPC and avoiding the veto were coming  
142 from delayed neutron emission from the PTFE field cage. Secondary particles in muon-induced  
143 cascade activated fluorine producing  $^{17}\text{N}$  from  $^{19}\text{F}$ .  $^{17}\text{N}$  undergoes  $\beta$  decay with half-life of 4.2 s  
144 with emission of a neutron.

145 We also made considerations of systematic uncertainties. We concluded that the major uncer-  
146 tainty was due to the neutron production modelling in Geant4. From comparisons with previous  
147 studies (see, for instance, Refs. [8, 10] for discussions of simulations and comparison of different  
148 models with experimental data) we conservatively assessed the uncertainty to be a factor of 2.  
149 Uncertainties in the muon flux are linked to an unknown location of the new cavern within the ex-  
150 isting level. Our estimate of 10% was based on measurements at different locations. We estimated  
151 the uncertainty of 20% for the deeper site where the flux was calculated based on the geophysical  
152 model of the Boulby mine but the exact location has not been determined. We approximated the  
153 muon spectrum at the deeper site to be the same as the one at the current laboratory. However, the  
154 calculated mean muon energies differ by about 9% (261 GeV and 282 GeV for the 2850 m w. e.  
155 and 3575 m w. e. sites, respectively) and we expect it would lead to a small increase in the neutron  
156 production yield of (6–7)%. This change is small compared with the other mentioned systematic  
157 uncertainties.

## 158 4 Conclusion

159 We have performed simulations for an experiment similar in design to the LZ detector, upscaled  
160 to 71 tonnes of LXe acting as a target. We conclude that, after applying a standard simplified  
161 analysis procedure and cuts, the event rate caused by cosmogenic activity stays below 1 event per  
162 10 years in the fiducial volume of the LXe-TPC (64 tonnes). This rate is well below the expected  
163 background of tens of events from beta decays from radon progeny and ERs/NRs from physics  
164 backgrounds such as two-neutrino double beta decay of  $^{136}\text{Xe}$  and solar/atmospheric neutrinos  
165 with ER events leaking into NR band due to limited discrimination. From the point of view of  
166 cosmogenic background, the depth of about 3 km w. e. or deeper is sufficient for a next generation  
167 dark matter experiment based on liquid xenon. Although there are significant systematic uncer-  
168 tainties related mainly to the modeling of neutron production, they cannot change our conclusion.  
169 The observed residual background of NR events comes from the production and delayed  $\beta - n$   
170 decay of  $^{17}\text{N}$  in PTFE (on fluorine) where only a single neutron scatter is detected. Our material  
171 budget contained about 2.8 t of PTFE. Although the residual background is very low, the design  
172 of a future experiment may need to limit PTFE usage to the necessary minimum.

173 We have also investigated two designs of veto system: a default one with instrumented liquid  
174 scintillator surrounding the cryostat, and an option without the scintillator. No significant differ-  
175 ence was observed between the two scenarios which lead us to the conclusion that the additional  
176 veto system is not required to suppress cosmogenic backgrounds for the sensitivity at the stud-  
177 ied depth. This conclusion, however, does not apply to other types of backgrounds (mainly from  
178 detector components) where liquid scintillator is particularly efficient in tagging neutron-induced  
179 events. The decision about the need for a scintillator veto can only be taken after a detailed simu-  
180 lation of all types of background from all components is completed.

## 181 Acknowledgments

182 We wish to thank UKRI-STFC for financial support of the Boulby case study and the whole team  
183 who worked on the Boulby Feasibility Study project.

184 **References**

- 185 [1] H. M. Araújo *et al.*, *Feasibility study for developing the Boulby underground laboratory into*  
186 *a facility for future major international projects*, Report to UKRI-STFC (2021).
- 187 [2] V. Pěč and V. A. Kudryavtsev, *Muon-induced background in a next-generation dark mat-*  
188 *ter experiment based on liquid xenon*, presentation file at [https://indico.cern.ch/event/](https://indico.cern.ch/event/922783/contributions/4892720/)  
189 [922783/contributions/4892720/](https://indico.cern.ch/event/922783/contributions/4892720/), 14th International Conference on Identification of Dark  
190 Matter (IDM2022) (2022).
- 191 [3] D. S. Akerib *et al.*, *The LUX-ZEPLIN (LZ) Experiment*, Nucl. Instrum. Meth. A **953**, 163047  
192 (2020), doi:[10.1016/j.nima.2019.163047](https://doi.org/10.1016/j.nima.2019.163047), [1910.09124](https://arxiv.org/abs/1910.09124).
- 193 [4] J. Aalbers *et al.*, *First dark matter search results from the LUX-ZEPLIN (LZ) experiment*  
194 (2022), [2207.03764](https://arxiv.org/abs/2207.03764).
- 195 [5] E. Aprile *et al.*, *Search for new physics in electronic recoil data from xenonnT* (2022), [2207.](https://arxiv.org/abs/2207.11330)  
196 [11330](https://arxiv.org/abs/2207.11330).
- 197 [6] P. Antonioli, C. Ghetti, E. V. Korolkova, V. A. Kudryavtsev and G. Sartorelli, *A Three-*  
198 *dimensional code for muon propagation through the rock: MUSIC*, Astropart. Phys. **7**, 357  
199 (1997), doi:[10.1016/S0927-6505\(97\)00035-2](https://doi.org/10.1016/S0927-6505(97)00035-2), [hep-ph/9705408](https://arxiv.org/abs/hep-ph/9705408).
- 200 [7] V. A. Kudryavtsev, *Muon simulation codes MUSIC and MUSUN for underground physics*,  
201 Comput. Phys. Commun. **180**, 339 (2009).
- 202 [8] L. Reichhart *et al.*, *Measurement and simulation of the muon-induced neutron yield in lead*,  
203 Astropart. Phys. **47**, 67 (2013).
- 204 [9] D. S. Akerib *et al.*, *Projected WIMP sensitivity of the LUX-ZEPLIN dark matter experiment*,  
205 Phys. Rev. D **101**(5), 052002 (2020), doi:[10.1103/physrevd.101.052002](https://doi.org/10.1103/physrevd.101.052002).
- 206 [10] L. Lindote *et al.*, *Simulation of neutrons produced by high-energy muons underground*,  
207 Astropart. Phys. **31**, 366 (2009).