

Comments on the Negative grade KdV Hierarchy

Y. F. Adans, J.F. Gomes* G.V. Lobo and A.H. Zimerman

Instituto de Física Teórica, IFT-Unesp
 Rua Dr. Bento Teobaldo Ferraz, 271, Bloco II,
 CEP 01140-070, São Paulo - SP, Brasil.
 * francisco.gomes@unesp.br

December 17, 2022



34th International Colloquium on Group Theoretical Methods in Physics
 Strasbourg, 18-22 July 2022
 doi:[10.21468/SciPostPhysProc.7](https://doi.org/10.21468/SciPostPhysProc.7)

Abstract

The construction of negative grade KdV hierarchy is proposed in terms of a Miura-gauge transformation. Such gauge transformation is employed within the zero curvature representation and maps the Lax operator of the mKdV into its counterpart in the KdV setting. Each odd negative KdV flow is obtained from an odd and its subsequent even negative mKdV flows. The negative KdV flows are shown to inherit the two different vacua structure that characterizes the associated mKdV flows.

Contents

1	Introduction	1
2	mKdV negative hierarchy	2
3	KdV negative hierarchy	4
4	Miura Transformation and Soliton Solutions	6
5	Conclusion	7
	References	8

1 Introduction

Integrable models have been focus of considerable attention in the past few years. These are very peculiar two dimensional field theories admitting an infinite number of conservation laws and soliton solutions. The algebraic construction of integrable models has provided a series of important achievements which allows its construction and classification in terms of the decomposition of the affine algebra into graded subspaces. Structural connection and the derivation

of many properties as the construction of conservation laws and soliton solutions can be set from the zero curvature representation [1], [2]. In particular the mKdV hierarchy, based on the affine $sl(2)$ algebra, provides the simplest example of systematic construction of a series of evolution equations associated to a universal object called Lax operator. For the mKdV case the relevant decomposition occurs according to the principal gradation. Explicit constructions for positive and negative graded sub-hierarchies have been obtained. The positive flows are known to be labelled by odd numbers whilst there are no restriction for the negative [3].

An interesting relation between the KdV and mKdV hierarchies can be realised by the Miura transformation which maps one hierarchy into the other. In ref. [4], [5] we have related the two hierarchies by a gauge transformation that maps one Lax operator into the other. Such *Miura-gauge transformation* acting upon the zero curvature maps the flows from one hierarchy into the other. For the positive sub-hierarchy the mapping is one to one, i.e., each flow equation of mKdV is mapped into its counterpart within the KdV hierarchy. However this is not true for the negative KdV sub-hierarchy. In sec. 3 we argue that only odd flows are consistent and hence since there are even and odd flows within the negative mKdV side, there shall be a mapping of a pair of mKdV flows into a single KdV flow. This is indeed true, in sect. 4 we construct these mappings and show that an odd and its subsequent even mKdV flows can be mapped into a single KdV flow. An interesting point to mention is that odd mKdV flows admit only *zero vacuum* whilst the even admit strictly *non-zero vacuum* solutions and the associated KdV flow ends up inheriting both types of structure.

2 mKdV negative hierarchy

In this section let us review the construction of mKdV hierarchy within the algebraic formalism. Consider the affine $\mathcal{G} = \hat{sl}(2)$ centerless Kac-Moody algebra generated by

$$h^{(m)} = \lambda^m h^{(0)}, \quad E_{\pm\alpha}^{(m)} = \lambda^m E_{\pm\alpha}^{(0)} \quad \text{with } \lambda \in \mathbb{C} \quad \text{and } n \in \mathbb{Z} \quad (1)$$

satisfying the following algebra

$$[h^{(m)}, E_{\pm\alpha}^{(n)}] = \pm 2E_{\pm\alpha}^{(m+n)}, \quad [E_{\alpha}^{(m)}, E_{-\alpha}^{(n)}] = h^{(m+n)}. \quad (2)$$

Introduce the principal grading operator

$$Q_p = 2\lambda \frac{d}{d\lambda} + \frac{1}{2}h \quad (3)$$

that decomposes the affine algebra into graded subspaces, i.e., $\mathcal{G} = \bigoplus_i \mathcal{G}_i$ with

$$[Q_p, \mathcal{G}_a] = a\mathcal{G}_a, \quad [\mathcal{G}_a, \mathcal{G}_b] \in \mathcal{G}_{a+b}, \quad a, b \in \mathbb{Z}, \quad (4)$$

where, for $\mathcal{G} = \hat{sl}(2)$,

$$\mathcal{G}_{2n} = \{h^{(n)} = \lambda^n h\}, \quad \mathcal{G}_{2n+1} = \{\lambda^n (E_{\alpha} + \lambda E_{-\alpha}), \lambda^n (E_{\alpha} - \lambda E_{-\alpha})\}. \quad (5)$$

A second important ingredient is the choice of a constant grade one element $E^{(1)} \in \mathcal{G}_1$

$$E^{(1)} = E_{\alpha}^{(0)} + E_{-\alpha}^{(1)} \quad (6)$$

such that it decomposes the affine algebra as $\hat{\mathcal{G}} = \mathcal{K} \oplus \mathcal{M}$, where \mathcal{K} is the *Kernel* of $E^{(1)}$:

$$\mathcal{K}_E = \{y \in \mathcal{K}, [y, E^{(1)}] = 0\} = \{E_{\alpha}^{(n)} + E_{-\alpha}^{(n+1)}\} \in \mathcal{G}_{2n+1} \quad (7)$$

and \mathcal{M} is its complement subspace. We now define the spatial Lax operator to be an universal algebraic object within the whole hierarchy to be

$$A_x^{\text{mKdV}}(\phi) = E^{(1)} + A^{(0)}(\phi) = E_\alpha^{(0)} + E_{-\alpha}^{(1)} + \partial_x \phi h^{(0)} = \begin{pmatrix} \partial_x \phi & 1 \\ \lambda & -\partial_x \phi \end{pmatrix} \quad (8)$$

where $v(x, t_{-N}) = \partial_x \phi$ is the field of the theory. We are interested in the negative time flows generated by the temporal Lax operator component of the form [3]

$$A_{t_{-N}}^{\text{mKdV}} = D^{(-N)} + D^{(-N+1)} + \dots + D^{(-1)}, \quad N = 1, 2, \dots \quad (9)$$

where $D^{(i)} \in \mathcal{G}_i$. Thus, for a given integer N , the zero curvature equation

$$[\partial_x + E^{(1)} + A^{(0)}, \partial_{t_{-N}} + D^{(-N)} + D^{(-N+1)} + \dots + D^{(-1)}] = 0 \quad (10)$$

decomposes according to the grading structure, i.e.,

$$[A^{(0)}, D^{(-N)}] + \partial_x D^{(-N)} = 0, \quad (11)$$

$$[A^{(0)}, D^{(-N+1)}] + [E^{(1)}, D^{(-N)}] + \partial_x D^{(-N+1)} = 0, \quad (12)$$

$$\vdots \quad \vdots$$

$$[E^{(1)}, D^{(-1)}] - \partial_{t_N} A^{(0)} = 0. \quad (13)$$

These eqns. can be solved grade by grade in order to determine $D^{(i)}$ and the evolution equation for $A^{(0)}(\phi)$ according to time t_{-N} is given by (13).

The simplest case is found by taking $N = 1$, leading to

$$A_{t_{-1}}^{\text{mKdV}} = e^{-2\phi} E_\alpha^{(-1)} + e^{2\phi} E_{-\alpha}^{(0)} = \begin{pmatrix} 0 & \lambda^{-1} e^{-2\phi} \\ e^{2\phi} & 0 \end{pmatrix} \quad (14)$$

associated with the well know sinh-Gordon equation,

$$\phi_{x, t_{-1}} = e^{2\phi} - e^{-2\phi}. \quad (15)$$

Notice that $v = \partial_x \phi = v_0 = \text{const.}$ is the vacuum solution of (15) only if $v_0 = 0 \rightarrow \phi = 0$. It therefore follows that the sinh-Gordon equation only admits zero vacuum solution.

Considering now $N = 2$, we find

$$\begin{aligned} A_{t_{-2}}^{\text{mKdV}} &= h^{(-1)} + (2e^{-2\phi} d^{-1}(e^{2\phi})) E_\alpha^{(-1)} - 2e^{2\phi} d^{-1}(e^{-2\phi}) E_{-\alpha}^{(0)} \\ &= \begin{pmatrix} \lambda^{-1} & \lambda^{-1} (2e^{-2\phi} d^{-1}(e^{2\phi})) \\ -2e^{2\phi} d^{-1}(e^{-2\phi}) & -\lambda^{-1} \end{pmatrix} \end{aligned} \quad (16)$$

where we have denoted $d^{-1} f = \int_0^x f dx'$. It leads to the following nonlocal equation of motion

$$\phi_{x, t_{-2}} = -2(e^{-2\phi} d^{-1}(e^{2\phi}) + e^{2\phi} d^{-1}(e^{-2\phi})). \quad (17)$$

Notice that $v = \partial_x \phi = v_0 = \text{const.}$ is the vacuum solution of (17) only if $v_0 \neq 0 \rightarrow \phi = v_0 x$. Such equation does not admit zero vacuum solution, but a constant vacuum solution $v = v_0$ ($\phi = v_0 x$), instead. In fact, it can be showed that all models associated to negative even values of N only admit non-zero vacuum solutions [3]. Let us consider the zero curvature equation in the vacuum regime, i.e.,

$$[E^{(1)} + v_0 h^{(0)}, D_{\text{vac}}^{(-N)} + D_{\text{vac}}^{(-N+1)} + \dots + D_{\text{vac}}^{(-1)}] = 0 \quad (18)$$

the lowest grade equation is

$$[v_0 h^{(0)}, D_{vac}^{(-N)}] = 0 \quad (19)$$

thus if $v_0 \neq 0$ $D_{vac}^{(-N)}$ must commute with $h^{(0)}$ and therefore $D_{vac}^{(-N)} \in \mathcal{G}_{-2n}$ and $N = 2n$. Conversely if $v_0 = 0$ the lowest grade eqn. becomes

$$[E^{(1)}, D_{vac}^{(-N)}] = 0 \quad (20)$$

thus $D_{vac}^{(-N)} \in \mathcal{K}_E$ and N is odd. Thus, the negative mKdV hierarchy splits in two sub-hierarchies: one even admitting strictly non-zero vacuum ($v_0 \neq 0$) and one odd admitting, only zero vacuum ($v_0 = 0$) solutions. The systematic construction of soliton solutions for the negative mKdV hierarchies was previously studied and can be written as follows (see [3]). For the odd sub-hierarchy the one soliton solution was constructed from dressing the zero vacuum solution leading to

$$v(x, t_{-2n+1}) = \partial_x \ln \left(\frac{1 - \beta e^{2kx + 2k^{-2n+1} t_{-2n+1}}}{1 + \beta e^{2kx + 2k^{-2n+1} t_{-2n+1}}} \right) \quad \text{with} \quad \omega_{-2n+1} = 2k^{-2n+1}. \quad (21)$$

For the even sub-hierarchy the constant value of the vacuum, v_0 introduces a deformation upon the dressing method. In [3] the solutions were constructed employing deformed vertex operators yielding for the one soliton,

$$v(x, t_{-2n}) = v_0 + \partial_x \ln \left(\frac{1 + \beta(v_0 - k)e^{2kx + \omega_{-2n} t_{-2n}}}{1 + \beta(v_0 + k)e^{2kx + \omega_{-2n} t_{-2n}}} \right) \quad \text{with} \quad \omega_{-2n} = \frac{2k}{v_0(k^2 - v_0^2)^n} \quad (22)$$

where in both cases β is a free parameter.

3 KdV negative hierarchy

For KdV hierarchy we employ the same algebraic structure of section 3, i.e., principal gradation, Q_p (3) and the constant grade one element $E^{(1)}$ (6). Propose the following Lax operator,

$$A_x^{\text{KdV}}(J) = E^{(1)} + A^{(-1)} = E_\alpha^{(0)} + E_{-\alpha}^{(1)} + J E_{-\alpha}^{(0)} = \begin{pmatrix} 0 & 1 \\ \lambda + J & 0 \end{pmatrix} \quad (23)$$

where $A^{(-1)} = J E_{-\alpha}^{(0)} \in \mathcal{G}_{-1}$ and $J = J(x, \tau_N)$ is the field of KdV hierarchy. For sub-hierarchy that leads to negative time-flow τ_{-N} , the temporal-part Lax operator is given by

$$A_{\tau_{-N}}^{\text{KdV}}(J) = \mathcal{D}^{(-N-2)} + \mathcal{D}^{(-N-1)} + \dots + \mathcal{D}^{(-1)} \quad (24)$$

where $\mathcal{D}^{(i)} \in \mathcal{G}_i$. The zero curvature decomposes according to the graded structure as

$$[A^{(-1)}, \mathcal{D}^{(-N-2)}] = 0 \quad (25)$$

$$\partial_x \mathcal{D}^{(-N-2)} + [A^{(-1)}, \mathcal{D}^{(-N-1)}] = 0 \quad (26)$$

$$\partial_x \mathcal{D}^{(-N-1)} + [E^{(1)}, \mathcal{D}^{(-N-2)}] + [A^{(-1)}, \mathcal{D}^{(-N)}] = 0 \quad (27)$$

\vdots

$$\partial_x \mathcal{D}^{(-1)} + [E^{(1)}, \mathcal{D}^{(-2)}] - \partial_{\tau_{-N}} A^{(-1)} = 0 \quad (28)$$

$$[E^{(1)}, \mathcal{D}^{(-1)}] = 0 \quad (29)$$

which allows solving for all $\mathcal{D}^{(i)}$ and determines the equation of motion (28) according to τ_{-N} . Notice that the lowest grade equation (25) implies that $\mathcal{D}^{(-N-2)}$ is proportional to $E_{-\alpha}^{(-m)}$ and

therefore $N = 2m - 1$. For this reason *all equations of motion for KdV hierarchy are associated with odd temporal flows*, in contrast to the mKdV case, where there are equations of motion associated to both, even and odd flows.

The equations of motion for KdV hierarchy are more conveniently expressed in terms of non-local field $J(x, \tau_N) = \partial_x \eta(x, \tau_N)$. The first negative flow is obtained from zero curvature with $N = 1$, leads to the following temporal Lax,

$$\begin{aligned} A_{\tau_{-1}}^{\text{KdV}} &= \frac{\eta_{\tau_{-1}}}{2} \left(E_{\alpha}^{(-1)} + E_{-\alpha}^{(0)} \right) + \frac{\eta_{x, \tau_{-1}}}{4} h^{(-1)} + \frac{2\eta_x \eta_{\tau_{-1}} - \eta_{2x, \tau_{-1}}}{4} E_{-\alpha}^{(-1)} \\ &= \left(\begin{array}{cc} \frac{\eta_{x, \tau_{-1}}}{4\lambda} & \frac{\eta_{\tau_{-1}}}{2\lambda} \\ \frac{2\eta_x \eta_{\tau_{-1}} - \eta_{2x, \tau_{-1}}}{4\lambda} + \frac{\eta_{\tau_{-1}}}{2} & -\frac{\eta_{x, \tau_{-1}}}{4\lambda} \end{array} \right) \end{aligned} \quad (30)$$

and equation of motion

$$4\eta_x \eta_{x, \tau_{-1}} + 2\eta_{2x} \eta_{\tau_{-1}} - \eta_{3x, \tau_{-1}} = 0. \quad (31)$$

This equation was first proposed in [6] using inverse of recursion operator. Later in [7], its hamiltonian and soliton solutions were discussed.

If we now take $N = 3$ in (24) and find for the associated temporal Lax,

$$\begin{aligned} A_{\tau_{-3}}^{\text{KdV}} &= \frac{\eta_{\tau_{-3}}}{2} \left(E_{\alpha}^{(-1)} + E_{-\alpha}^{(0)} \right) + \frac{\eta_{x, \tau_{-3}}}{4} h^{(-1)} - \frac{\mathcal{B}}{8} \left(E_{\alpha}^{(-2)} + E_{-\alpha}^{(-1)} \right) \\ &+ \frac{2\eta_{\tau_{-3}} \eta_x - \eta_{2x, \tau_{-3}}}{8} E_{-\alpha}^{(-1)} - \frac{\mathcal{B}_x}{16} h^{(-2)} + \frac{\mathcal{B}_{2x} - \eta_x \mathcal{B}}{8} E_{-\alpha}^{(-2)} \\ &= \left(\begin{array}{cc} \frac{\eta_{x, \tau_{-3}}}{4\lambda} - \frac{\mathcal{B}_x}{16\lambda^2} & \frac{\eta_{\tau_{-3}}}{2\lambda} - \frac{\mathcal{B}}{8\lambda^2} \\ \frac{1}{2}\eta_{\tau_{-3}} + \frac{2\eta_{\tau_{-3}} \eta_x - \eta_{2x, \tau_{-3}} - \mathcal{B}}{8\lambda} + \frac{\mathcal{B}_{2x} - \eta_x \mathcal{B}}{8\lambda^2} & -\frac{\eta_{x, \tau_{-3}}}{4\lambda} + \frac{\mathcal{B}_x}{16\lambda^2} \end{array} \right) \end{aligned} \quad (32)$$

where

$$\mathcal{B} = d^{-1}(4\eta_x \eta_{x, \tau_{-3}} + 2\eta_{2x} \eta_{\tau_{-3}} - \eta_{3x, \tau_{-3}}). \quad (33)$$

The corresponding equation of motion is given by

$$\begin{aligned} &-\frac{1}{2}\eta_{5x, \tau_{-3}} + 4\eta_x (-2\eta_{x, \tau_{-3}} \eta_x + \eta_{3x, \tau_{-3}} - \eta_{2x} \eta_{\tau_{-3}}) + 5\eta_{2x} \eta_{2x, \tau_{-3}} \\ &+ 4\eta_{x, \tau_{-3}} \eta_{3x} + \eta_{4x} \eta_{\tau_{-3}} + \eta_{2x} d^{-1}(4\eta_x \eta_{x, \tau_{-3}} + 2\eta_{2x} \eta_{\tau_{-3}} - \eta_{3x, \tau_{-3}}). \end{aligned} \quad (34)$$

Notice that vacuum solution $\eta = \eta_0 = \text{constant}$, either zero or non-zero, satisfy both equations of motion (31) and (34). Such behavior differs from the mKdV hierarchy where the equations of motion associated with odd-time flows are satisfied with zero vacuum and the even-time flows with non-zero vacuum (constant). This coalescence in vacuum solution presented in KdV hierarchy can be explained more generally from zero curvature projected around vacuum, i.e,

$$\left[A_x^{\text{KdV}} \Big|_{\text{vac}}, A_{\tau_{-N}}^{\text{KdV}} \Big|_{\text{vac}} \right] = \left[E^{(1)} + \eta_0 E_{-\alpha}^{(0)}, \mathcal{D}_{\text{vac}}^{(-N-2)} + \mathcal{D}_{\text{vac}}^{(-N-1)} + \dots + \mathcal{D}_{\text{vac}}^{(-1)} \right] = 0 \quad (35)$$

from lowest grade equation

$$\left[\eta_0 E_{-\alpha}^{(0)}, \mathcal{D}_{\text{vac}}^{(-N-2)} \right] = \left[\eta_0 E_{-\alpha}^{(0)}, a_{-N-2} E_{-\alpha}^{(-1/2(N+1))} \right] = 0 \quad (36)$$

this equation is automatically satisfied no matter η_0 is zero or non-zero if $N = 2n - 1$. It therefore follows that *the negative KdV hierarchy are associated to odd flows*, $\tau_{-N} = \tau_{-2n-1}$ and admit both, zero and non-zero vacuum solutions.

4 Miura Transformation and Soliton Solutions

In order to map the mKdV and KdV hierarchies let us consider the *Miura-gauge transformation* generated by (see [4], [5])

$$S_1 = e^{\phi_x E_{-\alpha}^{(0)}} = \begin{pmatrix} 1 & 0 \\ \phi_x & 1 \end{pmatrix} \quad (37)$$

which maps the two Lax operators, A_x^{mKdV} into A_x^{KdV} of eqns. (8) and (23) respectively, i.e.,

$$A_x^{\text{KdV}} = S_1 A_x^{\text{mKdV}} S_1^{-1} + S_1 \partial_x S_1^{-1} = E_{-\alpha}^{(0)} + E_{-\alpha}^{(1)} + J E_{-\alpha}^{(0)} \quad (38)$$

where

$$J(x, t) = \partial_x \eta(x, t) = (\phi_x)^2 - \phi_{2x}. \quad (39)$$

We now analyse how S_1 acts as local gauge transformation upon A_t^{mKdV} . Let us consider first its action on an even grade element $D^{(-2n)} = c_{-n} h^{(-n)}$:

$$\begin{aligned} D^{(-2n)} &\rightarrow e^{\phi_x E_{-\alpha}^{(0)}} (c_{-n} h^{(-n)}) e^{-\phi_x E_{-\alpha}^{(0)}} + e^{\phi_x E_{-\alpha}^{(0)}} \partial_t (e^{-\phi_x E_{-\alpha}^{(0)}}) \\ &= \underbrace{c_{-n} h^{(-n)}}_{\mathcal{G}_{-2n}} + \underbrace{2c_{-n} \phi_x E_{-\alpha}^{(-n)}}_{\mathcal{G}_{-2n-1}} - \underbrace{\partial_t \phi_x E_{-\alpha}^{(0)}}_{\mathcal{G}_{-1}}. \end{aligned} \quad (40)$$

On the other hand, if we consider $D^{(-2n+1)} = a_{-n} E_{-\alpha}^{(-n)} + b_{-n} E_{-\alpha}^{(-n+1)}$ under the local gauge generated by (37) we find

$$\begin{aligned} D^{(-2n+1)} &\rightarrow e^{\phi_x E_{-\alpha}^{(0)}} (a_{-n} E_{-\alpha}^{(-n)} + b_{-n} E_{-\alpha}^{(-n+1)}) e^{-\phi_x E_{-\alpha}^{(0)}} + e^{\phi_x E_{-\alpha}^{(0)}} \partial_t (e^{-\phi_x E_{-\alpha}^{(0)}}) \\ &= \underbrace{-a_{-n} (\phi_x)^2 E_{-\alpha}^{(-n)}}_{\mathcal{G}_{-2n-1}} - \underbrace{a_{-n} \phi_x h_1^{(-n)}}_{\mathcal{G}_{-2n}} + \underbrace{a_{-n} E_{-\alpha}^{(-n)} + b_{-n} E_{-\alpha}^{(-n+1)}}_{\mathcal{G}_{-2n+1}} - \underbrace{\partial_t \phi_x E_{-\alpha}^{(0)}}_{\mathcal{G}_{-1}}. \end{aligned} \quad (41)$$

Thus, any even negative mKdV time flow of the form $A_{t_{-2n}}^{\text{mKdV}} = D^{(-2n)} + D^{(-2n+1)} + \dots + D^{(0)}$ is mapped into its KdV counterpart with the following graded structure,

$$\begin{aligned} A_{\tau_{-2n+1}}^{\text{KdV}} &= e^{\phi_x E_{-\alpha}^{(0)}} (D^{(-2n)} + D^{(-2n+1)} + \dots + D^{(0)}) e^{-\phi_x E_{-\alpha}^{(0)}} - \phi_{x, t_{-2n}} E_{-\alpha}^{(0)} \\ &= \mathcal{D}^{(-2n-1)} + \mathcal{D}^{(-2n)} + \dots + \mathcal{D}^{(-1)}. \end{aligned} \quad (42)$$

For odd negative mKdV time flow of the form $A_{t_{-2n+1}}^{\text{mKdV}} = D^{(-2n+1)} + D^{(-2n+2)} + \dots + D^{(0)}$ will be mapped into

$$\begin{aligned} A_{\tau_{-2n+1}}^{\text{KdV}} &= e^{\phi_x E_{-\alpha}^{(0)}} (D^{(-2n+1)} + D^{(-2n+2)} + \dots + D^{(0)}) e^{-\phi_x E_{-\alpha}^{(0)}} - \phi_{x, t_{-2n+1}} E_{-\alpha}^{(0)} \\ &= \mathcal{D}^{(-2n-1)} + \mathcal{D}^{(-2n)} + \mathcal{D}^{(-2n+1)} + \dots + \mathcal{D}^{(-1)}. \end{aligned} \quad (43)$$

Thus, both $A_{t_{-2n+1}}^{\text{mKdV}}$ and $A_{t_{-2n}}^{\text{mKdV}}$ are transformed, by the Miura-gauge transformation (37), into a single graded KdV structure $A_{\tau_{-2n+1}}^{\text{KdV}}$ (42)-(43) (associated to flow τ_{-2n+1}). We therefore conclude that both *negative even and negative odd mKdV flows collapse within the same KdV odd flow*, i.e.,

$$t_{-2n+1}^{\text{mKdV}}, t_{-2n}^{\text{mKdV}} \xRightarrow{S_1} \tau_{-2n+1}^{\text{KdV}}. \quad (44)$$

Notice that this explains why each KdV negative flow admits both zero and non-zero vacuum solutions. They inherit the zero and the non-zero vacuum information from mKdV negative odd and its subsequent negative even flows respectively. Let us illustrate explicitly for the first

two negative mKdV flows, namely, t_{-1} and t_{-2} .

For t_{-1}^{mKdV} the field $\phi = \phi(x, t_{-1})$ satisfy the sinh-Gordon eqn (15). We then have

$$\begin{aligned} A_{\tau_{-1}}^{\text{KdV}} &= S_1 A_{t_{-1}}^{\text{mKdV}} S_1^{-1} + S_1 \partial_{t_{-1}} S_1^{-1} \\ &= e^{\phi_x E_{-\alpha}^{(0)}} \left(e^{-2\phi} E_{\alpha}^{(-1)} + e^{2\phi} E_{-\alpha}^{(0)} \right) e^{-\phi_x E_{-\alpha}^{(0)}} - \phi_{x, t_{-1}} E_{-\alpha}^{(0)} \end{aligned} \quad (45)$$

leading to

$$A_{\tau_{-1}}^{\text{KdV}} = e^{-2\phi} \left(E_{\alpha}^{(-1)} + E_{-\alpha}^{(0)} \right) + \frac{\eta_{x, t_{-1}}}{4} h^{(-1)} - (\phi_x)^2 e^{-2\phi} E_{-\alpha}^{(-1)} \quad (46)$$

where we used the Sinh-Gordon equation of motion, $\phi_{x, t_{-1}} = e^{2\phi} - e^{-2\phi}$ and *Miura* transformation, $\eta_x = (\phi_x)^2 - \phi_{2x}$ to simplify some terms. Note that in terms of zero curvature, we had already constructed $A_{\tau_{-1}}^{\text{KdV}}$ given in (30),

$$A_{\tau_{-1}}^{\text{KdV}} = \frac{\eta_{\tau_{-1}}}{2} \left(E_{\alpha}^{(-1)} + E_{-\alpha}^{(0)} \right) + \frac{\eta_{x, \tau_{-1}}}{4} h^{(-1)} + \frac{2\eta_x \eta_{\tau_{-1}} - \eta_{2x, \tau_{-1}}}{4} E_{-\alpha}^{(-1)} \quad (47)$$

From the condition for eqns (46) and (47) to agree we find

$$\eta_{\tau_{-1}} = 2 \cdot e^{-2\phi(x, t_{-1})}. \quad (48)$$

On the other hand, if we now use t_{-2}^{mKdV} with $\phi = \phi(x, t_{-2})$ satisfying (17), we get

$$\begin{aligned} A_{\tau_{-1}}^{\text{KdV}} &= S_1 A_{t_{-2}}^{\text{mKdV}} S_1^{-1} + S_1 \partial_{t_{-2}} S_1^{-1} \\ &= e^{\phi_x E_{-\alpha}^{(0)}} \left(h^{(-1)} + 2e^{-2\phi} d^{-1} (e^{2\phi}) E_{\alpha}^{(-1)} - 2e^{2\phi} d^{-1} (e^{-2\phi}) E_{-\alpha}^{(0)} \right) e^{-\phi_x E_{-\alpha}^{(0)}} - \phi_{x, t_{-2}} E_{-\alpha}^{(0)} \end{aligned} \quad (49)$$

leading to

$$A_{\tau_{-1}}^{\text{KdV}} = 2e^{-2\phi} d^{-1} (e^{2\phi}) \left(E_{\alpha}^{(-1)} + E_{-\alpha}^{(0)} \right) + \frac{\eta_{x, t_{-2}}}{4} h^{(-1)} + 8(\phi_x - \phi_x^2 e^{-2\phi} d_x^{-1} e^{2\phi}) E_{-\alpha}^{(-1)} \quad (50)$$

where we used the equation of motion for t_{-2}^{mKdV} (17) and *Miura transformation*. Thus, (50) only agrees with (47) if we set

$$\eta_{\tau_{-1}} = 2 \cdot 2e^{-2\phi(x, t_{-2})} d^{-1} (e^{2\phi(x, t_{-2})}) \quad (51)$$

Notice that the same $A_{\tau_{-1}}^{\text{KdV}}$ is written in two different ways, one in terms of the sinh-Gordon field $\phi(x, t_{-1})$ given by (46)-(48) and another, in terms of solution of eqn. (17) namely $\phi(x, t_{-2})$ in (50)-(51). This can be checked explicitly with solutions given in (21) and (22) for $n = 1$.

5 Conclusion

We have therefore concluded from the above simple example that solutions of the KdV equation associated to the time flow τ_{-1} inherits different vacuum structure from a pair of mKdV solutions (via *Miura transformation*). The first associated to mKdV flow t_{-1} , eqn. (15) (with zero vacuum) satisfying (48) and the second associated to mKdV flow t_{-2} , eqn. (17) (with non-zero vacuum) satisfying (51). The argument can be easily generalized for higher flows, each KdV flow admits both, zero and non-zero vacuum solutions. They are constructed from pair of subsequent of mKdV flows each of them admitting different vacuum structures. We expect to report in a future publication the generalization of our construction to the A_r - KdV hierarchy employing the gauge-Miura transformation proposed in [5]. We also expect to discuss the systematic construction of soliton (multisoliton) solutions and its vacuum structure in terms of vertex operators and its deformations in the lines of refs. [3], [4].

Acknowledgements

JFG and AHZ thank CNPq and Fapesp for support. YFA thanks São Paulo Research Foundation (FAPESP) for financial support under grant #2021/00623 – 4 and GVL is supported by Capes.

References

- [1] D. Olive and N. Turok, *Local Conserved Densities and Zero Curvature Conditions for Toda Lattice Field Theories*, Nucl. Phys. **B215** (1983), 470, DOI: 10.1016/0550-3213(85)90347-5
- [2] O. Babelon, D. Bernard and Talon, *Introduction to Classical Integrable Systems*, Cambridge Univ. Press (2009).
- [3] J.F. Gomes, G. Starvaggi França, G.R. de Melo and A. H. Zimerman, *Negative Even Grade mKdV Hierarchy and its Soliton Solutions*, J. Phys: Math. Theor. **42**(2009), 445204 arXiv:0906.5579, DOI: 10.1088/1751-8113/42/44/445204
- [4] J.F. Gomes, A.L. Retore and A.H. Zimerman, *Miura and generalized Bäcklund transformation for KdV hierarchy*, J. Phys. A: Math. Theor. **49** (2016) 504003, arXiv:1610.02303, DOI: 10.1088/1751-8113/49/50/504003
- [5] J. M. Carvalho Ferreira, J. F. Gomes, G. V. Lobo, A. H. Zimerman, *Gauge Miura and Bäcklund transformations for generalized A_n -KdV hierarchies*, J. Phys: Math. Theor. **54**(2021), 435201 arXiv:2106.00741, DOI: 10.1088/1751-8121/ac2718
- [6] J.M. Verosky, , *Negative powers of Olver recursion operators*, J. Math. Phys. **32**,1733 (1991), doi.org/10.1063/1.529234
- [7] Z. Qiao and Z. Fan, *Negative-order Korteweg–de Vries equations*, Physical Review E **86**, 016601 (2012), DOI: 10.1103/PhysRevE.86.016601