

Air showers and hadronic interactions with CORSIKA 8

Maximilian Reininghaus¹ for the CORSIKA 8 Collaboration^a

1 Karlsruher Institut für Technologie, Institut für Astroteilchenphysik,
Postfach 3640, 76021 Karlsruhe, Germany
reininghaus@kit.edu

December 23, 2022



51st International Symposium on Multiparticle Dynamics (ISMD2022)

Pitlochry, Scottish Highlands, 1-5 August 2022

[doi:10.21468/SciPostPhysProc.7](https://doi.org/10.21468/SciPostPhysProc.7)

Abstract

The CORSIKA 8 project is a collaborative effort aiming to develop a versatile C++ framework for the simulation of extensive air showers, intended to eventually succeed the long-standing FORTRAN version. I present an overview of its current capabilities, focusing on aspects concerning the hadronic and muonic shower components. In particular, I demonstrate the “cascade lineage” feature and its application to quantify the importance of certain phase-space regions in hadronic interactions for muon production. Additionally, I show first results using Pythia 8.3, which as of late is usable as interaction model in cosmic-ray applications and is currently being integrated into CORSIKA 8.

1 Introduction

A large part of the astroparticle physics community deals with the measurement of extensive air showers (EAS), particle cascades developing on macroscopically large scales of up to several 10 km. Making use of measurements of EAS observables in order to infer properties of the primary particle (high-energy cosmic rays, gamma rays, neutrinos) requires accurate predictions of these. Only Monte Carlo simulations are able to provide these at the necessary level of detail, relying on numerical methods and reliable models of the physical processes involved. The phase space covered in EAS is vast: Electromagnetic (EM), weak, and strong interactions play a role; dozens of particle species are involved; energy scales range from \sim keV up to ZeV. Software tools that can keep up with these requirements are a must-have and serve as a cornerstone of the field. For more than 30 years the FORTRAN code CORSIKA (Cosmic Ray Simulations for KASCADE) [1, 2] played that role and has become a de facto standard [3]. In recent years, however, it has become unfeasible to accommodate for the increasing needs of upcoming experiments by continuing to extend the existing “dinosource”.

Instead, efforts have been taken to develop a new C++ code from scratch, eventually termed CORSIKA 8 [4], with the goal to provide a modern, modular and flexible framework for simulations of particle showers. Since its inception in 2018 [5], CORSIKA 8 is developed as open-source project by an international collaboration. The code is publicly available at the [gitlab repository](#) and usable by early adopters. At the time of writing, a large fraction of features available in the legacy version are implemented in CORSIKA 8. Moreover, CORSIKA 8 has a number of unique features that are not available in other codes.

^afull author list available at <https://tinyurl.com/corsika8-202210>




Figure 1: Left: Distributions of the number of muon ancestor generations for primary energies of $E_0 = 10^{17}, 10^{18}, 10^{19}$ eV. Vertical dashed lines indicate the mean value of the distribution of the same colour. Taken from ref. [10]. Right: Mean number of muon ancestor generations as function of primary energy. Taken from ref. [11].

In this article I focus on the capabilities in the hadronic and muonic sectors of EAS, which to date remain not fully understood, as a number of discrepancies between experimental data and simulations suggest [6]. More general overviews of the current state of the project can be found in refs. [7, 8]. The structure of this article is as follows: In section 2 I present a study of the phase space of hadronic interactions relevant for muon production. Additionally, I show first results using Pythia 8 as hadronic interaction model in the context of air shower simulations in section 3, followed by concluding remarks.

2 Air shower genealogy

The muon component of EAS is a tracer of the hadronic interactions happening during the shower development. The bulk of muons observed at ground, having energies mostly in the 100 MeV to GeV range, stem from the decay of low-energy pions and kaons that form the last generation of the hadronic cascade. Measurements of the muon content of EAS induced by ultra-high energy cosmic rays (i.e., having energies ≥ 1 EeV) performed in several experiments show that there is a significant excess of muons in data compared to simulations using up-to-date versions of hadronic interaction models [9]. It is widely believed that this discrepancy, coined *muon puzzle* [6], is the result of a lack of understanding and mismodelling of hadronic interactions.

CORSIKA 8 is particularly well suited to shed more light onto the muon puzzle: It enables keeping the complete *lineage* of particles in memory so that particles can be related to any of their predecessor generations up to the primary particle. Details on the technical implementation are given in refs. [11, 12]. The first study to exploit this information is presented in ref. [10], whose results I summarize here. Figure 1 shows results regarding the number of generations N_{gen} , i.e. the number of hadronic interactions happening between the primary particle and the final muon that reaches ground. It is an important quantity because the number of muons grows exponentially with N_{gen} and small errors in the modelling of these interactions get amplified N_{gen} times, leading to a potentially large impact on the muon number [13]. The left plot shows the N_{gen} distributions of proton-induced EAS with energies of




Figure 2: Pseudorapidity distributions of $\pi^\pm + \text{air} \rightarrow \text{charged hadrons}$. The dashed line indicates generator-level distributions while the coloured and dotted lines show muon-weighted distributions of vertical, proton-induced showers of 10^{19} eV (in arbitrary units). Taken from ref. [10].

$E_0 = 10^{17}, 10^{18}, 10^{19}$ eV, simulated using the interaction model SIBYLL 2.3d [14] at high energies (> 63.1 GeV) together with the Hillas Splitting algorithm (HSA) [15] for low energies. With increasing primary energy, the distributions shift towards higher values of N_{gen} . The right plot shows the dependence of the mean $\langle N_{\text{gen}} \rangle$ on the primary energy. The logarithmic behaviour follows what is expected from the Heitler–Matthews toy model [16]. A more detailed analysis is given in ref. [11].

A second study deals with the importance of different phase-space regions in hadronic interactions with respect to muon production. Figure 2 shows pseudorapidity (η) distributions of charged hadrons in π^\pm -air collisions in four energy bins. Besides the pure generator-level spectra (dashed lines), which are in principle measurable in accelerator experiments, the *muon weighted* distributions are superimposed (coloured lines). This muon weight is given by the number of muons stemming directly or indirectly from the secondaries emitted at a given η , potentially after applying some selection criterion. The plot shows that at sufficiently high interaction energies ($\sqrt{s} \gtrsim 100$ GeV) essentially only particle production in the forward region plays a role, irrespective of the lateral distance of the muons. In this regime the muon weight can also be estimated well from the Heitler–Matthews model [6], indicated by the dotted line. At low interaction energies ($\sqrt{s} \lesssim 50$ GeV), however, the central region gains relevance




Figure 3: Mean shower maximum $\langle X_{\max} \rangle$ vs. number of muons N_{μ} for showers at $10^{17.5}$ eV and 60° inclination

especially for muons at distances larger than a few 100 m.

3 Towards Pythia 8 as interaction model in EAS simulations

A currently ongoing development is the inclusion of Pythia 8.3 [17] into CORSIKA 8 as hadronic interaction model. Until recently, this was technically unfeasible, but since the latest release (Pythia 8.307) a number of new features allow for an easy use in EAS simulations [18]: a) the ability to generate single events at arbitrary energies without a time-consuming re-initialization of the model, b) a much wider range of possible projectile species, c) an extended range of interaction energies down to 200 MeV (lab-frame), d) a simplified model of nuclear matter, allowing hadron-air collisions.

For a first comparison we have simulated showers with an energy of $10^{17.5}$ eV and an inclination of 60° . Hadrons and muons are fully propagated with CORSIKA 8, while EM particles are redirected into the CONEX code [19], which simulates the EM component of the shower using a numerical solution of the cascade equations describing the longitudinal development. We use QGSJet-II.04 [20], SIBYLL 2.3d and Pythia 8.307 as high-energy interaction models. In each case Pythia is used as low-energy interaction model. Figure 3 shows the results regarding the mean shower maximum $\langle X_{\max} \rangle$ and N_{μ} . While N_{μ} obtained with Pythia is in the same ballpark as the other models, a deviation of $\langle X_{\max} \rangle$ as large as the difference between proton and helium is apparent. Further studies to explain these differences, stemming from differences in hadron-air cross-sections, are ongoing and will be presented elsewhere [21]. Note that in our setup Pythia 8.307 cannot be used for nucleus-nucleus collisions at the moment so that we consider only proton-induced showers in that case.

4 Conclusions and outlook

CORSIKA 8 is a modern framework for the simulation of particle showers in air and other media. Over the past few years, a lot of progress has been made to make it a reliable, versa-

tile and future-proof tool. Although not yet feature-complete, in some aspects it can be used for studies that have previously been impossible. The availability of the complete particle lineage allows detailed studies of hadronic interactions and their relevance for muon production, which helps to shed light on the muon puzzle. It emphasizes the importance of dedicated accelerator measurements at both high and low energies that cover the relevant phase-space.

The ongoing integration of Pythia 8 as new hadronic interaction model in CORSIKA 8 offers a number of new perspectives for EAS simulations. On the one hand, the reduced energy threshold of only 200 MeV renders it suitable for both low and high energies. On the other hand, Pythia 8 can be tuned by the users, allowing to study the impact of internal model parameters on EAS observables, or possibly even to conduct combined fits to accelerator measurements and EAS data at the same time.

A first public release is anticipated for 2023 mid-year. Until then, the development focusses mainly on reaching a level of maturity sufficient to cover at least most mainstream applications with an accuracy on par with the legacy version CORSIKA 7. Issues currently being addressed include the integration of FLUKA [22, 23] as low-energy interaction model, proper handling of the LPM effect (see e.g. ref. [24]), photo-hadronic interactions with SOPHIA [25], and the introduction of thinning to speed up simulations at the highest energies (see e.g. ref. [26]). Furthermore, validation and further improvements of the already existing modules for simulating radio signals and Cherenkov light are work in progress. Besides these physics-driven aspects, efforts are made to provide a default application with an easy-to-use user interface for non-expert end users suitable for standard scenarios (in contrast to offering just the framework in which users need to create simulation setups on their own).

Acknowledgements

I thank Torbjörn Sjöstrand and Marius Utheim for providing example code that greatly simplified the integration of Pythia 8 in CORSIKA 8.

Simulations were performed on the bwForCluster BinAC of the University of Tübingen, supported by the state of Baden-Württemberg through bwHPC and the DFG through grant no. INST 37/935-1 FUGG.

References

- [1] J. Capdevielle, P. Gabriel, H. Gils, P. Grieder, D. Heck, N. Heide, J. Knapp, H. Mayer, J. Oehlschläger, H. Rebel, G. Schatz and T. Thouw, *Air shower simulations for KASCADE*, In *Proc. 21th Int. Cosmic Ray Conf.*, HE 7.3-3 (1990).
- [2] D. Heck, J. Knapp, J. N. Capdevielle, G. Schatz and T. Thouw, *CORSIKA: A Monte Carlo code to simulate extensive air showers*, Tech. Rep. FZKA-6019, Forschungszentrum Karlsruhe, doi:[10.5445/IR/270043064](https://doi.org/10.5445/IR/270043064) (1998).
- [3] K.-H. Kampert and A. A. Watson, *Extensive Air Showers and Ultra High-Energy Cosmic Rays: A Historical Review*, *Eur. Phys. J. H* **37**, 359 (2012), doi:[10.1140/epjh/e2012-30013-x](https://doi.org/10.1140/epjh/e2012-30013-x), [1207.4827](https://arxiv.org/abs/1207.4827).
- [4] M. Reininghaus and R. Ulrich, *CORSIKA 8 – Towards a modern framework for the simulation of extensive air showers*, *EPJ Web Conf.* **210**, 02011 (2019), doi:[10.1051/epjconf/201921002011](https://doi.org/10.1051/epjconf/201921002011), [1902.02822](https://arxiv.org/abs/1902.02822).

- [5] R. Engel, D. Heck, T. Huege, T. Pierog, M. Reininghaus, F. Riehn, R. Ulrich, M. Unger and D. Veberič, *Towards a Next Generation of CORSIKA: A Framework for the Simulation of Particle Cascades in Astroparticle Physics*, *Comput. Softw. Big Sci.* **3**(1), 2 (2019), doi:[10.1007/s41781-018-0013-0](https://doi.org/10.1007/s41781-018-0013-0), [1808.08226](https://arxiv.org/abs/1808.08226).
- [6] J. Albrecht, L. Cazon, H. Dembinski, A. Fedynitch, K.-H. Kampert, T. Pierog, W. Rhode, D. Soldin, B. Spaan, R. Ulrich and M. Unger, *The Muon Puzzle in cosmic-ray induced air showers and its connection to the Large Hadron Collider*, *Astrophys. Space Sci.* **367**, 27 (2022), doi:[10.1007/s10509-022-04054-5](https://doi.org/10.1007/s10509-022-04054-5), [2105.06148](https://arxiv.org/abs/2105.06148).
- [7] T. Huege, *CORSIKA 8 – the next-generation air shower simulation framework*, In *Proc. 21st ISVHECRI* (2022), [2208.14240](https://arxiv.org/abs/2208.14240).
- [8] J.-M. Alameddine *et al.*, *CORSIKA 8 – Contributions to the 37th International Cosmic Ray Conference in Berlin Germany (ICRC 2021)* (2021), [2112.11761](https://arxiv.org/abs/2112.11761).
- [9] D. Soldin, *Update on the Combined Analysis of Muon Measurements from Nine Air Shower Experiments*, *PoS ICRC2021*, 349 (2021), doi:[10.22323/1.395.0349](https://doi.org/10.22323/1.395.0349), [2108.08341](https://arxiv.org/abs/2108.08341).
- [10] M. Reininghaus, R. Ulrich and T. Pierog, *Air shower genealogy for muon production*, *PoS ICRC2021*, 463 (2021), doi:[10.22323/1.395.0463](https://doi.org/10.22323/1.395.0463), [2108.03266](https://arxiv.org/abs/2108.03266).
- [11] M. Reininghaus, *The air shower simulation framework CORSIKA 8: Development and first applications to muon production*, Ph.D. thesis, Karlsruhe Institut für Technologie (KIT) & Universidad Nacional de San Martín, doi:[10.5445/IR/1000152097](https://doi.org/10.5445/IR/1000152097) (2021).
- [12] A. A. Alves Jr., M. Reininghaus, A. Schmidt, R. Prechelt and R. Ulrich, *CORSIKA 8 - A novel high-performance computing tool for particle cascade Monte Carlo simulations*, *EPJ Web Conf.* **251**, 03038 (2021), doi:[10.1051/epjconf/202125103038](https://doi.org/10.1051/epjconf/202125103038).
- [13] L. Cazon, R. Conceição, M. A. Martins and F. Riehn, *Constraining the energy spectrum of neutral pions in ultra-high-energy proton-air interactions*, *Phys. Rev. D* **103**(2), 022001 (2021), doi:[10.1103/PhysRevD.103.022001](https://doi.org/10.1103/PhysRevD.103.022001), [2006.11303](https://arxiv.org/abs/2006.11303).
- [14] F. Riehn, R. Engel, A. Fedynitch, T. K. Gaisser and T. Stanev, *Hadronic interaction model Sibyll 2.3d and extensive air showers*, *Phys. Rev. D* **102**(6), 063002 (2020), doi:[10.1103/PhysRevD.102.063002](https://doi.org/10.1103/PhysRevD.102.063002), [1912.03300](https://arxiv.org/abs/1912.03300).
- [15] A. M. Hillas, *Two interesting techniques for Monte-Carlo simulation of very high energy hadron cascades*, In *Proc. 17th Int. Cosmic Ray Conf.*, p. 193 (1981).
- [16] J. Matthews, *A Heitler model of extensive air showers*, *Astropart. Phys.* **22**, 387 (2005), doi:[10.1016/j.astropartphys.2004.09.003](https://doi.org/10.1016/j.astropartphys.2004.09.003).
- [17] C. Bierlich *et al.*, *A comprehensive guide to the physics and usage of PYTHIA 8.3*, *SciPost Phys. Codebases* **8** (2022), doi:[10.21468/SciPostPhysCodeb.8](https://doi.org/10.21468/SciPostPhysCodeb.8), [2203.11601](https://arxiv.org/abs/2203.11601).
- [18] T. Sjöstrand and M. Utheim, *Hadron Interactions for Arbitrary Energies and Species, with Applications to Cosmic rays*, *Eur. Phys. J. C* **82**(1), 21 (2022), doi:[10.1140/epjc/s10052-021-09953-5](https://doi.org/10.1140/epjc/s10052-021-09953-5), [2108.03481](https://arxiv.org/abs/2108.03481).
- [19] T. Bergmann, R. Engel, D. Heck, N. N. Kalmykov, S. Ostapchenko, T. Pierog, T. Thouw and K. Werner, *One-dimensional Hybrid Approach to Extensive Air Shower Simulation*, *Astropart. Phys.* **26**, 420 (2007), doi:[10.1016/j.astropartphys.2006.08.005](https://doi.org/10.1016/j.astropartphys.2006.08.005), [astro-ph/0606564](https://arxiv.org/abs/astro-ph/0606564).

- [20] S. Ostapchenko, *Monte Carlo treatment of hadronic interactions in enhanced Pomeron scheme: I. QGSJET-II model*, Phys. Rev. D **83**, 014018 (2011), doi:[10.1103/PhysRevD.83.014018](https://doi.org/10.1103/PhysRevD.83.014018), [1010.1869](https://arxiv.org/abs/1010.1869).
- [21] M. Reininghaus, T. Sjöstrand and M. Uthmeim, In *Proc. 6th Int. Sympos. Ultra High Energy Cosmic Rays (UHECR2022)*, In preparation (2023).
- [22] A. Ferrari, P. R. Sala, A. Fasso and J. Ranft, *FLUKA: A multi-particle transport code (Program version 2005)*, Tech. Rep. CERN-2005-010, SLAC-R-773, INFN-TC-05-11, CERN-2005-10, doi:[10.2172/877507](https://doi.org/10.2172/877507) (2005).
- [23] G. Battistoni *et al.*, *Overview of the FLUKA code*, Annals Nucl. Energy **82**, 10 (2015), doi:[10.1016/j.anucene.2014.11.007](https://doi.org/10.1016/j.anucene.2014.11.007).
- [24] A. N. Cillis, H. Fanchiotti, C. A. García Canal and S. J. Sciutto, *Influence of the LPM effect and dielectric suppression on particle air showers*, Phys. Rev. D **59**, 113012 (1999), doi:[10.1103/PhysRevD.59.113012](https://doi.org/10.1103/PhysRevD.59.113012), [astro-ph/9809334](https://arxiv.org/abs/astro-ph/9809334).
- [25] A. Mücke, R. Engel, J. P. Rachen, R. J. Protheroe and T. Stanev, *SOPHIA: Monte Carlo simulations of photohadronic processes in astrophysics*, Comput. Phys. Commun. **124**, 290 (2000), doi:[10.1016/S0010-4655\(99\)00446-4](https://doi.org/10.1016/S0010-4655(99)00446-4), [astro-ph/9903478](https://arxiv.org/abs/astro-ph/9903478).
- [26] M. Kobal, *A thinning method using weight limitation for air-shower simulations*, Astropart. Phys. **15**, 259 (2001), doi:[10.1016/S0927-6505\(00\)00158-4](https://doi.org/10.1016/S0927-6505(00)00158-4).