

Entanglement dynamics with string measurement operators

Giulia Piccitto^{1*}, Angelo Russomanno², and Davide Rossini¹

¹ Dipartimento di Fisica dell'Università di Pisa and INFN,
Largo Pontecorvo 3, I-56127 Pisa, Italy

² Scuola Superiore Meridionale, Università di Napoli Federico II,
Largo San Marcellino 10, I-80138 Napoli, Italy

* giulia.piccitto@df.unipi.it

March 21, 2023

Abstract

We explain how to apply a Gaussian-preserving operator to a fermionic Gaussian state. We use this method to study the evolution of the entanglement entropy of an Ising spin chain following a Lindblad dynamics with string measurement operators, focusing on the quantum-jump unraveling of such Lindbladian. We find that the asymptotic entanglement entropy obeys an area law for finite-range string operators and a volume law for ranges of the string which scale with the system size. The same behavior is observed for the measurement-only dynamics, suggesting that measurements can play a leading role in this context.

Contents

1	Introduction	2
2	Fermionic Gaussian states	3
3	Evolving Gaussian states	5
3.1	Bogoliubov matrix evolution	6
3.2	QR decomposition	7
4	Number-preserving operators	9
4.1	Local operator	10
4.2	Two-site operator	10
5	Quantum jumps with string operators	10
5.1	Entanglement entropy	11
5.2	Measurement-only dynamics	12
5.3	Effect of the Hamiltonian	13
6	Conclusions	14
A	Rotation to the diagonal basis	15
B	Quantum-jump protocol	17
C	Evolution of the state: a different derivation	18

1 Introduction

Understanding the role of entanglement in quantum many-body systems is a subject under intensive study [1, 2]. While the ground-state entanglement properties have been widely investigated and are relatively well understood, there are still several open questions regarding the stabilization of quantum correlations during the non-equilibrium dynamics. By exploiting a quasi-particle description it has been shown that in integrable systems, after a sudden quench, the entanglement entropy linearly increases in time, and eventually settles to a stationary value which linearly scales with the system size (volume-law scaling) [3, 4]. A similar volume-law scaling occurs in ergodic short-range systems [5], while in the presence of a strong disorder the situation is more involved, due to the appearance of many-body localization (see Ref. [6] for a review).

A natural extension of the above scenario is for a quantum system coupled to an external environment, a setup which is crucial to applications in quantum technologies and quantum computing, since any experimental platform is subject to noise-induced decoherence [7, 8]. In the hypothesis of weak and Markovian coupling with the bath, a good modeling for this setup is to assume the evolution of the reduced density matrix of the system to be ruled by a master equation in a Lindblad form [9]. The system-bath framework can be also interpreted as a process in which the environment behaves as a classical stochastic process performing random measurements on the quantum system [10, 11]. The outcome of any realization of this process is a pure-state stochastic quantum trajectory and the density matrix obtained by averaging over such trajectories obeys a Lindblad-type dynamic evolution [12, 13].

Here we are interested in the so-called measurement-induced phase transitions (MIPTs), i.e., in the possibility to develop discontinuities as in the asymptotic entanglement properties that emerge from the interplay between the unitary dynamics (typically generating entanglement) and the measurement processes (typically destroying it). This kind of transitions have been first proposed and studied in unitary random circuits undergoing random measurements (see, e.g., Refs. [14–22] and citing references), which may display a transition between a phase in which the entanglement grows with the system size and another phase where it undergoes a saturation to a finite value (area-law scaling). Similarly, MIPTs may also arise in free-fermion random circuits with temporal randomness [23], Majorana random circuits [24, 25], Dirac fermions [26], and other Hamiltonian models undergoing random measurements at discrete times [27–30]. When measuring string operators, MIPTs can occur in the presence of a measurement-only dynamics, without the interplay with the Hamiltonian [31, 32].

Recently, an increasing number of works have been focusing on MIPTs in fermionic systems described by quadratic Hamiltonians on a lattice, undergoing quantum trajectories under the action of random measurements of onsite quadratic operators [33–45]. One of the reasons of this interest relies in the Gaussian structure of the states of such systems that, being entirely determined by two-point correlation functions, are suitable for a semi-analytical treatment up to large lattice sizes (of the order of hundreds of sites) [46–50]. In particular, it has been found that the asymptotic volume-law scaling of the entanglement following a unitary evolution [51, 52] is unstable under local measurements, independently

of the measurement strength [33, 35], and a MIPT may emerge. For example, the Ising chain exhibits a size-dependent crossover towards a subextensive or an area-law entanglement regime, depending on the system parameters, as well as on the measurement rate and the type of unraveling [36–40]. In a version of this model with temporal noise also in the Hamiltonian, this crossover has been analytically proved to be a true transition by means of the replica method [53]. In the same context of monitored fermionic models, a couple of recent works pointed out the importance of having long-range interactions to stabilize larger amounts of entanglement, if the system undergoes random onsite measurements [42, 43] (see [54, 55] for a similar phenomenon in quantum circuits).

Inspired by these latter findings, here we follow a complementary strategy and address the role of non-local string measurement operators in the dynamics of entanglement for a quadratic Hamiltonian system. The purpose of this paper is twofold. On the one hand, we collect and review all the technical details to implement the dynamics of Gaussian states in the presence of Gaussian-preserving measuring operators. On the other hand, we discuss the effect of non-local measurement operators on the dynamics of the entanglement entropy, while keeping the Hamiltonian with short-range interactions. To this aim, we implement a quantum-jump unraveling of the Lindblad dynamics [9, 10] for the Ising chain (mapped to the fermionic Kitaev chain) using strings of Pauli matrices as Lindblad operators (mapped to non-local two-point fermionic operators). The main result is that, when the string range r is comparable with the system size L , contrary to what happens in the presence of local jump operators, the measurement process can stabilize an asymptotic volume-law for the entanglement entropy. As a consequence, a MIPT is conjectured as a function of r . Quite interestingly, the same qualitative conclusion applies if we switch off the Hamiltonian and consider a measurement-only Lindblad dynamics induced by string operators, similarly to what has been studied in Ref. [31] for measurement-only models with nonlocal measurements of Pauli strings.

The paper is organized as follows. In Section 2 we briefly review the theory of quadratic fermionic Hamiltonians on a lattice. In Section 3 and Appendix A we present the technical details required to implement the dynamics of a Gaussian state under the effect of Gaussian-preserving exponentials of quadratic operators. We first set the general framework, and then shed light on the dynamical equations and on the normalization of the state. In Section 4 we apply such a machinery to string operators, a particular class of operators which generalize the one discussed in Ref. [38, 39]. Subsequently, in Section 5 we present and discuss numerical results for the Ising chain with non-local string measurement operators. Finally, our conclusions are drawn in Section 6. The other appendices contain some details on the quantum-jump measurement protocol (Appendix B) and an alternative approach to determine the evolved state with the string operator (Appendix C).

2 Fermionic Gaussian states

We consider a generic free-fermion system on a lattice with L sites, described by the quadratic Hamiltonian

$$\hat{H} = \sum_{i,j=1}^L \left(Q_{i,j} \hat{c}_i^\dagger \hat{c}_j + P_{i,j} \hat{c}_i^\dagger \hat{c}_j^\dagger + \text{h.c.} \right), \quad (1)$$

where $\hat{c}_j^{(\dagger)}$ denotes the anticommuting annihilation (creation) operator on the j -th site. To ensure the Hermiticity of \hat{H} , the coefficient matrices obey $Q^\dagger = Q$ and $P^T = -P$.

In this context, a Gaussian state has the generic form ¹

$$|\psi\rangle = \mathcal{N} \exp \left\{ \frac{1}{2} \sum_{p,q} Z_{p,q} \hat{c}_p^\dagger \hat{c}_q^\dagger \right\} |0\rangle_{\hat{c}}, \quad (2)$$

with

$$Z = -(u^\dagger)^{-1} v^\dagger, \quad (3)$$

and \mathcal{N} a normalization constant. Here u and v are $L \times L$ matrices that obey the following symplectic unitarity conditions

$$u^\dagger u + v^\dagger v = I, \quad u v^\dagger + v^* u^T = 0, \quad (4)$$

where I denotes the $L \times L$ identity matrix. These conditions are equivalent to say that the matrix

$$\mathbb{U} = \begin{pmatrix} u & v^* \\ v & u^* \end{pmatrix} \quad (5)$$

is unitary and preserves the fermionic anticommutation relations: In fact it defines a new set of fermionic anticommuting operators

$$\hat{\gamma}_\kappa = \sum_j \left(u_{j,\kappa}^* \hat{c}_j + v_{j,\kappa}^* \hat{c}_j^\dagger \right), \quad (6)$$

that enjoy the property

$$\hat{\gamma}_\kappa |\psi\rangle = 0, \quad \forall \kappa \in \{1, \dots, L\}, \quad (7)$$

meaning that $|\psi\rangle$ is the vacuum of the $\hat{\gamma}$ fermions, $|\psi\rangle \equiv |0\rangle_{\hat{\gamma}}$ (the matrix \mathbb{U} is the so-called Bogoliubov transformation. For more details see, e.g., Refs. [50, 56]).

We remark that, because of the fermionic nature of the \hat{c} -operators, the matrix Z in Eq (3) is antisymmetric, as can be proved by exploiting Eqs. (4) [56]. In this context, the Nambu spinor notation is useful. Defining $\hat{\Psi} \equiv (\hat{c}_1, \dots, \hat{c}_L, \hat{c}_1^\dagger, \dots, \hat{c}_L^\dagger)^T$, and $\hat{\Phi} \equiv (\hat{\gamma}_1, \dots, \hat{\gamma}_L, \hat{\gamma}_1^\dagger, \dots, \hat{\gamma}_L^\dagger)^T$, one can rewrite Eq. (6) as

$$\hat{\Phi} = \mathbb{U}^\dagger \hat{\Psi}. \quad (8)$$

The state in Eq. (2) is Gaussian, enjoys the Wick's theorem, and thus it is completely determined by two-point correlation functions. Using Eq. (7) one can show that these correlation functions have a simple expression in terms of the \mathbb{U} matrix:

$$G_{i,j} \equiv \langle \hat{c}_i^\dagger \hat{c}_j \rangle = [v v^\dagger]_{i,j}, \quad F_{i,j} \equiv \langle \hat{c}_i \hat{c}_j \rangle = [v u^\dagger]_{i,j}. \quad (9)$$

An important case where the state has the Gaussian form in Eq. (2) is the ground state of the Hamiltonian (1). The Hamiltonian reads

$$\hat{H} = \frac{1}{2} \hat{\Psi}^\dagger \mathbb{H} \hat{\Psi} + \text{const.}, \quad \text{with} \quad \mathbb{H} = \begin{pmatrix} Q & P \\ -P^* & -Q^* \end{pmatrix}. \quad (10)$$

The $2L \times 2L$ Hermitian matrix \mathbb{H} is called the Bogoliubov-de Gennes Hamiltonian matrix, which can be diagonalized through the implementation of a Bogoliubov transformation

¹Hereafter we omit extrema in all the summations running from 1 to L .

\mathbb{U} of the form (5), such that $\mathbb{U}^{-1} \mathbb{H} \mathbb{U} = \text{diag}[\omega_\kappa, -\omega_\kappa]_{\kappa=1, \dots, L}$. The above transformation defines a new set of fermionic quasi-particles $\hat{\gamma}_\kappa$ [Eq. (6)], according to which the Hamiltonian of Eq. (1) can be written as

$$\hat{H} = \sum_{\kappa} \omega_\kappa \hat{\gamma}_\kappa^\dagger \hat{\gamma}_\kappa + \text{const.} \quad (11)$$

Its ground state is a Gaussian state of the form in Eq. (2), that corresponds to the vacuum of the $\hat{\gamma}_\kappa$ operators [see Eq. (7)].

We note that the above formalism can be easily generalized to the case of time-varying coefficients $Q_{i,j}(t)$ and $P_{i,j}(t)$ in Eq. (1), by admitting a time dependence of the Bogoliubov transformation matrix $\mathbb{U} = \mathbb{U}(t)$. Moreover, as we shall see below in Section 3, the form of a generic Gaussian state Eq. (2) is preserved by the application of the exponential of any operator quadratic in the \hat{c} fermions.

3 Evolving Gaussian states

We now discuss the evolution of a generic Gaussian state $|\psi\rangle$ under the effect of an operator of the form

$$\hat{M}(s) = e^{\xi s \hat{A}}, \quad (12a)$$

being $s \in \mathbb{R}$, $\xi = \{1, i\}$, and

$$\hat{A} = \sum_{i,j} \left(D_{i,j} \hat{c}_i^\dagger \hat{c}_j + O_{i,j} \hat{c}_i^\dagger \hat{c}_j^\dagger + \text{h.c.} \right), \quad (12b)$$

a generic Hermitian quadratic operator. For simplicity of notations, we assume all coefficients in \hat{A} to be real, but our analysis can be easily generalized to the complex case.

Due to the fermionic anticommutation rules, we have $D^T = D$, $O^T = -O$. Notice that, if $\xi = i$, the operator in Eqs. (12) describes the real-time evolution according to a (pseudo)-Hamiltonian \hat{A} . On the other hand, if $\xi = 1$, it describes an imaginary-time evolution (related to the construction of the thermal ensemble at inverse temperature $\beta = 1/s$), that can be thought of as the analytic continuation of the purely imaginary evolution. As discussed in Section 4, in some particular cases, the action played by such kind of operator on a given state can be intended as part of a measurement process.

In what follows we discuss how to evaluate, for any (real) symmetric \hat{A} operator, the s -evolved state

$$|\psi(s)\rangle = \hat{M}(s) |\psi\rangle,$$

taking as initial state $|\psi\rangle$ a generic Gaussian state of the form (2). In Appendix A we show that, because of the quadratic structure of \hat{A} , the state $|\psi(s)\rangle$ keeps the Gaussian form of Eq. (2):

$$|\psi(s)\rangle \propto \exp \left\{ \frac{1}{2} \sum_{p,q} Z_{p,q}(s) \hat{c}_p^\dagger \hat{c}_q^\dagger \right\} |0\rangle_{\hat{c}}, \quad (13)$$

with

$$Z(s) = -[u^\dagger(s)]^{-1} v^\dagger(s) \quad (14)$$

and $u(s), v(s)$ submatrices of the s -evolved Bogoliubov matrix $\mathbb{U}(s)$. In the next Subsection we are going to show how to evaluate the matrix $\mathbb{U}(s)$ (an alternative derivation is provided in Appendix A).

3.1 Bogoliubov matrix evolution

To characterize the evolved state $|\psi(s)\rangle$, one only needs to know how the Bogoliubov matrix $\mathbb{U}(s)$ evolves with $\hat{M}(s)$. Exploiting the identity $[\hat{M}(s)]^{-1} = \hat{M}(-s)$, we can write

$$|\psi(s)\rangle = \hat{M}(s) |\psi\rangle \propto e^{\hat{Z}(s)} \hat{M}(s) |0\rangle_{\hat{c}}, \quad (15)$$

where $|\psi\rangle$ is a generic Gaussian state given by Eq. (2). The operator

$$\hat{Z}(s) = \frac{1}{2} \sum_{p,q} Z_{p,q} \hat{c}_p^\dagger(s) \hat{c}_q^\dagger(s) \quad (16)$$

is defined through the evolved fermions

$$\hat{c}_q^{(\dagger)}(s) = \hat{M}(s) \hat{c}_q^{(\dagger)} \hat{M}(-s), \quad (17)$$

obeying anticommutation relations $\{\hat{c}_q(s), \hat{c}_p^\dagger(s)\} = \delta_{p,q}$. Note that, for $\xi = 1$, the transformation $\hat{M}(s)$ is not unitary and the conjugation operation does not commute with the evolution one, hence $\hat{c}_q^\dagger(s) \neq [\hat{M}(s) \hat{c}_q \hat{M}(-s)]^\dagger$. Since $\hat{M}(s) |0\rangle_{\hat{c}} \propto |0\rangle_{\hat{c}}$, to determine the evolved state, one has to calculate $\hat{Z}(s)$. This can be done by evaluating

$$\partial_s \hat{c}_q^{(\dagger)}(s) = \xi \hat{M}(s) [\hat{A}, \hat{c}_q^{(\dagger)}] \hat{M}(-s). \quad (18)$$

Given the quadratic structure of the operator \hat{A} , the above commutators read

$$[\hat{A}, \hat{c}_q^\dagger] = 2 \sum_i (D_{i,q} \hat{c}_i^\dagger + O_{q,i} \hat{c}_i), \quad (19a)$$

$$[\hat{A}, \hat{c}_q] = -2 \sum_i (D_{i,q} \hat{c}_i + O_{q,i} \hat{c}_i^\dagger). \quad (19b)$$

Therefore

$$\partial_s \hat{c}_q^\dagger(s) = 2\xi \sum_i (O_{q,i} \hat{c}_i(s) + D_{i,q} \hat{c}_i^\dagger(s)), \quad (20a)$$

$$\partial_s \hat{c}_q(s) = -2\xi \sum_i (D_{i,q} \hat{c}_i(s) + O_{q,i} \hat{c}_i^\dagger(s)). \quad (20b)$$

At this point, we make the Ansatz that the fermionic operators depend on s only through a Bogoliubov-like matrix

$$\hat{\Psi}(s) = \mathbb{U}^{\text{ev}}(s) \hat{\Phi}, \quad \text{with} \quad \mathbb{U}^{\text{ev}}(s) = \begin{pmatrix} u^{\text{ev}}(s) & \bar{v}^{\text{ev}}(s) \\ v^{\text{ev}}(s) & \bar{u}^{\text{ev}}(s) \end{pmatrix}. \quad (21)$$

The expression for the submatrices of the $\mathbb{U}^{\text{ev}}(s)$ is given in Eq. (29). Note that, while for $\xi = i$ one has $\bar{u}^{\text{ev}}(s) = [u^{\text{ev}}(s)]^*$ and $\bar{v}^{\text{ev}}(s) = [v^{\text{ev}}(s)]^*$ (so that $\mathbb{U}^{\text{ev}}(s)$ is a true Bogoliubov transformation), in general this is not true.

The above Ansatz is equivalent to assume that the s -evolved state $|\psi(s)\rangle$ is a Bogoliubov vacuum $\hat{\gamma}_\mu(s) |\psi(s)\rangle = 0$, $\forall \mu$. By substituting this Ansatz in Eq. (20), we obtain a set of differential equations for the elements of the evolved matrix:

$$\partial_s u_{q,\mu}^{\text{ev}}(s) = -2\xi \sum_i (D_{q,i} u_{i,\mu}^{\text{ev}}(s) + O_{q,i} v_{i,\mu}^{\text{ev}}(s)), \quad (22a)$$

$$\partial_s \bar{v}_{q,\mu}^{\text{ev}}(s) = -2\xi \sum_i (D_{q,i} \bar{v}_{i,\mu}^{\text{ev}}(s) + O_{q,i} \bar{u}_{i,\mu}^{\text{ev}}(s)), \quad (22b)$$

$$\partial_s v_{q,\mu}^{\text{ev}}(s) = 2\xi \sum_i (D_{q,i} v_{i,\mu}^{\text{ev}}(s) + O_{q,i} u_{i,\mu}^{\text{ev}}(s)), \quad (22c)$$

$$\partial_s \bar{u}_{q,\mu}^{\text{ev}}(s) = 2\xi \sum_i (D_{q,i} \bar{u}_{i,\mu}^{\text{ev}}(s) + O_{q,i} \bar{v}_{i,\mu}^{\text{ev}}(s)), \quad (22d)$$

with the initial condition: $u_{q,\mu}^{\text{ev}}(0) = [\bar{u}_{q,\mu}^{\text{ev}}(0)]^* = u_{q,\mu}$, and $v_{q,\mu}^{\text{ev}}(0) = [\bar{v}_{q,\mu}^{\text{ev}}(0)]^* = v_{q,\mu}$. In a compact form, these equations read

$$\partial_s \mathbb{U}^{\text{ev}}(s) = -2\xi \mathbb{A} \mathbb{U}^{\text{ev}}(s), \quad \text{with } \mathbb{A} = \begin{pmatrix} D & O \\ -O & -D \end{pmatrix}, \quad (23)$$

being $D^T = D$, $O^T = -O$. This leads to the formal solution

$$\mathbb{U}^{\text{ev}}(s) = e^{-2\xi \mathbb{A} s} \mathbb{U}. \quad (24)$$

The matrix $\mathbb{U}^{\text{ev}}(s)$ is not *a priori* characterizing a fermionic state in that it should be unitary, to guarantee the normalization of the state, and should obey Eqs. (4), to impose the anticommutation relations for the $\hat{\Psi}(s)$. While the former requirement is automatically fulfilled for $\xi = i$ (because of the unitarity of the evolution operator), in the case $\xi = 1$ it must be forced by performing a QR-decomposition, as discussed in details in the next subsection. Once the unitarity of the matrix is imposed, Eqs. (4) are satisfied if the Bogoliubov matrix preserves the symplectic structure of Eq. (6), and this happens for \hat{A} being Hermitian.

By substituting the normalized s-evolved Bogoliubov matrix in $Z(s)$ [cf. Eq. (14)], we obtain the expression for $|\psi(s)\rangle$. If the above conditions are satisfied, this is a Gaussian fermionic state, as the one in Eq. (13). A more detailed discussion about this point is provided in Appendix A.

3.2 QR decomposition

We now discuss how to restore the unitarity of the s-evolved Bogoliubov matrix $\mathbb{U}^{\text{ev}}(s)$, when considering a norm-non preserving evolution with $\xi = 1$, that is in general obtained by QR-decomposing the s-evolved Bogoliubov matrix $\mathbb{U}^{\text{ev}}(s)$. This procedure keeps the $Z(s)$ matrix in Eq. (14) – and then the Gaussian state – invariant, as we are going to see below. Nevertheless, it is necessary to apply the known formulae [56] for evaluating the entanglement entropy and the local observables.

The QR-decomposition is a procedure that allows one to decompose a generic matrix as $K = U_Q R$, with U_Q a unitary matrix and R an upper triangular one. In what follows, we provide an argument to show that, if $\hat{A} = \hat{A}^\dagger$, the unitary matrix obtained by QR-decomposing the s-evolved Bogoliubov matrix $\mathbb{U}^{\text{ev}}(s)$ describes a fermionic state, i.e., it has the symplectic form of Eq. (6). The straightforward way would be to check it by construction, a procedure that requires some involved calculations. Here we choose a different strategy. Since R is positive definite (we have checked it numerically), the QR decomposition must be unique. Therefore, we assume U_Q to have the right structure and then we show that it exists a decomposition that is compatible with this assumption.

Let us consider a symmetric operator \hat{A} defined in Eq. (12b) and the associated coefficient matrix \mathbb{A} defined in Eq. (23). Because of the symmetries of D and O , we have that $\{D, O\} = 0$.³ We notice that

$$\mathbb{A}^{2n} = \begin{pmatrix} X^{2n} & 0 \\ 0 & X^{2n} \end{pmatrix}, \quad \text{with } X^2 = D^2 - O^2. \quad (25)$$

By exploiting the block diagonal form of Eq. (25) and the identities

$$\sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!} = \cosh(x), \quad \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n+1)!} = \frac{\sinh(x)}{x}, \quad (26)$$

²For $s = it$ and $\hat{A} = \hat{H}$, it returns the usual Hamiltonian time evolution of the system [56].

³This simply follows from the fact that $(\{D, O\})^\dagger = -\{D, O\}$. Being D and O real matrices, it implies that the anticommutator vanishes.

we can write

$$e^{-2\xi\mathbb{A}s} = \sum_{n=0}^{\infty} \left[\frac{(-2\xi\mathbb{A}s)^{2n}}{(2n)!} + \frac{-2\xi\mathbb{A}s (-2\xi\mathbb{A}s)^{2n}}{(2n+1)!} \right] = \begin{pmatrix} \text{ch}^+(s) & \text{sh}(s) \\ -\text{sh}(s) & \text{ch}^-(s) \end{pmatrix}, \quad (27)$$

where, for shortness of notation, we have posed

$$\text{ch}^{\pm}(s) = I \cosh(-2\xi X s) \pm D \sinh(-2\xi X s) X^{-1}, \quad \text{sh}(s) = O \sinh(-2\xi X s) X^{-1}. \quad (28)$$

Notice that $[\text{ch}^{\pm}(s)]^{\dagger} = \text{ch}^{\pm}(s)$, while $\text{sh}^{\dagger}(s) = -\text{sh}(s)$. Moreover, since $\{D, O\} = 0$, we have $[D, \text{ch}^{\pm}(s)] = [O, \text{ch}^{\pm}(s)] = 0$ and $[D, \text{sh}(s)] = [O, \text{sh}(s)] = 0$. This means that

$$\begin{pmatrix} u^{\text{ev}}(s) & \bar{v}^{\text{ev}}(s) \\ v^{\text{ev}}(s) & \bar{u}^{\text{ev}}(s) \end{pmatrix} = \begin{pmatrix} \text{ch}^+(s)u + \text{sh}(s)v & \text{ch}^+(s)v^* + \text{sh}(s)u^* \\ -\text{sh}(s)u + \text{ch}^-(s)v & -\text{sh}(s)v^* + \text{ch}^-(s)u^* \end{pmatrix}. \quad (29)$$

We can QR-decompose $\mathbb{U}^{\text{ev}}(s)$

$$\mathbb{U}^{\text{ev}}(s) = \mathbb{U}(s) R(s), \quad (30)$$

making the Ansatz that the $\mathbb{U}(s)$ preserves the symplectic form of Eq. (6), so that (30) can be rewritten as

$$\begin{pmatrix} u^{\text{ev}}(s) & \bar{v}^{\text{ev}}(s) \\ v^{\text{ev}}(s) & \bar{u}^{\text{ev}}(s) \end{pmatrix} = \begin{pmatrix} u(s) & v^*(s) \\ v(s) & u^*(s) \end{pmatrix} \begin{pmatrix} r_1(s) & r_3(s) \\ 0 & r_2(s) \end{pmatrix}. \quad (31)$$

In what follows, we show that there exists a decomposition compatible with this Ansatz, by imposing Eq. (4) and that, because of the unicity, it must be the only possible. Let us assume to have derived $r_1(s)$ by construction. We can explicitly write $\mathbb{U}(s) = \mathbb{U}^{\text{ev}}(s)R(s)^{-1}$, exploiting the fact that the inverse of an upper block triangular matrix remains an upper block triangular matrix:

$$\begin{pmatrix} u(s) & v^*(s) \\ v(s) & u^*(s) \end{pmatrix} = \begin{pmatrix} u^{\text{ev}}(s) r_1^{-1}(s) & u^{\text{ev}}(s) r_3^{-1}(s) + \bar{v}^{\text{ev}}(s) r_2^{-1}(s) \\ v^{\text{ev}}(s) r_1^{-1}(s) & v^{\text{ev}}(s) r_3^{-1}(s) + \bar{v}^{\text{ev}}(s) r_2^{-1}(s) \end{pmatrix}. \quad (32)$$

We can substitute these expressions in Eq. (4) to obtain

$$u^{\dagger}u + v^{\dagger}v = [r_2^T(s)]^{-1} \left([\bar{u}^{\text{ev}}]^T(s) u^{\text{ev}}(s) + [\bar{v}^{\text{ev}}]^T(s) v^{\text{ev}}(s) \right) [r_1(s)]^{-1} = I. \quad (33)$$

Since $[\bar{u}^{\text{ev}}]^T(s) u^{\text{ev}}(s) + [\bar{v}^{\text{ev}}]^T(s) v^{\text{ev}}(s) = I$, the above equation provides a relation between two of the blocks of the matrix $R(s)$:

$$r_2(s) = [r_1^T(s)]^{-1}. \quad (34)$$

By imposing the second constraint in Eq. (4), we find the relation for $r_3(s)$

$$r_3(s) = -r_1^T \left([u^{\text{ev}}]^{\dagger}(s) \bar{v}^{\text{ev}}(s) + [v^{\text{ev}}]^{\dagger}(s) \bar{u}^{\text{ev}}(s) \right)^{-1} \left([u^{\text{ev}}]^{\dagger}(s) u^{\text{ev}}(s) + [v^{\text{ev}}]^{\dagger}(s) v^{\text{ev}}(s) \right). \quad (35)$$

As a last comment we notice that the matrix $Z(s)$ in Eq. (14) – and then the Gaussian state – are left invariant by the application of the QR decomposition:

$$Z(s) = [u^{\dagger}(s)]^{-1} v(s)^{\dagger} = \left\{ [u^{\text{ev}}(s) r_1^{-1}(s)]^{\dagger} \right\}^{-1} [v^{\text{ev}}(s) r_1^{-1}(s)]^{\dagger} = \left\{ [u^{\text{ev}}]^{\dagger}(s) \right\}^{-1} [v^{\text{ev}}]^{\dagger}(s). \quad (36)$$

4 Number-preserving operators

We now consider a generic number-preserving operator

$$\hat{A}_{\mathcal{I},\mathcal{J}} = \sum_{i \in \mathcal{I}} \sum_{j \in \mathcal{J}} \hat{c}_i^\dagger \hat{c}_j, \quad (37)$$

where \mathcal{I}, \mathcal{J} are two sets of indices labeling certain sites on the system, and apply

$$\hat{M}_{\mathcal{I},\mathcal{J}} = e^{\beta \hat{A}_{\mathcal{I},\mathcal{J}}} \quad (38)$$

to a Gaussian state. We focus on this particular example of the theory presented in Section 3.1, since it generalizes the construction discussed in Ref. [38,39]. By exploiting the anticommutation relations, it is possible to show that $\hat{A}_{\mathcal{I},\mathcal{J}}^n = \alpha^{n-1} \hat{A}_{\mathcal{I},\mathcal{J}}$, with $\alpha = |\mathcal{I} \cap \mathcal{J}|$. In fact

$$\begin{aligned} \hat{A}_{\mathcal{I},\mathcal{J}}^2 &= \sum_{i,l \in \mathcal{I}} \sum_{j,m \in \mathcal{J}} \hat{c}_i^\dagger \hat{c}_j \hat{c}_l^\dagger \hat{c}_m = \sum_{i,l \in \mathcal{I}} \sum_{j,m \in \mathcal{J}} (\hat{c}_i^\dagger \hat{c}_j \delta_{l,m} + \hat{c}_i^\dagger \hat{c}_j \hat{c}_m \hat{c}_l^\dagger) \\ &= \sum_{i,l \in \mathcal{I}} \sum_{j,m \in \mathcal{J}} \left[\hat{c}_i^\dagger \hat{c}_j \delta_{l,m} + \frac{1}{2} \hat{c}_i^\dagger (\hat{c}_j \hat{c}_m + \hat{c}_m \hat{c}_j) \hat{c}_l^\dagger \right] = |\mathcal{I} \cap \mathcal{J}| \sum_{i \in \mathcal{I}} \sum_{j \in \mathcal{J}} \hat{c}_i^\dagger \hat{c}_j, \end{aligned} \quad (39)$$

where the last equality holds, since j and m run in the same set. Therefore, if $\mathcal{I} \cap \mathcal{J} = \emptyset$, we have $\hat{A}_{\mathcal{I},\mathcal{J}}^2 = 0$ and $\hat{M}_{\mathcal{I},\mathcal{J}} = \hat{1} + \hat{A}_{\mathcal{I},\mathcal{J}}$, where $\hat{1}$ denotes the identity operator. Otherwise

$$\hat{M}_{\mathcal{I},\mathcal{J}} = \sum_{n=0}^{\infty} \frac{(\beta \hat{A}_{\mathcal{I},\mathcal{J}})^n}{n!} = \hat{1} + \frac{(e^{\alpha\beta} - 1) \hat{A}_{\mathcal{I},\mathcal{J}}}{\alpha}. \quad (40)$$

In what follows, without loss of generality, we set $\beta \equiv \log(\alpha+1)/\alpha$, so that $\hat{M}_{\mathcal{I},\mathcal{J}} = \hat{1} + \hat{A}_{\mathcal{I},\mathcal{J}}$ reduces to that studied in Ref. [38,39] for $\mathcal{I} = \mathcal{J} = \{i\}$. As discussed in Appendix B, the application of such a $\hat{M}_{\mathcal{I},\mathcal{J}}$ can be thought as the measurement step of a quantum-jump dynamics.

We want to evaluate the effect of this operator on a Gaussian state

$$|\psi\rangle_{\hat{M}_{\mathcal{I},\mathcal{J}}} \equiv \hat{M}_{\mathcal{I},\mathcal{J}} |\psi\rangle = (\hat{1} + \hat{A}_{\mathcal{I},\mathcal{J}}) |\psi\rangle. \quad (41)$$

Following the procedure discussed in Section 3.1, we write the evolved state as

$$|\psi\rangle_{\hat{M}_{\mathcal{I},\mathcal{J}}} \propto \exp \left\{ \frac{1}{2} \sum_{p,q} Z_{p,q} \hat{k}_p^\dagger \hat{k}_q^\dagger \right\} |0\rangle, \quad (42)$$

where we introduced the transformed fermionic operators $\hat{k}_q^{(\dagger)} = \hat{M}_{\mathcal{I},\mathcal{J}} \hat{c}_q^{(\dagger)} \hat{M}_{\mathcal{I},\mathcal{J}}^{-1}$. By exploiting the fact that $\hat{M}_{\mathcal{I},\mathcal{J}}^{-1} = \hat{1} - (\alpha+1)^{-1} \hat{A}_{\mathcal{I},\mathcal{J}}$, we derive the exact expression of $\hat{k}^{(\dagger)}$

$$\hat{k}_q^\dagger = \hat{c}_q^\dagger + \sum_{j \in \mathcal{J}} \delta_{q,j} \sum_{i \in \mathcal{I}} \hat{c}_i^\dagger, \quad (43a)$$

$$\hat{k}_q = \hat{c}_q - (\alpha+1)^{-1} \sum_{j \in \mathcal{J}} \delta_{q,j} \sum_{i \in \mathcal{I}} \hat{c}_i. \quad (43b)$$

This leads to the following expressions for the evolved Bogoliubov matrix:

$$u_{q,\mu}^{\text{ev}} = u_{q,\mu} - (\alpha+1)^{-1} \sum_{j \in \mathcal{J}} \delta_{q,j} \sum_{i \in \mathcal{I}} u_{i,\mu}, \quad \bar{u}_{q,\mu}^{\text{ev}} = u_{q,\mu}^* + \sum_{j \in \mathcal{J}} \delta_{q,j} \sum_{i \in \mathcal{I}} u_{i,\mu}^*, \quad (44a)$$

$$\bar{v}_{q,\mu}^{\text{ev}} = v_{q,\mu}^* - (\alpha+1)^{-1} \sum_{j \in \mathcal{J}} \delta_{q,j} \sum_{i \in \mathcal{I}} v_{i,\mu}^*, \quad v_{q,\mu}^{\text{ev}} = v_{q,\mu} + \sum_{j \in \mathcal{J}} \delta_{q,j} \sum_{i \in \mathcal{I}} v_{i,\mu}. \quad (44b)$$

As discussed in Appendix C, an alternative derivation can be obtained from the evolved antisymmetric matrix Z .

4.1 Local operator

The above results can be applied straightforwardly to local (l) operators, as for the one used in Ref. [38, 39], $\hat{A}_{\mathcal{I},\mathcal{J}}^l \equiv \hat{n}_i = \hat{c}_i^\dagger \hat{c}_i$, such that $\hat{M}_{\mathcal{I},\mathcal{J}}^l \equiv e^{\log(2)\hat{A}^l} = \hat{1} + \hat{c}_i^\dagger \hat{c}_i$. The Bogoliubov matrix, evolves according to Eqs. (44), which read

$$u_{q,\mu}^{\text{ev},1} = (1 - \delta_{q,i}/2)u_{q,\mu}, \quad \bar{u}_{q,\mu}^{\text{ev},1} = (1 + \delta_{q,i})u_{q,\mu}^*, \quad (45a)$$

$$\bar{v}_{q,\mu}^{\text{ev},1} = (1 - \delta_{q,i}/2)v_{q,\mu}^*, \quad v_{q,\mu}^{\text{ev},1} = (1 + \delta_{q,i})v_{q,\mu}. \quad (45b)$$

This evolution is implemented by applying to the \mathbb{U} a block-diagonal matrix T with entries $T_{i+L,i+L} = T_{i,i}^{-1} = 2$ (see Eq. (1) of Ref. [39]).

4.2 Two-site operator

We now consider a non-local (nl) operator with support on two sites ($\mathcal{I} = \mathcal{J} = \{i, j\}$):

$$\hat{A}_{\mathcal{I},\mathcal{J}}^{\text{nl}} = (\hat{c}_i^\dagger + \hat{c}_j^\dagger)(\hat{c}_i + \hat{c}_j), \quad (46)$$

Since $\alpha = |\mathcal{I} \cap \mathcal{J}| = 2$, we have $\hat{M}_{\mathcal{I},\mathcal{J}}^{\text{nl}} \equiv e^{\frac{\log(3)}{2}\hat{A}^{\text{nl}}} = \hat{1} + \hat{A}_{\mathcal{I},\mathcal{J}}^{\text{nl}}$. Without losing in generality, in the following we will focus on one dimensional systems and set $j = i + r$ (with $r > 0$). As we shall see in the next Section, a measurement operator as the one in Eq. (46) may unveil a richer physics than local ones. In fact, in the spin-1/2 language, it corresponds to a string operator. The evolved Bogoliubov matrix can be derived from Eqs. (44) which, in this case, read

$$u_{q,\mu}^{\text{ev},\text{nl}} = u_{q,\mu} - \frac{1}{3}(\delta_{q,i} + \delta_{q,i+r})(u_{i,\mu} + u_{i+r,\mu}), \quad (47a)$$

$$\bar{v}_{q,\mu}^{\text{ev},\text{nl}} = v_{q,\mu}^* - \frac{1}{3}(\delta_{q,i} + \delta_{q,i+r})(v_{i,\mu}^* + v_{i+r,\mu}^*), \quad (47b)$$

$$\bar{u}_{q,\mu}^{\text{ev},\text{nl}} = u_{q,\mu}^* + (\delta_{q,i} + \delta_{q,i+r})(u_{i,\mu}^* + u_{i+r,\mu}^*), \quad (47c)$$

$$v_{q,\mu}^{\text{ev},\text{nl}} = v_{q,\mu} + (\delta_{q,i} + \delta_{q,i+r})(v_{i,\mu} + v_{i+r,\mu}). \quad (47d)$$

5 Quantum jumps with string operators

In this section we use the machinery presented above to describe the evolution of a quantum Ising chain under the effect of a quantum-jump dynamics generated by $\hat{M}_{\mathcal{I},\mathcal{J}}^{\text{nl}}$ [cf. Eq. (46) and below], with emphasis on the evolution of the entanglement entropy.

We consider the Hamiltonian

$$\hat{H} = -J \sum_j \hat{\sigma}_j^x \hat{\sigma}_{j+1}^x - h \sum_j \hat{\sigma}_j^z, \quad (48)$$

where $\hat{\sigma}_j^\alpha$ ($\alpha = x, y, z$) are the usual spin-1/2 Pauli matrices acting on the j -th site, while $J > 0$ denotes the ferromagnetic spin-spin coupling and h the transverse magnetic field. In the thermodynamic limit, the ground state exhibits a zero-temperature continuous transition at $|h/J| = 1$, separating a paramagnetic (symmetric) phase, for $|h/J| > 1$, from a ferromagnetic (symmetry-broken) phase, for $|h/J| < 1$ [57, 58]. Through a Jordan-Wigner transformation [59, 60], the Ising model (48) can be mapped into the so-called Kitaev chain

$$\hat{H} = -J \sum_j \left(\hat{c}_j^\dagger \hat{c}_{j+1} + \hat{c}_j^\dagger \hat{c}_{j+1}^\dagger + \text{h.c.} \right) - 2h \sum_j \hat{c}_j^\dagger \hat{c}_j. \quad (49)$$

This Hamiltonian is of the form (1), with

$$Q_{j,j} = h, \quad Q_{j,j+1} = Q_{j+1,j} = -J/2, \quad P_{j,j+1} = -P_{j+1,j} = -J/2. \quad (50)$$

Hereafter we work in units of $\hbar = 1$, set $J = 1$ as the energy scale, and assume periodic boundary conditions for fermions.

We implement the quantum-jump protocol described in Appendix B on the fermionic model above, by initializing the system in the ground state of the Hamiltonian (49) with a given value of h and then Trotterizing the time evolution in steps $\delta t \propto (4N\gamma)^{-1}$, to ensure convergence, being γ the strength of the measurements. The emerging dynamics is one of the possible unravelings [10] of the Lindblad master equation [9]

$$\dot{\rho}(t) = -i[\hat{H}, \rho(t)] - \gamma \sum_j \left(\hat{A}_j(r) \rho \hat{A}_j(r) - \frac{1}{2} \{ \hat{A}_j^2(r), \rho \} \right), \quad (51)$$

being $\hat{A}_j(r) = (\hat{c}_j^\dagger + \hat{c}_{j+r}^\dagger)(\hat{c}_j + \hat{c}_{j+r})$ the same operator as in Eq. (46), with $i \equiv j + r$.⁴ In the spin language, this operator is a sum of strings of Pauli matrices, of length r :

$$\hat{A}_j(r) = \prod_{k < j} \hat{\sigma}_k^z \hat{\sigma}_j^- \prod_{k' < j+r} \hat{\sigma}_{k'}^z \hat{\sigma}_{j+r}^+ + \prod_{k < j+r} \hat{\sigma}_k^z \hat{\sigma}_{j+r}^- \prod_{k' < j} \hat{\sigma}_{k'}^z \hat{\sigma}_j^+ + \frac{1}{2} (\hat{\sigma}_j^z + \hat{\sigma}_{j+r}^z - 2), \quad (52)$$

with $\hat{\sigma}_j^\pm = \hat{\sigma}_j^x \pm i\hat{\sigma}_j^y$. Hereafter, we refer to it as the string operator.

We point out that our framework is similar to the one presented in Ref. [31], although with some important qualitative differences. The former is described by a random sequence of stabilizer-formalism measurements of Pauli string operators. These operators have a fixed range r and are taken from a random distribution of strings, located around different sites, and made of different sets of onsite Pauli operators.

In this work, in contrast, we are following an unraveling of the quantum master equation Eq. (51). As detailed in Appendix B, we apply random quantum-jump measurements of string operators $\hat{1} + \hat{A}_j(r)$, located at different sites, but with the structure fixed as the one in Eq. (52). This is sufficient to achieve a measurement-induced nontrivial dynamics, with important consequences on the entanglement generation.

It is important to emphasize that the operators $\hat{1} + \hat{A}_j(r)$ have the exponential form given by Eqs. (38) and (40) (see also Sec. 4). In this way, the state keeps its Gaussian form under the application of the quantum jumps, and a numerical analysis of quite large system sizes is possible.

5.1 Entanglement entropy

Let us focus our attention on the entanglement entropy. Given a pure state $|\psi\rangle$, there is an unambiguous way to quantify its quantum correlations. We divide the system in two subchains, one with length ℓ and the other with length $L - \ell$. After evaluating the density matrix $\rho_\ell = \text{Tr}_{L-\ell} [|\psi\rangle\langle\psi|]$ reduced to the subsystem ℓ , the entanglement entropy is quantified by its von Neumann entropy [7, 8]:

$$S_\ell[\rho_\ell] = -\rho_\ell \log \rho_\ell. \quad (53)$$

For the Ising model we are discussing herewith, it is known that after, a sudden quench in a purely Hamiltonian dynamics, the entanglement entropy approaches a stationary value that increases linearly with ℓ , thus obeying a volume-law behavior [3, 4]. As discussed in Refs. [33–40], this dynamical volume law is generally destroyed by any arbitrary small

⁴Note that, with our convention for the boundary conditions, we have $\hat{c}_{j+r}^{(\dagger)} \equiv \hat{c}_{j+r-L}^{(\dagger)}$ for any $j + r > L$.

local weak measurement process. In this scenario, the asymptotic entanglement entropy is conjectured to exhibit either a logarithmic scaling or an area-law behavior, depending on the strength of measurements and of the Hamiltonian parameters. We are going to show that the situation is quite different for the case of the nonlocal operators defined in Eq. (52).

In practice, to avoid spurious contributions from classical correlations, we evaluate Eq. (53) on each stochastic quantum trajectory $|\psi_t\rangle$ (defined in Appendix B), and then average over different stochastic realizations.⁵ For simplicity of notation, we denote the ensemble-averaged entanglement entropy as

$$S_\ell(t) \equiv \langle\langle S_\ell[\rho_\ell^{\text{stoch}}(t)] \rangle\rangle, \quad \text{where} \quad \rho_\ell^{\text{stoch}}(t) = \text{Tr}_{L-\ell}[|\psi_t\rangle\langle\psi_t|], \quad (54)$$

and where the symbol $\langle\langle \dots \rangle\rangle$ denotes the average over different stochastic realizations. We also define the asymptotic entanglement entropy as

$$\bar{S}_\ell = \lim_{T \rightarrow \infty} \int_{t^*}^T dt' S_\ell(t'). \quad (55)$$

We remark that, in all the simulations shown here, we keep fixed the partition to $\ell = L/4$ and study the scaling with L . Below we present two different scenarios: the measurement-only dynamics (Section 5.2) and the full quantum-jump dynamics (Section 5.3).

5.2 Measurement-only dynamics

We first neglect the Hamiltonian contribution, i.e., we implement the quantum-jump protocol described in Appendix B with $\hat{H} = 0$. The Hamiltonian is only used to set the initial state as its ground state with $h = 0.5$. However, we have carefully checked that this choice does not affect the asymptotic values of $S_\ell(t)$, but only the transient dynamics.⁷ The measurement-only dynamics allows to isolate the effects of measuring the string operators, without considering the interplay with the unitary dynamics.

The upper panels of Figure 1 report the ensemble-averaged entanglement entropy $S_\ell(t)$ versus the rescaled time γt , for three ranges of the string measurement operator: $r = 1$ (panel a), $r = 4$ (panel b), and $r = L/2$ (panel c), and for different system sizes L . For $r = 1$, the measurement-only dynamics cannot generate entanglement and the system stabilizes on a low-entangled asymptotic state. This is confirmed by Fig. 1(d), that shows the behavior of the asymptotic entanglement entropy \bar{S}_ℓ versus L , which tends to saturate to a constant value, thus obeying an area-law behavior. As expected, this result is analogous to the case where onsite operators are measured [38, 39].

When the correlation range of the dissipative string operator is increased, as for $r = 4$, we may identify three regimes: the short-time transient, the metastable plateau, and the asymptotic dynamics. At short times the entanglement entropy [cf. inset of Fig. 1(b)] grows to reach a maximum, that saturates to a constant value for large L . This is clarified by the red curve in Fig. 1(e), showing the maximum value of the entanglement entropy versus the system size (notice the logarithmic scale of the x -axis). After the initial transient, the entanglement entropy attains a metastable plateau where the entanglement entropy exhibits an area-law behavior [green curve in Fig. 1(e)]. Finally, in a time $\tau \sim L^2$, the

⁵For details on how to evaluate the von Neumann entropy of the reduced density matrix of Gaussian states see, e.g., Ref. [56].

⁶Ensemble-averaged entanglement entropies have then been obtained with $N_{\text{avg}} = 10^2$ different quantum trajectories. Subsequently, to perform time averages, we fix a given evolution time T and average in the interval $t' \in [0.7T, T]$.

⁷Note that the perfectly paramagnetic state with spins pointing along the z direction ($h = \infty$) is a dark state of the dissipation, therefore, initializing the system close to such state leads to slower convergence.




Figure 1: Top panels: plots of $S_\ell(t)$ versus the rescaled time γt , during the quantum-jump process with $\hat{H} = 0$. The different panels refer to $r = 1$ (a), $r = 4$ (b), and $r = L/2$ (c). The various curves are for different sizes L [see color code in (a)]. The inset in (b) is a magnification at small times, where a maximum emerges in correspondence of the vertical dashed line. Bottom panels: scaling of \bar{S}_ℓ versus L , for $r = 1$ (d), $r = 4$ (e), $r = L/2$ (f). The three curves in (e) refer to the short-time peak (red), the metastable plateau (green), and the long-time value (blue). The dashed line in (f) extrapolates to the asymptotic value for $L \rightarrow \infty$.

entanglement entropy approaches a non-vanishing asymptotic value still obeying an area law [blue curve in Fig. 1(e)].

This scenario dramatically changes when r is forced to be a thermodynamic fraction of L (e.g., $r = L/2$): In that case, the measurement-only dynamics leads to a stationary state, whose entanglement obeys a volume-law behavior. This is clarified by Fig. 1(f), which shows the rescaled asymptotic entanglement entropy \bar{S}_ℓ/L versus L^{-1} . In this plot, the saturation of \bar{S}_ℓ/L to a finite value for $L^{-1} \rightarrow 0$ clearly suggests a linear growth $\bar{S}_\ell \propto L$ for large system sizes.

5.3 Effect of the Hamiltonian

We now switch on the Kitaev Hamiltonian \hat{H} and follow the full quantum-jump dynamics for $h = 0.1$, starting from the ground state with $h_i = 0.5$. As for the case with no Hamiltonian, we have checked that the asymptotic values reached by the entanglement entropy are independent of h_i , while they may depend on h (although the qualitative behaviors with L are unaffected by this choice).

Figure 2 displays $S_\ell(t)$ versus γt (upper panels) and the asymptotic value \bar{S}_ℓ versus L , for the three ranges of $r = 1$ (left) $r = 4$ (middle), and $r = L/2$ (right). In panel (a) we show the local case $r = 1$. The long-time averaged entanglement entropy attains a stationary value which is almost independent of L , as clearly visible from Fig. 2(d), thus reflecting an area-law behavior. Such value is generally larger than the one obtained in the measurement-only dynamics, without the entangling action of \hat{H} [compare with Fig. 1(a)].

In Fig. 2(b) we show the case $r = 4$, which displays a convergence in time faster than that for a measurement-only dynamics [compare with Fig. 1(b)]. The asymptotic data in




Figure 2: Same plots as in Figure 1, but in the simultaneous presence of the Hamiltonian and the string measurement process. We observe qualitatively similar behaviors, although with different time scales. Here we fix $\gamma = 0.5$, $h = 0.1$, and start from the ground state with $h_i = 0.5$.

Fig. 2(e) (x -axis in logarithmic scale) suggest that \bar{S}_ℓ initially grows sub-extensively with L , to eventually attain an area-law regime. Physically, this can be understood by the fact that for L much larger than r , the dissipation becomes effectively local and the extensive growth is destroyed.

In contrast with that, if the range r of the jump operator is comparable with the total size L , the measurements can stabilize a volume-law scaling of the entanglement, as we can see in Fig. 1(c), that refers to $r = L/2$. Here \bar{S}_ℓ increases in time and quickly attains an asymptotic value that follows a volume-law scaling [see the data in Fig. 1(f) for \bar{S}_ℓ/L versus L^{-1} , which display convergence to a finite value, in the $L \rightarrow \infty$ limit].

6 Conclusions

We have unveiled some aspects of the entanglement-entropy dynamics of fermionic many-body Gaussian states, in the presence of a Gaussian-preserving evolution, with a special focus on the possible emergence of measurement-induced phase transitions in such systems. The first part of the paper presents a detailed technical discussion on how to maintain and treat Gaussianity, when exponentials of quadratic fermion operators are applied to a Gaussian state. Our purpose is to provide the reader with a comprehensive guide to implement fermionic Gaussian evolutions and study the dynamics (not necessarily unitary) of such systems. Among all the possible treatable evolutions, we focused on the quantum-jump unraveling of a Lindblad dynamics induced by a particular class of quadratic fermion operators, having a simple definition in the fermionic language, but a non-trivial string structure in terms of the Pauli matrices. We called them string operators.

In the second part of the paper, after having derived the equation of motions for a quantum-jump dynamics induced by the string operators, we have solved it for the quantum Ising chain, focusing on the time behavior of the entanglement entropy. Our main result is that the scaling of the entanglement entropy with the system size L strongly depends on the range r of the string operators. For short strings (r finite), we see that the asymptotic entanglement entropy obeys a non-extensive scaling behavior. In this case,

the measurement operator is effectively local in space and thus it asymptotically destroys quantum correlations that may be generated by the unitary dynamics. For string ranges comparable with the system size ($r \sim L$), the measurement process may become highly entangling and is able to stabilize a volume-law scaling in the asymptotic entanglement entropy.

Remarkably, the picture above is valid both for a measurement-only dynamics and for a dynamics where also the effect of a short-range Hamiltonian is considered. So, the measurement dynamics of the string operators is strong enough to induce different entanglement scalings by itself.

A natural extension of our work could be to understand how the MIPT scenario would change in the mutual presence of unitary dynamics and long-range measurement operators (e.g., by considering some power-law decay of dissipators with the distance). It would be also interesting to go beyond free-fermion models and address the role of interactions in the entanglement dynamics under variable-range monitoring.

Acknowledgements

We acknowledge fruitful discussions with V. Alba, M. Collura, R. Fazio, M. Schirò, and X. Turkeshi.

Funding information This work has been partly funded by the Italian MIUR through PRIN Project No. 2017E44HRF.

A Rotation to the diagonal basis

Here we discuss a different derivation of the s -evolved state $|\psi(s)\rangle = \hat{M}(s)|\psi\rangle$, where $|\psi\rangle$ is a generic Gaussian state [Eq. (2)], and the $\hat{M}(s) = e^{\xi s \hat{A}}$ is defined in Eqs. (12).

The operator \hat{A} [Eq. (12b)] is quadratic in the \hat{c} fermions and can be written compactly as $\hat{A} = \hat{\Psi}^\dagger \mathbb{A} \hat{\Psi}$, where $\hat{\Psi} = (\hat{c}_1, \dots, \hat{c}_L, \hat{c}_1^\dagger, \dots, \hat{c}_L^\dagger)^T$ is the Nambu spinor associated to the \hat{c} fermions and \mathbb{A} the Bogoliubov-de Gennes matrix associated to \hat{A} , as defined in Eq. (23). This matrix can be diagonalized by a Bogoliubov transformation \mathbb{W} , such that

$$\mathbb{W}^\dagger \mathbb{A} \mathbb{W} = \text{diag} [\epsilon_\nu, -\epsilon_\nu]_{\nu=1, \dots, L}, \quad \text{and} \quad \hat{A} = \sum_\nu \epsilon_\nu \left(\hat{\eta}_\nu^\dagger \hat{\eta}_\nu - \hat{\eta}_\nu \hat{\eta}_\nu^\dagger \right). \quad (56)$$

The unitary matrix \mathbb{W} defines a new set of $\hat{\eta}$ fermions, whose associated Nambu spinor is $\hat{\Theta} \equiv (\hat{\eta}_1, \dots, \hat{\eta}_L, \hat{\eta}_1^\dagger, \dots, \hat{\eta}_L^\dagger)^T = \mathbb{W}^\dagger \hat{\Psi}$.

Let us now evaluate $|\psi(s)\rangle$. We remind that the initial state $|\psi\rangle$ is the vacuum of the $\hat{\gamma}$ fermions, defined through the Bogoliubov transformation in Eq. (6): $\hat{\Phi} = \mathbb{U}^\dagger \hat{\Psi}$, where $\hat{\Phi} = (\hat{\gamma}_1, \dots, \hat{\gamma}_L, \hat{\gamma}_1^\dagger, \dots, \hat{\gamma}_L^\dagger)^T$. These fermions can be transformed into those diagonalizing \mathbb{A} by means of a rotation

$$\hat{\Phi} = \mathbb{U}^\dagger \mathbb{W} \hat{\Theta} = \mathbb{Q}^\dagger \hat{\Theta}, \quad \text{being} \quad \mathbb{Q} = \begin{pmatrix} u_\eta & v_\eta^* \\ v_\eta & u_\eta^* \end{pmatrix}. \quad (57)$$

Therefore we have the relation

$$\mathbb{U} = \mathbb{W} \mathbb{Q}, \quad (58)$$

with \mathbb{U} given in Eq. (5). In the $\hat{\eta}$ representation, the state (2) takes the form

$$|\psi\rangle \propto \exp\left\{\frac{1}{2}\sum_{\mu,\nu} Z_{\mu,\nu}^{(\eta)} \hat{\eta}_\mu^\dagger \hat{\eta}_\nu^\dagger\right\} |0\rangle_{\hat{\eta}}, \quad (59)$$

with $Z^{(\eta)} = -[u_\eta^\dagger]^{-1} v_\eta^\dagger$.

Going on analogously as in Sec. 3.1, we can write the s -evolved state as

$$|\psi'\rangle \propto \exp\left\{\frac{1}{2}\sum_{\mu,\nu} Z_{\mu,\nu}^{(\eta)} \hat{\eta}_\mu^\dagger(s) \hat{\eta}_\nu^\dagger(s)\right\} \hat{M}(s) |0\rangle_{\hat{\eta}}, \quad (60)$$

with $\hat{\eta}_\nu^\dagger(s) \equiv \hat{M}(s) \hat{\eta}_\nu^\dagger \hat{M}(-s)$. The operator $\hat{M}(s)$ is number-preserving, therefore we can exploit the methods of Section 4. We first notice that $\hat{M}(s) |0\rangle_{\hat{\eta}} \propto |0\rangle_{\hat{\eta}}$, in fact

$$e^{\xi s \sum_\nu \epsilon_\nu (\hat{\eta}_\nu^\dagger \hat{\eta}_\nu - \hat{\eta}_\nu \hat{\eta}_\nu^\dagger)} |0\rangle_{\hat{\eta}} = e^{-\xi s \sum_\nu \epsilon_\nu} |0\rangle_{\hat{\eta}}, \quad (61)$$

because $\hat{\eta}_\nu |0\rangle_{\hat{\eta}} = 0, \forall \nu \in \{1, \dots, L\}$. Then we notice that, since $(\hat{\eta}_\nu^\dagger \hat{\eta}_\nu)^n = \hat{\eta}_\nu^\dagger \hat{\eta}_\nu^n$ (cf. Section 4), we have

$$\begin{aligned} \hat{M}(\pm s) &= e^{\pm \xi s \hat{A}} = e^{\mp \xi s \sum_\nu \epsilon_\nu} \prod_\nu \left[1 + \sum_{n=1}^{\infty} \frac{(\pm 2\xi s \epsilon_\nu)^n}{n!} \hat{\eta}_\nu^\dagger \hat{\eta}_\nu \right] \\ &= e^{\mp \xi s \sum_\nu \epsilon_\nu} \prod_\nu \left[1 + (e^{\pm 2\xi s \epsilon_\nu} - 1) \hat{\eta}_\nu^\dagger \hat{\eta}_\nu \right], \end{aligned} \quad (62)$$

where we have applied the diagonal form of \hat{A} , in Eq. (56). Therefore, according to the fact that $[\hat{\eta}_\mu^\dagger \hat{\eta}_\mu, \hat{\eta}_\nu^\dagger] = \delta_{\mu,\nu} \hat{\eta}_\mu^\dagger$, we can write

$$\hat{\eta}_\nu^\dagger(s) = [1 + (e^{2\xi s \epsilon_\nu} - 1) \hat{\eta}_\nu^\dagger \hat{\eta}_\nu] \hat{\eta}_\nu^\dagger [1 + (e^{-2\xi s \epsilon_\nu} - 1) \hat{\eta}_\nu^\dagger \hat{\eta}_\nu] = e^{2\xi s \epsilon_\nu} \hat{\eta}_\nu^\dagger. \quad (63)$$

Substituting into Eq. (60) we get

$$|\psi(s)\rangle \propto \exp\left\{\frac{1}{2}\sum_{\mu,\nu} \tilde{Z}_{\mu,\nu}^{(\eta)} \hat{\eta}_\mu^\dagger \hat{\eta}_\nu^\dagger\right\} |0\rangle_{\hat{\eta}}, \quad \text{with } \tilde{Z}_{\mu,\nu}^{(\eta)} \equiv Z_{\mu,\nu}^{(\eta)} e^{2\xi s (\epsilon_\mu + \epsilon_\nu)}. \quad (64)$$

We can also write $\tilde{Z}^{(\eta)} = -[u_\eta^{\text{ev}\dagger}(s)]^{-1} v_\eta^{\text{ev}\dagger}(s)$, with

$$\begin{pmatrix} u_\eta^{\text{ev}}(s) \\ v_\eta^{\text{ev}}(s) \end{pmatrix} = \begin{pmatrix} e^{-2\xi s \epsilon} & 0 \\ 0 & e^{2\xi s \epsilon} \end{pmatrix} \begin{pmatrix} u_\eta \\ v_\eta \end{pmatrix}, \quad \text{being } \epsilon = \text{diag}(\epsilon_\nu). \quad (65)$$

At this point, if $\xi = 1$, we apply the QR decomposition to $\begin{pmatrix} u_\eta^{\text{ev}}(s) \\ v_\eta^{\text{ev}}(s) \end{pmatrix}$, as explained in Sec. 3.2. In this way, the matrix $\tilde{Z}^{(\eta)}$ remains unchanged, but can be written as $\tilde{Z}^{(\eta)} = -[u_\eta^\dagger(s)]^{-1} v_\eta^\dagger(s)$, with $u_\eta(s), v_\eta(s)$ obeying the symplectic unitarity conditions Eq. (4) and written as

$$\begin{pmatrix} u_\eta^{\text{ev}}(s) \\ v_\eta^{\text{ev}}(s) \end{pmatrix} = \begin{pmatrix} u_\eta(s) \\ v_\eta(s) \end{pmatrix} r_1, \quad (66)$$

with r_1 some upper triangular matrix $L \times L$.

We have therefore reduced to the standard Gaussian form for the evolved state $|\psi(s)\rangle$ in the $\hat{\eta}$ representation, with the Z matrix given by $\tilde{Z}^{(\eta)} = -[u_\eta^\dagger(s)]^{-1} v_\eta^\dagger(s)$, and the symplectic unitary matrix

$$\mathbb{Q}(s) = \begin{pmatrix} u_\eta(s) & v_\eta^*(s) \\ v_\eta(s) & u_\eta^*(s) \end{pmatrix} \quad (67)$$

providing the set of fermionic operators $\hat{\gamma}_\nu(s)$ annihilating $|\psi(s)\rangle$ [we have $\hat{\Phi}(s) = \mathbb{Q}^\dagger(s) \hat{\Theta}$, with $\hat{\Phi}(s) \equiv (\hat{\gamma}_1(s), \dots, \hat{\gamma}_L(s), \hat{\gamma}_1^\dagger(s), \dots, \hat{\gamma}_L^\dagger(s))^T$]. At this point it is possible to come back to the \hat{c} representation. We can use Eq. (58), and obtain the Bogoliubov matrix $\mathbb{U}(s)$ [the one such that $\hat{\Phi}(s) = \mathbb{U}^\dagger(s) \hat{\Psi}$] as $\mathbb{U}(s) = \mathbb{W} \mathbb{Q}(s)$. Applying \mathbb{W} on the left of both sides of Eq. (65) and writing

$$\mathbb{U}(s) = \begin{pmatrix} u(s) & v^*(s) \\ v(s) & u^*(s) \end{pmatrix},$$

we obtain

$$\begin{pmatrix} u(s) \\ v(s) \end{pmatrix} r_1 = e^{-2\xi s \mathbb{A}} \begin{pmatrix} u \\ v \end{pmatrix}. \quad (68)$$

[To get this result, we used the relation $\mathbb{U}(s) = \mathbb{W} \mathbb{Q}(s)$, as well as Eqs. (56), (58), and (66)]. The Z matrix of the evolved state in this representation is thus $Z(s) = -[u^\dagger(s)]^{-1} v^\dagger(s)$. Notice that Eq. (68) is the same result of Eq. (24), when the QR decomposition discussed in Sec. 3.2 is applied.

As a last comment we point out that Eq. (68) refers to the first two blocks of the Bogoliubov matrix \mathbb{U} only. This is consistent with the fact that, in the Nambu spinor notation, the degrees of freedom are doubled and there is some redundant information that can be neglected.

B Quantum-jump protocol

We start from the following master equation in the Lindblad form:

$$\dot{\rho}(t) = \mathcal{L}[\rho(t)] \equiv -i[\hat{H}, \rho(t)] - \gamma \sum_j \left(\hat{m}_j \rho \hat{m}_j^\dagger - \frac{1}{2} \{ \hat{m}_j^\dagger \hat{m}_j, \rho \} \right), \quad (69)$$

where $\rho(t)$ is the density matrix describing the state of the system at time t and $\hat{m}_j^{(\dagger)}$ are the Lindblad jump operators, while \mathcal{L} denotes the Lindbladian superoperator acting on ρ . The solution of Eq. (69) is, in general, a mixed state that can be thought of as an average over many pure-state quantum trajectories of a suitable stochastic process. Given a Lindbladian, there are many stochastic processes (or unravelings) that produce the same averaged density matrix [10]. Among them, in this paper we focus on the so-called quantum-jump unraveling [61], whose implementation is described below.

We define the non-Hermitian Hamiltonian

$$\hat{H}_{\text{eff}} = \hat{H} - \frac{i\gamma}{2} \sum_j \hat{m}_j^\dagger \hat{m}_j, \quad (70)$$

then we rewrite Eq. (69)

$$\dot{\rho}(t) = -i[\hat{H}_{\text{eff}}, \rho(t)] - \gamma \sum_j \hat{m}_j \rho \hat{m}_j^\dagger. \quad (71)$$

In this form, the Lindblad dynamics ruled by Eq. (69) can be interpreted as a deterministic non-Hermitian time evolution of a pure quantum state $|\psi_t\rangle$ generated by \hat{H}_{eff} , plus a stochastic part given by the possibility of applying \hat{m}_j to the state $|\psi_t\rangle$ [cf. the second term in the right-hand-side of Eq. (71)]. For \hat{m}_j Hermitian, this dynamics can be thought of as an occasional, yet abrupt, measurement process in which, at any time, the system has a chance to be POVM-measured [7, 8], i.e., to undergo the action of one of the operators \hat{m}_j , taken from a distribution (see more details below).

In this framework, Eq. (69) can be solved by implementing the following stochastic process. The time evolution is discretized in time slices of step dt . Then, at each step:

1. With probability $\pi_j = \gamma \langle \hat{m}_j^\dagger \hat{m}_j \rangle dt$, the jump operator is applied:

$$|\psi_{t+dt}\rangle \mapsto \frac{\hat{m}_j |\psi_t\rangle}{\|\hat{m}_j |\psi_t\rangle\|}; \quad (72)$$

2. With probability $\pi = 1 - \sum_j \pi_j$ the non-Hermitian evolution occurs:

$$|\psi_{t+dt}\rangle \mapsto \frac{e^{-i\hat{H}_{\text{eff}} dt} |\psi_t\rangle}{\|e^{-i\hat{H}_{\text{eff}} dt} |\psi_t\rangle\|}. \quad (73)$$

Note that the Lindblad master equation in Eq. (69) is invariant under a shift of the jump operators $\hat{m}_j \mapsto \hat{1} + \hat{m}_j$. In practice, one can implement the protocol described above with the Lindblad operators $\hat{1} + \hat{m}_j$ (as we actually do in Sec. 5), instead of \hat{m}_j , and obtain the same averaged density matrix, although the single trajectories $|\psi_t\rangle$ may behave differently.

We remark that, since the von Neumann entropy of Eq. (53) quantifies the amount of genuine quantum correlations of a pure state, we first evaluate it over the single pure-state quantum trajectory and then perform the ensemble averaging. It is important to fix the order of the two operations of measuring and ensemble averaging, since they are commuting only for linear functions of the state ρ (as for expectation values of observables), but not for quantities as the one in Eq. (53). For a more detailed discussion see, e.g., Refs. [33, 38, 39].

C Evolution of the state: a different derivation

In this appendix we provide a different derivation of the analytic expression for the evolved state $|\psi\rangle_{\hat{M}_{\mathcal{I},\mathcal{J}}}$, by evaluating the operator $\hat{\mathcal{Z}}(1) = \frac{1}{2} \sum_{p,q} Z_{p,q} \hat{c}_p^\dagger \hat{c}_q^\dagger(1)$, in Eq. (16). To this purpose we notice that

$$[\hat{A}_{\mathcal{I},\mathcal{J}}, \hat{\mathcal{Z}}] = \frac{1}{2} \sum_{p,q} \sum_{i \in \mathcal{I}} (Z_{p,\mathcal{J}} \delta_{q,i} + Z_{q,\mathcal{J}} \delta_{p,i}) \hat{c}_p^\dagger \hat{c}_q^\dagger, \quad (74)$$

where we have defined $\hat{\mathcal{Z}} = \frac{1}{2} \sum_{p,q} Z_{p,q} \hat{c}_p^\dagger \hat{c}_q^\dagger$ and $Z_{p,\mathcal{J}} = \sum_{j \in \mathcal{J}} Z_{p,j}$. Consequently $[\hat{\mathcal{Z}}, [\hat{M}_{\mathcal{I},\mathcal{J}}, \hat{\mathcal{Z}}]] = 0$. This allows to write

$$\hat{M}_{\mathcal{I},\mathcal{J}} e^{\hat{\mathcal{Z}}} = e^{\hat{\mathcal{Z}}} \left(\hat{M}_{\mathcal{I},\mathcal{J}} + [\hat{M}_{\mathcal{I},\mathcal{J}}, \hat{\mathcal{Z}}] \right), \quad (75)$$

that, together with $\hat{M}_{\mathcal{I},\mathcal{J}} |0\rangle = |0\rangle$, leads to

$$\hat{M}_{\mathcal{I},\mathcal{J}} |\psi\rangle_{\hat{M}_{\mathcal{I},\mathcal{J}}} = \mathcal{N} e^{\hat{\mathcal{Z}}} \left(\hat{1} + [\hat{A}_{\mathcal{I},\mathcal{J}}, \hat{\mathcal{Z}}] \right) |0\rangle. \quad (76)$$

Since $[\hat{A}_{\mathcal{I},\mathcal{J}}, \hat{\mathcal{Z}}]^2 = 0$, we can write $\exp\{[\hat{A}_{\mathcal{I},\mathcal{J}}, \hat{\mathcal{Z}}]\} = \hat{1} + [\hat{A}_{\mathcal{I},\mathcal{J}}, \hat{\mathcal{Z}}]$ and, therefore,

$$\hat{M}_{\mathcal{I},\mathcal{J}} |\psi\rangle_{\hat{M}_{\mathcal{I},\mathcal{J}}} = \mathcal{N} e^{\hat{\mathcal{Z}}} e^{[\hat{A}_{\mathcal{I},\mathcal{J}}, \hat{\mathcal{Z}}]} |0\rangle \equiv \mathcal{N} e^{\hat{\mathcal{Z}}'} |0\rangle, \quad (77)$$

being

$$\hat{\mathcal{Z}}' = \frac{1}{2} \sum_{i \in \mathcal{I}} \sum_{p,q} \left[Z_{p,q} + (Z_{p,\mathcal{J}} \delta_{q,i} + Z_{q,\mathcal{J}} \delta_{p,i}) \right] \hat{c}_p^\dagger \hat{c}_q^\dagger. \quad (78)$$

For the local case $\hat{M}_{\mathcal{I},\mathcal{J}}^1$ in Subsec 4.1, the jumped state $|\psi\rangle_{\hat{M}_{\mathcal{I},\mathcal{J}}^1}$ is described by the antisymmetric matrix $[Z^1]'$, with

$$[Z^1]_{p,q}' = (1 + \delta_{q,i} + \delta_{p,i}) Z_{p,q}. \quad (79)$$

On the other hand, for the non-local case $\hat{M}_{\mathcal{I},\mathcal{J}}^{\text{nl}}$ in Subsec. 4.2, the antisymmetric matrix $[Z^{\text{nl}}]'$ reads

$$[Z^{\text{nl}}]_{p,q}' = Z_{p,q} + (\delta_{p,i} + \delta_{p,i+r})(Z_{i,q} + Z_{i+r,q}). \quad (80)$$

References

- [1] L. Amico, R. Fazio, A. Osterloh and V. Vedral, *Entanglement in many-body systems*, Rev. Mod. Phys. **80**, 517 (2008), doi:[10.1103/RevModPhys.80.517](https://doi.org/10.1103/RevModPhys.80.517).
- [2] P. Calabrese and J. Cardy, *Entanglement entropy and conformal field theory*, J. Phys. A: Math. Theor. **42**, 504005 (2009), doi:[10.1088/1751-8113/42/50/504005](https://doi.org/10.1088/1751-8113/42/50/504005).
- [3] V. Alba and P. Calabrese, *Entanglement and thermodynamics after a quantum quench in integrable systems*, Proc. Natl. Acad. Sci. U.S.A. **114**, 7947 (2017), doi:[10.1073/pnas.1703516114](https://doi.org/10.1073/pnas.1703516114).
- [4] V. Alba and P. Calabrese, *Entanglement dynamics after quantum quenches in generic integrable systems*, SciPost Phys. **4**, 017 (2018), doi:[10.21468/SciPostPhys.4.3.017](https://doi.org/10.21468/SciPostPhys.4.3.017).
- [5] R. Singh, J. H. Bardarson and F. Pollmann, *Signatures of the many-body localization transition in the dynamics of entanglement and bipartite fluctuations*, New J. Phys. **18**(2), 023046 (2016), doi:[10.1088/1367-2630/18/2/023046](https://doi.org/10.1088/1367-2630/18/2/023046).
- [6] D. A. Abanin, E. Altman, I. Bloch and M. Serbyn, *Colloquium: Many-body localization, thermalization, and entanglement*, Rev. Mod. Phys. **91**, 021001 (2019), doi:[10.1103/RevModPhys.91.021001](https://doi.org/10.1103/RevModPhys.91.021001).
- [7] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information: 10th Anniversary Edition*, Cambridge University Press, New York, USA, doi:[10.1017/CBO9780511976667](https://doi.org/10.1017/CBO9780511976667) (2010).
- [8] G. Benenti, G. Casati, D. Rossini and G. Strini, *Principles of Quantum Computation and Information: A Comprehensive Textbook*, World Scientific, Singapore, doi:[10.1142/10909](https://doi.org/10.1142/10909) (2018).
- [9] H.-P. Breuer and F. Petruccione, *The theory of open quantum systems*, Oxford University Press, doi:[10.1093/acprof:oso/9780199213900.001.0001](https://doi.org/10.1093/acprof:oso/9780199213900.001.0001) (2002).
- [10] M. B. Plenio and P. L. Knight, *The quantum-jump approach to dissipative dynamics in quantum optics*, Rev. Mod. Phys. **70**, 101 (1998), doi:[10.1103/RevModPhys.70.101](https://doi.org/10.1103/RevModPhys.70.101).
- [11] A. J. Daley, *Quantum trajectories and open many-body quantum systems*, Adv. Phys. **63**(2), 77 (2014), doi:[10.1080/00018732.2014.933502](https://doi.org/10.1080/00018732.2014.933502).
- [12] A. Barchielli and M. Gregoratti, *Quantum trajectories and measurements in continuous time: the diffusive case*, vol. 782, Springer Berlin, Heidelberg, doi:[10.1007/978-3-642-01298-3](https://doi.org/10.1007/978-3-642-01298-3) (2009).
- [13] H. M. Wiseman and G. J. Milburn, *Quantum measurement and control*, Cambridge University Press, doi:[10.1017/CBO9780511813948](https://doi.org/10.1017/CBO9780511813948) (2009).
- [14] Y. Li, X. Chen and M. P. A. Fisher, *Quantum zeno effect and the many-body entanglement transition*, Phys. Rev. B **98**, 205136 (2018), doi:[10.1103/PhysRevB.98.205136](https://doi.org/10.1103/PhysRevB.98.205136).

- [15] Y. Li, X. Chen and M. P. A. Fisher, *Measurement-driven entanglement transition in hybrid quantum circuits*, Phys. Rev. B **100**, 134306 (2019), doi:[10.1103/PhysRevB.100.134306](https://doi.org/10.1103/PhysRevB.100.134306).
- [16] A. Chan, R. M. Nandkishore, M. Pretko and G. Smith, *Unitary-projective entanglement dynamics*, Phys. Rev. B **99**, 224307 (2019), doi:[10.1103/PhysRevB.99.224307](https://doi.org/10.1103/PhysRevB.99.224307).
- [17] B. Skinner, J. Ruhman and A. Nahum, *Measurement-induced phase transitions in the dynamics of entanglement*, Phys. Rev. X **9**, 031009 (2019), doi:[10.1103/PhysRevX.9.031009](https://doi.org/10.1103/PhysRevX.9.031009).
- [18] M. Szyniszewski, A. Romito and H. Schomerus, *Entanglement transition from variable-strength weak measurements*, Phys. Rev. B **100**, 064204 (2019), doi:[10.1103/PhysRevB.100.064204](https://doi.org/10.1103/PhysRevB.100.064204).
- [19] M. J. Gullans and D. A. Huse, *Scalable probes of measurement-induced criticality*, Phys. Rev. Lett. **125**, 070606 (2020), doi:[10.1103/PhysRevLett.125.070606](https://doi.org/10.1103/PhysRevLett.125.070606).
- [20] C.-M. Jian, Y.-Z. You, R. Vasseur and A. W. W. Ludwig, *Measurement-induced criticality in random quantum circuits*, Phys. Rev. B **101**, 104302 (2020), doi:[10.1103/PhysRevB.101.104302](https://doi.org/10.1103/PhysRevB.101.104302).
- [21] A. Zabalo, M. J. Gullans, J. H. Wilson, S. Gopalakrishnan, D. A. Huse and J. H. Pixley, *Critical properties of the measurement-induced transition in random quantum circuits*, Phys. Rev. B **101**, 060301 (2020), doi:[10.1103/PhysRevB.101.060301](https://doi.org/10.1103/PhysRevB.101.060301).
- [22] A. C. Potter and R. Vasseur, *Entanglement dynamics in hybrid quantum circuits, in "Entanglement in Spin Chains. Quantum Science and Technology"*, Springer, Cham, doi:[10.1007/978-3-031-03998-0_9](https://doi.org/10.1007/978-3-031-03998-0_9) (2022).
- [23] X. Chen, Y. Li, M. P. A. Fisher and A. Lucas, *Emergent conformal symmetry in nonunitary random dynamics of free fermions*, Phys. Rev. Res. **2**(3), 033017 (2020), doi:[10.1103/physrevresearch.2.033017](https://doi.org/10.1103/physrevresearch.2.033017).
- [24] Y. Bao, S. Choi and E. Altman, *Symmetry enriched phases of quantum circuits*, Ann. Phys. **435**, 168618 (2021), doi:[10.1016/j.aop.2021.168618](https://doi.org/10.1016/j.aop.2021.168618).
- [25] A. Nahum and B. Skinner, *Entanglement and dynamics of diffusion-annihilation processes with majorana defects*, Phys. Rev. Res. **2**, 023288 (2020), doi:[10.1103/PhysRevResearch.2.023288](https://doi.org/10.1103/PhysRevResearch.2.023288).
- [26] M. Buchhold, Y. Minoguchi, A. Altland and S. Diehl, *Effective theory for the measurement-induced phase transition of dirac fermions*, Phys. Rev. X **11**, 041004 (2021), doi:[10.1103/PhysRevX.11.041004](https://doi.org/10.1103/PhysRevX.11.041004).
- [27] O. Lunt and A. Pal, *Measurement-induced entanglement transitions in many-body localized systems*, Phys. Rev. Res. **2**, 043072 (2020), doi:[10.1103/PhysRevResearch.2.043072](https://doi.org/10.1103/PhysRevResearch.2.043072).
- [28] Q. Tang and W. Zhu, *Measurement-induced phase transition: A case study in the nonintegrable model by density-matrix renormalization group calculations*, Phys. Rev. Res. **2**, 013022 (2020), doi:[10.1103/PhysRevResearch.2.013022](https://doi.org/10.1103/PhysRevResearch.2.013022).
- [29] D. Rossini and E. Vicari, *Measurement-induced dynamics of many-body systems at quantum criticality*, Phys. Rev. B **102**, 035119 (2020), doi:[10.1103/PhysRevB.102.035119](https://doi.org/10.1103/PhysRevB.102.035119).

- [30] P. Sierant, G. Chiriacò, F. M. Surace, S. Sharma, X. Turkeshi, M. Dalmonte, R. Fazio and G. Pagano, *Dissipative floquet dynamics: from steady state to measurement induced criticality in trapped-ion chains*, Quantum **6**, 638 (2022), doi:[10.22331/q-2022-02-02-638](https://doi.org/10.22331/q-2022-02-02-638).
- [31] M. Ippoliti, M. J. Gullans, S. Gopalakrishnan, D. A. Huse and V. Khemani, *Entanglement phase transitions in measurement-only dynamics*, Phys. Rev. X **11**, 011030 (2021), doi:[10.1103/PhysRevX.11.011030](https://doi.org/10.1103/PhysRevX.11.011030).
- [32] A. Sriram, T. Rakovszky, V. Khemani and M. Ippoliti, *Topology, criticality, and dynamically generated qubits in a stochastic measurement-only kitaev model*, doi:[10.48550/arXiv.2207.07096](https://doi.org/10.48550/arXiv.2207.07096) (2022), [2207.07096](https://arxiv.org/abs/2207.07096).
- [33] X. Cao, A. Tilloy and A. D. Luca, *Entanglement in a fermion chain under continuous monitoring*, SciPost Phys. **7**, 24 (2019), doi:[10.21468/SciPostPhys.7.2.024](https://doi.org/10.21468/SciPostPhys.7.2.024).
- [34] N. Lang and H. P. Büchler, *Entanglement transition in the projective transverse field ising model*, Phys. Rev. B **102**, 094204 (2020), doi:[10.1103/PhysRevB.102.094204](https://doi.org/10.1103/PhysRevB.102.094204).
- [35] O. Alberton, M. Buchhold and S. Diehl, *Entanglement transition in a monitored free-fermion chain: From extended criticality to area law*, Phys. Rev. Lett. **126**(17), 170602 (2021), doi:[10.1103/physrevlett.126.170602](https://doi.org/10.1103/physrevlett.126.170602).
- [36] X. Turkeshi, A. Biella, R. Fazio, M. Dalmonte and M. Schirò, *Measurement-induced entanglement transitions in the quantum ising chain: From infinite to zero clicks*, Phys. Rev. B **103**(22), 224210 (2021), doi:[10.1103/physrevb.103.224210](https://doi.org/10.1103/physrevb.103.224210).
- [37] X. Turkeshi, M. Dalmonte, R. Fazio and M. Schirò, *Entanglement transitions from stochastic resetting of non-hermitian quasiparticles*, Phys. Rev. B **105**, L241114 (2021), doi:[10.1103/PhysRevB.105.L241114](https://doi.org/10.1103/PhysRevB.105.L241114).
- [38] G. Piccitto, A. Russomanno and D. Rossini, *Entanglement transitions in the quantum ising chain: A comparison between different unravelings of the same lindbladian*, Phys. Rev. B **105**, 064305 (2022), doi:[10.1103/PhysRevB.105.064305](https://doi.org/10.1103/PhysRevB.105.064305).
- [39] G. Piccitto, A. Russomanno and D. Rossini, *Erratum: Entanglement transitions in the quantum ising chain: A comparison between different unravelings of the same lindbladian*, Phys. Rev. B **106**, 219901(E) (2022), doi:[10.1103/PhysRevB.106.219901](https://doi.org/10.1103/PhysRevB.106.219901).
- [40] M. Coppola, E. Tirrito, D. Karevski and M. Collura, *Growth of entanglement entropy under local projective measurements*, Phys. Rev. B **105**, 094303 (2022), doi:[10.1103/PhysRevB.105.094303](https://doi.org/10.1103/PhysRevB.105.094303).
- [41] E. Tirrito, A. Santini, R. Fazio and M. Collura, *Full counting statistics as probe of measurement-induced transitions in the quantum ising chain*, doi:[10.48550/arXiv.2212.09405](https://doi.org/10.48550/arXiv.2212.09405) (2022), [2212.09405](https://arxiv.org/abs/2212.09405).
- [42] T. Minato, K. Sugimoto, T. Kuwahara and K. Saito, *Fate of measurement-induced phase transition in long-range interactions*, Phys. Rev. Lett. **128**, 010603 (2022), doi:[10.1103/PhysRevLett.128.010603](https://doi.org/10.1103/PhysRevLett.128.010603).
- [43] T. Müller, S. Diehl and M. Buchhold, *Measurement-induced dark state phase transitions in long-ranged fermion systems*, Phys. Rev. Lett. **128**, 010605 (2022), doi:[10.1103/PhysRevLett.128.010605](https://doi.org/10.1103/PhysRevLett.128.010605).

- [44] C. Zerba and A. Silva, *Measurement phase transitions in the no-click limit as quantum phase transitions of a non-hermitean vacuum*, doi:[10.48550/arXiv.2301.07383](https://doi.org/10.48550/arXiv.2301.07383) (2023), [2301.07383](https://arxiv.org/abs/2301.07383).
- [45] A. Paviglianiti and A. Silva, *Multipartite entanglement in the measurement-induced phase transition of the quantum ising chain*, doi:[10.48550/arXiv.2302.06477](https://doi.org/10.48550/arXiv.2302.06477) (2023), [2302.06477](https://arxiv.org/abs/2302.06477).
- [46] S. L. Braunstein and P. van Loock, *Quantum information with continuous variables*, Rev. Mod. Phys. **77**, 513 (2005), doi:[10.1103/RevModPhys.77.513](https://doi.org/10.1103/RevModPhys.77.513).
- [47] C. Weedbrook, S. Pirandola, R. García-Patrón, N. J. Cerf, T. C. Ralph, J. H. Shapiro and S. Lloyd, *Gaussian quantum information*, Rev. Mod. Phys. **84**, 621 (2012), doi:[10.1103/RevModPhys.84.621](https://doi.org/10.1103/RevModPhys.84.621).
- [48] S. Bravyi, *Lagrangian representation for fermionic linear optics*, Quantum Inf. Comput. **5**, 216 (2005), doi:[10.5555/2011637.2011640](https://doi.org/10.5555/2011637.2011640).
- [49] D. P. DiVincenzo and B. M. Terhal, *Fermionic linear optics revisited*, Found. Phys. **35**, 1967 (2005), doi:[10.1007/s10701-005-8657-0](https://doi.org/10.1007/s10701-005-8657-0).
- [50] M.-w. Xiao, *Theory of transformation for the diagonalization of quadratic hamiltonians*, doi:[10.48550/arXiv.0908.0787](https://doi.org/10.48550/arXiv.0908.0787) (2009), [0908.0787](https://arxiv.org/abs/0908.0787).
- [51] M. Fagotti and P. Calabrese, *Evolution of entanglement entropy following a quantum quench: Analytic results for the xy chain in a transverse magnetic field*, Phys. Rev. A **78**(1), 010306 (2008), doi:[10.1103/physreva.78.010306](https://doi.org/10.1103/physreva.78.010306).
- [52] P. Calabrese and J. Cardy, *Evolution of entanglement entropy in 1d systems*, J. Stat. Mech. **2005**(04), P04010 (2005), doi:[10.1088/1742-5468/2005/04/p04010](https://doi.org/10.1088/1742-5468/2005/04/p04010).
- [53] M. Fava, L. Piroli, T. Swann, D. Bernard and A. Nahum, *Nonlinear sigma models for monitored dynamics of free fermions* (2023), [2302.12820](https://arxiv.org/abs/2302.12820).
- [54] M. Block, Y. Bao, S. Choi, E. Altman and N. Yao, *The measurement-induced transition in long-range interacting quantum circuits*, Phys. Rev. Lett. **128**, 010604 (2022), doi:[10.1103/PhysRevLett.128.010604](https://doi.org/10.1103/PhysRevLett.128.010604).
- [55] S. Sharma, X. Turkeshi, R. Fazio and M. Dalmonte, *Measurement-induced criticality in extended and long-range unitary circuits*, SciPost Physics Core **5**(2), 023 (2022), doi:[10.21468/scipostphyscore.5.2.023](https://doi.org/10.21468/scipostphyscore.5.2.023).
- [56] G. B. Mbeng, A. Russomanno and G. E. Santoro, *The quantum ising chain for beginners*, doi:[10.48550/arXiv.2009.09208](https://doi.org/10.48550/arXiv.2009.09208) (2020), [2009.09208](https://arxiv.org/abs/2009.09208).
- [57] S. Sachdev, *Quantum Phase Transitions*, Cambridge University Press, doi:[10.1017/CBO9780511973765](https://doi.org/10.1017/CBO9780511973765) (2011).
- [58] D. Rossini and E. Vicari, *Coherent and dissipative dynamics at quantum phase transitions*, Phys. Rep. **936**, 1 (2021), doi:[10.1016/j.physrep.2021.08.003](https://doi.org/10.1016/j.physrep.2021.08.003).
- [59] P. Pfeuty, *The one-dimensional ising model with a transverse field*, Ann. Phys. **57**, 79 (1970), doi:[10.1016/0003-4916\(70\)90270-8](https://doi.org/10.1016/0003-4916(70)90270-8).
- [60] E. Lieb, T. Schultz and D. Mattis, *Two soluble models of an antiferromagnetic chain*, Ann. Phys. **16**, 407 (1961), doi:[10.1016/0003-4916\(61\)90115-4](https://doi.org/10.1016/0003-4916(61)90115-4).

- [61] J. Dalibard, Y. Castin and K. Mølmer, *Wave-function approach to dissipative processes in quantum optics*, Phys. Rev. Lett. **68**, 580 (1992), doi:[10.1103/PhysRevLett.68.580](https://doi.org/10.1103/PhysRevLett.68.580).