

GRIFFIN: A C++ library for electroweak radiative corrections in fermion scattering and decay processes

Lisong Chen^{1,2} and Ayres Freitas¹

¹Pittsburgh Particle-physics Astro-physics & Cosmology Center (PITT-PACC), Department of Physics & Astronomy, University of Pittsburgh, Pittsburgh, PA 15260, USA

²Institut für Theoretische Teilchenphysik, Karlsruhe Institute of Technology (KIT), Wolfgang-Gaede Straße 1, 76128 Karlsruhe, Germany

Abstract

This paper describes a modular framework for the description of electroweak scattering and decay processes, including but not limited to Z-resonance physics. The framework consistently combines a complex-pole expansion near an s-channel resonance with a regular fixed-order perturbative description away from the resonance in a manifestly gauge-invariant scheme. Leading vertex correction contributions are encapsulated in form factors that can be predicted or treated as numerical fit parameters.

This framework has been implemented in the publicly available object-oriented C++ library GRIFFIN. Version 1.0 of this library provides Standard Model predictions for the IR-subtracted matrix elements for the process $f\bar{f} \rightarrow f'\bar{f}'$ with full NNLO and leading higher-order contributions on the Z-resonance, and with NLO corrections off-resonance. The library can straightforwardly be extended to include higher-order corrections, should they become available, or predictions for new physics models. It can be interfaced with Monte-Carlo programs to account for QED and QCD initial-state and final-state radiation.

1 Introduction

Studies of fermion scattering, $f\bar{f} \rightarrow f'\bar{f}'$ for center-of-mass energies near the Z-boson resonance, $\sqrt{s} \sim M_Z$, have played a crucial role in elucidating the structure of the Standard Model (SM) and putting constraints in potential new physics beyond the SM (BSM). These include precision measurements at LEP and SLC (where $f = e$) [1], as well as Drell-Yan production at the TeVatron and LHC (where $f = u, d, s, c, b$) [2–4]. Even higher levels of precision can be achieved at the high-luminosity run of the LHC (HL-LHC) [5] and one of several proposed future e^+e^- colliders: FCC-ee [6], CEPC [7], ILC [8, 9], CLIC [10, 11].

The relevant outcomes of these experiments are typically presented in terms of a set of so-called electroweak pseudo-observables (EWPOs) that encapsulate the dominant radiative

corrections in the SM and are most sensitive to BSM physics. Examples of EWPOs are effective Z-fermion couplings, partial Z-boson widths, the effective weak mixing angle $\sin^2 \theta_{\text{eff}}^f$, and the asymmetry parameters A_f ; see *e.g.* Ref. [1] for the definition of these quantities. For the full description of the observable cross-sections, however, a number of other ingredients besides the EWPOs are needed, such as contributions from diagrams without s-channel Z-bosons (*i.e.* s-channel photon exchange and box diagrams) and initial- and final-state QED and QCD radiation effects. A number of software packages provide these ingredients with complete next-to-leading order (NLO) and some partial higher-order corrections included [12–29]. Among these, the ZFITTER [14,15] and TOPAZ0 [13] packages also provide extensive formulae for real photon radiation contributions, including certain selection cuts. They have been widely used in experimental studies. One can refer to [30] for the most recent updates on such analytical methods. Alternatively, QED radiation can be simulated with Monte-Carlo (MC) methods. For example, the electroweak corrections provided by the package DIZET [12], which is a component of ZFITTER, have been linked to the MC programs KoralZ [31] and KKMC [32]¹

However, despite the tremendous success of these software tools, they may not be easily adaptable to future applications that require a higher level of precision. Such applications call for a setup that enables the incorporation of higher-order corrections (NNLO and beyond) in a well-controlled and manifestly gauge-invariant way, as well as a modular object-oriented structure for the straightforward implementation of new SM or BSM contributions. In this article, the new software package GRIFFIN (Gauge-invariant Resonance In Four-Fermion INteractions) is introduced, which aims to provide a framework with these desirable features. It is written in C++ and defines a class hierarchy that can be extended with new results (both in the SM and beyond) without modifying its interface to external users (such as MC or fitting programs). While its object-oriented structure is, in principle, general enough to implement any arbitrary physics process, the current version is focused on 4-fermion processes, *i.e.* scattering processes of the form $f\bar{f} \rightarrow f'\bar{f}'$ or decay processes like muon decay. The relevant matrix elements for these processes are implemented in IR-subtracted form, which can be interfaced with the MC program to treat QED and QCD radiation. To describe the Z-boson resonance, it uses a Laurent expansion of the hard matrix elements² about the complex pole $s_0 \equiv M_Z^2 - iM_Z\Gamma_Z$. Since this pole is an analytical property of the S-matrix, both the pole location and the coefficients of the expansion are individually gauge-invariant [34–39].

The paper is organized as follows. Section 2 introduces the formalism for the complex-pole expansion and discusses what building blocks are required to describe the Z resonance at NNLO precision. On the other hand, outside of the resonance region, no pole expansion is needed. In section 3, it is discussed how on- and off-resonance predictions can be consistently matched to obtain reliable results for $f\bar{f} \rightarrow f'\bar{f}'$ at any center-of-mass energy. The implementation of these elements within the GRIFFIN library is described in section 4, and

¹Similar functionality, without using the DIZET package, is also provided by a number of other MC tools. See Ref. [33] for a broader overview.

²Here “hard” refers to the matrix element without initial-state and final-state QED/QCD radiation since the former would produce a deformation of the resonance lineshape.

numerical results and comparisons with ZFITTER/DIZET are shown in section 5. Finally, a summary is provided in section 6.

2 Fermion pair production on the Z resonance

The matrix element for the process $f\bar{f} \rightarrow f'\bar{f}'$ can be decomposed into four different chirality structures, which here will be delineated according to their vector or axial-vector couplings:

$$\mathcal{M} = [M_{VV}\gamma^\mu \otimes \gamma_\mu - M_{VA}\gamma^\mu \otimes \gamma_\mu\gamma^5 - M_{AV}\gamma^\mu\gamma^5 \otimes \gamma_\mu + M_{AA}\gamma^\mu\gamma^5 \otimes \gamma_\mu\gamma^5], \quad (1)$$

where the \otimes stands for the outer product of two fermion chains. In terms of these quantities, the differential cross-section is given by

$$\frac{d\sigma}{d\cos\theta} = \frac{N_c}{32\pi s} |\mathcal{M}|^2 \quad (2)$$

$$\begin{aligned} &= \frac{N_c s}{32\pi} \left[(1 + c_\theta^2) (|M_{VV}|^2 + |M_{VA}|^2 + |M_{AV}|^2 + |M_{AA}|^2) \right. \\ &\quad + 4c_\theta \operatorname{Re}\{M_{VV}M_{AA}^* + M_{VA}M_{AV}^*\} \\ &\quad - 2P_f(1 + c_\theta^2) \operatorname{Re}\{M_{VV}M_{AV}^* + M_{VA}M_{AA}^*\} \\ &\quad \left. - 4P_f c_\theta \operatorname{Re}\{M_{VV}M_{VA}^* + M_{AV}M_{AA}^*\} \right], \quad (3) \end{aligned}$$

where $c_\theta = \cos\theta$, θ is the scattering angle in the center-of-mass frame, and s is the center-of-mass energy, P_f is the polarization of the incoming fermion f , and the masses of f and f' have been neglected. For Z-boson exchange at tree level, the four chiral matrix elements read

$$M_{VV}^{(0)} = \frac{v_{f(0)}^Z v_{f'(0)}^Z}{s - s_0}, \quad M_{VA}^{(0)} = \frac{v_{f(0)}^Z a_{f'(0)}^Z}{s - s_0}, \quad M_{AV}^{(0)} = \frac{a_{f(0)}^Z v_{f'(0)}^Z}{s - s_0}, \quad M_{AA}^{(0)} = \frac{a_{f(0)}^Z a_{f'(0)}^Z}{s - s_0}, \quad (4)$$

where $s_0 \equiv M_Z^2 - iM_Z\Gamma_Z$ and

$$v_{f(0)}^Z = \frac{eI_f^3(1 - 4|Q_f|s_w^2)}{2s_w c_w}, \quad a_{f(0)}^Z = \frac{eI_f^3}{2s_w c_w} \quad (5)$$

are the vector and axial-vector couplings of the Z-boson to the fermion f ($f = \ell, \nu, u, d, \dots$). Furthermore, s_w and c_w stand for the sine and cosine of weak-mixing angle.

Note that throughout this document, M_Z and Γ_Z refer to the mass and width of the Z-boson in the complex-pole scheme, which is theoretically well-defined and gauge-invariant [34–39]. However, most experimental measurements are typically reported in terms of the so-called running-width scheme, leading to different values for the mass and width, which we denote as M_Z^{exp} and Γ_Z^{exp} . The two definitions are related via [40]³

$$M_Z = M_Z^{\text{exp}} (1 + (\Gamma_Z^{\text{exp}}/M_Z^{\text{exp}})^2)^{-1/2}, \quad \Gamma_Z = \Gamma_Z^{\text{exp}} [1 + (\Gamma_Z^{\text{exp}}/M_Z^{\text{exp}})^2]^{-1/2}. \quad (6)$$

³See Ref. [41] for a general review of the treatment of electroweak gauge-boson resonances.

In general, when including photon-exchange diagrams and higher-order contributions, the matrix elements can be written as Laurent expansion about the complex pole s_0 ,

$$M_{ij} = \frac{R_{ij}}{s - s_0} + S_{ij} + (s - s_0)S'_{ij} + \dots \quad (i, j = V, A). \quad (7)$$

Note that the scattering angle θ is kept fixed when expanding $M_{ij}(s, \theta)$. To construct explicit expressions for R, S, S' , we introduce the following quantities:

$$Z_{Vf}(s) \equiv v_f^Z(s) + v_f^\gamma(s) \frac{\Sigma_{\gamma Z}(s)}{s + \Sigma_{\gamma\gamma}(s)}, \quad G_{Vf}(s) \equiv v_f^\gamma(s), \quad (8)$$

$$Z_{Af}(s) \equiv a_f^Z(s) + a_f^\gamma(s) \frac{\Sigma_{\gamma Z}(s)}{s + \Sigma_{\gamma\gamma}(s)}, \quad G_{Af}(s) \equiv a_f^\gamma(s), \quad (9)$$

$$\Sigma_Z(s) \equiv \Sigma_{ZZ}(s) - \frac{[\Sigma_{\gamma Z}(s)]^2}{s + \Sigma_{\gamma\gamma}(s)}. \quad (10)$$

Here v_f^V (a_f^V) is the vector (axial-vector) form factor for the vertex between the gauge boson V ($V=Z, \gamma$) and the fermion f , including loop contributions. $\Sigma_{V_1 V_2}$ is the self-energy for incoming V_1 and outgoing V_2 ($V_{1,2}=Z, \gamma$). Furthermore, we denote

$$B_{ij}(s, t) : \text{Contribution of } \gamma\gamma, ZZ \text{ and } WW \text{ box diagrams for initial-state} \quad (11) \\ \text{vector/axial-vector current } (i = V, A) \text{ and final-state vector/axial-} \\ \text{vector current } (j = V, A);$$

$$B_{\gamma Z, ij}(s, t) = \frac{B_{\gamma Z, ij}^R}{s - s_0} + B_{\gamma Z, ij}^S + (s - s_0)B_{\gamma Z, ij}^{S'} + \dots : \quad (12) \\ \text{Contribution of } \gamma Z \text{ box diagrams, which can also contribute to the} \\ \text{leading pole term } R_{ij}.$$

It should be noted that the coefficients $B_{\gamma Z, ij}^{R, S, S', \dots}$ contain additional logarithms $\ln(1 - \frac{s}{s_0})$ that become singular on the pole and need to be accounted for in the Laurent expansion. Up to one-loop order, one thus has $B_{\gamma Z, ij}^X = B_{\gamma Z, ij}^{X, 1} \ln(1 - \frac{s}{s_0}) + B_{\gamma Z, ij}^{X, 0}$ ($X = R, S, S', \dots$), where $B_{\gamma Z, ij}^{X, i}$ are independent of s . In practice, these contributions are calculated by expanding the full analytical expression for the one-loop γZ box diagrams while tracking both polynomial and logarithmic singularities.

In terms of the above quantities, the coefficients of the complex-pole expansion are read

$$R_{ij} = \left[\frac{Z_{if} Z_{jf'}}{1 + \Sigma'_Z} \right]_{s=s_0} + B_{\gamma Z, ij}^R, \quad (13)$$

$$S_{ij} = \left[\frac{Z_{if} Z'_{jf'} + Z'_{if} Z_{jf'}}{1 + \Sigma'_Z} - \frac{Z_{if} Z_{jf'} \Sigma''_Z}{2(1 + \Sigma'_Z)^2} + \frac{G_{if} G_{jf'}}{s + \Sigma_{\gamma\gamma}} + B_{ij} \right]_{s=s_0} + B_{\gamma Z, ij}^S, \quad (14)$$

$$S'_{ij} = \left[\frac{Z_{if} Z''_{jf'} + Z''_{if} Z_{jf'} + 2Z'_{if} Z'_{jf'}}{2(1 + \Sigma'_Z)} - \frac{(Z_{if} Z'_{jf'} + Z'_{if} Z_{jf'}) \Sigma''_Z + \frac{1}{3} Z_{if} Z_{jf'} \Sigma'''_Z}{2(1 + \Sigma'_Z)^2} + \frac{Z_{if} Z_{jf'} (\Sigma''_Z)^2}{4(1 + \Sigma'_Z)^3} \right. \\ \left. + \frac{G_{if} G'_{jf'} + G'_{if} G_{jf'}}{s + \Sigma_{\gamma\gamma}} - \frac{G_{if} G_{jf'} (1 + \Sigma'_{\gamma\gamma})}{(s + \Sigma_{\gamma\gamma})^2} + B'_{ij} \right]_{s=s_0} + B_{\gamma Z, ij}^{S'}, \quad (15)$$

Here X' denotes the derivative of X with respect to s .

The vertex form factors and box diagrams can contain infrared (IR) divergencies from QED and (in the case of external quarks) QCD corrections. When interfacing the matrix elements with a Monte-Carlo (MC) program, these IR divergent contributions and the corresponding real emission contributions will be produced by the MC phase-space generator and showering algorithm. Thus they must be excluded from the hard matrix elements encoded in GRIFFIN.

For the vertex form factors, the IR-divergent contributions can be factorized, $Z_{if}^{\text{tot}} = R_f^i \times Z_{if}$, where R_f^i ($i = V, A$) contain the QED/QCD corrections to the $f\bar{f}$ pair (see *e.g.* Ref. [42]). They are defined via the matrix elements for the decay of a vector boson into $f\bar{f}$:

$$R_f^V(s) \equiv \frac{\mathcal{M}_{V^* \rightarrow f\bar{f}}^{\text{QED/QCD}}}{\mathcal{M}_{V^* \rightarrow f\bar{f}}^{\text{Born}}}, \quad R_f^A(s) \equiv \frac{\mathcal{M}_{A^* \rightarrow f\bar{f}}^{\text{QED/QCD}}}{\mathcal{M}_{A^* \rightarrow f\bar{f}}^{\text{Born}}}, \quad (16)$$

where V^* (A^*) denotes a generic vector boson with invariant mass s that couples to the $f\bar{f}$ fermion current with a pure vector (axial-vector) coupling, and the superscript ‘‘QED/QCD’’ indicates that all QED and QCD to the desired order are included. The factorization is not perfect, but the remaining non-factorizable contributions [43, 44] are IR-finite and can be incorporated into Z_{if} order by order. Here and in the following, we adopt the notation that Z_{if} is the IR-finite vertex form factor after the IR-divergent QED/QCD contributions have been factored off, whereas Z_{if}^{tot} is the vertex form factor including all QED/QCD corrections.

The subtraction of these contributions is less straightforward for the box diagrams, which contain IR-divergent initial-final interference (IFI) terms. We here restrict ourselves to a discussion at NLO, where one encounters IR-divergent IFI terms from two sources, the $\gamma\gamma$ boxes, and the γZ boxes. Following the CEEX MC scheme of Ref. [45], they can be removed with the following subtraction terms:

$$\gamma\gamma \text{ box:} \quad B_{\text{VV}(1)} = B_{\text{VV}(1)}^{\text{tot}} - S_{\text{VV}}^{(0)} \frac{\alpha}{\pi} Q_f Q_{f'} f_{\text{IR}}(m_\gamma, t, u), \quad (17)$$

$$\gamma Z \text{ box:} \quad B_{\gamma Z, ij(1)} = B_{\gamma Z, ij(1)}^{\text{tot}} - \frac{R_{ij}^{(0)}}{s - s_0} \frac{\alpha}{\pi} Q_f Q_{f'} [f_{\text{IR}}(m_\gamma, t, u) + \delta_G(s, t, u)], \quad (18)$$

$$f_{\text{IR}}(m_\gamma, t, u) = \frac{2\pi}{\alpha} [R_{e(1)}(t) - R_{e(1)}(u)] = \ln\left(\frac{1 - c_\theta}{1 + c_\theta}\right) \left[\ln\left(\frac{2m_\gamma^2}{s\sqrt{1 - c_\theta^2}}\right) + \frac{1}{2} \right],$$

$$\delta_G(s, t, u) = -2 \ln\left(\frac{1 - c_\theta}{1 + c_\theta}\right) \ln\left(\frac{s_0 - s}{s_0}\right). \quad (19)$$

Here the subscripts (n) indicate the loop order. $R_{e(1)}$ is the radiative factor defined in eq. (16) with one-loop QED corrections (the upper indices for denoting vector/axial vector are suppressed here since the QED correction is chiral-blind). When matching our IR-subtracted results to an MC generator, the R factors can be implemented with any IR regularization scheme in the MC program since they are based on a physical process and thus scheme independent. Only for illustration, we show their form when using a small photon mass m_γ as a regulator. The current version of GRIFFIN uses this subtraction

scheme, but other schemes for removing the IR-divergent IFI contributions could also be easily implemented.

Near the Z resonance, when aiming for a description at $N^n\text{LO}$ precision, it is typically sufficient to compute only the leading coefficient R to n -loop order, whereas $(n-1)$ -loop and $(n-2)$ -loop precision is adequate for S and S' , respectively⁴.

Furthermore, the ratio $\Gamma_Z/M_Z = \mathcal{O}(\alpha)$, where $\mathcal{O}(\alpha)$ denotes electroweak NLO corrections, which implies that one can perform expansions in the perturbative order, α , and Γ_Z/M_Z in parallel. For example, $f(s_0) = f(M_Z^2) - iM_Z\Gamma_Z f'(M_Z^2) - \frac{M_Z^2\Gamma_Z^2}{2} f''(M_Z^2) + \dots$. Thus, in summary, we adopt the power counting $(s-s_0)/M_Z^2 \sim \Gamma_Z/M_Z \sim \alpha$ for the expansion of the matrix element near the Z pole.

If we wish to expand up to NNLO for the leading pole term, one would in principle, also need the γZ box to two-loop order, which is currently unknown. However, it was shown in Refs. [46–48] that at NLO the total contribution of IFI terms to R_{ij} vanishes when adding up the virtual γZ boxes and real photon radiation (see also Refs. [49, 50]). This argument, of course, only holds for sufficiently inclusive observables. Furthermore, for quarks in either the initial or final state, Ref. [51] demonstrated that the resonance pole also cancels in the IFI contributions for mixed electroweak-QCD NNLO corrections. A similar argument should apply to the $\gamma\gamma Z$ boxes at electroweak NNLO, although a more careful analysis of this issue would be desirable. Assuming that this argument holds, one only needs to include $B_{\gamma Z(m)}^R$, $m = 1, \dots, n-1$ for the computation of $R_{ij}^{(n)}$.

Based on the above considerations, the result for an expansion up to NNLO for the leading pole term R (which implies NLO precision for S and LO for S') reads

$$R_{ij}^{(0)} = Z_{if(0)}Z_{jf'(0)}, \quad (20)$$

$$R_{ij}^{(1)} = [Z_{if(0)}Z_{jf'(1)} + Z_{if(1)}Z_{jf'(0)} - Z_{if(0)}Z_{jf'(0)}\Sigma'_{Z(1)}]_{s=M_Z^2}, \quad (21)$$

$$R_{ij}^{(2)} = [Z_{if(0)}Z_{jf'(2)} + Z_{if(2)}Z_{jf'(0)} + Z_{if(1)}Z_{jf'(1)} - Z_{if(0)}Z_{jf'(0)}\Sigma'_{Z(2)} - \Sigma'_{Z(1)}R_{ij}^{(1)} - iM_Z\Gamma_Z(Z_{if(0)}Z'_{jf'(1)} + Z'_{if(1)}Z_{jf'(0)} - Z_{if(0)}Z_{jf'(0)}\Sigma''_{Z(1)})]_{s=M_Z^2} + B_{\gamma Z,ij(1)}^R, \quad (22)$$

$$S_{ij}^{(0)} = \frac{1}{M_Z^2}G_{if(0)}G_{jf'(0)}, \quad (23)$$

$$S_{ij}^{(1)} = \left[Z_{if(0)}Z'_{jf'(1)} + Z'_{if(1)}Z_{jf'(0)} - \frac{1}{2}Z_{if(0)}Z_{jf'(0)}\Sigma''_{Z(1)} + \frac{1}{M_Z^2}(G_{if(0)}G_{jf'(1)} + G_{if(1)}G_{jf'(0)}) + \frac{iM_Z\Gamma_Z - \Sigma_{\gamma\gamma(1)}}{M_Z^4}G_{if(0)}G_{jf'(0)} + B_{ij(1)} \right]_{s=M_Z^2} + B_{\gamma Z,ij(1)}^S, \quad (24)$$

$$S'_{ij}{}^{(0)} = -\frac{1}{M_Z^4}G_{if(0)}G_{jf'(0)}, \quad (25)$$

where the subscripts (n) again indicate the loop order.

⁴This power counting can be extended to more terms, beyond S' , in the Laurent expansion.

As mentioned in the introduction, electroweak pseudo-observables (EWPOs) are used as an intermediate step when comparing experimental data to theory expectations. The EWPOs can be expressed in terms of the form factors $F_{V,A}^f$ defined in Ref. [52]⁵ and in terms of the effective weak mixing angle $\sin^2 \theta_{\text{eff}}^f$ (as defined, *e.g.*, in Ref. [54]). Up to NNLO, and using the power counting $\alpha \sim \Gamma_Z/M_Z$, they are given by

$$\sin^2 \theta_{\text{eff}}^f = \frac{1}{4|Q_f|} \left[1 - \text{Re} \frac{Z_{Vf}}{Z_{Af}} \right]_{s=M_Z^2}, \quad (26)$$

$$F_A^f = \left[\frac{|Z_{Af}|^2}{1 + \text{Re} \Sigma'_Z} - \frac{1}{2} M_Z \Gamma_Z |a_{f(0)}^Z|^2 \text{Im} \Sigma''_Z \right]_{s=M_Z^2} + \mathcal{O}(\alpha^3), \quad (27)$$

$$F_V^f = \left[\frac{|Z_{Vf}|^2}{1 + \text{Re} \Sigma'_Z} - \frac{1}{2} M_Z \Gamma_Z |v_{f(0)}^Z|^2 \text{Im} \Sigma''_Z \right]_{s=M_Z^2} + \mathcal{O}(\alpha^3) \quad (28)$$

$$= F_A^f \left[(1 - 4|Q_f| \sin^2 \theta_{\text{eff}}^f)^2 + \left(\text{Im} \frac{Z_{Vf}}{Z_{Af}} \right)^2 \right] \quad (29)$$

For $f = \nu$ the effective weak mixing angle is ill-defined and irrelevant, and only F_A^ν is needed.

The matrix elements for the process $f\bar{f} \rightarrow f'f'$ can be expressed in terms of these form factors. In fact, they only enter the leading pole coefficient, R , as follows:

$$\begin{aligned} R_{ij}^{(0+1+2)} &= 4I_f^3 I_{f'}^3 \sqrt{F_A^f F_A^{f'}} \left[\tilde{Q}_i^f \tilde{Q}_j^{f'} (1 + i r_{AA}^I - \frac{1}{2} (r_{AA}^I)^2 + \frac{1}{2} \delta \bar{X}_{(2)}) \right. \\ &\quad \left. + (\tilde{Q}_i^f I_{j,f'} + \tilde{Q}_j^{f'} I_{i,f}) (i - r_{AA}^I) - I_{i,f} I_{j,f'} \right] \\ &\quad + M_Z \Gamma_Z Z_{if(0)} Z'_{jf'(0)} x_{ij}^I, \end{aligned} \quad (30)$$

where

$$\tilde{Q}_V^f = 1 - 4|Q_f| \sin^2 \theta_{\text{eff}}^f, \quad \tilde{Q}_A^f = 1, \quad (31)$$

$$I_{V,f} = \frac{1}{(a_{f(0)}^Z)^2} [a_{f(0)}^Z \text{Im} Z_{Vf(1)} - v_{f(0)}^Z \text{Im} Z_{Af(1)}], \quad I_{A,f} = 0, \quad (32)$$

$$\delta \bar{X}_{(2)} = -(\text{Im} \Sigma'_{Z(1)})^2 + 2 \frac{B_{\gamma Z, ij(1)}^R}{R_{ij}^{(0)}}, \quad (33)$$

$$r_{AA}^I = \frac{\text{Im} Z_{Af(1)}}{a_{f(0)}^Z} + \frac{\text{Im} Z_{Af'(1)}}{a_{f'(0)}^Z} - \text{Im} \Sigma'_{Z(1)}, \quad (34)$$

$$x_{ij}^I = \frac{\text{Im} Z'_{if(1)}}{Z_{if(0)}} + \frac{\text{Im} Z'_{jf'(1)}}{Z_{jf'(0)}} - \frac{1}{2} \text{Im} \Sigma''_{Z(1)}. \quad (35)$$

3 Combination of on- and off-resonance fermion-pair production

The Laurent series (7) is only a good approximation in a window of a few GeV about the Z resonance. For values of \sqrt{s} outside of this window, a non-expanded version of the matrix

⁵ F_A^f is related to ρ_f introduced in Ref. [53], up to a normalization factor.

element provides a more accurate description. A unified formulation that works for a wide range of center-of-mass energies near and far from the Z resonance is given by the following prescription:

$$M_{ij} = M_{ij}^{\text{exp},s_0} + M_{ij}^{\text{noexp}} - M_{ij}^{\text{exp},M_Z^2}, \quad (36)$$

where M_{ij}^{exp,s_0} is the matrix element expanded about the complex pole s_0 as in (7), and M_{ij}^{noexp} is the matrix element without any expansion in s and Dyson summation. In other words, it is a straightforward fixed-order matrix element for which the full NLO electroweak corrections are known (see *e.g.* Ref. [55,56]). To avoid double counting, the expanded version of the latter, $M_{ij}^{\text{exp},M_Z^2}$ must be subtracted. Since M_{ij}^{noexp} has a pole at $s = M_Z^2$, the expansion for $M_{ij}^{\text{exp},M_Z^2}$ must be performed about that point⁶:

$$M_{ij}^{\text{exp},M_Z^2} = \frac{\overline{R}'_{ij}}{(s - M_Z^2)^2} + \frac{\overline{R}_{ij}}{s - M_Z^2} + \overline{S}_{ij} + (s - M_Z^2)\overline{S}'_{ij} + \dots \quad (37)$$

Up to NLO, the coefficients are given by

$$\overline{R}'_{ij}{}^{(0)} = 0, \quad \overline{R}'_{ij}{}^{(1)} = -Z_{if(0)}Z_{jf'(0)}\Sigma_{Z(1)}\Big|_{s=M_Z^2}, \quad (38)$$

$$\overline{R}_{ij}{}^{(0)} = Z_{if(0)}Z_{jf'(0)}, \quad (39)$$

$$\overline{R}_{ij}{}^{(1)} = [Z_{if(0)}Z_{jf'(1)} + Z_{if(1)}Z_{jf'(0)} - Z_{if(0)}Z_{jf'(0)}\Sigma'_{Z(1)}]_{s=M_Z^2} + B_{\gamma Z,ij(1)}^{\overline{R}}, \quad (40)$$

$$\overline{S}_{ij}{}^{(0)} = \frac{1}{M_Z^2}G_{if(0)}G_{jf'(0)}, \quad (41)$$

$$\begin{aligned} \overline{S}_{ij}{}^{(1)} = & \left[Z_{if(0)}Z'_{jf'(1)} + Z'_{if(1)}Z_{jf'(0)} - \frac{1}{2}Z_{if(0)}Z_{jf'(0)}\Sigma''_{Z(1)} + \frac{1}{M_Z^2}(G_{if(0)}G_{jf'(1)} + G_{if(1)}G_{jf'(0)}) \right. \\ & \left. - \frac{\Sigma_{\gamma\gamma(1)}}{M_Z^4}G_{if(0)}G_{jf'(0)} + B_{ij(1)} \right]_{s=M_Z^2} + B_{\gamma Z,ij(1)}^{\overline{S}}, \end{aligned} \quad (42)$$

$$\overline{S}'_{ij}{}^{(0)} = -\frac{1}{M_Z^4}G_{if(0)}G_{jf'(0)}, \quad (43)$$

Note the presence of the double-pole term \overline{R}' , which is purely imaginary and does not exist for the complex-pole expansion. Similar to eq. (12), the γZ box diagrams also contribute to

⁶ M_{ij}^{exp,s_0} contains expansion terms $(s - s_0)^i$ for $i \leq 1$, whereas an expansion of $M_{ij}^{\text{noexp}} - M_{ij}^{\text{exp},M_Z^2}$ would have terms $(s - M_Z^2)^i$ with $i > 1$. Since the two series have different expansion points, the match between them is not perfect, but the mismatch is of order $\mathcal{O}(\Gamma_Z^2/M_Z^2)$ or $\mathcal{O}(\alpha\Gamma_Z/M_Z)$, which is beyond the level of accuracy of our results.

We also want to point out that our matching scheme for combining resonant and off-resonant regions is not unique. One could, for instance, use the complex-mass scheme [57, 58] to calculate the off-resonant matrix elements, but a more careful investigation of this would be needed. At NLO our prescriptions is equivalent to the one in Ref. [20].

the single-pole term \bar{R} ,

$$B_{\gamma Z,ij}(s,t) = \frac{B_{\gamma Z,ij}^{\bar{R}}}{s - M_Z^2} + B_{\gamma Z,ij}^{\bar{S}} + \dots \quad (44)$$

The difference $M_{ij}^{\text{noexp}} - M_{ij}^{\text{exp},M_Z^2}$ is free of any poles at $s = M_Z^2$ and in fact it vanishes in the limit $s \rightarrow M_Z^2$. All three terms in eq. (36) are separately finite and gauge-invariant. With currently available results, M_{ij}^{exp,s_0} can be evaluated to NNLO order near the Z pole, as described in the previous section. The current state of the art for M_{ij}^{noexp} is NLO, so that, for consistency, $M_{ij}^{\text{exp},M_Z^2}$ should also be computed to NLO.

Both the matrix element coefficients (30) and (23)–(25), as well as the complete matrix element (36) are implemented in the GRIFFIN library.

4 Structure of the GRIFFIN library

The GRIFFIN package provides a framework for a hierarchy of C++ classes to compute in principle, any electroweak observable or pseudo-observable within a given model. The current version implements SM predictions for EWPOs and matrix elements for the process $f\bar{f} \rightarrow f'\bar{f}'$, with $f \neq f'$. Still, it is straightforward to include other items as well, including but not limited to:

- matrix elements for $f\bar{f} \rightarrow f\bar{f}$, with the same fermion type in the initial and final state, which includes Bhabha scattering;
- matrix elements for radiation of additional photons ($f\bar{f} \rightarrow f'\bar{f}'\gamma$) and/or fermion pairs ($f\bar{f} \rightarrow f'\bar{f}'f''\bar{f}''$), with appropriate subtraction of IR singularities (which is broadly equivalent to the concept of “electroweak pseudo-parameters” (EWPP) in section C.3 of Ref. [59]);
- predictions for EWPOs in BSM theories or in terms of effective theory extensions of the SM with higher-dimensional operators;

The library contains two base classes:

- class `inval`, which contains user-provided input parameters for a given model (such as the SM or some extension thereof);
- class `psobs`, which returns a numerical prediction for an observable or pseudo-observable, for the input parameters provided by an `inval` object.

In its basic form, `inval` simply has some basic methods for setting and retrieving the values of some input parameters. However, one can define extended classes derived from `inval` to perform computations of input parameters, such as translating between masses in the complex-pole scheme and the running-width scheme, see eq. (6), or computing the W-boson mass from the Fermi constant [60].

Boson masses and widths		Fermion masses			Couplings
M_W	Γ_W	m_e^{OS}	$m_d^{\overline{\text{MS}}}(M_Z)$	$m_u^{\overline{\text{MS}}}(M_Z)$	$\alpha(0)$
M_Z	Γ_Z	m_μ^{OS}	$m_s^{\overline{\text{MS}}}(M_Z)$	$m_c^{\overline{\text{MS}}}(M_Z)$	$\Delta\alpha \equiv 1 - \alpha(0)/\alpha(M_Z^2)$
M_H		m_τ^{OS}	$m_b^{\overline{\text{MS}}}(M_Z)$	m_t^{OS}	$\alpha_s^{\overline{\text{MS}}}(M_Z)$
					G_μ

Table 1: SM input parameters used in GRIFFIN. Here OS and $\overline{\text{MS}}$ refer to the on-shell and $\overline{\text{MS}}$ scheme, respectively. For most quantities currently encoded in GRIFFIN, fermion masses besides the top-quark mass are being ignored. The CKM matrix is taken to be the unit matrix. $\alpha(0)$ refers to the electromagnetic coupling in the Thomson limit, and G_μ is the Fermi constant of muon decay.

The base version of GRIFFIN defines a set of input parameters for SM calculations, listed in Tab. 1. Most of these parameters are defined within the on-shell (OS) renormalization scheme, with the exception of light quark masses and the strong coupling, for which the $\overline{\text{MS}}$ scheme is assumed (at the scale $\mu = M_Z$). Additional input parameters for flavor physics or BSM scenarios can be easily added.

The user has the option to choose between input classes that either use $\alpha(0)$, M_W , M_Z or $\alpha(0)$, G_μ , M_Z as inputs to define the electroweak couplings. Here $\alpha(0)$ is the electromagnetic coupling in the Thomson limit, and G_μ is the Fermi constant of muon decay. An additional input is the shift $\Delta\alpha$ between the running electromagnetic couplings at the scales $q^2 = 0$ and $q^2 = M_Z^2$. $\Delta\alpha$ receives contributions from leptons, which has been computed to four-loop order [61], and from quarks or hadrons, which can be extracted from data [62–64].

A child class descending from `psobs` can in principle encode predictions for any observable or pseudo-observable within any given model. The base version of GRIFFIN includes SM predictions for form factors, such as $\sin^2 \theta_{\text{eff}}^f$ and $F_{V,A}^f$, and for matrix elements for the process $f\bar{f} \rightarrow f'\bar{f}'$ near the Z resonance, using the complex pole expansion described in the previous section.

GRIFFIN version 1.0 contains the following SM corrections:

- Complete one-loop corrections for $\sin^2 \theta_{\text{eff}}^f$ [53,65] are implemented in the class `SW_SMNLO`. On top of this, electroweak [66–71] and mixed electroweak-QCD [72–76] two-loop corrections, as well as partial higher-order corrections are available in the class `SW_SMNNLO`. The latter include $\mathcal{O}(\alpha_t \alpha_s^2)$ [77,78], $\mathcal{O}(\alpha_t^2 \alpha_s)$, $\mathcal{O}(\alpha_t^3)$ [79,80] and $\mathcal{O}(\alpha_t \alpha_s^3)$ [81–83] corrections in the limit of a large top Yukawa coupling y_t , where $\alpha_t \equiv y_t^2/(4\pi)$, and leading fermionic three-loop corrections of orders $\mathcal{O}(\alpha^3)$ and $\mathcal{O}(\alpha^2 \alpha_s)$ [84,85]. In addition, non-factorizable $\mathcal{O}(\alpha \alpha_s)$ $Zq\bar{q}$ vertex contributions [43,86–90] are also implemented in `SW_SMNNLO`.
- Similarly, the classes `FA_SMNLO` and `FV_SMNLO` provide one-loop corrections [53] for the form factors $F_{V,A}^f$, whereas `FA_SMNNLO` and `FV_SMNNLO` contain electroweak [52,91–93]

and mixed electroweak-QCD [72–76] two-loop corrections, as well as the partial higher-order corrections and non-factorizable contributions mentioned in the previous bullet point.

- For the process $f\bar{f} \rightarrow f'\bar{f}'$: The class `mat_SMNNLO` computes the matrix element according to section 3 with the following ingredients:
 - All contributions needed to compute the matrix element coefficient R to NNLO accuracy according to (30), and the coefficients S and S' to NLO and LO, respectively, see eqs. (23)–(25). These are also separately available in the member functions `coeffR`, `coeffS`, `coeffSp` of `mat_SMNNLO`.
 - The off-resonance contribution, $M_{ij}^{\text{noexp}} - M_{ij}^{\text{exp}, M_Z^2}$, to NLO precision, see section 3, which is also separately available via the member function `resoffZ`.
- When using the input parameter set $\alpha(0), G_\mu, M_Z$, one needs to compute M_W from these quantities according to

$$G_\mu = \frac{\pi\alpha}{\sqrt{2}M_W^2(1 - M_W^2/M_Z^2)}(1 + \Delta r). \quad (45)$$

Here Δr accounts for radiative corrections. The class `dr_SMNNLO` contains all higher-order corrections discussed in Ref. [60], plus the leading fermionic three-loop corrections of orders $\mathcal{O}(\alpha^3)$ and $\mathcal{O}(\alpha^2\alpha_s)$ [84, 85]. These corrections are used in the input classes `invalGmu` and `SMvalGmu`.

For any of these quantities, QED and QCD corrections on the external legs have been factored out, as explained in detail in section 2. The logic is that QED/QCD effects depend on detector acceptance and selection cuts and are best simulated with MC methods. GRIFFIN could be interfaced with suitable MC tools to provide the hard electroweak matrix elements.

5 Sample results and comparisons

In this section, we show numerical comparisons between GRIFFIN and the DIZET library of EW radiative corrections [14, 15, 94] for the EWPOs and the differential cross-section. For the latter, we use some of the computational frameworks of the KKMCEE project⁷ [95].

We first perform a benchmark test of the EWPOs in comparison with DIZET v 6.45 [94], including NNLO and leading NNNLO corrections. In DIZET, the form factor is defined as in Eq. 2.4.9 and Eq. 2.4.10 of Ref. [14].

$$\Gamma_{Z \rightarrow f\bar{f}} = \Gamma_0 c_f |\rho_Z^f| (|g_Z^f|^2 R_V^f + R_A^f) + \delta_{\alpha\alpha_s}, \quad (46)$$

where we have neglected all lepton masses, $c_f = N_c^f$ is the number of colors, and

$$\Gamma_0 = \frac{G_\mu M_Z^3}{24\sqrt{2}\pi}. \quad (47)$$

⁷The authors are grateful to S. Jadach for sharing a suitable test program with us.

In eq. (46), g_Z^f is a complex-valued variables, which in our notation from section 2 is given by

$$g_Z^f = \frac{Z_{Vf}^f}{Z_{Af}^f}, \quad (48)$$

On the other hand, in GRIFFIN, we define the partial width of Z-boson as

$$\Gamma_{Z \rightarrow f\bar{f}} = \frac{N_c^f M_Z}{12\pi} (F_V^f R_V^f + F_A^f R_A^f). \quad (49)$$

By setting this equal to eq. (46) we obtain the relation

$$\frac{N_c^f M_Z}{12\pi} F_A^f \left(\frac{F_V^f}{F_A^f} R_V^f + R_A^f \right) = \frac{N_c^f M_Z}{12\pi} F_A^f (|g_Z^f|^2 R_V^f + R_A^f) = \Gamma_0 c_f |\rho_Z^f| (|g_Z^f|^2 R_V^f + R_A^f), \quad (50)$$

This implies the following relation between $|\rho_Z^f|$ and F_A^f :

$$|\rho_Z^f| = \frac{2\sqrt{2}F_A^f}{G_\mu M_Z^2} \quad (51)$$

One should notice that the non-factorizable mixed QCD-EW corrections are considered as an additive part to the Z widths in DIZET, whereas in GRIFFIN, they are absorbed in the form factors $F_{V,A}$. This will cause a small numerical mismatch when comparing the form factor F_A to ρ_Z^f . Besides, one also has to notice that eq. 51 is the modulus of ρ_Z^f instead of $\text{Re } \rho_Z^f$. Hence to compare these observables, we need to use both the $\text{Im } \rho_Z^f$ and $\text{Re } \rho_Z^f$ output from DIZET to reconstruct $|\rho_Z^f|$.

The flags used by DIZET v.6.45 are listed as follows:

IHVP=5	IAMT4=8	IQCD=3	IMOMS=1	IMASS=0
ISCRE=0	IALEM=0	IMASK=0	ISCAL=0	IBARB=2
IFTJR=1	IFACR=2	IFACT=0	IHIGS=0	IAFMT=3
IEWLC=0	ICZAK=1	IHIGS=1	IALE2=3	IGFER=2
IDDZZ=1	IAMW2=1	ISFSR=1	IDMWW=0	IDSWW=0

Due to the limited options of EW input schemes offered by subroutines in DIZET, we have to set G_μ , M_Z as inputs and use the DIZET outputs for M_W and $\Gamma_{W,Z}$ as inputs for GRIFFIN, as shown in the following table:

GRIFFIN input parameters	
DIZET input parameters	DIZET output
$\alpha_s(M_Z^2) = 0.118$, $\alpha = 1/137.035999084$	$\Gamma_Z = 2.495890$ GeV
$\Delta\alpha = 0.059$, $M_Z = 91.1876$ GeV, $G_\mu = 1.166137 \times 10^{-5}$	$M_W = 80.3599$ GeV
$m_t = 173.0$ GeV, $M_H = 125.0$ GeV, $m_{e,\mu,\tau,u,d,s,c,b} = 0$ GeV	$\Gamma_W = 2.090095$ GeV

	DIZET 6.45	GRIFFIN all orders	GRIFFIN $\mathcal{O}(\alpha, \alpha^2, \alpha_t \alpha_s, \alpha_t \alpha_s^2)$
Δr	3.63947×10^{-2}	3.68836×10^{-2}	3.63987×10^{-2}

	$ \rho_Z^f $		$\sin^2 \theta_{\text{eff}}^f$		$\Gamma_{Z \rightarrow f\bar{f}}$	
	DIZET 6.45	GRIFFIN	DIZET 6.45	GRIFFIN	DIZET 6.45	GRIFFIN
$\nu\bar{\nu}$	1.00800	1.00814	0.231119	NAN	0.167206	0.167197
$\ell\bar{\ell}$	1.00510	1.00519	0.231500	0.231534	0.083986	0.083975
$u\bar{u}$	1.00578	1.00573	0.231393	0.231420	0.299938	0.299958
$d\bar{d}$	1.00675	1.00651	0.231266	0.231309	0.382877	0.382846
$b\bar{b}$	0.99692	0.99420	0.232737	0.23292	0.376853	0.377432

Table 2: The numerical comparison of the EWPOs and form factors ρ between DIZET and GRIFFIN. The partial width results are for a single fermion family. See text for details.

With these inputs, we find the numerical results for the form factors Δr , $|\rho_Z^f|$ and $\sin^2 \theta_{\text{eff}}^f$, as well as the partial Z width $\Gamma_{Z \rightarrow f\bar{f}}$ shown in Tab. 2. Both the DIZET and GRIFFIN results for the form factors include full $\mathcal{O}(\alpha^2)$ corrections, $\mathcal{O}(\alpha\alpha_s)$ and $\mathcal{O}(\alpha\alpha_s^2)$ QCD corrections, and leading higher-order corrections in an expansion in m_t^2 of $\mathcal{O}(\alpha_t^3)$, $\mathcal{O}(\alpha_t^2\alpha_s)$, and $\mathcal{O}(\alpha_t\alpha_s^3)$, but with the following differences: (a) the GRIFFIN result for $|\rho_Z^f|$ additionally includes non-factorizable EW-QCD corrections; (b) the DIZET result for Δr does not include the $\mathcal{O}(\alpha_t\alpha_s^3)$ contributions, and the $\mathcal{O}(\alpha\alpha_s, \alpha\alpha_s^2)$ terms are computed only in a large- m_t approximation.

A better agreement for Δr is obtained when adjusting GRIFFIN to match the order of Δr in DIZET, by only summing corrections of $\mathcal{O}(\alpha, \alpha^2, \alpha_t\alpha_s, \alpha_t\alpha_s^2)$. In this case, one finds a 4-digit agreement, as shown in Tab. 2.

Most predictions given by both programs for $|\rho_Z^f|$, $\sin^2 \theta_{\text{eff}}^f$ and $\Gamma_{Z \rightarrow f\bar{f}}$ agree with each other by at least four decimal points. As aforementioned, the definition of the effective weak-mixing angle at $f = \nu$ is ill-defined. Owing to an alternative definition of $\sin^2 \theta_{\text{eff}}^f$ by DIZET (see eq. 5.6 in Ref. [15]), a number is yet produced without phenomenological implications. The discrepancy is mildly larger for the form factors $|\rho_Z^d|$ and $|\rho_Z^b|$ for quark final states, which reflects the different implementations of the non-factorizable EW-QCD corrections (as mentioned above, these are encapsulated in the form factors in GRIFFIN, but treated separately in DIZET). This is especially true for $|\rho_Z^b|$, where the top quark comes into play for these types of corrections. However, these implementation differences do not affect the predictions for the partial widths at the given order, and indeed one can see from the table the numbers for $\Gamma_{Z \rightarrow q\bar{q}}$ agree better.

Let us now move on to comparisons of predictions for the differential cross-section, where for concreteness, we focus on the process $e^+e^- \rightarrow \mu^+\mu^-$. Within GRIFFIN, these predictions have been computed using the class `mat_SMNLO`, whereas for `Dizet 6.45`, they are based on outputs of the subroutine `ROKANC`, which have been assembled into predictions for the differential cross-section using the `KKMCee` framework. Given that the γZ box contribution is not included in DIZET, we also turned off the γZ contributions in GRIFFIN for this




Figure 1: Ratios of the differential cross-section for $e^+e^- \rightarrow \mu^+\mu^-$ using predictions from GRIFFIN v1.0 and DIZET 6.45, for three choices of the scattering angle θ . The left plot is focused on the Z-pole region, while the right plot shows a wider range of center-of-mass energies.

comparison for consistency. The results are shown in Fig. 1.

The left plot in the figure shows that there is very good agreement between GRIFFIN v1.0 and DIZET 6.45 in the Z-pole region, with deviations of $\mathcal{O}(10^{-3})$ or less. When analyzing a larger range of center-of-mass energies, as shown in the right plot, one finds larger discrepancies at the level of 0.5–2%. This is not surprising since away from the Z resonances, both codes only deliver NLO precision, and differences in the implementation in GRIFFIN and DIZET would be of NNLO. In particular, DIZET does not use the manifestly gauge-invariant pole expansion scheme described in sections 2 and 3. Note that the relative corrections in some kinematic regions (*e.g.* $\cos\theta = +0.8$ below the resonance and $\cos\theta = -0.8$ above the resonance) are enhanced due to strong cancellations between the s-channel photon and Z exchange contributions, which render the tree-level matrix element small. In these regions, the NLO corrections reach 20–30%, so that $\mathcal{O}(\%)$ discrepancies from missing NNLO contributions are perfectly consistent with expectations.

We also wish to note that GRIFFIN provides a framework that can be systematically extended to higher orders, and NNLO corrections for $e^+e^- \rightarrow f\bar{f}$ can be included in the library once they become available. With this improvement, the theory uncertainty for the differential cross-section will likely be reduced significantly below 1%.

6 Summary

The GRIFFIN library provides a consistent description of the IR-subtracted matrix elements for fermion scattering for a wide range of center-of-mass energies, including close to and far away from the Z-boson resonance. This is achieved by merging a complex-pole expansion, which provides an accurate description near the Z peak, with an unexpanded fixed-order calculation, which is more adequate outside of the Z resonance region.

Version 1.0 of the library includes all currently available higher-order SM corrections for the leading Z-pole term, and NLO SM corrections for the remainder. It also includes two input parameter schemes, which either use the Fermi constant G_μ or the W mass M_W as inputs and all available higher-order SM corrections for the translation between the two. The results have been validated and compared to the DIZET 6.45 library.

The structure of GRIFFIN is modular and object-oriented to easily facilitate future extensions. Possible such extensions are:

- Higher-order corrections for the Z-pole form factors and matrix elements;
- Matrix elements for Bhabha scattering and/or final states with additional partons;
- Predictions for BSM scenarios, including effective theory frameworks such as SMEFT;
- Implementation of different schemes for factorization of initial- and final-state radiation;
- Implementation of other processes, such as charged-current Drell-Yan production with a W-boson resonance, or W-boson decays.

The authors invite the community to contact them with feedback, suggestions for improvement, or to contribute new modules to include in the GRIFFIN system.

Acknowledgments

The authors are indebted to S. Jadach and J. Holeczek for providing us with a test program within the KKMCEE framework for producing comparisons, as well as for numerous discussions. We also thank Z. Wąs for helpful communications. This work has been supported in part by the U.S. National Science Foundation under grant no. PHY-2112829.

A Appendix: Download information

The GRIFFIN source code is available at <https://github.com/lisongc/GRIFFIN/releases>. The manual is also available at https://github.com/lisongc/GRIFFIN_manual.

References

- [1] ALEPH, DELPHI, L3, OPAL, SLD, LEP ELECTROWEAK WORKING GROUP, SLD ELECTROWEAK GROUP, SLD HEAVY FLAVOUR GROUP collaboration, S. Schael et al., *Precision electroweak measurements on the Z resonance*, *Phys. Rept.* **427** (2006) 257–454, [[hep-ex/0509008](#)].
- [2] CDF, D0 collaboration, T. A. Aaltonen et al., *Tevatron Run II combination of the effective leptonic electroweak mixing angle*, *Phys. Rev. D* **97** (2018) 112007, [[1801.06283](#)].
- [3] ATLAS collaboration, *Measurement of the effective leptonic weak mixing angle using electron and muon pairs from Z-boson decay in the ATLAS experiment at $\sqrt{s} = 8$ TeV*, ATLAS-CONF-2018-037.
- [4] CMS collaboration, A. M. Sirunyan et al., *Measurement of the weak mixing angle using the forward-backward asymmetry of Drell-Yan events in pp collisions at 8 TeV*, *Eur. Phys. J. C* **78** (2018) 701, [[1806.00863](#)].
- [5] ATLAS, CMS collaboration, *Addendum to the report on the physics at the HL-LHC, and perspectives for the HE-LHC: Collection of notes from ATLAS and CMS*, *CERN Yellow Rep. Monogr.* **7** (2019) Addendum, [[1902.10229](#)].
- [6] FCC collaboration, A. Abada et al., *FCC-ee: The Lepton Collider: Future Circular Collider Conceptual Design Report Volume 2*, *Eur. Phys. J. ST* **228** (2019) 261–623.
- [7] CEPC STUDY GROUP collaboration, CEPC Study Group, J. Guimarães da Costa, Y. Gao, S. Jin, J. Qian, C. Tully, C. Young, L. Wang, M. Ruan, H. Zhu, Q. Ouyang et al., *CEPC Conceptual Design Report: Volume 2 - Physics & Detector*, [1811.10545](#).
- [8] H. Baer, T. Barklow, K. Fujii, Y. Gao, A. Hoang, S. Kanemura et al., *The International Linear Collider Technical Design Report - Volume 2: Physics*, , [1306.6352](#).
- [9] P. Bambade et al., *The International Linear Collider: A Global Project*, , [1903.01629](#).
- [10] *Physics and Detectors at CLIC: CLIC Conceptual Design Report*, [1202.5940](#).
- [11] CLICDP, CLIC collaboration, T. Charles et al., *The Compact Linear Collider (CLIC) - 2018 Summary Report*, [1812.06018](#).
- [12] D. Y. Bardin, M. S. Bilenky, T. Riemann, M. Sachwitz and H. Vogt, *Dizet: A Program Package for the Calculation of Electroweak One Loop Corrections for the Process $e^+ e^- \rightarrow f^+ f^-$ Around the Z0 Peak*, *Comput. Phys. Commun.* **59** (1990) 303–312.

- [13] G. Montagna, O. Nicosini, F. Piccinini and G. Passarino, *TOPAZ0 4.0: A New version of a computer program for evaluation of deconvoluted and realistic observables at LEP-1 and LEP-2*, *Comput. Phys. Commun.* **117** (1999) 278–289, [[hep-ph/9804211](#)].
- [14] D. Y. Bardin, P. Christova, M. Jack, L. Kalinovskaya, A. Olchevski, S. Riemann et al., *ZFITTER v.6.21: A Semianalytical program for fermion pair production in e^+e^- annihilation*, *Comput. Phys. Commun.* **133** (2001) 229–395, [[hep-ph/9908433](#)].
- [15] A. B. Arbuzov, M. Awramik, M. Czakon, A. Freitas, M. W. Grunewald, K. Monig et al., *ZFITTER: A Semi-analytical program for fermion pair production in e^+e^- annihilation, from version 6.21 to version 6.42*, *Comput. Phys. Commun.* **174** (2006) 728–758, [[hep-ph/0507146](#)].
- [16] D. Y. Bardin, L. Kalinovskaya and G. Nanava, *An Electroweak library for the calculation of EWRC to $e^+e^- \rightarrow f\bar{f}$ within the topfit project*, [hep-ph/0012080](#).
- [17] U. Baur, S. Keller and W. K. Sakumoto, *QED radiative corrections to Z boson production and the forward backward asymmetry at hadron colliders*, *Phys. Rev. D* **57** (1998) 199–215, [[hep-ph/9707301](#)].
- [18] U. Baur, O. Brein, W. Hollik, C. Schappacher and D. Wackerth, *Electroweak radiative corrections to neutral current Drell-Yan processes at hadron colliders*, *Phys. Rev. D* **65** (2002) 033007, [[hep-ph/0108274](#)].
- [19] S. Dittmaier and M. Krämer, *Electroweak radiative corrections to W boson production at hadron colliders*, *Phys. Rev. D* **65** (2002) 073007, [[hep-ph/0109062](#)].
- [20] S. Dittmaier and M. Huber, *Radiative corrections to the neutral-current Drell-Yan process in the Standard Model and its minimal supersymmetric extension*, *JHEP* **01** (2010) 060, [[0911.2329](#)].
- [21] S. Dittmaier, T. Schmidt and J. Schwarz, *Mixed NNLO QCD \times electroweak corrections of $\mathcal{O}(N_f\alpha_s\alpha)$ to single-W/Z production at the LHC*, *JHEP* **12** (2020) 201, [[2009.02229](#)].
- [22] J. Gluza, A. Lorca and T. Riemann, *Automated use of DIANA for two-fermion production at colliders*, *Nucl. Instrum. Meth. A* **534** (2004) 289–292, [[hep-ph/0409011](#)].
- [23] C. M. Carloni Calame, G. Montagna, O. Nicosini and A. Vicini, *Precision electroweak calculation of the charged current Drell-Yan process*, *JHEP* **12** (2006) 016, [[hep-ph/0609170](#)].
- [24] C. M. Carloni Calame, G. Montagna, O. Nicosini and A. Vicini, *Precision electroweak calculation of the production of a high transverse-momentum lepton pair at hadron colliders*, *JHEP* **10** (2007) 109, [[0710.1722](#)].

- [25] A. Arbuzov, D. Bardin, S. Bondarenko, P. Christova, L. Kalinovskaya, G. Nanava et al., *One-loop corrections to the Drell–Yan process in SANC. (II). The Neutral current case*, *Eur. Phys. J. C* **54** (2008) 451–460, [0711.0625].
- [26] D. Y. Bardin et al., *Precision Description of Processes at Colliders in the SANC System*, *Phys. Part. Nucl.* **50** (2019) 395–432.
- [27] V. A. Zykunov, *Electroweak corrections to the Drell-Yan process in the high dimuon mass range*, hep-ph/0702203.
- [28] W. Płaczek, S. Jadach and M. W. Krasny, *Drell-Yan processes with WINHAC*, *Acta Phys. Polon. B* **44** (2013) 2171–2178, [1310.5994].
- [29] L. Barze, G. Montagna, P. Nason, O. Nicrosini, F. Piccinini and A. Vicini, *Neutral current Drell-Yan with combined QCD and electroweak corrections in the POWHEG BOX*, *Eur. Phys. J. C* **73** (2013) 2474, [1302.4606].
- [30] V. Bertone, M. Cacciari, S. Frixione and G. Stagnitto, *The partonic structure of the electron at the next-to-leading logarithmic accuracy in QED*, *JHEP* **03** (2020) 135, [1911.12040].
- [31] S. Jadach, B. F. L. Ward and Z. Was, *The Monte Carlo program KORALZ, for the lepton or quark pair production at LEP / SLC energies: From version 4.0 to version 4.04*, *Comput. Phys. Commun.* **124** (2000) 233–237, [hep-ph/9905205].
- [32] A. Arbuzov, S. Jadach, Z. Was, B. F. L. Ward and S. A. Yost, *The Monte Carlo Program KKMC, for the Lepton or Quark Pair Production at LEP/SLC Energies—Updates of electroweak calculations*, *Comput. Phys. Commun.* **260** (2021) 107734, [2007.07964].
- [33] S. Frixione et al., *Initial state QED radiation aspects for future e^+e^- colliders*, in *2022 Snowmass Summer Study*, 3, 2022, 2203.12557.
- [34] S. Willenbrock and G. Valencia, *On the definition of the Z boson mass*, *Phys. Lett. B* **259** (1991) 373–376.
- [35] A. Sirlin, *Theoretical considerations concerning the Z_0 mass*, *Phys. Rev. Lett.* **67** (1991) 2127–2130.
- [36] R. G. Stuart, *Gauge invariance, analyticity and physical observables at the Z_0 resonance*, *Phys. Lett. B* **262** (1991) 113–119.
- [37] H. G. J. Veltman, *Mass and width of unstable gauge bosons*, *Z. Phys. C* **62** (1994) 35–52.
- [38] P. Gambino and P. A. Grassi, *The Nielsen identities of the SM and the definition of mass*, *Phys. Rev. D* **62** (2000) 076002, [hep-ph/9907254].

- [39] P. A. Grassi, B. A. Kniehl and A. Sirlin, *Width and partial widths of unstable particles in the light of the Nielsen identities*, *Phys. Rev. D* **65** (2002) 085001, [[hep-ph/0109228](#)].
- [40] D. Y. Bardin, A. Leike, T. Riemann and M. Sachwitz, *Energy Dependent Width Effects in $e^+ e^-$ Annihilation Near the Z Boson Pole*, *Phys. Lett. B* **206** (1988) 539–542.
- [41] A. Denner and S. Dittmaier, *Electroweak Radiative Corrections for Collider Physics*, *Phys. Rept.* **864** (2020) 1–163, [[1912.06823](#)].
- [42] K. G. Chetyrkin, J. H. Kuhn and A. Kwiatkowski, *QCD corrections to the e^+e^- cross-section and the Z boson decay rate*, *Phys. Rept.* **277** (1996) 189–281, [[hep-ph/9503396](#)].
- [43] A. Czarnecki and J. H. Kuhn, *Nonfactorizable QCD and electroweak corrections to the hadronic Z boson decay rate*, *Phys. Rev. Lett.* **77** (1996) 3955–3958, [[hep-ph/9608366](#)].
- [44] R. Harlander, T. Seidensticker and M. Steinhauser, *Complete corrections of Order α_s^2 to the decay of the Z boson into bottom quarks*, *Phys. Lett. B* **426** (1998) 125–132, [[hep-ph/9712228](#)].
- [45] S. Jadach, B. F. L. Ward and Z. Was, *Coherent exclusive exponentiation CEEEX: The Case of the resonant $e^+ e^-$ collision*, *Phys. Lett. B* **449** (1999) 97–108, [[hep-ph/9905453](#)].
- [46] K. Melnikov and O. I. Yakovlev, *Final state interaction in the production of heavy unstable particles*, *Nucl. Phys. B* **471** (1996) 90–120, [[hep-ph/9501358](#)].
- [47] W. Beenakker, A. P. Chapovsky and F. A. Berends, *Nonfactorizable corrections to W pair production*, *Phys. Lett. B* **411** (1997) 203–210, [[hep-ph/9706339](#)].
- [48] W. Beenakker, A. P. Chapovsky and F. A. Berends, *Nonfactorizable corrections to W pair production: Methods and analytic results*, *Nucl. Phys. B* **508** (1997) 17–63, [[hep-ph/9707326](#)].
- [49] M. Greco, G. Panzeri-Srivastava and Y. Srivastava, *Radiative Corrections to $e^+ e^- \rightarrow \mu^+ \mu^-$ Around the Z0*, *Nucl. Phys. B* **171** (1980) 118.
- [50] S. Jadach and Z. Was, *Suppression of QED Interference Contributions to the Charge Asymmetry at the Z⁰ Resonance*, *Phys. Lett. B* **219** (1989) 103–106.
- [51] S. Dittmaier, A. Huss and C. Schwinn, *Mixed QCD-electroweak $\mathcal{O}(\alpha_s\alpha)$ corrections to Drell-Yan processes in the resonance region: pole approximation and non-factorizable corrections*, *Nucl. Phys. B* **885** (2014) 318–372, [[1403.3216](#)].
- [52] A. Freitas, *Higher-order electroweak corrections to the partial widths and branching ratios of the Z boson*, *JHEP* **04** (2014) 070, [[1401.2447](#)].

- [53] A. A. Akhundov, D. Y. Bardin and T. Riemann, *Electroweak One Loop Corrections to the Decay of the Neutral Vector Boson*, *Nucl. Phys. B* **276** (1986) 1–13.
- [54] M. Awramik, M. Czakon and A. Freitas, *Electroweak two-loop corrections to the effective weak mixing angle*, *JHEP* **11** (2006) 048, [[hep-ph/0608099](#)].
- [55] M. Consoli, W. Hollik and F. Jegerlehner, *Electroweak Radiative Corrections for Z Physics*, in *LEP Physics Workshop*, CERN-TH-5527-89, 1989.
- [56] F. Boudjema et al., *Standard model processes*, in *AGS / RHIC Users Annual Meeting*, 1, 1996, [hep-ph/9601224](#).
- [57] A. Denner, S. Dittmaier, M. Roth and L. H. Wieders, *Electroweak corrections to charged-current $e^+ e^- \rightarrow 4$ fermion processes: Technical details and further results*, *Nucl. Phys. B* **724** (2005) 247–294, [[hep-ph/0505042](#)].
- [58] A. Denner and S. Dittmaier, *The Complex-mass scheme for perturbative calculations with unstable particles*, *Nucl. Phys. B Proc. Suppl.* **160** (2006) 22–26, [[hep-ph/0605312](#)].
- [59] A. Blondel et al., *Standard model theory for the FCC-ee Tera-Z stage*, in *Mini Workshop on Precision EW and QCD Calculations for the FCC Studies : Methods and Techniques*, vol. 3/2019 of *CERN Yellow Reports: Monographs*, (Geneva), CERN, 9, 2018, [1809.01830](#), DOI.
- [60] M. Awramik, M. Czakon, A. Freitas and G. Weiglein, *Precise prediction for the W boson mass in the standard model*, *Phys. Rev. D* **69** (2004) 053006, [[hep-ph/0311148](#)].
- [61] C. Sturm, *Leptonic contributions to the effective electromagnetic coupling at four-loop order in QED*, *Nucl. Phys. B* **874** (2013) 698–719, [[1305.0581](#)].
- [62] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, *A new evaluation of the hadronic vacuum polarisation contributions to the muon anomalous magnetic moment and to $\alpha(\mathbf{m}_Z^2)$* , *Eur. Phys. J. C* **80** (2020) 241, [[1908.00921](#)].
- [63] A. Keshavarzi, D. Nomura and T. Teubner, *$g - 2$ of charged leptons, $\alpha(M_Z^2)$, and the hyperfine splitting of muonium*, *Phys. Rev. D* **101** (2020) 014029, [[1911.00367](#)].
- [64] F. Jegerlehner, *$\alpha_{QED,eff}(s)$ for precision physics at the FCC-ee/ILC*, in *Theory for the FCC-ee: Report on the 11th FCC-ee Workshop Theory and Experiments* (A. Blondel, J. Gluza, S. Jadach, P. Janot and T. Riemann, eds.), vol. 3/2020 of *CERN Yellow Reports: Monographs*, pp. 9–37, 2020, DOI.
- [65] W. J. Marciano and A. Sirlin, *Radiative corrections to neutrino induced neutral current phenomena in the $SU(2)_L \times U(1)$ theory*, *Phys. Rev.* **D22** (1980) 2695.

- [66] M. Awramik, M. Czakon, A. Freitas and G. Weiglein, *Complete two-loop electroweak fermionic corrections to $\sin^2 \theta_{\text{eff}}^{\text{lept}}$ and indirect determination of the Higgs boson mass*, *Phys. Rev. Lett.* **93** (2004) 201805, [hep-ph/0407317].
- [67] W. Hollik, U. Meier and S. Uccirati, *The effective electroweak mixing angle $\sin^2 \theta^{\text{eff}}$ with two-loop fermionic contributions*, *Nucl. Phys.* **B731** (2005) 213–224, [hep-ph/0507158].
- [68] M. Awramik, M. Czakon and A. Freitas, *Bosonic corrections to the effective weak mixing angle at $O(\alpha^2)$* , *Phys. Lett.* **B642** (2006) 563–566, [hep-ph/0605339].
- [69] W. Hollik, U. Meier and S. Uccirati, *The effective electroweak mixing angle $\sin^2 \theta^{\text{eff}}$ with two-loop bosonic contributions*, *Nucl. Phys.* **B765** (2007) 154–165, [hep-ph/0610312].
- [70] M. Awramik, M. Czakon, A. Freitas and B. Kniehl, *Two-loop electroweak fermionic corrections to $\sin^2 \theta_{\text{eff}}^{\text{bb}}$* , *Nucl. Phys.* **B813** (2009) 174–187, [0811.1364].
- [71] I. Dubovyk, A. Freitas, J. Gluza, T. Riemann and J. Usovitsch, *The two-loop electroweak bosonic corrections to $\sin^2 \theta_{\text{eff}}^{\text{bb}}$* , *Phys. Lett.* **B762** (2016) 184–189, [1607.08375].
- [72] A. Djouadi and C. Verzegnassi, *Virtual very heavy top effects in LEP/SLC precision measurements*, *Phys. Lett.* **B195** (1987) 265–271.
- [73] A. Djouadi, *$O(\alpha_s)$ vacuum polarization functions of the standard model gauge bosons*, *Nuovo Cim.* **A100** (1988) 357.
- [74] B. A. Kniehl, *Two loop corrections to the vacuum polarizations in perturbative QCD*, *Nucl. Phys.* **B347** (1990) 86–104.
- [75] B. A. Kniehl and A. Sirlin, *Dispersion relations for vacuum polarization functions in electroweak physics*, *Nucl. Phys.* **B371** (1992) 141–148.
- [76] A. Djouadi and P. Gambino, *Electroweak gauge boson selfenergies: Complete QCD corrections*, *Phys. Rev.* **D49** (1994) 3499–3511, [hep-ph/9309298].
- [77] L. Avdeev, J. Fleischer, S. Mikhailov and O. Tarasov, *$O(\alpha_s^2)$ correction to the electroweak ρ parameter*, *Phys. Lett.* **B336** (1994) 560–566, [hep-ph/9406363].
- [78] K. Chetyrkin, J. H. Kühn and M. Steinhauser, *Corrections of order $O(G_F M_t^2 \alpha_s^2)$ to the ρ parameter*, *Phys. Lett.* **B351** (1995) 331–338, doi:10.1016/0370-2693(95)00380-4, [hep-ph/9502291].
- [79] J. J. van der Bij, K. G. Chetyrkin, M. Faisst, G. Jikia and T. Seidensticker, *Three loop leading top mass contributions to the ρ parameter*, *Phys. Lett.* **B498** (2001) 156–162, [hep-ph/0011373].

- [80] M. Faisst, J. H. Kühn, T. Seidensticker and O. Veretin, *Three loop top quark contributions to the ρ parameter*, *Nucl. Phys.* **B665** (2003) 649–662, [[hep-ph/0302275](#)].
- [81] Y. Schröder and M. Steinhauser, *Four-loop singlet contribution to the ρ parameter*, *Phys. Lett.* **B622** (2005) 124–130, [[hep-ph/0504055](#)].
- [82] K. G. Chetyrkin, M. Faisst, J. H. Kühn, P. Maierhöfer and C. Sturm, *Four-loop QCD corrections to the ρ parameter*, *Phys. Rev. Lett.* **97** (2006) 102003, [[hep-ph/0605201](#)].
- [83] R. Boughezal and M. Czakon, *Single scale tadpoles and $O(G_F m_t^2 \alpha_s^3)$ corrections to the ρ parameter*, *Nucl. Phys.* **B755** (2006) 221–238, [[hep-ph/0606232](#)].
- [84] L. Chen and A. Freitas, *Leading fermionic three-loop corrections to electroweak precision observables*, *JHEP* **07** (2020) 210, [[2002.05845](#)].
- [85] L. Chen and A. Freitas, *Mixed EW-QCD leading fermionic three-loop corrections at $\mathcal{O}(\alpha_s \alpha^2)$ to electroweak precision observables*, *JHEP* **03** (2021) 215, [[2012.08605](#)].
- [86] R. Harlander, T. Seidensticker and M. Steinhauser, *Complete corrections of order $\alpha \alpha_s$ to the decay of the Z boson into bottom quarks*, *Phys. Lett.* **B426** (1998) 125–132, [[hep-ph/9712228](#)].
- [87] J. Fleischer, O. Tarasov, F. Jegerlehner and P. Raczka, *Two loop $O(\alpha_s G_\mu m_t^2)$ corrections to the partial decay width of the Z^0 into $b\bar{b}$ final states in the large top mass limit*, *Phys. Lett.* **B293** (1992) 437–444.
- [88] G. Buchalla and A. J. Buras, *QCD corrections to the $\bar{s}dZ$ vertex for arbitrary top quark mass*, *Nucl. Phys.* **B398** (1993) 285–300.
- [89] G. Degrassi, *Current algebra approach to heavy top effects in $Z \rightarrow b + \bar{b}$* , *Nucl. Phys.* **B407** (1993) 271–289, [[hep-ph/9302288](#)].
- [90] K. Chetyrkin, A. Kwiatkowski and M. Steinhauser, *Leading top mass corrections of order $O(\alpha \alpha_s m_t^2 / m_W^2)$ to partial decay rate $\Gamma(Z \rightarrow b\bar{b})$* , *Mod. Phys. Lett.* **A8** (1993) 2785–2792.
- [91] A. Freitas, *Two-loop fermionic electroweak corrections to the Z-boson width and production rate*, *Phys. Lett.* **B730** (2014) 50–52, [[1310.2256](#)].
- [92] I. Dubovyk, A. Freitas, J. Gluza, T. Riemann and J. Usovitsch, *Complete electroweak two-loop corrections to Z boson production and decay*, *Phys. Lett.* **B783** (2018) 86–94, [[1804.10236](#)].
- [93] I. Dubovyk, A. Freitas, J. Gluza, T. Riemann and J. Usovitsch, *Electroweak pseudo-observables and Z-boson form factors at two-loop accuracy*, *JHEP* **08** (2019) 113, [[1906.08815](#)].

- [94] “ZFITTER 6.42/DIZET 6.45.” <http://sanc.jinr.ru/users/zfitter/>.
- [95] S. Jadach, B. F. L. Ward, Z. Was, S. A. Yost and A. Siodmok, *Multi-photon Monte Carlo event generator KKMCEE for lepton and quark pair production in lepton colliders*, *Comput. Phys. Commun.* **283** (2023) 108556, [2204.11949].