

Quantum Monte Carlo Simulation of the 3D Ising Transition on the Fuzzy Sphere

Johannes S. Hofmann^{1*}, Florian Goth², Wei Zhu³, Yin-Chen He⁴, and Emilie Huffman^{4†}

¹ Department of Condensed Matter Physics, Weizmann Institute of Science, Rehovot, 76100, Israel

² Institute for Theoretical Physics, University of Würzburg, Würzburg, 97074, Germany

³ School of Science, Westlake University, Hongzhou, 310024, P. R. China

⁴ Perimeter Institute for Theoretical Physics, Waterloo, Ontario N2L 2Y5, Canada

* johannes-stephan.hofmann@weizmann.ac.il, † ehuffman@perimeterinstitute.ca

Abstract

We present a numerical quantum Monte Carlo (QMC) method for simulating the 3D phase transition on the recently proposed fuzzy sphere [Phys. Rev. X 13, 021009 (2023)]. By introducing an additional $SU(2)$ layer degree of freedom, we reformulate the model into a form suitable for sign-problem-free QMC simulation. From the finite-size-scaling, we show that this QMC-friendly model undergoes a quantum phase transition belonging to the 3D Ising universality class, and at the critical point we compute the scaling dimensions from the state-operator correspondence, which largely agrees with the prediction from the conformal field theory. These results pave the way to construct sign-problem-free models for QMC simulations on the fuzzy sphere, which could advance the future study on more sophisticated criticalities.

Copyright attribution to authors.

This work is a submission to SciPost Physics Core.

License information to appear upon publication.

Publication information to appear upon publication.

Received Date

Accepted Date

Published Date

1

2 Contents

3	1 Introduction	2
4	2 Model and Method	3
5	2.1 Review of fuzzy sphere regularization	3
6	2.2 The density form of interaction	5
7	2.3 Four component fuzzy sphere model	5
8	3 Results	7
9	3.1 QMC Simulations	7
10	3.2 Finite-Size-Scaling	8
11	3.3 Dimensionless two-point correlator	9
12	3.4 Energy gaps and state-operator correspondence	10
13	4 Conclusions	11
14	A Order Parameter	12

15	B Extracting Energy Gaps	13
16	C Finite size scaling of energy gaps	15
17	References	17

18

20 1 Introduction

21 Critical phenomena emerging at classical and quantum phase transitions are of great interest
 22 due to their experimental relevance and theoretical significance [1, 2]. Many critical phe-
 23 nomena are believed to be described by conformal field theories (CFTs), which are strongly-
 24 interacting and pose challenges for studies in higher space-time dimensions beyond 2D (i.e.,
 25 1+1D). A recent method known as fuzzy (non-commutative) sphere regularization [3] has
 26 been invented to investigate 3D (i.e., 2+1D) critical phenomena governed by 3D CFTs on a
 27 cylindrical geometry represented as $S^2 \times \mathbb{R}$. Compared to traditional lattice regularization,
 28 the fuzzy sphere regularization offers numerous advantages in the study of 3D CFTs, primarily
 29 due to the utilization of radial quantization in $S^2 \times \mathbb{R}$ [4, 5] as well as the exact preservation
 30 of sphere $SO(3)$ symmetry [6, 7], as convincingly demonstrated recently [3, 8–11].

31 Firstly, the fuzzy sphere enables direct access to information regarding the emergent con-
 32 formal symmetry in the critical state [3, 10]. Secondly, it allows for the direct extraction of
 33 various data of the CFTs, including numerous scaling dimensions of conformal primary op-
 34 erators [3, 10], operator product expansion coefficients [8], and four-point correlators [9].
 35 For instance, scaling dimensions can be computed directly from excitation energies of the sys-
 36 tem, and their accuracy can be further improved using conformal perturbation [12]. Thirdly,
 37 the fuzzy sphere scheme is applicable to a variety of 3D CFTs, including Ising [3], $O(N)$
 38 Wilson-Fisher, $SO(5)$ deconfined phase transition [10], critical gauge theories [10], and de-
 39 fect CFTs [11]. Lastly, the fuzzy sphere regularization exhibits an incredibly small finite-size
 40 effect when the Hamiltonian is reasonably fine-tuned. These advantageous features of fuzzy
 41 sphere regularization present an exciting opportunity to explore 3D CFTs with high efficiency,
 42 accuracy, and comprehensiveness.

43 The fuzzy sphere regularization considers a microscopic quantum Hamiltonian modeling
 44 fermions (with multiple flavors) on continuous spherical space and projecting fermions into the
 45 lowest spherical Landau level [3, 6, 13]. In comparison with the regular lattice model, the fuzzy
 46 sphere model preserves the continuous rotational symmetry exactly in the UV limit. Thanks to
 47 the extremely small finite-size effect achieved through fine-tuning, numerical algorithms such
 48 as exact diagonalization (ED) and density matrix renormalization group (DMRG) methods are
 49 highly effective in studying the fuzzy sphere model of the 3D Ising CFT and $SO(5)$ deconfined
 50 phase transition. However, the computational cost of these two algorithms will eventually
 51 grow exponentially with the system size. More importantly, for cases involving a large number
 52 of fermion flavors, the computational costs of ED and DMRG quickly surpass practical resource
 53 and time limitations. In these cases, it would be helpful to be able to study models on the fuzzy
 54 sphere using a method that scales polynomially in time, such as quantum Monte Carlo (QMC).

55 The aim of this paper is to utilize the 3D Ising CFT as an example to demonstrate the
 56 application of the QMC approach in studying 3D CFTs on the fuzzy sphere. A similar discussion
 57 for the fuzzy torus model can be found in Ref. [13, 14]. In contrast to the fuzzy sphere Ising
 58 model introduced in Ref. [3], we introduce an additional flavor index to the fermions, which
 59 results in no sign problem for the QMC simulations. As a benchmark, we provide numerical

60 results of finite-size scaling, indicating that this model also belongs to the 3D Ising universality
 61 class. Furthermore, we introduce observables that enable the extraction of energy gaps in the
 62 spectrum corresponding to specific symmetry quantum numbers. This allows us to investigate
 63 the presence of conformal symmetry at criticality and extract scaling dimensions through the
 64 state-operator correspondence. Our numerical results for energy gaps are consistent with the
 65 universality of the 3D CFT Ising model, albeit with a larger finite-size effect compared to the
 66 previously studied fuzzy sphere Ising model [3]. In summary, we believe the QMC enriches
 67 the arsenal to study the fuzzy sphere model.

68 This paper is organized as follows: in Section II we introduce the model and its sym-
 69 metries, and we discuss how it can be implemented in auxiliary-field QMC simulations. We
 70 also argue for why the simulations are sign-problem-free. In Section III we discuss finite-size-
 71 scaling results and give evidence that the model is in the 3D Ising universality class, and we
 72 discuss energy spectrum results and give evidence for emergent conformal symmetry. Section
 73 IV contains our conclusions.

74 2 Model and Method

75 2.1 Review of fuzzy sphere regularization

76 The fuzzy sphere regularization considers fermions moving on a sphere in the presence of
 77 a magnetic monopole with $4\pi s$ flux sitting in the center of the sphere. In general, we can
 78 consider multi-flavor fermions ψ_α with the flavor index α , described by a Hamiltonian,

$$H = H_{\text{kin}} + H_{\text{int}}. \quad (1)$$

79 Here H_{kin} is the kinetic term of fermions, and H_{int} is an interaction which takes forms such as
 80 a density-density interaction,

$$\int d^2\vec{r}_1 d^2\vec{r}_2 U(\vec{r}_1 - \vec{r}_2) n^a(\vec{r}_1) n^b(\vec{r}_2), \quad (2)$$

81 where $n^a(\vec{r}) = \psi^\dagger(\vec{r})_\alpha \psi(\vec{r})_\beta M_{\alpha\beta}^a$ and M^a is a matrix defined in the fermion flavor space.
 82 $U(\vec{r}_1 - \vec{r}_2)$ is a rotationally invariant interaction, and we take it to be short ranged such as
 83 $\delta(\vec{r}_1 - \vec{r}_2)$ and $\nabla^2 \delta(\vec{r}_1 - \vec{r}_2)$.

84 The energy levels of H_{kin} form quantized Landau levels, whose wave-functions are de-
 85 scribed by the monopole Harmonics (i.e. spin-weighted spherical Harmonics) $Y_{n+s,m}^{(s)}(\theta, \varphi)$ [15],
 86 with $\mathbf{n} = \mathbf{0}, \mathbf{1}, \dots$ as the Landau level index and (θ, φ) as the spherical coordinates. Each Lan-
 87 dau level has an energy $E_n = [(\mathbf{n} + \mathbf{1}/2) + \mathbf{n}(\mathbf{n} + \mathbf{1})/2s] \omega_c / 2\pi$, with the cyclotron frequency
 88 ω_c [6, 16]. The states of each Landau level have an angular momentum $L = s + \mathbf{n}$, hence
 89 they are $(2s + 2\mathbf{n} + 1)$ -fold degenerate, which can be labeled by the quantum number of the
 90 z -component of the angular momentum L^z , $m = -s - \mathbf{n}, -s - \mathbf{n} + 1, \dots, s + \mathbf{n}$. Because we
 91 may set the scale of H_{kin} as large as we like ($\omega_c \rightarrow \infty$) relative to our scale of interest for
 92 H_{int} , we may enforce that there are no fluctuations out of the lowest Landau level (LLL). The
 93 physics of interest will come from the terms that make up H_{int} . Hence, we consider the limit
 94 that $H_{\text{kin}} \gg H_{\text{int}}$ such that we can project the system into the LLL. The annihilation operator
 95 $\psi(\theta, \varphi)$ on the LLL can be written as

$$\psi_\alpha(\theta, \varphi) = \frac{1}{\sqrt{N}} \sum_{m=-s}^s \bar{Y}_{s,m}^{(s)}(\theta, \varphi) c_{m,\alpha}, \quad (3)$$

96 where $c_{m,\alpha}$ stands for the annihilation operator of Landau orbital m , and it is independent of
 97 coordinates (θ, φ) . $N = 2s + 1$ is the number of orbitals, playing the role of area of the 2D

98 space. The prefactor $1/\sqrt{N}$ ensures that the density operator,

$$n^a(\theta, \varphi) = \frac{1}{N} \sum_{m_1, m_2} Y_{s, m_1}^{(s)} \bar{Y}_{s, m_2}^{(s)} c_{m_1, \alpha}^\dagger c_{m_2, \beta} M_{\alpha\beta}^a, \quad (4)$$

99 is an intensive quantity.

100 Under this LLL projection Eq. (3), any rotation invariant density-density interaction in the
101 form of eq. (2) can be written as the Haldane pseudopotentials [6] in terms of second quantized
102 fermion operators,

$$\sum_{m_1, m_2, m_3, m_4} V_{m_1, m_2, m_3, m_4} (c_{m_1, \alpha}^\dagger M_{\alpha\beta}^a c_{m_4, \beta}) (c_{m_2, \eta}^\dagger M_{\eta\gamma}^b c_{m_3, \gamma}) \quad (5)$$

103 with

$$V_{m_1, m_2, m_3, m_4} = \delta_{m_1+m_2, m_3+m_4} \sum_{l=0}^{2s} V_l (4s - 2l + 1) \times \begin{pmatrix} s & s & 2s-l \\ m_1 & m_2 & -m_1-m_2 \end{pmatrix} \begin{pmatrix} s & s & 2s-l \\ m_4 & m_3 & -m_3-m_4 \end{pmatrix}, \quad (6)$$

104 where $\begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix}$ is the Wigner $3j$ -symbol. V_l are numbers whose values are specifically
105 depending on the form of the interaction $U(\vec{r}_1 - \vec{r}_1)$. For the remainder of this work, we focus
106 on $U(\vec{r}_1 - \vec{r}_1) = U(\Omega_{12}) = \frac{g_0}{N} \delta(\Omega_{12}) + \frac{g_1}{N^2} \nabla^2 \delta(\Omega_{12})$ with $\delta(\Omega_{12}) = \delta(\varphi_1 - \varphi_2) \delta(\cos \theta_1 - \cos \theta_2)$.
107 The corresponding Haldane pseudo-potentials are

$$V_0 = \frac{2s+1}{4s+1} g_0 - \frac{s}{4s+1} g_1, \quad V_1 = \frac{s}{4s-1} g_1, \quad V_{l \geq 2} = 0. \quad (7)$$

108 To realize the $2+1$ D Ising transition, Ref. [3] introduced a Hamiltonian that has two flavors
109 of fermions $\psi^\dagger = (\psi_\uparrow^\dagger, \psi_\downarrow^\dagger)$ with their interaction,

$$H_{\text{int}} = \int N^2 d\Omega_1 d\Omega_2 U(\Omega_{12}) [n^0(\theta_1, \varphi_1) n^0(\theta_2, \varphi_2) - n^z(\theta_1, \varphi_1) n^z(\theta_2, \varphi_2)] - h \int N d\Omega n^x(\theta, \varphi), \quad (8)$$

110 where $\Omega = (\theta, \varphi)$ is a spherical coordinate and $n^a(\theta, \varphi) = \psi^\dagger(\theta, \varphi) \sigma^a \psi(\theta, \varphi)$ is a local
111 density operator with $\sigma^{x,y,z}$ being Pauli matrices, $\sigma^0 = I_{2 \times 2}$. The first term behaves like an
112 Ising ferromagnetic interaction, while the second term is the transverse field.

113 It is straightforward to solve the second quantized Hamiltonian Eq. (5) using unbiased
114 numerical algorithm such as the ED and the DMRG, although their computational costs grow
115 exponentially with the system size $N = 2s + 1$. So it is highly desirable to develop QMC
116 method for the simulation of a fuzzy sphere model, and it is the focus of this paper.

117 It is worth mentioning why the LLL projection leads to a fuzzy sphere. We can consider the
118 projection of the coordinates of a unit sphere, denoted as $x = (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta)$.
119 After the projection, the coordinates are transformed into $(2s + 1) \times (2s + 1)$ matrices, where
120 $(X)_{m_1, m_2} = \int d\Omega x \bar{Y}_{s, m_1}^{(s)}(\Omega) Y_{s, m_2}^{(s)}(\Omega)$. These matrices satisfy the following relations:

$$[X_i, X_j] = \frac{1}{s+1} i \epsilon_{ijk} X_k, \quad \sum_{i=1}^3 X_i X_i = \frac{s}{s+1} \mathbf{1}_{2s+1}. \quad (9)$$

121 The fact that the three coordinates satisfy the $SO(3)$ algebra formally defines a fuzzy sphere
122 [7]. It is interesting to note that in the limit as $s \rightarrow \infty$, the fuzziness disappears and a unit
123 sphere is recovered.

124 2.2 The density form of interaction

125 To facilitate QMC simulation, we would like to write the Hamiltonian in terms of the density
126 operator in the angular momentum space $n_{l,m}^a$, defined as,

$$n^a(\theta, \varphi) = \frac{1}{N} \sum_{l,m} n_{l,m}^a Y_l^m(\theta, \varphi). \quad (10)$$

127 Here $Y_l^m(\theta, \varphi)$ is the spherical harmonics, with $m = -l, -l+1, \dots, l$ and $l \in \mathbb{Z}$. $n_{l,m}^a$ can be
128 obtained using the spherical harmonic transformation,

$$\begin{aligned} n_{l,m}^a &= N \int d\Omega \bar{Y}_l^m(\theta, \varphi) n^a(\theta, \varphi) \\ &= N \sqrt{\frac{2l+1}{4\pi}} \sum_{m_1=-s}^s (-1)^{3s+m_1} \begin{pmatrix} s & l & s \\ -m_1 & m & m_1-m \end{pmatrix} \\ &\quad \times \begin{pmatrix} s & l & s \\ -s & 0 & s \end{pmatrix} c_{m_1,\alpha}^\dagger c_{m_1-m,\beta} M_{\alpha\beta}^a \end{aligned} \quad (11)$$

129 To have the term $\begin{pmatrix} s & l & s \\ -m_1 & m & m_1-m \end{pmatrix}$ non-vanishing, we should have $l \leq 2s$. One can show
130 $n_{l,m}^\dagger = (-1)^m n_{l,-m}$.

131 In this context, it is convenient to decompose the potential $U(\theta_{12}) = \sum_l \frac{2l+1}{4\pi} U_l P_l(\cos \theta_{12})$
132 using the Legendre polynomials, $P_l(\cos \theta_{12}) = \frac{4\pi}{2l+1} \sum_{m=-l}^l \bar{Y}_l^m(\Omega_1) Y_l^m(\Omega_2)$, such that the
133 interaction terms take the form

$$\begin{aligned} &\int N^2 d\Omega_1 d\Omega_2 U(\theta_{12}) n^a(\theta_1, \varphi_1) n^b(\theta_2, \varphi_2) \\ &= \sum_{l=0}^{2s} U_l \sum_{m=-l}^l (n_{l,m}^a)^\dagger n_{l,m}^b, \end{aligned} \quad (12)$$

134 with the coefficients $U_l = g_0/N - l(l+1)g_1/N^2$.

135 2.3 Four component fuzzy sphere model

136 In comparison to Ref. [3], we consider four flavors of fermions, $\psi^\dagger = (\psi_{\uparrow,+}^\dagger, \psi_{\uparrow,-}^\dagger, \psi_{\downarrow,+}^\dagger, \psi_{\downarrow,-}^\dagger)$,
137 i.e., we introduce an additional ‘‘layer’’ degree of freedom (+, -). The Pauli-matrices σ^i and
138 τ^i act on the spin (\uparrow, \downarrow) and layer indices, respectively. Let us define the operators $n_{l,m}^0$ and
139 $n_{l,m}^z$ according to Eq. (11) with $M^0 = \sigma^0 \tau^0$ and $M^z = \sigma^z \tau^0$, respectively. The Hamiltonian
140 reads

$$\begin{aligned} H_{\text{int}} &= \sum_{l=0}^{2s} U_l \sum_{m=-l}^l [(n_{l,m}^0)^\dagger n_{l,m}^0 - (n_{l,m}^z)^\dagger n_{l,m}^z] \\ &\quad + h \sum c_m^\dagger \sigma^x \tau^0 c_m, \end{aligned} \quad (13)$$

141 and the interaction favors a ferromagnetic state for $g_0, g_1 > 0$.

142 There are four symmetries of this model, which are

143 1. Ising \mathbb{Z}_2 symmetry: $c_m \rightarrow \sigma^x \tau^0 c_m$.




Figure 1: Order parameter data for $V_0 = 0.5564$ and $V_1 = 0.1$, which shows evidence for a continuous phase transition consistent with that of the 3D Ising Model. (a) Binder ratio data showing a crossing that drifts somewhat in system size: $N = 12$ and $N = 14$ cross around $h = 0.21$, whereas $N = 8$ and $N = 10$ cross closer to $h = 0.20$. (b) Magnetization data showing consistent crossing between $h = 0.21$ and $h = 0.22$ for $N = 10, 12, 14$. Ising $\Delta_\sigma = 0.518$ is assumed. (c) Magnetization data plotted along with a universal scaling function fit. Fixing $\Delta_\sigma = 0.518$ and $\Delta_\epsilon = 1.41$ (as shown in this figure) yields a good fit ($\chi^2 = 1.305$) with $h_c = 0.2129(8)$, consistent with 3D Ising universality. Fitting using an $h = 0.21$ estimate consistent with the Binder ratio and magnetization crossing data gives $\Delta_\sigma = 0.49(2)$ and $\Delta_\epsilon = 1.28(6)$.

- 144 2. $SO(3)$ sphere rotation symmetry: $\mathbf{c}_{m=-s,\dots,s}$ form the spin- s representation of $SO(3)$.
 145 3. Particle-hole symmetry: $\mathbf{c}_m \rightarrow i\sigma^y \tau^0 \mathbf{c}_m^\dagger$, and $i \rightarrow -i$.
 146 4. Layer $SU(2)$: generated by $\mathbf{c}_m \rightarrow \sigma^0 \tau^{x,y,z} \mathbf{c}_m$.

147 The first three of these symmetries are the same as those of the two-flavor model studied
 148 in [3]. The layer symmetry is an additional symmetry for the four flavors, which allows for
 149 sign-problem-free QMC simulations of this model. At the Ising transition, the layer $SU(2)$
 150 degrees of freedom need to be gapped. We have verified this in Appendix C.

151 Before moving on, we remark that the four component fuzzy sphere model Eq. 13 is not a
 152 simple product of the two-component model Eq. 8, so the phase transition point h_c of the four
 153 component model is distinct from that of the two-component model. While the interaction is
 154 chosen to be of density form in the layer degree of freedom, i.e., using τ^0 , to disfavor layer-
 155 magnetism, the two layers are coupled and spontaneous layer-symmetry breaking cannot ruled
 156 out in general. Nevertheless, we ensure the universality of the four component model falls in
 157 the 3D Ising class, as shown below.

158 3 Results

159 3.1 QMC Simulations

160 We simulate the model, eq. (13), using projector auxiliary Quantum Monte Carlo (AFQMC).
 161 To fit this goal, we rewrite the sums of quartic terms in the following way

$$\begin{aligned}
 U_l \sum_{m=-l}^l (n_{l,m}^a)^\dagger n_{l,m}^a &= U_l \sum_{m=-l}^l (-1)^m (n_{l,-m}^a) n_{l,m}^a \\
 &= \frac{U_l}{4} \sum_{m=-l}^l \left((1+i)n_{l,m}^a + (1-i)(-1)^m n_{l,-m}^a \right)^2,
 \end{aligned}
 \tag{14}$$

162 where in the first equality we make use of the density operator identity

$$n_{l,m}^{\dagger} = (-1)^m n_{l,-m}. \tag{15}$$

163 The squared operators in the second line of (14) are Hermitian, and thus AFQMC as imple-
 164 mented in [17] is applicable. The projector we use is the half-filled solution to the model when
 165 $\mathbf{g}_0 = \mathbf{g}_1 = \mathbf{0}$, where the Ising spins are polarized by the transverse field term $\mathbf{h} \sum \mathbf{c}_m^\dagger \boldsymbol{\sigma}^x \boldsymbol{\tau}^0 \mathbf{c}_m$.

166 Now we show, that the QMC simulation of this model is sign-problem-free. After the
 167 Hubbard-Stratonovich transformation, we have a prefactor of $\sqrt{-\Delta \tau U_l / 4}$. If $\mathbf{g}_0 - \mathbf{g}_1 l(l+1)/(2s+1)$
 168 of the expression in (13) is always positive, then we get an extra factor of i for the \mathbf{n}^0 terms,
 169 which picks up a sign under antiunitary transformations. The antiunitary particle-hole trans-
 170 formation \mathcal{P} ,

$$\begin{aligned}
 \mathbf{c}_m^\dagger &\rightarrow i \boldsymbol{\sigma}^y \boldsymbol{\tau}^0 \mathbf{c}_m, & i &\rightarrow -i \\
 n_{l,m}^z &\rightarrow (-1)^m n_{l,-m}^z \\
 n_{l,m}^0 &\rightarrow -(-1)^m n_{l,-m}^0,
 \end{aligned}
 \tag{16}$$

171 is a symmetry with \mathcal{P}^2 . Combined with the $SU(2)$ layer symmetry, it guarantees the absence of
 172 the sign-problem in this model [18]. Hence, the computational complexity scales polynomially
 173 with system size N . However, in this basis, the auxiliary fields couple to the operators $\mathbf{n}_{l,m}^a$ of
 174 typical rank N , compared to the usual rank of $\mathcal{O}(1)$ in lattice models. Therefore, the compute
 175 time of the algorithm scales as N^4 instead of the conventional N^3 .

176 In this particular model, we focus on the critical point that occurs in a regime where both
 177 $\mathbf{g}_0, \mathbf{g}_1 > \mathbf{0}$. We have not proven in the discussion above that the absence of a sign problem
 178 occurs when instances of $\mathbf{g}_0 - \mathbf{g}_1 l(l+1)/(2s+1)$ is positive for small l but is negative for
 179 large l , yet we encountered no sign problem in our simulations. One explanation may be that
 180 the Wigner-3j prefactor $\begin{pmatrix} s & l & s \\ -s & 0 & s \end{pmatrix}$ decays exponentially in l , causing a suppression of terms
 181 that change the overall prefactor signs in (13), and so the smallness of the couplings of these
 182 terms may be important.

183 Due to the nonlocal nature of the operators in (13), controlling Trotter discretization errors
 184 becomes a more demanding task, as observed in [14]. We alleviate some of these effects
 185 by adopting a stabilized second-order Trotter decomposition developed by Blanes et al., as
 186 discussed in [19]. The effectiveness of these alternate splitting schemes in the realm of AFQMC
 187 was shown in [20]. Furthermore, to implement the Wigner-3j prefactors efficiently, we utilize
 188 the software package detailed in [21].

189 Here we utilize QMC to compute the evolution of the order parameter and CFT dimen-
 190 sionless two-point correlators across the transition point, as well as extract energy gaps for
 191 the lowest lying states using time-displaced correlation functions (see Appendix B). While the

192 Lowest Landau level basis has already been used for a QMC study in [22], the CFT-inspired
 193 use of time-displaced correlation functions and two-point correlators on the fuzzy sphere is
 194 new for QMC studies.

195 Below we will set $V_1 = 0.1$, $V_0 = 0.5564$, and tune h to realize a 2+1D Ising transition.
 196 We find this choice of (V_0, V_1) has a smaller finite size effect. By inverting the equations in
 197 (7), we have

$$\begin{aligned} g_0 &= \frac{4s+1}{2s+1}V_0 + \frac{4s-1}{2s+1}V_1 \\ g_1 &= \frac{4s-1}{s}V_1, \end{aligned} \quad (17)$$

198 so this corresponds to the region $g_0, g_1 > 0$.

199 3.2 Finite-Size-Scaling

200 To look for a phase transition, we begin with the order parameter for the Ising phase transition,
 201 which in the Landau Level basis (see Appendix A) is given by

$$M = \sum_{m=-s}^s c_m^\dagger \sigma^z c_m. \quad (18)$$

202 In Figure 1(a), we have plotted the Binder cumulant, given by

$$U_4 = 1 - \frac{\langle M^4 \rangle}{3 \langle M^2 \rangle^2}, \quad (19)$$

203 and we see a crossing in the vicinity of $0.20 - 0.21$, that drifts to larger couplings with larger
 204 $N = 2s + 1$. With this evidence of there being a quantum phase transition, we can find further
 205 evidence that the phase transition is in the Ising universality class by assuming that Δ_σ is equal
 206 to 0.518 , as is known for the universality class [23], and checking the the magnetization data,
 207 as seen in Figure 1(b). Here we see a good crossing for $N = 10, 12, 14$, which is consistent
 208 with the choice of Δ_σ . The crossing is at a larger h than the Binder cumulant crossing, that
 209 is because the Binder cumulant suffers from larger finite size effects. Similar finite size effects
 210 have also been observed in the two-layer model [3].

211 To see that the data is consistent with both $\eta (= 2\Delta_\sigma - 1)$ and $\nu (= 1/(3 - \Delta_\epsilon))$, critical ex-
 212 ponents in the 3D Ising universality class, we have performed a data collapse to a universal scal-
 213 ing function, assuming that $\langle M^2 \rangle / (\sqrt{N})^{4-2\Delta_\sigma}$ has a functional form of $f_0 + f_1x + f_2x^2 + f_3x^3$,
 214 where $x = (h - h_c) (\sqrt{N})^{3-\Delta_\epsilon}$. In addition to the fit parameters, we pay attention to the quan-
 215 tity χ^2/dof , with numerator χ^2 given by

$$\chi^2 = \frac{\sum_n (O_n - F_n)^2}{\sigma_n^2}, \quad (20)$$

216 where O_n is a measurement, F_n is the expected value from the fit, and σ_n is the variance
 217 for the measurement. The quantity χ^2/dof is reduced by the number of degrees of freedom
 218 (dof) for the fitted data, which is the number of measurements minus the number of free
 219 parameters. A good fit that captures the features of the data but does not overfit is close to 1.
 220 When we fix $\Delta_\sigma = 0.518$ and $\Delta_\epsilon = 1.41$ and leave the other five parameters free, we get a
 221 fit for the data $N = 10, 12, 14$, with $\chi^2/\text{dof} = 1.305$ and estimate for the critical coupling of
 222 $h_c = 0.2129(8)$, as seen in Figure 1(c). If instead we fix $h = 0.21$, as suggested by the Binder
 223 ratio, and leave six parameters including the critical exponents free, a fitting ($\chi^2/\text{dof} = 1.491$)
 224 gives us $\Delta_\sigma = 0.49(2)$ and $\Delta_\epsilon = 1.28(6)$, values consistent with Ising universality for these
 225 relatively small system sizes.




Figure 2: Data from CFT dimensionless two-point correlators. (a) Values of $\Delta_\sigma^{\text{proxy}}$ from $G_{\sigma\sigma}$ using the equal-time correlators. The crossing of the $\Delta_\sigma^{\text{proxy}}$ values through the $\Delta_\sigma = 0.518$ critical exponent value occurs at larger h -values as N increases, consistent with the finite-size-scaling drift that was observed earlier. (b) Extrapolation to infinite lattice size Δ_σ from $G_{\sigma\sigma}$ correlation functions for different values of h and $N = 6, 8, 10, 12, 14$. Linear fits suggest that, from this dataset, $h = 0.212$ is closest to criticality. (c) Plotting of the angle-dependence of the $G_{\sigma\sigma}$ correlator with data from $N = 6, 8, 10, 12, 14$ at $h = 0.212$.

226 3.3 Dimensionless two-point correlator

227 To take the advantage of fuzzy sphere regularization, below we compute CFT dimensionless
228 two-point correlators on a sphere [9] at equal time,

$$\begin{aligned} G_{\phi\phi}(\theta) &= \langle \phi(\theta = \varphi = 0) \phi(\theta, \varphi = 0) \rangle \\ &= \frac{1}{(2 \sin(\theta/2))^{2\Delta_\phi}}, \end{aligned} \quad (21)$$

229 where ϕ is a CFT primary operator, and (θ, φ) are the spherical coordinates specifying the
230 positions of the two operators. We mainly focus on the lowest \mathbb{Z}_2 -odd primary σ , which can
231 be well approximated by the UV operator n^z [8, 9], up to a non-universal normalization (say
232 \sqrt{A}) and higher order corrections $O(1/\sqrt{N})$ from operators with higher scaling dimensions.
233 So we can first compute the equal-time two-point correlator,

$$\begin{aligned} f(\theta) &= \langle n^z(\theta = \varphi = 0) n^z(\theta, \varphi = 0) \rangle \\ &= \sum_{l=0}^{2s} \bar{Y}_{l,m=0}(\theta, 0) Y_{l,m=0}(0, 0) \langle n_{l,0}^z n_{l,0}^z \rangle, \end{aligned} \quad (22)$$

234 and then

$$G_{\sigma\sigma}(\theta) = A f(\theta) + O(1/\sqrt{N}), \quad (23)$$

235 where A is a nonuniversal number. Because we have an explicit expression for f , we know
236 the exact values for $f(\theta = \pi)$ and $f''(\theta)|_{\theta=\pi}$, where the derivatives are taken in θ . Then by
237 assuming that $G_{\sigma\sigma}$ has the critical scaling form of (21), we can solve the following system,

$$\begin{aligned} A f(\pi) &= 1/(2 \sin(\pi/2))^{2\Delta_\sigma^{\text{proxy}}} \\ A f''(\theta)|_{\theta=\pi} &= \frac{\partial^2}{\partial \theta^2} \left(1/(2 \sin(\theta/2))^{2\Delta_\sigma^{\text{proxy}}} \right) \Big|_{\theta=\pi}, \end{aligned} \quad (24)$$

238 $\Delta_\sigma^{\text{proxy}}$ is a number that will extrapolate to the universal Δ_σ at the critical point as $N \rightarrow \infty$.
239 Second derivatives are used for the second equation in (24) because the first derivatives in θ
240 are zero for both f and critical $G_{\phi\phi}$ at $\theta = \pi$.

241 Figure 2(a) shows the extracted $\Delta_\sigma^{\text{proxy}}$ values for different values of h in the vicinity of
 242 the h_c determined by finite-size-scaling. Here we see that the $\Delta_\sigma^{\text{proxy}}$ indeed crosses through
 243 the 3D Ising $\Delta_\sigma = 0.518$ value in this region, and furthermore we see that the h at which this
 244 crossing occurs increases with system size N , which is consistent with the drift that we saw in
 245 the finite-size-scaling. Moreover, the drift appears to be slowing with increasing N , another
 246 consistency with finite-size scaling.

247 We can see more consistencies with finite-size-scaling from the results in Figure 2(b), which
 248 linearly extrapolate the values of $\Delta_\sigma^{\text{proxy}}$ as $N \rightarrow \infty$ for different values of h . Here we see that
 249 for the infinite N extrapolation, the $h = 0.212$ data is closest to the critical Δ_σ , which is
 250 consistent with the earlier universal scaling fit of $h_c = 0.2129(8)$. We use this $h = 0.212$ data
 251 to show the calculation of $G_{\sigma\sigma}(\theta)$ as a function of θ for QMC data at $N = 6, 8, 10, 12, 14$ and
 252 how it approaches the exact expression as N increases. The QMC data-derived expressions for
 253 $G_{\sigma\sigma}(\theta)$ are given in Figure 2(c) and are rescaled so that they are equal to the exact expression
 254 of $G_{\sigma\sigma}(\theta = \pi)$.

255 3.4 Energy gaps and state-operator correspondence

256 Next we turn to the state-operator correspondence [4, 5] on the sphere, namely, the scaling
 257 dimensions Δ_n are related to energy gaps by

$$\delta E_n = E_n - E_0 = \frac{v}{R} \Delta_n, \quad (25)$$

258 where R is the radius of the sphere and v is the model-dependent velocity of light.

259 While we are unable to get the full low lying energy spectrum directly using QMC, we
 260 are able to obtain energy gaps for the lowest lying states in each symmetry quantum number
 261 sector by using time-displaced correlation functions (see Appendix B). For an operator O_S with
 262 the quantum number S , we have:

$$\langle O_S(\tau) O_S(0) \rangle = \sum_n a_n^2 e^{-\tau(E_{S,n} - E_0)}, \quad (26)$$

263 where E_0 is the ground state energy, $E_{S,n}$ represents the energies of eigenstates $|\psi_n\rangle$ in the
 264 quantum number sector S , a_n is an operator O_S and state $|\psi_n\rangle$ dependent non-universal factor.
 265 At long time $\tau \gg 1$, the lowest energy will dominate and can be extracted by fitting the
 266 exponential decay.

267 In the data that follows, we will use density operators $n_{l,m}^i$ to measure energy gaps in
 268 different quantum number sectors. Specifically,

269 1) $n_{l,m}^z$ can measure gaps in the \mathbb{Z}_2 -odd, parity-even, and angular momentum (i.e. Lorentz
 270 spin) l sector;

271 2) $n_{l,m}^x$ can measure gaps in the \mathbb{Z}_2 -even, parity-even, and angular momentum l sector;

272 3) $n_{l,m}^0$ can measure gaps in the \mathbb{Z}_2 -even, parity-odd, and angular momentum l sector.

273 Figure 3 shows QMC data at the critical point $h = 0.212$. The energy gaps are scaled such
 274 that the gap measured from $n_{l=2,m=0}^0$ is rescaled to 4, the scaling dimension of the lowest
 275 parity-odd descendent of the energy-momentum tensor, $\Delta_{\epsilon_{\nu\rho\eta}\partial_\rho T_{\mu\nu}}$. In doing so, we find the
 276 gaps measured from other operators to be consistent with the scaling dimensions of primary
 277 and descendant operators of the 3D Ising CFT. The density operator ($n_{l=2,m=0}^x$) we measured
 278 does not seem to have an overlap with the state of stress tensor (with $\Delta_{T_{\mu\nu}} = 3$). Instead it
 279 gives the level-2 descendant of ϵ primary, i.e., $\partial_\mu \partial_\nu \epsilon$.




Figure 3: Comparison between 3D Ising CFT data and rescaled energy gaps for $N = 8$ and $h = 0.212$ measured by density operators. The energy gaps are rescaled by the factor that makes the lowest \mathbb{Z}_2 -even, parity-odd gap at $l = 2$ equal to the lowest parity-odd descendent of the energy-momentum tensor, **4.0**.

280 4 Conclusions

281 We have introduced a model that is amenable to using sign-problem free quantum Monte
 282 Carlo to simulate the $(2+1)\text{-D}$ transverse Ising model on a fuzzy sphere. Through finite-size
 283 scaling we have found data consistent with the model's phase transition being in the 3D Ising
 284 universality class, and we also have shown that we can recover the same critical exponents
 285 from the model's energy spectra, which is evidence of emergent conformal symmetry.

286 While these calculations are not competitive with ED and DMRG for small lattices, this
 287 work opens the door to larger scale calculations for models where there are too many degrees
 288 of freedom to make ED/DMRG calculations infeasible, or for when large sizes are desired
 289 for more accurate determination of critical exponents. Here, we were able to introduce an
 290 additional layer degree of freedom to avoid the sign-problem, and chose the interaction to
 291 disfavor spontaneous layer-symmetry breaking energetically. However, such instabilities can-
 292 not be ruled out in general. However, there are many interesting critical phenomena which
 293 naturally involve multiple flavors; one interesting target is the critical gauge theories proposed
 294 in Ref. [10].

295 Acknowledgements

296 We thank Chao Han and Zheng Zhou for stimulating discussions.

297 **Funding information** Research at Perimeter Institute is supported in part by the Govern-
 298 ment of Canada through the Department of Innovation, Science and Industry Canada and
 299 by the Province of Ontario through the Ministry of Colleges and Universities. WZ is sup-
 300 ported by National Natural Science Foundation of China (No. 92165102) and foundation of
 301 the Westlake University. JH was supported by the European Research Council (ERC) under

302 grant HQMAT (Grant Agreement No. 817799), the Israel-US Binational Science Foundation
 303 (BSF), and by a Research grant from Irving and Cherna Moskowitz. FG acknowledges fi-
 304 nancial support through the German Research Foundation, project-id 258499086 - SFB 1170
 305 ‘ToCoTronics’. This work used Bridges 2 at the Pittsburgh Supercomputing Center through
 306 allocation PHY170036 from the Advanced Cyberinfrastructure Coordination Ecosystem: Ser-
 307 vices & Support (ACCESS) program, which is supported by National Science Foundation grants
 308 #2138259, #2138286, #2138307, #2137603, and #2138296.

309 A Order Parameter

310 The Ising order parameter is n^z , and for QMC simulations we measure the two-point correla-
 311 tion function,

$$\begin{aligned} \langle n^z(\theta, \varphi) n^z(\theta', \varphi') \rangle &= \sum_{l,m,l',m'} \langle n_{l,m}^z n_{l',m'}^z \rangle Y_l^m(\theta, \varphi) Y_{l'}^{m'}(\theta', \varphi') \\ &= \sum_{l,m} (-1)^m \langle n_{l,0}^z n_{l,0}^z \rangle Y_l^m(\theta, \varphi) Y_l^{-m}(\theta', \varphi'). \end{aligned} \quad (\text{A.1})$$

312 The last equation comes from the conservation of angular momentum,

$$\begin{aligned} \langle n_{l,m}^z n_{l',m'}^z \rangle &= \begin{pmatrix} l & l' & 0 \\ m & m' & 0 \end{pmatrix} \mathcal{O}_l \\ &= \delta_{l,l'} \delta_{m,-m'} \begin{pmatrix} l & l & 0 \\ m & -m & 0 \end{pmatrix} \langle n_{l,0}^z n_{l,0}^z \rangle / \begin{pmatrix} l & l & 0 \\ 0 & 0 & 0 \end{pmatrix} \\ &= (-1)^m \delta_{l,l'} \delta_{m,-m'} \langle n_{l,0}^z n_{l,0}^z \rangle. \end{aligned} \quad (\text{A.2})$$

313 Therefore, we need to evaluate $\langle n_{l,0}^z n_{l,0}^z \rangle$ for each l . To do the finite-size-scaling, we cal-
 314 culate the order parameter $\langle M^2 \rangle$, with $M = \int d\Omega n^z(\theta, \varphi)$,

$$\begin{aligned} \langle M^2 \rangle &= \int d\Omega d\Omega' \langle n^z(\theta, \varphi) n^z(\theta', \varphi') \rangle \\ &= \sum_{l,m} (-1)^m \langle n_{l,0}^z n_{l,0}^z \rangle \int d\Omega d\Omega' Y_l^m(\theta, \varphi) Y_l^{-m}(\theta', \varphi') \\ &= 4\pi \langle n_{0,0}^z n_{0,0}^z \rangle \end{aligned} \quad (\text{A.3})$$

315 The last equation comes from $\int d\Omega Y_l^m(\theta, \varphi) = \sqrt{4\pi} \delta_{l,0} \delta_{m,0}$. Using Wigner-3j identities, we
 316 find that,

$$\begin{aligned} n_{0,0}^z &= (2s+1) \sqrt{\frac{1}{4\pi}} \sum_{m_1=-s}^s (-1)^{3s+m_1} \begin{pmatrix} s & 0 & s \\ -m_1 & 0 & m_1 \end{pmatrix} \begin{pmatrix} s & 0 & s \\ -s & 0 & s \end{pmatrix} c_{m_1}^\dagger \sigma^z c_{m_1} \\ &= \sqrt{\frac{1}{4\pi}} \sum_{m_1=-s}^s c_{m_1}^\dagger \sigma^z c_{m_1}. \end{aligned} \quad (\text{A.4})$$

317 Therefore, the order parameter M^2 is

$$\langle M^2 \rangle = \sum_{m_1, m_2=-s}^s \langle (c_{m_1}^\dagger \sigma^z c_{m_1}) (c_{m_2}^\dagger \sigma^z c_{m_2}) \rangle. \quad (\text{A.5})$$

318 B Extracting Energy Gaps

319 In projector QMC, we are able to get the energy gap between the first excited state in symmetry
 320 sector S and the ground state in the following way. Assuming a trial wavefunction, $|\psi_0\rangle$, an
 321 operator that creates overlap between the states in symmetry sector S and the ground state,
 322 O_S , and a complete set of states $\sum_n |n\rangle \langle n|$, where $|n\rangle$ is an eigenstate with energy eigenvalue
 323 E_n , we have that

$$\begin{aligned}
 \langle O_S(\tau) O_S(0) \rangle &= \frac{\langle \psi_0 | e^{-(\beta-\tau)H} O_S e^{-\tau H} O_S | \psi_0 \rangle}{\langle \psi_0 | \psi_0 \rangle} \\
 &= \sum_n \frac{\langle \psi_0 | e^{-(\beta-\tau)H} O_S | n \rangle \langle n | e^{-\tau H} O_S | \psi_0 \rangle}{\langle \psi_0 | \psi_0 \rangle} \\
 &= \sum_n \frac{\langle \psi_0 | O_S | n \rangle \langle n | O_S | \psi_0 \rangle}{\langle \psi_0 | \psi_0 \rangle} e^{-\beta E_0} e^{-\tau(E_n - E_0)}.
 \end{aligned} \tag{B.1}$$

324 This term has contributions from all eigenstates that have the symmetry S . The higher energy
 325 states will have gaps that will be suppressed relative to the smallest energy gap, and so we can
 326 approximate

$$\langle O_S(\tau) O_S(0) \rangle = C_1 e^{-\tau(E_S^0 - E_0)} + C_2 e^{-\tau(E_S^1 - E_0)}, \tag{B.2}$$

327 where E_S^0 and E_S^1 are the lowest energy and second lowest energy corresponding to states in
 328 symmetry sector S , respectively.

329 In practice, we found that in order to extract E_S^0 , sometimes a fit to the two exponentials
 330 with prefactors C_1 and C_2 in (5) is necessary, but sometimes a fit to a single exponential (which
 331 assumes $C_2 = 0$) is more appropriate. The procedure for fitting to one versus two exponentials
 332 involves the following steps:

- 333 1. Find the τ interval where the data is distinguishable from zero according to error bars.
 334 The largest time in this interval is the initial ‘‘endpoint’’ guess.
- 335 2. Initially guess that the ‘‘midpoint’’ in time—where one exponential versus the other ex-
 336 ponential dominates—is 30% of the full time interval.
- 337 3. Test a fit to a single exponential—if the initial data point is smaller than the $t = 0$ value
 338 for the fitted function, gradually adjust the midpoint and endpoint guesses down until
 339 this is not the case.
- 340 4. If the initial data point is still within errors of the $t = 0$ value for the single-exponential
 341 fitted function, fit the data to a single exponential. If not move on to a two-exponential
 342 fit and then revise the midpoint guess such that the value of the larger exponential in
 343 the fit is negligible compared to that of the smaller exponential at the midpoint.

344 For the operators $n_{1,0}^z$, $n_{2,0}^0$, and $n_{2,0}^x$, the fit to a single exponential ends up being more ap-
 345 propriate. Figure 4 shows the fits to a single exponential (in the cases of $n_{2,0}^2$ and $n_{2,0}^x$, we
 346 chose a single exponential because there was very little small τ data to fit to a higher expo-
 347 nential). However, the $n_{0,0}^x$, $n_{1,0}^x$ observables require two exponentials. Figure 5 shows the
 348 two exponential fits for these operators at coupling $h = 0.212$ and $s = 3.5$.

349 For the two-exponential fits, we first use the midpoint data to iteratively fit one exponential
 350 at a time: midpoint to endpoint is the fit for the lower energy and then the startpoint to the
 351 midpoint is the fit for the higher energy. We alternate fitting one exponential versus the other
 352 while fixing the parameters of the nonfitted exponential according to the previous fit. This
 353 gives us the ‘‘restricted’’ fit listed as ‘‘fit’’ in Figure 5. We then use these fitted energy values as




Figure 4: Fits to one exponential using semilog plots. The vertical lines show the locations of the endpoint and startpoints for the data. Plots (a), (b), and (c) show the data from $n_{0,0}^z$, $n_{1,0}^z$, and $n_{2,0}^z$, respectively. Plot (d) shows the data from $n_{2,0}^0$ and plot (e) shows the data from $n_{2,0}^x$.




Figure 5: Fits to two exponentials using semilog plots. The vertical solid lines show the locations of the endpoint and midpoint for a restricted fit—where the C_1 exponential and the C_2 exponential are fitted separately but iteratively using information from previous results to arrive at a final answer. The dashed lines give an interval of midpoints that were tested in order to estimate the error bar. The solid diagonal lines give the two-exponential restricted and unrestricted fits (“fit” and “fit-unrestricted”). The dashed diagonal lines are fits for each of the two exponentials, both for restricted (“fit-part”) and unrestricted (“fit-part_ur”) fits. Plot (a) shows the data from $n_{0,0}^x$, and plot (b) shows the data from $n_{1,0}^x$.

354 initial guesses for an “unrestricted” fit that fits both exponentials at once. This gives the “fit-
 355 unrestricted” in Figure 5. The value of the energy estimate is the mean of these restricted and
 356 unrestricted fit energies. Finally, we obtain error bars by performing fits to a single exponential
 357 for the smaller energy from a midpoint to the endpoint, where we calculate the midpoint as

$$\tau = -\frac{1}{(E_s^1 - E_s^0) \ln\left(\frac{\epsilon C_1}{C_2}\right)}, \quad (\text{B.3})$$

358 where ϵ is a small number representing the time when the value of $C_2 e^{-\tau(E_s^1 - E_0)}$ is ϵ times
 359 $C_1 e^{-\tau(E_s^0 - E_0)}$. We take a range of $\epsilon \in \{0.01, 0.1\}$ to fit E_s^0 and use this range of E_s^0 values to
 360 estimate the error for the energy. The boundaries of this range of midpoints are given by the
 361 dashed vertical lines in Figure 5.

362 C Finite size scaling of energy gaps

363 Because the QMC model studied has an additional $SU(2)$ -layer symmetry, one check to make
 364 is whether the degrees of freedom in the layer $SU(2)$ non-singlet representations are gapped
 365 at the phase transition. All the operators in the Hamiltonian are of the form $\sigma^i \tau^0$, which are
 366 the singlets of the layer $SU(2)$ symmetry. Figure 6 shows the energy gaps obtained from these
 367 operators as a function of $1/\sqrt{N}$ at $h = 0.2$, which is in the vicinity of the critical point. All
 368 gaps are decreasing with system size and the gaps for the \mathbb{Z}_2 -odd sector seem to be trending
 369 linearly towards the origin, as required by the state-operator correspondence. The \mathbb{Z}_2 -even
 370 sector gaps are also decreasing with increasing system size, but they have larger error bars
 371 due to interference with other higher energy descendants in their spectrum, and the details of
 372 their fits are given in Appendix B of the Supplementary Material. On the other hand, the layer
 373 $SU(2)$ non-singlet, e.g., layer triplet gaps should be finite in the thermodynamic limit. Figure
 374 7 shows layer triplet energy gaps measured by $\sigma^i \tau^{x,y,z}$, as a function of $1/\sqrt{N}$, and from here
 375 we see that these excitations appear to be gapped.




Figure 6: Data showing the energy gaps obtained from the $SU(2)$ singlet operators as a function of $1/\sqrt{N}$ for $N = 8, 10, 14$. This data is in the vicinity of the critical point at $V_1 = 0.1, V_0 = 0.5564, h = 0.2$. The gaps decrease with system size, as expected. Extrapolations are shown for the \mathbb{Z}_2 -odd sector, but not the even sector since the error bars are so large.




Figure 7: Data showing the energy gaps obtained from operators of the form $\sigma^i \tau^z$ as a function of $1/\sqrt{N}$ for $N = 8, 10, 14$ (operators of the form $\sigma^i \tau^{x,y}$ would give the same states, making this the $SU(2)$ triplet symmetry class). This data is in the vicinity of the critical point at $V_1 = 0.1, V_0 = 0.5564, h = 0.2$. These energies appear to be gapped out.

376 **References**

- 377 [1] S. Sachdev, *Quantum Phase Transitions*, Cambridge University Press, Cambridge, Eng-
378 land, ISBN 9780511973765 (2011).
- 379 [2] J. Cardy, *Scaling and Renormalization in Statistical Physics*, Cambridge University Press,
380 Cambridge, England, ISBN 9781316036440 (1996).
- 381 [3] W. Zhu, C. Han, E. Huffman, J. S. Hofmann and Y.-C. He, *Uncovering Conformal Symmetry*
382 *in the 3D Ising Transition: State-Operator Correspondence from a Quantum Fuzzy Sphere*
383 *Regularization*, Phys. Rev. X **13**(2), 021009 (2023), doi:[10.1103/PhysRevX.13.021009](https://doi.org/10.1103/PhysRevX.13.021009),
384 [2210.13482](https://doi.org/10.1103/PhysRevX.13.021009).
- 385 [4] J. L. Cardy, *Conformal invariance and universality in finite-size scaling*, Journal of Physics
386 A: Mathematical and General **17**(7), L385 (1984), doi:[10.1088/0305-4470/17/7/003](https://doi.org/10.1088/0305-4470/17/7/003).
- 387 [5] J. L. Cardy, *Universal amplitudes in finite-size scaling: generalisation to arbitrary di-*
388 *mensionality*, Journal of Physics A: Mathematical and General **18**(13), L757 (1985),
389 doi:[10.1088/0305-4470/18/13/005](https://doi.org/10.1088/0305-4470/18/13/005).
- 390 [6] F. D. M. Haldane, *Fractional quantization of the hall effect: A hierarchy of incompressible*
391 *quantum fluid states*, Phys. Rev. Lett. **51**, 605 (1983), doi:[10.1103/PhysRevLett.51.605](https://doi.org/10.1103/PhysRevLett.51.605).
- 392 [7] J. Madore, *The fuzzy sphere*, Class. Quantum Gravity **9**(1), 69 (1992),
393 doi:[10.1088/0264-9381/9/1/008](https://doi.org/10.1088/0264-9381/9/1/008).
- 394 [8] L. Hu, Y.-C. He and W. Zhu, *Operator Product Expansion Coefficients of the 3D Ising Criti-*
395 *cality via Quantum Fuzzy Sphere* (2023), [2303.08844](https://arxiv.org/abs/2303.08844).
- 396 [9] C. Han, L. Hu, W. Zhu and Y.-C. He, *Conformal four-point correlators of the 3D Ising*
397 *transition via the quantum fuzzy sphere* (2023), [2306.04681](https://arxiv.org/abs/2306.04681).
- 398 [10] Z. Zhou, L. Hu, W. Zhu and Y.-C. He, *The $SO(5)$ Deconfined Phase Transi-*
399 *tion under the Fuzzy Sphere Microscope: Approximate Conformal Symmetry, Pseudo-*
400 *Criticality, and Operator Spectrum*, arXiv e-prints arXiv:2306.16435 (2023),
401 doi:[10.48550/arXiv.2306.16435](https://doi.org/10.48550/arXiv.2306.16435), [2306.16435](https://arxiv.org/abs/2306.16435).
- 402 [11] L. Hu, Y.-C. He and W. Zhu, *Solving Conformal Defects in 3D Conformal Field*
403 *Theory using Fuzzy Sphere Regularization*, arXiv e-prints arXiv:2308.01903 (2023),
404 doi:[10.48550/arXiv.2308.01903](https://doi.org/10.48550/arXiv.2308.01903), [2308.01903](https://arxiv.org/abs/2308.01903).
- 405 [12] B.-X. Lao and S. Rychkov, *3D Ising CFT and Exact Diagonalization on Icosahedron*, arXiv
406 e-prints arXiv:2307.02540 (2023), doi:[10.48550/arXiv.2307.02540](https://doi.org/10.48550/arXiv.2307.02540), [2307.02540](https://arxiv.org/abs/2307.02540).
- 407 [13] M. Ippoliti, R. S. K. Mong, F. F. Assaad and M. P. Zaletel, *Half-filled landau levels: A*
408 *continuum and sign-free regularization for three-dimensional quantum critical points*, Phys.
409 Rev. B **98**, 235108 (2018), doi:[10.1103/PhysRevB.98.235108](https://doi.org/10.1103/PhysRevB.98.235108).
- 410 [14] Z. Wang, M. P. Zaletel, R. S. K. Mong and F. F. Assaad, *Phases of the (2+1) dimen-*
411 *sional $SO(5)$ nonlinear sigma model with topological term*, Physical Review Letters **126**(4)
412 (2021), doi:[10.1103/physrevlett.126.045701](https://doi.org/10.1103/physrevlett.126.045701).
- 413 [15] T. T. Wu and C. N. Yang, *Dirac monopole without strings: monopole harmonics*, Nuclear
414 Physics B **107**(3), 365 (1976).

- 415 [16] M. Greiter, *Landau level quantization on the sphere*, Physical Review B **83**, 115129
416 (2011), doi:[10.1103/PhysRevB.83.115129](https://doi.org/10.1103/PhysRevB.83.115129), [1101.3943](https://arxiv.org/abs/1101.3943).
- 417 [17] F. F. Assaad, M. Bercx, F. Goth, A. Götz, J. S. Hofmann, E. Huffman, Z. Liu, F. P. Toldin,
418 J. S. E. Portela and J. Schwab, *The ALF (Algorithms for Lattice Fermions) project release 2.0.*
419 *Documentation for the auxiliary-field quantum Monte Carlo code*, SciPost Phys. Codebases
420 p. 1 (2022), doi:[10.21468/SciPostPhysCodeb.1](https://doi.org/10.21468/SciPostPhysCodeb.1).
- 421 [18] C. Wu and S.-C. Zhang, *Sufficient condition for absence of the sign problem in*
422 *the fermionic quantum monte carlo algorithm*, Phys. Rev. B **71**, 155115 (2005),
423 doi:[10.1103/PhysRevB.71.155115](https://doi.org/10.1103/PhysRevB.71.155115).
- 424 [19] S. Blanes, F. Casas and J. M. Sanz-Serna, *Numerical integrators for the hybrid*
425 *monte carlo method*, SIAM Journal on Scientific Computing **36**(4), A1556 (2014),
426 doi:[10.1137/130932740](https://doi.org/10.1137/130932740), <https://doi.org/10.1137/130932740>.
- 427 [20] F. Goth, *Higher order auxiliary field quantum monte carlo methods* (2020), [2009.04491](https://arxiv.org/abs/2009.04491).
- 428 [21] H. T. Johansson and C. Forssén, *Fast and accurate evaluation of wigner $3j$, $6j$, and $9j$*
429 *symbols using prime factorization and multiword integer arithmetic*, SIAM Journal on
430 Scientific Computing **38**(1), A376 (2016), doi:[10.1137/15m1021908](https://doi.org/10.1137/15m1021908).
- 431 [22] B.-B. Chen, X. Zhang, Y. Wang, K. Sun and Z. Y. Meng, *Phases of $(2+1)d$ $so(5)$ non-linear*
432 *sigma model with a topological term on a sphere: multicritical point and disorder phase*
433 (2023), [2307.05307](https://arxiv.org/abs/2307.05307).
- 434 [23] M. Hasenbusch, *Finite size scaling study of lattice models in the three-dimensional ising*
435 *universality class*, Phys. Rev. B **82**, 174433 (2010), doi:[10.1103/PhysRevB.82.174433](https://doi.org/10.1103/PhysRevB.82.174433).