

Pairing Symmetry and Fermion Projective Symmetry Groups

Xu Yang^{1*}, Sayak Biswas^{1†}, Shuangyuan Lu^{1†}, Mohit Randeria¹ and Yuan-Ming Lu¹

¹ Department of Physics, The Ohio State University, Columbus, OH 43210, USA

* yang.6309@osu.edu, † These two authors contributed equally.

Abstract

The Ginzburg-Landau (GL) theory is very successful in describing the pairing symmetry, a fundamental characterization of the broken symmetries in a paired superfluid or superconductor. However, GL theory does not describe fermionic excitations such as Bogoliubov quasiparticles or Andreev bound states that are directly related to topological properties of the superconductor. In this work, we show that the symmetries of the fermionic excitations are captured by a Projective Symmetry Group (PSG), which is a group extension of the bosonic symmetry group in the superconducting state. We further establish a correspondence between the pairing symmetry and the fermion PSG. When the normal and superconducting states share the same spin rotational symmetry, there is a simpler correspondence between the pairing symmetry and the fermion PSG, which we enumerate for all 32 crystalline point groups. We also discuss the general framework for computing PSGs when the spin rotational symmetry is spontaneously broken in the superconducting state. This PSG formalism leads to experimental consequences, and as an example, we show how a given pairing symmetry dictates the classification of topological superconductivity.

Copyright attribution to authors.

This work is a submission to SciPost Physics.

License information to appear upon publication.

Publication information to appear upon publication.

Received Date

Accepted Date

Published Date

1

2 Contents

3	1 Introduction	2
4	2 Characterization of broken symmetries in a superconductor	4
5	2.1 Projective Symmetry Group and Projective Representation	4
6	2.2 Pairing Symmetry and Projective Representations	5
7	3 Constraints on the pairing symmetry by the PSG	7
8	4 Examples	8
9	4.1 Tetragonal Symmetry	8
10	4.2 Hexagonal Symmetry	11
11	4.3 Trigonal Symmetry	13
12	4.4 Physical consequences of the PSG	14
13	4.4.1 3d SCs with mirror reflection symmetry M_x	15
14	4.4.2 2d SCs with 2-fold rotational symmetry C_{2z}	16

15	5 General framework	16
16	5.1 Group extension and pairing symmetry in a generic superconductor	17
17	5.2 Examples: superfluid A and B phases in Helium-3	18
18	5.2.1 Superfluid B phase of Helium-3	19
19	5.2.2 Superfluid A phase of Helium-3	19
20	6 Conclusion	20
21	A A short introduction to projective representation and 2-cocycle	21
22	B How PSG constrains the pairing symmetry for all crystalline point groups	24
23	C GAP program for PSG calculation	24
24	References	29
25		
26		

27 1 Introduction

28 One of the most fundamental characterizations of a superconductor or a paired superfluid is
 29 the symmetry of its pair wavefunction. The standard way of describing pairing symmetry is
 30 in terms of the irreducible representations (irreps) of the *normal state* symmetry group \mathcal{G}_0
 31 which constrains the form of the Ginzburg-Landau (GL) free energy functional [1–4]. \mathcal{G}_0 can
 32 be written as

$$\mathcal{G}_0 = \mathbf{G}_0 \times U(1) = \begin{cases} \mathbf{X}_0 \times \mathbf{SO}(3)_{\text{spin}} \times U(1) & \text{Weak SOC} \\ \mathbf{X}_0 \times U(1) & \text{Strong SOC} \end{cases} \quad (1)$$

33 where \mathbf{X}_0 is the crystalline point group, and SOC denotes spin-orbit coupling. At a second or-
 34 der phase transition, the superconductor spontaneously breaks global charge $U(1)$ symmetry
 35 as the system condenses into a particular irrep of the normal state symmetry group. In gen-
 36 eral, the group of unbroken symmetries in the superconducting phase, $\mathbf{G} \subseteq \mathbf{G}_0$. For example,
 37 $\mathbf{G} = \mathbf{X} \times \mathbf{SO}(3)_{\text{spin}}$ for a singlet superconductor with weak SOC, where $\mathbf{X} \subseteq \mathbf{X}_0$ is the point
 38 group symmetry preserved in the superconductor. In the presence of a strong SOC we have
 39 $\mathbf{G} = \mathbf{X}$ with $\mathbf{X} \subseteq \mathbf{X}_0$ being the unbroken point group of the superconductor.

40 Essentially all of the phonon-mediated superconductors (SCs) exhibit singlet “s-wave” pair-
 41 ing, where the superconducting (SC) state transforms according to the trivial representation of
 42 \mathbf{X}_0 . But superfluid ^3He [5] and many quantum materials, including the heavy fermion SCs [6],
 43 the high T_c cuprates [7], and Sr_2RuO_4 [8], condense into nontrivial irreps.

44 In this paper, we wish to focus on the relation between pairing symmetry and the symmetry
 45 of the Hamiltonian describing the *fermionic excitations in the superconducting state*. At the mean
 46 field level, one focuses on the Bogoliubov-de Gennes (BdG) Hamiltonian, but the fermionic
 47 symmetry analysis applies equally beyond the BdG framework where one needs to take into
 48 account interactions between quasiparticles. The approach we develop here will allow us to
 49 gain new insights that go beyond the (bosonic) GL theory.

50 Examples of questions which this formalism would shed light on include: (a) the rela-
 51 tion between pairing symmetry and topology, as the K-theory classification [9–11] of non-
 52 interacting topological SCs is based on the BdG Hamiltonians, (b) how interactions between
 53 quasiparticles for various pairing symmetries impacts the classification of interacting topolog-
 54 ical SC phases [11–14], (c) the relation between pairing symmetry and excitations in topo-

55 logical defects such as Majorana zero modes trapped in vortices [15–19], and (d) whether
 56 new probes of electronic excitations can provide insight into the pairing symmetry [20]. We
 57 discussed question (a) in section 4.4 of the manuscript. We will return to other questions in
 58 subsequent papers.

59 Here, we first show how starting with the pairing symmetry, together with the crystalline
 60 symmetries that dictate the normal state electronic structure, we can derive the projective
 61 symmetry group (PSG) [21] for the fermionic excitations in the SC state. We first focus on
 62 the cases where the superconductor shares the same spin rotational symmetry as the normal
 63 state, we present an exhaustive classification of the SC state PSG corresponding to every al-
 64 lowable pairing symmetry for the 32 crystalline point groups with and without SOC. When
 65 confronted with a new superconductor, we would like to use these results in the “reverse” di-
 66 rection, namely, how can we deduce the possible pairing symmetry, given fermionic properties
 67 in the SC state. Mathematically, the map from the pairing symmetry to the SC state PSG is,
 68 in general, neither injective nor surjective, and thus it cannot be inverted. Nevertheless, we
 69 show below that the SC state PSG does constrain to a considerable extent the possible pairing
 70 symmetries. We also present numerous examples that serve to illustrate our general results.

71 To describe the symmetries of the fermionic Hamiltonian we need (i) to focus on the *super-*
 72 *conducting state* symmetry group \mathbf{G} as distinct from the *normal state* \mathbf{G}_0 relevant for GL theory,
 73 and (ii) to take into account fermion parity $(-1)^{\hat{F}}$, where \hat{F} is the total number of fermions in
 74 the system. Let us discuss each of these points in turn.

75 On general grounds, the SC state symmetry group \mathbf{G} is a subgroup of the normal state \mathbf{G}_0 .
 76 If the irrep into which the GL theory condenses is one-dimensional, then in fact $\mathbf{G} = \mathbf{G}_0$. While
 77 this is obvious for the trivial \mathbf{A}_1 representation, an example may be useful to illustrate why
 78 this is true quite generally. Consider the $\mathbf{d}_{x^2-y^2}$ pairing state in the cuprates that transforms
 79 according to the \mathbf{B}_{1g} irrep of the tetragonal symmetry group \mathbf{D}_{4h} . The pair wavefunction
 80 changes sign under a $\pi/2$ rotation, and one might naively think that this breaks \mathbf{C}_4 down to
 81 \mathbf{C}_2 . However, one can compensate for this minus sign by having the fermion operators pick up
 82 an $e^{i\pi/2}$ phase under \mathbf{C}_4 and thus have the electronic Hamiltonian retain the full symmetry of
 83 the normal state. We will see a generalization of this at play in the analysis later in section 2.

84 On the other hand, if the irrep has a dimension > 1 , then one needs to solve the GL
 85 equations to find the SC state that minimizes the free energy. Then the SC state symmetry
 86 is lower than that in the normal state, and \mathbf{G} is a proper subgroup of \mathbf{G}_0 . For example,
 87 ^3He is a \mathbf{p} -wave, triplet superfluid, corresponding to the $\mathbf{L} = \mathbf{1}, \mathbf{S} = \mathbf{1}$ irrep of the normal state
 88 symmetry group $\mathbf{G}_0 = \mathbf{SO}(\mathbf{3})_{\text{orbital}} \times \mathbf{SO}(\mathbf{3})_{\text{spin}}$. Depending on external parameters various
 89 superfluid states are stabilized, and in the \mathbf{B} -phase of ^3He , for instance, \mathbf{G}_0 is broken down to
 90 $\mathbf{G} = \mathbf{SO}(\mathbf{3})_{\mathbf{L}+\mathbf{S}}$ [22]. We will discuss a general framework to understand the PSG of fermion
 91 excitations in any superconductor in Section 5, where the superconductor can spontaneously
 92 break the normal-state spin rotational symmetry.

93 The second point above related to fermion parity may seem trivial: it enforces that a Hamil-
 94 tonian can only have terms with an even number of fermion operators. It leads, however, to
 95 the important mathematical structure of a projective symmetry group (PSG) \mathbf{G}_f acting on the
 96 many-body Hilbert space. In Section 2, we discuss in detail how \mathbf{G}_f is built as a central exten-
 97 sion of \mathbf{G} by the fermion parity group \mathbb{Z}_2^F .

98 The rest of the paper is organized as follows. In Section 3 we show how the fermion PSG
 99 \mathbf{G}_f can constrain the pairing symmetry of the SC state, applying the framework to all 32 point
 100 groups (see Table 7) and demonstrating it by a few examples in section 4. We further discuss
 101 how the PSG determines topological properties of the SC in section 4.4. While sections 2-3
 102 focus on the cases where the normal state and the SC state shares the same spin rotational
 103 symmetries, in section 5.1 we describe a generic theory framework that applies to all supercon-
 104 ductors, and further demonstrate its power in the examples of A- and B-phases of superfluid

105 ^3He in section 5.2. Finally we conclude in section 6 with a discussion on how the fermion PSG
 106 in SCs discussed here differs from the PSG first introduced in quantum spin liquids [21, 23],
 107 and an outlook to future studies.

108 2 Characterization of broken symmetries in a superconductor

109 2.1 Projective Symmetry Group and Projective Representation

110 Any Hamiltonian must conserve fermion parity $(-1)^{\hat{F}}$ even if it does not conserve particle
 111 number \hat{F} , as, for instance, in the presence of pairing. The fermion symmetry group G_f acting
 112 on the many-body Hilbert space of fermions is a projective symmetry group (PSG). Mathemat-
 113 ically, G_f is a central extension of the bosonic symmetry group G in the SC state by the fermion
 114 parity group $\mathbb{Z}_2^F = \{(\pm 1)^{\hat{F}}\}$. This may be written as a short exact sequence

$$1 \rightarrow \mathbb{Z}_2^F \rightarrow G_f \rightarrow G \rightarrow 1 \quad (2)$$

115 where \mathbb{Z}_2^F is in the center of G_f . Thus fermion parity commutes with all elements of G_f and
 116 the quotient group $G_f/\mathbb{Z}_2^F = G$.

117 Let us denote by \hat{g} the operator corresponding to the group element $g \in G$ that acts on
 118 Hilbert space. In general it could be unitary or anti-unitary. The group G_f is then the set
 119 $\{(\pm 1)^{\hat{F}} \hat{g} \mid g \in G\}$ with the product rule between $(\eta_1)^{\hat{F}} \hat{g}$ and $(\eta_2)^{\hat{F}} \hat{h}$ (with $\eta_i = \pm 1$) given
 120 by

$$[(\eta_1)^{\hat{F}} \hat{g}][(\eta_2)^{\hat{F}} \hat{h}] = [\eta_1 \eta_2 \omega(g, h)]^{\hat{F}} \widehat{g h} \quad (3)$$

121 ω called the 2-cocycle is a function $\omega : G \times G \rightarrow \{+1, -1\}$ that satisfies

$$\omega(g, h) \omega(g h, k) = \omega(g, h k) \omega(h, k), \quad (4)$$

122 so that the multiplication is associative, and $\omega(e_G, e_G) = 1$, so that the identity element is well
 123 defined. Each inequivalent cocycle furnishes a distinct projective symmetry group. Thus PSGs
 124 are characterized by classes of inequivalent cocycles $[\omega]$ which form the second cohomology
 125 group $\mathcal{H}^2(G, \mathbb{Z}_2)$.

126 As an example, consider time reversal symmetry where $G = \mathbb{Z}_2^T = \{1, T\}$. In this case,
 127 $\mathcal{H}^2(\mathbb{Z}_2, \mathbb{Z}_2) = \mathbb{Z}_2$ and there are two PSGs characterized by the two inequivalent cocycles: (1)
 128 $\omega(T, T) = 1$ in which case $\hat{T}^2 = 1$, and (2) $\omega(T, T) = -1$ where $\hat{T}^2 = (-1)^{\hat{F}}$. In the first case
 129 $G_f = \mathbb{Z}_2 \times \mathbb{Z}_2$ while in the second $G_f = \mathbb{Z}_4$. Physically, the action of the different PSGs on
 130 the even particle number subspace is the same as that of the bosonic group G . The distinction
 131 appears in how G_f acts on the odd particle number subspace, in particular, the single particle
 132 subspace.

133 In general, one could have both unitary and anti-unitary symmetries but in this paper
 134 we will focus on *unitary* operators $\hat{g} \in G_f$, under which the fermion annihilation operator
 135 transforms as

$$\hat{g} \hat{c}_{k\alpha} \hat{g}^{-1} = [U^g(\mathbf{k})]_{\alpha\beta}^{\dagger} \hat{c}_{g\mathbf{k}\beta} \quad (5)$$

136 where \mathbf{k} is the (crystal) momentum, and the α labels spin, orbital/sublattice/band degrees of
 137 freedom (d.o.f.). Using $(-1)^{\hat{F}} \hat{c}_{k\alpha} (-1)^{\hat{F}} = -\hat{c}_{k\alpha}$ and eq. (3), we find that

$$U^g(h\mathbf{k}) U^h(\mathbf{k}) = \omega(g, h) U^{gh}(\mathbf{k}). \quad (6)$$

138 The U^g 's thus form a projective representation of G with coefficients in $\{\pm 1\}$. Equivalently,
 139 one can regard $\{\pm U^g \mid g \in G\}$ as a linear representation of G_f with $(-1)^{\hat{F}}$ represented by -1 .

140 2.2 Pairing Symmetry and Projective Representations

141 To be concrete, let us focus on the BdG Hamiltonian

$$\hat{H} = \hat{H}_0 + (\hat{H}_{\text{pair}} + \text{h.c.}) \quad (7)$$

142 where

$$\hat{H}_0 = \sum_{\alpha\beta;\mathbf{k}} \hat{c}_{\mathbf{k}\alpha}^\dagger h_{\alpha\beta}(\mathbf{k}) \hat{c}_{\mathbf{k}\beta} \quad (8)$$

143 is the kinetic energy that describes the normal state electronic dispersion, and

$$\hat{H}_{\text{pair}} = \sum_{\alpha\beta;\mathbf{k}} \hat{c}_{\mathbf{k}\alpha}^\dagger \Delta_{\alpha\beta}(\mathbf{k}) \hat{c}_{-\mathbf{k}\beta}^\dagger \quad (9)$$

144 describes the pairing. Fermi statistics dictates that $\Delta_{\alpha\beta}(\mathbf{k}) = -\Delta_{\beta\alpha}(-\mathbf{k})$.

145 Initially, we restrict ourselves for simplicity to situations where $\mathbf{SO}(3)_{\text{spin}}$ is *not* broken
146 spontaneously in the SC state. In this case, the SC state symmetry group G is of the form

$$G = \begin{cases} X \times \mathbf{SO}(3)_{\text{spin}} & \text{Weak SOC} \\ X & \text{Strong SOC} \end{cases} \quad (10)$$

147 where X is the point group of crystalline symmetries. In either case the pairing order parameter
148 $\Delta(\mathbf{k})$ forms a 1d linear representation of crystalline point group X . Moreover the relevant
149 fermionic PSGs are of the form $G_f \simeq (X_f \times \mathbf{SU}(2))/\mathbb{Z}_2$ and $G_f \simeq X_f$ for the weak and strong
150 SOC cases respectively where X_f is itself a central extension of X with respect to fermion parity.
151 In the first case, we get an $\mathbf{SU}(2)$ as a \mathbb{Z}_2 central extension of $\mathbf{SO}(3)_{\text{spin}}$ and a quotient by \mathbb{Z}_2
152 is required to take into account the "double-counting" of \mathbb{Z}_2^F . It is thus sufficient to look at
153 the central extensions of X . Later, in Section 5, we shall present a more general treatment and
154 discuss the case of ${}^3\text{He}$ where spin rotation is spontaneously broken in the SC state. In such
155 cases, the fermion symmetry group might have a more complicated form and it is no longer
156 sufficient to look at central extensions of the spatial part alone.

157 We now discuss *three* different projective representations of X and explore how these are
158 related. First, we begin with $X_f^0 = \{(\pm 1)^{\hat{F}} \hat{g}_0 \mid g \in X\}$ the PSG of X that preserves the kinetic
159 part of the BdG hamiltonian i.e., $\hat{g}_0 \hat{H}_0 \hat{g}_0^{-1} = \hat{H}_0$. The fermion operators then transform
160 according to the corresponding projective representation $U_0^g(\mathbf{k})$, defined by

$$\hat{g}_0 c_{\mathbf{k}\alpha} \hat{g}_0^{-1} = [U_0^g(\mathbf{k})]_{\alpha\beta}^\dagger \hat{c}_{g\mathbf{k}\beta}, \quad (11)$$

161 which preserves the normal state band structure

$$U_0^g(\mathbf{k}) h(\mathbf{k}) [U_0^g(\mathbf{k})]^\dagger = h(g\mathbf{k}). \quad (12)$$

162 We shall call X_f^0 the *normal state PSG* and denote the corresponding 2-cocycle by ω_0 . For
163 systems with weak SOC, crystalline symmetries do not act on the spin degrees of freedom and
164 the PSG is trivial in this case $\omega_0(g, h) = 1$ for any elements $g, h \in X$. In the presence of
165 strong SOC the projective representation is non-trivial with operations like two fold rotations
166 and mirror reflections now squaring to fermion parity, $\omega_0(C_2, C_2) = \omega_0(M, M) = -1$. This
167 becomes evident by looking at the forms of the projective representations in the two cases.

$$U_0^g(\mathbf{k}) = \begin{cases} u_{\text{orbital}}^g(\mathbf{k}) \otimes \mathbb{1}_{\text{spin}} & \text{weak SOC} \\ u_{\text{orbital}}^g(\mathbf{k}) \otimes e^{i\frac{\theta_g}{2} \hat{n}_g \cdot \vec{\sigma}} & \text{strong SOC} \end{cases} \quad (13)$$

168 where \hat{n}_g and θ_g are the rotation axis and angle associated with crystalline symmetry operation
169 $g \in X$.

170 Next, we note that the normal state PSG preserves the pairing term only up to a phase,
171 namely

$$\hat{g}_0 \hat{H}_{\text{pair}} \hat{g}_0^{-1} = e^{i\Phi_g} \hat{H}_{\text{pair}} \quad (14)$$

172 The phases $\{e^{i\Phi_g} \mid g \in X\}$ form a 1D *linear* representation of X , which we call the *pairing*
173 *symmetry* $\mathcal{R}_{\text{pair}}$. The phases Φ_g 's satisfy the equation

$$\Phi_g + \Phi_h = \Phi_{gh} + 2n\pi \quad (n \in \mathbb{Z}). \quad (15)$$

174 The pairing matrix $\Delta(\mathbf{k})$ satisfies

$$U_0^g(\mathbf{k})\Delta(\mathbf{k})[U_0^g(-\mathbf{k})]^T = e^{i\Phi_g} \Delta(g\mathbf{k}). \quad (16)$$

175 We see from eq. (14) that the PSG X_f^0 that leaves \hat{H}_0 invariant, fails to preserve the pairing
176 term. However the situation can be fixed as follows. We modify the transformation of the
177 fermions $\hat{g}' c_{k\alpha} \hat{g}'^{-1} = [\tilde{U}(\mathbf{k})]_{\alpha\beta}^\dagger \hat{c}_{k\beta}$ with

$$\tilde{U}^g(\mathbf{k}) = e^{-i\Phi_g/2} U_0^g(\mathbf{k}) \quad (17)$$

178 The kinetic part \hat{H}_0 , which is invariant under $U(1)$ phase rotations, is preserved by the modified
179 transformations as can be seen from (12). The new transformations are also symmetries of
180 the pairing term \hat{H}_{pair} as $\tilde{U}_g(\mathbf{k})$'s lead to eq. (16) without the phase factor $e^{i\Phi_g}$ appearing on
181 the right-hand side.

182 We thus define *SC state* PSG \tilde{X}_f that preserves the full BdG Hamiltonian by

$$\tilde{X}_f = \left\{ (\pm 1)^{\hat{f}} \hat{g}' = (\pm 1)^{\hat{f}} e^{-i(\Phi_g/2)\hat{f}} \hat{g} \mid g \in X \right\} \quad (18)$$

183 This PSG is characterized by the 2-cocycle $\tilde{\omega}$.

184 The last step here is to look at the relation between the normal and the superconducting
185 state PSGs, or equivalently, between their cocycles ω_0 and $\tilde{\omega}$. The phases $\{e^{-i\Phi_g/2} \mid g \in X\}$
186 form a 1D *projective* representation of X , which we call \mathcal{R}_Φ . This follows from (15) by observ-
187 ing that $e^{-i\Phi_g/2} e^{-i\Phi_h/2} = (-1)^n e^{-i\Phi_{gh}/2}$. From eqn.(17) one concludes that the cocycle ω_Φ
188 associated with \mathcal{R}_Φ satisfies

$$\tilde{\omega}(g, h) = \omega_\Phi(g, h) \omega_0(g, h) \quad (19)$$

189 To summarize, we encountered the following projective representations and their associ-
190 ated cocycles which define the corresponding PSG's:

$$\text{Normal state: } U_0^g(h\mathbf{k}) U_0^h(\mathbf{k}) = \omega_0(g, h) U_0^{gh}(\mathbf{k}) \quad (20a)$$

$$\mathcal{R}_\Phi: e^{-i\Phi_g/2} e^{-i\Phi_h/2} = \omega_\Phi(g, h) e^{-i\Phi_{gh}/2} \quad (20b)$$

$$\text{SC state: } \tilde{U}^g(h\mathbf{k}) \tilde{U}^h(\mathbf{k}) = \tilde{\omega}(g, h) \tilde{U}^{gh}(\mathbf{k}) \quad (20c)$$

191 Eq. (17) relates the three projective representations and eq. (19) relates their cocycles.

192 Given the normal state PSG and the pairing symmetry of the SC state, one can use the for-
193 malism described above to determine the SC state PSG. This is achieved in the following steps.
194 Pairing symmetry being a 1D linear representation, $\mathcal{R}_{\text{pair}}$ can be read off from the character

195 table of X . Taking the square roots of the characters one obtains the 1D projective representa-
 196 tion \mathcal{R}_Φ and its cocycle ω_Φ . With the normal state PSG known eq. (19) gives the SC state PSG
 197 while eq. (17) gives the SC state projective representation explicitly. Thus knowing the pairing
 198 symmetry enables us to find the SC state PSG that preserves the BdG Hamiltonian. In the next
 199 Section we turn to the inverse problem of constraining the pairing symmetry, knowing the SC
 200 state PSG.

201 3 Constraints on the pairing symmetry by the PSG

202 One longstanding experimental challenge in the field of superconductivity is how to unambigu-
 203 ously determine the pairing symmetry of a superconductor material, based on experimental
 204 measurements. Since all fermionic excitations in the superconductor form a linear represen-
 205 tation of the SC state PSG \tilde{X}_f , the low-temperature physical properties of the superconductor
 206 completely depend on the PSG. For example, as will be discussed in section 4.4, the topological
 207 properties of the SC phase are determined by the PSG. As a result, it seems plausible to detect
 208 the SC state PSG \tilde{X}_f using various experimental probes, which we will clarify in future publi-
 209 cations. This observation motivates us to answer the following question: given a SC state PSG
 210 \tilde{X}_f , what are the pairing symmetries compatible with \tilde{X}_f ? In other words, how does a given
 211 PSG constrain the possible pairing symmetry in a superconductor? The answer to this question
 212 will allow us to constrain or even determine the pairing symmetry of a SC, by experimentally
 213 detecting its PSG.

214 Based on the discussions in section 2.2, we can readily derive the constraints on the pairing
 215 symmetry by the PSG from relations (17) and (19). Specifically, given a SC state PSG \tilde{X}_f and
 216 its associated 2-cocycle $\tilde{\omega}$, we can follow the steps listed below to obtain the possible pairing
 217 symmetries \mathcal{R}_{pair} in (14)-(16):

218 (1) Given the crystalline point group X , determine the normal state PSG X_f^0 and associated
 219 2-cocycle $\{\omega_0\}$ of the normal-state symmetry transformations $\{U_0^g | g \in X\}$. This only depends
 220 on the strength of SOCs in the system.

221 (2) Compute the 2-cocycle $\{\omega_\Phi\}$ from $\{\omega_0\}$ and $\{\tilde{\omega}\}$ from relation (19).

222 (3) Obtain all one-dimensional (1d) projective representations $\{\mathcal{R}_\Phi(g) | g \in X\}$ of the crys-
 223 talline symmetry group X compatible with 2-cocycle $\{\omega_\Phi\}$ obtained in step (2), satisfying

$$\mathcal{R}_\Phi(g)\mathcal{R}_\Phi(h) = \omega_\Phi(g, h)\mathcal{R}_\Phi(gh) \quad (21)$$

224 (5) For each 1d projective representation $\mathcal{R}_\Phi(g)$ obtained in step (3), compute the 1d
 225 linear representation

$$\mathcal{R}_{pair}(g) = [\mathcal{R}_\Phi(g)]^{-2} \quad (22)$$

226 of the pairing order parameter. The collection of all results $\{\mathcal{R}_{pair}\}$ correspond to all the pos-
 227 sible pairing symmetries compatible with the PSG \tilde{X}_f .

228
 229 We have applied our general computational scheme to the case of 32 crystalline point
 230 groups for both strong SOCs and negligible (weak) SOCs. Our analysis focuses on SC order
 231 parameters that are spatially uniform, where the Cooper pairs condense in a state with zero
 232 center of mass momentum. Lattice translations then act trivially on the SC state and leave
 233 the BdG Hamiltonian invariant, and it is sufficient for us to focus on point group symmetry
 234 alone. Most experimentally relevant systems exhibit spatially uniform pairing (in the absence
 235 of strong disorder). It is only in exceptional circumstances – under very limited range of ex-
 236 ternal parameters in a few systems – that one expects the SC order parameter to sponta-
 237 neously break translational symmetry, e.g., in FFLO or pair density wave state. In such cases,

238 we would need to investigate space group symmetries which we leave for future investiga-
 239 tion. Group cohomology and projective representation calculations are performed using the
 240 GAP computer algebra program [24]. The correspondence between fermion PSGs \mathbf{G}_f and the
 241 representations \mathcal{R}_{pair} of the superconducting order parameter is established for all 32 point
 242 groups, and the results are summarized in Table. 7 in Appendix B.

243 4 Examples

244 We now demonstrate the above formalism for different point groups. In section 4.1 we con-
 245 sider systems with tetragonal symmetry. Cuprates and ruthenates which belong to this cate-
 246 gory have point group D_{4h} . But for instance in cuprates, only the Cu-O plane is relevant for
 247 superconductivity and it suffices to consider the point group C_{4v} for the purpose of illustration.
 248 In section 4.2 we treat systems with hexagonal symmetry. A discussion of superconductivity
 249 on a honeycomb lattice is followed by a remark on how our formalism can be applied to the
 250 case of magic angle twisted bilayer graphene. In section 4.3 we discuss superconductivity in
 251 transition metal dichalcogenides with trigonal point group C_{3v} .

252 The purpose of these examples is two-fold. First, we present a detailed account of how
 253 the table in appendix B is constructed and what information can be extracted from it. Sec-
 254 ond, we make a direct connection with real physical systems by producing examples of order
 255 parameters $\Delta_{\alpha\beta}(\mathbf{k})$ for each 1D irrep (pairing symmetry) of the relevant point group.

256 As mentioned earlier we shall restrict ourselves to cases where there is no additional break-
 257 ing of spin rotation symmetry when going from the normal to the SC phase. Examples which
 258 do not fit in this category, like superfluid He^3 , are discussed in the section 5.2.

259 4.1 Tetragonal Symmetry

260 To be concrete, consider a two dimensional square lattice in the xy plane. The relevant crys-
 261 talline point group is $X = C_{4v}$. The group is generated by a rotation by $\pi/2$ about the z -axis,
 262 C_4 and reflection about a vertical mirror in the yz plane, σ_v . The action of these operations
 263 can be summarized as

$$(x, y, z) \xrightarrow{C_4} (-y, x, z) \quad (23a)$$

$$(x, y, z) \xrightarrow{\sigma_v} (-x, y, z) \quad (23b)$$

264 The group law is captured by the relations $C_4^4 = e$, $\sigma_v^2 = e$ and $C_4^3\sigma_v = \sigma_v C_4$. Equivalently
 265 the group is generated by the vertical mirror σ_v and the diagonal mirror $\sigma_d = \sigma_v C_4$. Since
 266 $\sigma_v^2 = \sigma_d^2 = e$, these could have either $+1$ or -1 characters in a 1D irrep. Consequently there
 267 are four 1D irreps for this group, each labeled uniquely by a tuple of σ_v and σ_d characters,
 268 $(e^{i\Phi_{\sigma_v}}, e^{i\Phi_{\sigma_d}})$ taking values $(\pm 1, \pm 1)$. The characters for the other group elements can then be
 269 obtained using the group laws. In particular, it follows from $C_2 = (\sigma_d \sigma_v)^2$ that the character
 270 for the two-fold rotation in the four 1D irreps is $+1$.

271 Let us now turn our attention to the possible fermion PSGs for this group. From the group
 272 cohomology calculation we have $\mathcal{H}^{(2)}(C_{4v}, \mathbb{Z}_2) = \mathbb{Z}_2^3$, corresponding to eight inequivalent
 273 classes of 2-cocycles for this group characterized by the 3-tuple

$$(\omega(C_2, C_2), \omega(\sigma_v, \sigma_v), \omega(\sigma_d, \sigma_d)) = (\pm 1, \pm 1, \pm 1). \quad (24)$$

274 The eight PSGs are thus distinguished on the basis of whether the two fold rotation, C_2 and
 275 the two mirrors σ_v and σ_d square to ± 1 .

276 We are now in a position to explore the connection between the pairing symmetries and
 277 fermion PSGs for this group. First consider the case when because of weak spin-orbit coupling

278 there is spin rotation invariance in the normal state. The symmetry operations that preserve
 279 the kinetic energy act only on the momentum label, keeping the spin label unaltered. Denoted
 280 by superscript $\mathbf{0}$ these are

$$\hat{C}_2^{\mathbf{0}} \hat{c}_{k\alpha} (\hat{C}_2^{\mathbf{0}})^{-1} = \hat{c}_{C_2 k\alpha} \quad (25a)$$

$$\hat{\sigma}_v^{\mathbf{0}} \hat{c}_{k\alpha} (\hat{\sigma}_v^{\mathbf{0}})^{-1} = \hat{c}_{\sigma_v k\alpha} \quad (25b)$$

$$\hat{\sigma}_d^{\mathbf{0}} \hat{c}_{k\alpha} (\hat{\sigma}_d^{\mathbf{0}})^{-1} = \hat{c}_{\sigma_d k\alpha} \quad (25c)$$

281 Consequently, the normal state PSG is trivial and

$$(\omega_0(C_2, C_2), \omega_0(\sigma_v, \sigma_v), \omega_0(\sigma_d, \sigma_d)) = (+1, +1, +1). \quad (26)$$

282 Given the assumption that pairing does not break spin rotation invariance in the superconduct-
 283 ing phase, condensation takes place in the singlet channel. This enforces the pair wavefunction
 284 to be of the form

$$\Delta_{\alpha\beta}(\mathbf{k}) = \Psi(\mathbf{k})(i\sigma_y)_{\alpha\beta} \quad (27)$$

285 where α, β are spin labels and Pauli exclusion constrains the orbital part of the pair wavefunc-
 286 tion to obey $\Psi(-\mathbf{k}) = \Psi(\mathbf{k})$. As has been discussed in detail in previous sections, the phases
 287 $\{e^{i\Phi_s}\}$ acquired by the pairing term in (9), when acted upon by the operations in (25), consti-
 288 tute a 1D linear irrep of C_{4v} which we refer to as pairing symmetry $\mathcal{R}_{\text{pair}}$. We also learnt that
 289 (25) must be modified by compensating phase rotations so as to make the new transformations
 290 symmetries of the BdG hamiltonian.

291 Different pairing symmetries modify the normal state transformations in (25) differently.
 292 When the pairing symmetry is A_1 , which is the case when say $\Psi(\mathbf{k})$ is a constant Ψ_0 indepen-
 293 dent of \mathbf{k} , the normal state transformations already preserve the pairing term and no modifi-
 294 cation is necessary. The normal and the SC state PSGs are the same in this case. If however
 295 $\Psi(\mathbf{k}) = \Psi_0(k_x^2 - k_y^2)$, σ_v keeps the pairing term unchanged whereas under σ_d (or equiva-
 296 lently under C_4) it acquires a negative sign. The pairing symmetry in this case is B_1 , labeled
 297 by $(e^{i\Phi_{\sigma_v}}, e^{i\Phi_{\sigma_d}}) = (+1, -1)$. Eqn. (25c) now has to be modified by a factor of i appearing on
 298 the right hand side, i.e, the modified σ_d must take $\hat{c}_{k\alpha}$ to $i\hat{c}_{\sigma_d k\alpha}$.

299 For a generic irrep, when the orbital part transforms as

$$\Psi(\mathbf{k}) = e^{i\Phi_s} \Psi(g\mathbf{k}) \quad (28)$$

300 the compensating phases are the square roots of the characters of the relevant irrep. Denoted
 301 with primes, the transformations that preserve the BdG hamiltonian are then

$$\hat{C}'_2 \hat{c}_{k\alpha} (\hat{C}'_2)^{-1} = e^{i\Phi_{C_2}/2} \hat{c}_{C_2 k\alpha} \quad (29a)$$

$$\hat{\sigma}'_v \hat{c}_{k\alpha} (\hat{\sigma}'_v)^{-1} = e^{i\Phi_{\sigma_v}/2} \hat{c}_{\sigma_v k\alpha} \quad (29b)$$

$$\hat{\sigma}'_d \hat{c}_{k\alpha} (\hat{\sigma}'_d)^{-1} = e^{i\Phi_{\sigma_d}/2} \hat{c}_{\sigma_d k\alpha} \quad (29c)$$

302 For instance, for A_1 and B_1 pairing symmetries, $(e^{i\Phi_{C_2}/2}, e^{i\Phi_{\sigma_v}/2}, e^{i\Phi_{\sigma_d}/2})$ can be chosen to be
 303 $(1, 1, 1)$ and $(1, 1, i)$ respectively.

304 The resulting SC state PSGs are different across pairing symmetries. For the A_1 irrep, the
 305 SC state PSG is trivial. With the diagonal mirror now squaring to fermion parity, the SC state
 306 PSG for B_1 becomes

$$(\tilde{\omega}(C_2, C_2), \tilde{\omega}(\sigma_v, \sigma_v), \tilde{\omega}(\sigma_d, \sigma_d)) = (+1, +1, -1). \quad (30)$$

307 As elaborated in previous sections, the reason for this is best understood once we recognize
 308 that the compensating phases, $\{e^{-i\Phi_g/2} \mid g \in X\}$ form a 1D projective representation, \mathcal{R}_Φ

$\mathcal{R}_{\text{pair}}$	$\Psi(\mathbf{k})$	\mathcal{R}_{Φ}	ω_{Φ}	$\tilde{\omega}$
$A_1 : (+1, +1)$	1	$(\pm 1, \pm 1, \pm 1)$	$(+1, +1, +1)$	$(+1, +1, +1)$
$A_2 : (-1, -1)$	$k_x k_y (k_x^2 - k_y^2)$	$(\pm 1, \pm i, \pm i)$	$(+1, -1, -1)$	$(+1, -1, -1)$
$B_1 : (+1, -1)$	$k_x^2 - k_y^2$	$(\pm 1, \pm 1, \pm i)$	$(+1, +1, -1)$	$(+1, +1, -1)$
$B_2 : (-1, +1)$	$k_x k_y$	$(\pm 1, \pm i, \pm 1)$	$(+1, -1, +1)$	$(+1, -1, +1)$

Table 1: Tetragonal Symmetry ($X = C_{4v}$) with weak SOC. Here $\mathcal{R}_{\text{pair}} \equiv (e^{i\Phi_{\sigma_v}}, e^{i\Phi_{\sigma_d}})$ and $\mathcal{R}_{\Phi} \equiv (e^{-i\Phi_{C_2}/2}, e^{-i\Phi_{\sigma_v}/2}, e^{-i\Phi_{\sigma_d}/2})$

309 of X . The corresponding cocycle given by ω_{Φ} could be different for the different pairing
310 symmetries. For example,

$$(\omega_{\Phi}(C_2, C_2), \omega_{\Phi}(\sigma_v, \sigma_v), \omega_{\Phi}(\sigma_d, \sigma_d)) = (1^2, 1^2, 1^2) \text{ and } (1^2, 1^2, i^2 = -1) \quad (31)$$

311 for the A_1 and B_1 irreps respectively. Pairing symmetry thus dictates ω_{Φ} which through (19) in
312 turn decides the SC state PSG. Table 1 summarizes the results of the above analysis for C_{4v} with
313 weak SOC. For each irrep, we give an example of $\Psi(\mathbf{k})$, show the 1D projective representation
314 of the compensating phases \mathcal{R}_{Φ} , the cocycle ω_{Φ} and finally the SC state PSG $\tilde{\omega}$.

315

316 In the presence of strong spin orbit coupling, the transformations that preserve the kinetic
317 energy are combined spatial and spin rotation. A rotation by angle θ about $\hat{\mathbf{n}}$ transforms
318 the spinor by $e^{-i\frac{\theta}{2}(\hat{\mathbf{n}} \cdot \boldsymbol{\sigma})}$ while inversion leaves it unaffected. A mirror could be viewed as a
319 combination of inversion and a two fold rotation about an axis perpendicular to the mirror
320 plane. For instance, reflection about the yz mirror plane is then effectively a two-fold rotation
321 about the x axis and would be implemented by $-i\sigma_x$ in the spinor basis. The transformations
322 that preserve kinetic energy are

$$\hat{C}_2^0 \hat{c}_{k\alpha} (\hat{C}_2^0)^{-1} = [-i\sigma_z]_{\alpha\beta} \hat{c}_{C_2 k\beta} \quad (32a)$$

$$\hat{\sigma}_v^0 \hat{c}_{k\alpha} (\hat{\sigma}_v^0)^{-1} = [-i\sigma_x]_{\alpha\beta} \hat{c}_{\sigma_v k\beta} \quad (32b)$$

$$\hat{\sigma}_d^0 \hat{c}_{k\alpha} (\hat{\sigma}_d^0)^{-1} = [-i\hat{\mathbf{n}}' \cdot \boldsymbol{\sigma}]_{\alpha\beta} \hat{c}_{\sigma_d k\beta} \quad (32c)$$

323 Where $\hat{\mathbf{n}}' = (\hat{x} - \hat{y})/\sqrt{2}$ and the Einstein summation convention is implied. With two fold
324 rotations and hence mirrors now squaring to fermion parity, the normal state PSG is

$$(\omega_0(C_2, C_2), \omega_0(\sigma_v, \sigma_v), \omega_0(\sigma_d, \sigma_d)) = (-1, -1, -1). \quad (33)$$

325 In the absence of spin rotation invariance in the normal state, the pair wavefunction is an
326 admixture of singlet and triplet parts and takes the form

$$\Delta_{\alpha\beta}(\mathbf{k}) = \Psi(\mathbf{k}) [i\sigma_y]_{\alpha\beta} + \mathbf{d}(\mathbf{k}) \cdot [\vec{\sigma}(i\sigma_y)]_{\alpha\beta} \quad (34)$$

327 where Pauli exclusion now requires the three component complex vector \mathbf{d} to obey $\mathbf{d}(\mathbf{k}) = -\mathbf{d}(-\mathbf{k})$.
328 Since the C_2 character in all the one dimensional irreps is $+1$, we must have

$$(i\sigma_z)\Delta(\mathbf{k})(i\sigma_z)^T = \Delta(C_2\mathbf{k}) = \Delta(-\mathbf{k}), \quad (35)$$

329 where the last equality follows from the fact that we are in two spatial dimensions. It is
330 immediately seen that this implies $\mathbf{d}_z(\mathbf{k}) = \mathbf{d}_z(-\mathbf{k})$ and the only way this could be consistent
331 with the constraint imposed by Pauli exclusion is when $\mathbf{d}_z(\mathbf{k}) = \mathbf{0}$. Similarly, by effecting

$\mathcal{R}_{\text{pair}}$	$\mathbf{d}(\mathbf{k})$	\mathcal{R}_{Φ}	$\boldsymbol{\omega}_{\Phi}$	$\tilde{\boldsymbol{\omega}}$
$A_1 : (+1, +1)$	$k_y \hat{x} - k_x \hat{y}$	$(\pm 1, \pm 1, \pm 1)$	$(+1, +1, +1)$	$(-1, -1, -1)$
$A_2 : (-1, -1)$	$k_x \hat{x} + k_y \hat{y}$	$(\pm 1, \pm i, \pm i)$	$(+1, -1, -1)$	$(-1, +1, +1)$
$B_1 : (+1, -1)$	$k_y \hat{x} + k_x \hat{y}$	$(\pm 1, \pm 1, \pm i)$	$(+1, +1, -1)$	$(-1, -1, +1)$
$B_2 : (-1, +1)$	$k_x \hat{x} - k_y \hat{y}$	$(\pm 1, \pm i, \pm 1)$	$(+1, -1, +1)$	$(-1, +1, -1)$

Table 2: Tetragonal Symmetry ($X = C_{4v}$) with strong SOC. Here $\mathcal{R}_{\text{pair}} \equiv (e^{i\Phi_{\sigma_v}}, e^{i\Phi_{\sigma_d}})$ and $\mathcal{R}_{\Phi} \equiv (e^{-i\Phi_{C_2}/2}, e^{-i\Phi_{\sigma_v}/2}, e^{-i\Phi_{\sigma_d}/2})$

332 transformations for σ_v and σ_d on the pairing term we conclude that to transform as a 1D irrep
 333 labeled by the characters $(e^{i\Phi_{\sigma_v}}, e^{i\Phi_{\sigma_d}})$, the non-zero components of the \mathbf{d} vector, must satisfy
 334

$$(+d_x(\mathbf{k}), -d_y(\mathbf{k})) = e^{i\Phi_{\sigma_v}} (d_x(\sigma_v \mathbf{k}), d_y(\sigma_v \mathbf{k})) \quad (36a)$$

$$(-d_y(\mathbf{k}), -d_x(\mathbf{k})) = e^{i\Phi_{\sigma_d}} (d_x(\sigma_d \mathbf{k}), d_y(\sigma_d \mathbf{k})) \quad (36b)$$

335 and $\Psi(\mathbf{k})$, like in the case for weak SOC, satisfies (28). Table 2 provides examples of the $\mathbf{d}(\mathbf{k})$
 336 vector for each pairing symmetry. All of these examples belong to a $(p+ip) \uparrow + (p-ip) \downarrow$ type
 337 SC. As before, square roots of the characters of the 1D irrep form the compensating phases
 338 which modify the transformations in (32) and different SC state PSGs are obtained for the four
 339 pairing symmetries as outlined in table 2.

340

341 A few comments are in order. First, comparing the two tables we observe that since the 1D
 342 projective representation \mathcal{R}_{Φ} formed by the compensating phases and the corresponding co-
 343 cycle $\boldsymbol{\omega}_{\Phi}$ depend solely on the pairing symmetry, the correspondence between $\mathcal{R}_{\text{pair}}$ and $\boldsymbol{\omega}_{\Phi}$
 344 is identical irrespective of the strength of SOC. The difference in the normal state PSG $\boldsymbol{\omega}_0$
 345 accounts for the difference in the SC state PSG $\tilde{\boldsymbol{\omega}}$ between the corresponding rows of tables 1
 346 and 2.

347 Second, a question arises as to why only four of the eight PSGs appear in each of the two ta-
 348 bles. The answer is apparent once we observe that the $\boldsymbol{\omega}_{\Phi}$ column only contains the four PSGs
 349 with $\boldsymbol{\omega}_{\Phi}(C_2, C_2) = +1$. This is easily seen as follows. Group law tells us that $\sigma_v \sigma_d = C_2 \sigma_d \sigma_v$.
 350 Then for any 1D projective representation ϕ , we must have $\phi(\sigma_v) \phi(\sigma_d) = \pm \phi(C_2) \phi(\sigma_d) \phi(\sigma_v)$.
 351 Since ϕ 's are all non-zero complex numbers, dividing both sides by $\phi(\sigma_v) \phi(\sigma_d)$ gives $\phi(C_2) = \pm 1$
 352 and hence $\boldsymbol{\omega}_{\Phi}(C_2, C_2) = +1$. In other words PSGs with $\boldsymbol{\omega}(C_2, C_2) = -1$ cannot have a 1D rep-
 353 resentation.

354 Finally, both tables show a one-one correspondence between the four pairing symmetries
 355 and four out of the eight possible PSGs. Knowledge of the SC state PSG (from topological or
 356 spectroscopic properties) thus uniquely determines the pairing symmetry.

357 4.2 Hexagonal Symmetry

358 Consider a two dimensional honeycomb lattice in the xy plane with a plaquet center chosen
 359 as the origin and the x -axis passing through a bond center. A six fold rotation about the z -axis,
 360 C_6 and a reflection about a vertical mirror σ_v in the yz plane then transform the coordinates
 361 as

$$(x, y) \xrightarrow{C_6} \left(\frac{1}{2}x - \frac{\sqrt{3}}{2}y, \frac{1}{2}y + \frac{\sqrt{3}}{2}x \right) \quad (37a)$$

$$(x, y) \xrightarrow{\sigma_v} (-x, y) \quad (37b)$$

362 C_6 and σ_v generate the point group C_{6v} . It comprises of six rotations and six mirror reflections
 363 and the group law is captured by the relations $C_6^6 = e$, $\sigma_v^2 = e$ and $C_6\sigma_v C_6 = \sigma_v$. From these
 364 relations it is evident that the C_6 and σ_v characters in a 1D linear irrep of C_{6v} could only be
 365 ± 1 . Indeed, there are four 1D irreps for this group labeled by $(e^{i\Phi_{C_6}}, e^{i\Phi_{\sigma_v}}) = (\pm 1, \pm 1)$. Here
 366 we note that not only is the group D_6 isomorphic to C_{6v} , but has indistinguishable action in
 367 two spatial dimensions. In D_6 , the two-fold rotation about the in-plane y -axis, C_{2y} assumes
 368 the role of σ_v in C_{6v} . Thus, when we are strictly in two spatial dimensions, C_{6v} and D_6 can
 369 be used interchangeably.

370 Since $\mathcal{H}^{(2)}(C_{6v}, \mathbb{Z}_2) = \mathbb{Z}_2^3$, there are eight possible PSGs distinguished on the basis of
 371 whether C_2 and σ_v square to $+1$ or -1 and whether they commute or anti-commute. The
 372 classes of 2-cocycles are labeled by

$$\left(\omega(C_2, C_2), \omega(\sigma_v, \sigma_v), \frac{\omega(C_2, \sigma_v)}{\omega(\sigma_v, C_2)} \right) = (\pm 1, \pm 1, \pm 1). \quad (38)$$

373 We discuss the case when the normal and the SC states have spin rotation invariance.
 374 Denoted by the superscript $\mathbf{0}$, the transformations that preserve the kinetic energy are

$$\hat{C}_6^{\mathbf{0}} \hat{c}_{k\alpha s} (\hat{C}_6^{\mathbf{0}})^{-1} = (\tau_x)_{\alpha\beta} \hat{c}_{C_6 k \beta s} \quad (39a)$$

$$\hat{\sigma}_v^{\mathbf{0}} \hat{c}_{k\alpha s} (\hat{\sigma}_v^{\mathbf{0}})^{-1} = \hat{c}_{\sigma_v k \alpha s} \quad (39b)$$

375 Where α, β are sub-lattice labels, s labels spin and $\vec{\tau}$ denotes Pauli matrices in the sub-lattice
 376 space. The momentum \mathbf{k} is measured from the Γ point of the Brillouin zone. The normal state
 377 PSG is trivial with

$$\left(\omega_0(C_2, C_2), \omega_0(\sigma_v, \sigma_v), \frac{\omega_0(C_2, \sigma_v)}{\omega_0(\sigma_v, C_2)} \right) = (+1, +1, +1). \quad (40)$$

378 Here we consider a generic situation where both the bands participate in pairing and we
 379 express the pair wavefunction in the sub-lattice basis. If however we have a weak coupling
 380 scenario in which only a single band takes part in pairing, it is more convenient to express
 381 the pair wavefunction in the active band basis. For the present case, consistent with Pauli
 382 exclusion, the spin singlet wave function has the form

$$[\Delta(\mathbf{k})]_{\alpha s \beta s'} = \Psi_{\alpha\beta}(\mathbf{k})(i\sigma_y)_{ss'} \quad (41)$$

383 where $\Psi_{\alpha\beta}(\mathbf{k}) = \Psi_{\beta\alpha}(-\mathbf{k})$. For the pairing term to transform as the irrep $(e^{i\Phi_{C_6}}, e^{i\Phi_{\sigma_v}})$ under
 384 (39), $\Psi_{\alpha\beta}(\mathbf{k})$ satisfies

$$(\tau_x)_{\alpha\gamma} \Psi_{\gamma\delta}(\mathbf{k}) (\tau_x)_{\beta\delta} = e^{i\Phi_{C_6}} \Psi_{\alpha\beta}(C_6 \mathbf{k}) \quad (42a)$$

$$\Psi_{\alpha\beta}(\mathbf{k}) = e^{i\Phi_{\sigma_v}} \Psi_{\alpha\beta}(\sigma_v \mathbf{k}) \quad (42b)$$

385 In table 3 we provide examples of $\Psi_{\alpha\beta}(\mathbf{k})$ satisfying (42) for each pairing symmetry. The
 386 compensating phases $(e^{-i\Phi_{C_6}/2}, e^{-i\Phi_{\sigma_v}/2})$ forming the 1D projective representation \mathcal{R}_{Φ} and
 387 the corresponding 2-cocycle ω_{Φ} are also tabulated. A product of ω_{Φ} and ω_0 then gives the
 388 SC state PSG $\tilde{\omega}$. The four pairing symmetries correspond to four distinct $\tilde{\omega}$ s. The SC state
 389 PSG thus uniquely determines the pairing symmetry for this point group. Like in the previous
 390 case, only four out of the eight possible PSGs appear in table 3. Inspecting the ω_{Φ} column
 391 we observe that it only has entries with $\omega_{\Phi}(C_2, \sigma_v)/\omega_{\Phi}(\sigma_v, C_2) = +1$. Since complex num-
 392 bers always commute, it is impossible to get a 1D projective representation of C_{6v} where
 393 $\omega_{\Phi}(C_2, \sigma_v)/\omega_{\Phi}(\sigma_v, C_2) = -1$.

\mathcal{R}_{pair}	$\Psi_{AA}(\mathbf{k})$	$\Psi_{BB}(\mathbf{k})$	$\Psi_{AB}(\mathbf{k})$	\mathcal{R}_Φ	ω_Φ	$\tilde{\omega}$
$A_1 = (+1, +1)$	Δ_0	Δ_0	Δ'_0	$(\pm 1, \pm 1)$	$(+1, +1, +1)$	$(+1, +1, +1)$
$A_2 = (+1, -1)$	$\Delta_0 f(\mathbf{k})$	$\Delta_0 f(\mathbf{k})$	$\Delta'_0 g(\mathbf{k})$	$(\pm 1, \pm i)$	$(+1, -1, +1)$	$(+1, -1, +1)$
$B_1 = (-1, +1)$	Δ_0	$-\Delta_0$	0	$(\pm i, \pm 1)$	$(-1, +1, +1)$	$(-1, +1, +1)$
$B_2 = (-1, -1)$	$\Delta_0 f(\mathbf{k})$	$-\Delta_0 f(\mathbf{k})$	0	$(\pm i, \pm i)$	$(-1, -1, +1)$	$(-1, -1, +1)$

Table 3: Hexagonal symmetry ($X = C_{6v}$) with weak SOC. Here $\mathcal{R}_{pair} = (e^{i\Phi_{c_6}}, e^{i\Phi_{\sigma_v}})$, $\mathcal{R}_\Phi = (e^{-i\Phi_{c_6}/2}, e^{-i\Phi_{\sigma_v}/2})$. Also, $f(\mathbf{k}) = k_x k_y (k_x^2 - 3k_y^2)(k_y^2 - 3k_x^2)$ and $g(\mathbf{k}) = k_x(3k_y^2 - k_x^2)$

394 We end this subsection discussing superconductivity in magic angle twisted bilayer graphene
395 (MATBG) where the pairing symmetry is still not known although there has been some theo-
396 retical proposals [25]. The experimental observation of nematicity in the SC state [26], shows
397 that the normal state D_6 symmetry, is spontaneously broken in the SC state. Thus condensa-
398 tion must take place either in the E_1 or the E_2 irrep of D_6 . As pointed out in the introduction,
399 if it were any of the 1D irreps, the pair wavefunction would be invariant under D_6 up-to a
400 phase rotation, and the SC state would not show the observed nematicity. This corresponds
401 to the orbital part being a p -wave for the E_1 irrep or a d -wave for the E_2 irrep in the pair
402 wavefunction proposed in [25]. The residual symmetry in the SC state is the two-fold rotation
403 about z -axis, $X = C_{2z}$. Since the E_1 irrep (p -wave) has a C_2 character -1 and the E_2 irrep
404 (d -wave) has a C_2 character $+1$, these correspond to the two 1D irreps of X . There is a one
405 to one correspondence between \mathcal{R}_{pair} and PSGs for X as shown in table B and thus the two
406 possible pairing symmetries would give two distinct SC state PSGs.

407 4.3 Trigonal Symmetry

408 Like in the previous subsection, we consider the honeycomb lattice in the xy plane except
409 now two different species occupy the A and B sub-lattices. Such is the case, for example, in a
410 mono-layer transition metal dichalcogenide (TMD). The resulting point group C_{3v} is generated
411 by a three-fold rotation about the z -axis (C_3) and reflection about a vertical mirror in the yz
412 plane (σ_v) which act on the coordinates as

$$(x, y) \xrightarrow{C_3} \left(-\frac{1}{2}x - \frac{\sqrt{3}}{2}y, -\frac{1}{2}y + \frac{\sqrt{3}}{2}x \right) \quad (43a)$$

$$(x, y) \xrightarrow{\sigma_v} (-x, y) \quad (43b)$$

413 The relations $C_3^3 = \sigma_v^2 = e$ and $C_3\sigma_v C_3 = \sigma_v$ capture the group law. There are two 1D irreps
414 for this group with $e^{i\Phi_{\sigma_v}} = \pm 1$ and two PSGs with σ_v squaring to unity in one and to the
415 fermion parity in the other, $\omega(\sigma_v, \sigma_v) = \pm 1$

416 In TMDs, the presence of strong Ising SOC breaks spin rotation invariance [27]. Hole dop-
417 ing away from charge neutrality creates small Fermi surface pockets at the K and K' valleys.
418 Denoted by superscript 0 , the symmetry operations that preserve the kinetic energy act on the
419 fermion operator $\hat{c}_{\mathbf{k}\nu s}$ for the active band as

$$\hat{C}_3^0 \hat{c}_{\mathbf{k}\nu s} (\hat{C}_3^0)^{-1} = \left[e^{-i\frac{\pi}{3}\sigma_z} \right]_{ss'} \hat{c}_{C_3\mathbf{k}\nu s'} \quad (44a)$$

$$\hat{\sigma}_v^0 \hat{c}_{\mathbf{k}\nu s} (\hat{\sigma}_v^0)^{-1} = [\tau_x]_{\nu\nu'} [i\sigma_x]_{ss'} \hat{c}_{\sigma_v\mathbf{k}\nu' s'} \quad (44b)$$

420 where ν is the valley and s is the spin label and momentum \mathbf{k} is measured from the K or K'
421 point. Pauli matrices $\vec{\sigma}$ and $\vec{\tau}$ act on spin and valley spaces respectively. The normal state PSG
422 is thus described by the cocycle $\omega_0(\sigma_v, \sigma_v) = -1$.

$\mathcal{R}_{pair} \equiv e^{i\Phi_{\sigma_v}}$	$\Psi(\mathbf{k})$	$\mathcal{R}_{\Phi} \equiv e^{-i\Phi_{\sigma_v}/2}$	ω_{Φ}	$\tilde{\omega}$
$A_1 = +1$	Δ_0	± 1	$+1$	-1
$A_2 = -1$	$\Delta_0 k_y (3k_x^2 - k_y^2)$	$\pm i$	-1	$+1$

Table 4: Trigonal Symmetry ($X = C_{3v}$) with strong SOC

423 To ensure the Cooper pair has a zero center of mass momentum, pairing must be inter-
 424 valley. Because of time reversal invariance, the Fermi surface pockets at opposite valleys have
 425 oppositely polarized spins. If the spin polarization is $\sigma_z = +1$ in the K valley ($\tau_z = +1$),
 426 then it is along $\sigma_z = -1$ in the K' valley ($\tau_z = -1$). Therefore, in addition to Pauli ex-
 427 clusion, the order parameter matrix $\Delta(\mathbf{k})$ in the spin-valley space must satisfy the constraint
 428 $\mathcal{P}^T \Delta(\mathbf{k}) = \Delta(\mathbf{k}) \mathcal{P} = \Delta(\mathbf{k})$ where $\mathcal{P} = \frac{1}{2}(\mathbb{1} + \sigma_z \tau_z)$ projects onto the $\sigma_z \tau_z = +1$ space.
 429 Consistent with these requirements, $\Delta(\mathbf{k})$ takes the form

$$\Delta(\mathbf{k}) = \left(\Psi(\mathbf{k})\tau_+ - \Psi(-\mathbf{k})\tau_- \right) (\hat{\mathbf{z}} \cdot \vec{\sigma})(i\sigma_y) + \left(\Psi(\mathbf{k})\tau_+ + \Psi(-\mathbf{k})\tau_- \right) (i\sigma_y) \quad (45)$$

430 As expected, the absence of spin rotation invariance in the normal state results in a pair wave-
 431 function which is a superposition of singlet and triplet parts. For the pairing term to transform
 432 as a 1D irrep of C_{3v} under (44), $\Psi(\mathbf{k})$ must satisfy

$$\Psi(\mathbf{k}) = \Psi(C_3 \mathbf{k}) \quad (46a)$$

$$\Psi(-\mathbf{k}) = e^{i\Phi_{\sigma_v}} \Psi(\sigma_v \mathbf{k}) \quad (46b)$$

433 Table 4 shows that the two 1D irreps are in a one-one correspondence with the two SC state
 434 PSGs. It also gives an example of $\Psi(\mathbf{k})$ for each pairing symmetry.

435 4.4 Physical consequences of the PSG

436 The projective symmetry group G_f of the BdG Hamiltonian has effects on all fermionic ex-
 437 citations of the superconductor, since the Bogoliubov quasiparticles as excitations of the BdG
 438 Hamiltonian form a linear representation of the PSG G_f . In particular, the topological prop-
 439 erties of the superconductor is determined by the PSG, as different PSGs give rise to different
 440 classifications of fermion topological superconductors (TSCs) [11, 14, 28]. This is a well-known
 441 fact in the classification of gapped fermion topological phases, both in the 10-fold way [29]
 442 classification of non-interacting topological superconductors [11, 30], and in the interacting
 443 classification of fermion symmetry protected topological phases [14, 31]. For example, in the
 444 case of time reversal symmetry \mathcal{T} , it is well known that two- and three-dimensional topolog-
 445 ical insulators only exist for spinful electrons with $\hat{T}^2 = (-1)^{\hat{F}}$ and $G_f = U(1) \rtimes Z_4^{\mathcal{T}}$, which
 446 is a different symmetry class (class AII in the 10-fold way [29]) than spinless case (class AI in
 447 the 10-fold way [29]), with $\hat{T}^2 = \mathbb{1}$ and $G_f = U(1) \rtimes Z_2^{\mathcal{T}}$. In addition to topological classifi-
 448 cations, these two distinct symmetry classes have many other different properties, such as the
 449 presence vs. absence of Kramers degeneracy of fermion excitations. Below we illustrate how
 450 different PSGs, and hence different pairing symmetries, give rise to different classifications of
 451 TSCs, in the case of crystalline symmetries [28, 32–37]. Our classification scheme applies to
 452 gapped topological SCs. Thus, weak pairing unconventional SC with gap nodes are not part
 453 of the classification. However, there are fully gapped unconventional SCs, like the $(\mathbf{p} + i\mathbf{p})$
 454 state in 2D and the B-phase of He3, which are topologically non-trivial. Our analysis focus
 455 on understanding how, in the presence of additional crystalline symmetries, pairing symmetry
 456 through the PSG affects the classification of such states. We use 3d SCs with mirror reflection
 457 symmetry M_x , and 2d SCs with 2-fold rotational symmetry C_{2z} as two known examples to
 458 demonstrate this fact.

SOC strength	pairing symmetry	G_f	K-theory classification [38, 39]
Weak	A'	$SU(2) \times \mathbb{Z}_2^{\hat{M}_x}$	\mathbb{Z}
	A''	$SU(2) \times \mathbb{Z}_4^{\hat{M}_x} / \mathbb{Z}_2$	\mathbb{Z}_2
Strong	A'	$\mathbb{Z}_4^{\hat{M}_x}$	0
	A''	$\mathbb{Z}_2^{\hat{M}_x} \times \mathbb{Z}_2^F$	\mathbb{Z}

Table 5: Classification of class D topological superconductor in 3d with mirror reflection M_x . The fermion projective symmetry groups G_f are listed for superconductors with weak/strong SOC and A'/A'' pairing symmetries. Note that the topological classification is solely determined by G_f .

SOC strength	pairing symmetry	G_f	K-theory classification [40, 41]
Weak	A	$SU(2) \times \mathbb{Z}_2^{\hat{C}_{2z}}$	\mathbb{Z}
	B	$SU(2) \times \mathbb{Z}_4^{\hat{C}_{2z}} / \mathbb{Z}_2$	\mathbb{Z}^2
Strong	A	$\mathbb{Z}_4^{\hat{C}_{2z}}$	\mathbb{Z}^2
	B	$\mathbb{Z}_2^{\hat{C}_{2z}} \times \mathbb{Z}_2^F$	\mathbb{Z}

Table 6: Classification of class D topological superconductor in 2d with C_{2z} rotation perpendicular to the 2d x - y plane. The fermion PSGs G_f are listed for superconductors with weak/strong SOC and A/B pairing symmetries. Note that the topological classification is solely determined by G_f .

4.4.1 3d SCs with mirror reflection symmetry M_x

Our first example is the classification of TSCs in three dimension (3d) in the presence of only mirror reflection symmetry M_x which reverses the x coordinate. From the group cohomology $\mathcal{H}^{(2)}(\mathbb{Z}_2^{\hat{M}_x}, \mathbb{Z}_2) = \mathbb{Z}_2$, we find two possible fermion PSGs in the presence of strong SOC: $G_f = \mathbb{Z}_2^{\hat{M}_x} \times \mathbb{Z}_2^F$ with $\hat{M}_x^2 = +1$, and $G_f = \mathbb{Z}_4^{\hat{M}_x}$ with $\hat{M}_x^2 = (-1)^{\hat{F}}$. Similarly, in the presence of weak SOC and spin rotational symmetry, the two possible PSGs are given by $G_f = SU(2) \times \mathbb{Z}_2^{\hat{M}_x}$ with $\hat{M}_x^2 = +1$, and $G_f = SU(2) \times \mathbb{Z}_4^{\hat{M}_x} / \mathbb{Z}_2$ with $\hat{M}_x^2 = (-1)^{\hat{F}}$.

For weakly interacting systems, K-theory [10, 11, 30, 38, 39] can be used to classify distinct TSCs described by BdG Hamiltonians. In the presence of strong SOC, it gives rise to a \mathbb{Z} classification of TSCs for the case of $\hat{M}_x^2 = +1$, and a trivial classification for the case of $\hat{M}_x^2 = (-1)^{\hat{F}}$ [38, 39]. In the presence of a weak SOC and $SU(2)$ spin rotational symmetry, there is a \mathbb{Z} classification of TSCs for the case of $\hat{M}_x^2 = +1$, and a \mathbb{Z}_2 classification for the case of $\hat{M}_x^2 = (-1)^{\hat{F}}$ [38, 39]. With this result we can now readily bridge the gap between pairing symmetry and the K-theory classification of TSC via the projective symmetry group G_f .

A mirror symmetry satisfying $\hat{M}_x^2 = +1$ is preserved either in a singlet superconductor with pairing symmetry A' in the presence of a weak SOC, or in a superconductor with pairing symmetry A'' in the presence of a strong SOC. The classifications of weakly-interacting TSCs in these two cases are both \mathbb{Z} .

To compare, a mirror symmetry satisfying $\hat{M}_x^2 = (-1)^{\hat{F}}$ corresponds to either a singlet superconductor with pairing symmetry A'' in the presence of a weak SOC, or a pairing symmetry A' in the presence of a strong SOC. For these two cases the classifications of TSCs are \mathbb{Z}_2 and trivial, respectively. The results are summarized in Table 5.

481 4.4.2 2d SCs with 2-fold rotational symmetry C_{2z}

482 Our second example is the classification of TSCs in two dimensions (2d) with a C_{2z} rotation
 483 perpendicular to the 2d plane. In this case $\mathcal{H}^{(2)}(C_{2z}, \mathbb{Z}_2) = \mathbb{Z}_2$, which yields two different
 484 fermion PSGs in the presence of a strong (weak) SOC: one with $\hat{C}_{2z}^2 = +1$ and the other with
 485 $\hat{C}_{2z}^2 = (-1)^{\hat{f}}$, as shown in Table 6. Accordingly, the K -theory classification of C_{2z} symmetric
 486 TSCs [40] are given by \mathbb{Z} for $\hat{C}_{2z}^2 = +1$ and \mathbb{Z}^2 for $\hat{C}_{2z}^2 = (-1)^{\hat{f}}$.

487 From the relationship between pairing symmetry and projective symmetry group, we find
 488 that the $\hat{C}_{2z}^2 = +1$ case corresponds to either a singlet SC with pairing symmetry A or a SC
 489 with a strong SOC and pairing symmetry B . Then for these two cases the classifications of
 490 topological superconductors are both \mathbb{Z} .

491 The $\hat{C}_{2z}^2 = (-1)^{\hat{f}}$ case corresponds to either a singlet superconductor with pairing symme-
 492 try B or a superconductor with a strong SOC and pairing symmetry A . For these two cases the
 493 classifications of TSCs are both \mathbb{Z}^2 . The results are summarized in Table 6.

494 From these two examples, we see that BdG Hamiltonians with different PSGs generally
 495 give rise to different topological classifications. Based on the correspondence between the
 496 fermion PSG and the pairing symmetry discussed in sections 2-3, the classification of TSCs is
 497 therefore directly related to the pairing symmetry, as demonstrated in Table 5-6. For TSCs of
 498 all the possible pairing symmetries associated with a magnetic point group symmetry, Ref. [28]
 499 summarizes a full list of K -theory classification for both the cases of spinless (weak SOC) and
 500 spinful (strong SOC) electrons.

501 5 General framework

502 So far we have only focused on cases where the normal and the SC states have the same spin
 503 rotational symmetry. This simplifies the the form of the fermion PSG as explained below. For
 504 systems with weak SOC the physical symmetry group is $G = X \times SO(3)_{\text{spin}}$. When we take a
 505 central extension by the fermion parity group to obtain the fermion PSG, both in the normal
 506 and SC state PSGs, the $SO(3)_{\text{spin}}$ becomes an $SU(2)_{\text{spin}}$. The spatial part however undergoes
 507 different central extensions: X_f^0 in the normal state PSG and \tilde{X}_f in the SC state PSG. Thus,
 508 the fermion PSG preserving the kinetic energy is $(X_f^0 \times SU(2))/\mathbb{Z}_2$ and that preserving the
 509 BdG is $(\tilde{X}_f \times SU(2))/\mathbb{Z}_2$ (taking a quotient by \mathbb{Z}_2 takes care of the ‘‘double-counting’’ of \mathbb{Z}_2^F).
 510 Thus, the difference between the normal and SC state PSGs is completely captured by different
 511 central extensions of X by \mathbb{Z}_2^F . This holds true for systems with strong SOC where spin rotation
 512 symmetry is altogether absent and with $G = X$, the fermion PSG is synonymous with the central
 513 extension of X by \mathbb{Z}_2^F .

514 When spin rotation is spontaneously broken in the SC state, the fermion PSG no longer
 515 admits such a simple description in terms of central extensions of the spatial part. When the
 516 physical symmetry group in the SC state is $G = X \times S$ where S is a subgroup of the normal
 517 state spin rotation group, as we show in section 5.1, the fermion PSG could be a generic
 518 group extension of X by the fermion spin rotation symmetry group S_f . S_f in turn is a central
 519 extension of S by the fermion parity group and could in general be non-Abelian.

520 In superfluid ^3He , condensation into the spin triplet channel spontaneously breaks the
 521 continuous spin rotation symmetry present in the normal state. We discuss it in section 5.2 in
 522 the light of this general framework.

5.1 Group extension and pairing symmetry in a generic superconductor

Let the normal state spin rotational symmetry group $S_0 \subseteq SO(3)_{\text{spin}}$ be spontaneously broken down to $S \subseteq S_0$ in the SC state. With the charge $U(1)$ symmetry in the normal state completely broken, the SC state physical (bosonic) symmetry group G takes the form $G = X \times S$, where X denotes the spatial symmetry group preserved by the SC state. We now describe the structure of the fermion symmetry group G_f in such cases. Some of the relevant mathematical details can be found in Appendix A.

First of all, the fermion spin rotation (or internal) symmetry group in the SC-state, S_f is a subgroup of G_f and given by a *central* extension of the physical spin rotation symmetry group S :

$$1 \rightarrow \mathbb{Z}_2^F \rightarrow S_f \rightarrow S \rightarrow 1 \quad (47)$$

S_f has the form $S_f = \{(\pm 1)^{\hat{F}} \hat{s}' \mid s \in S\}$ where under a spin rotation \hat{s}' , the fermion operator transforms as

$$\hat{s}' \hat{c}_{k\alpha} \hat{s}'^{-1} = [\tilde{U}^s]_{\alpha\beta}^\dagger \hat{c}_{k\beta} \quad (48a)$$

$$\tilde{U}^s = e^{-i\phi_s} U_0^s \quad (48b)$$

The transformation is a combination of an $SU(2)$ spin rotation U_0^s that preserves the kinetic energy and a compensating phase rotation $e^{-i\phi_s}$ required to make \hat{s}' a symmetry of the BdG hamiltonian. Being an internal (on-site) symmetry, \hat{s}' leaves the momentum label unchanged on both sides of (48a). For a given S , the possible choices for S_f is captured by $\mathcal{H}^{(2)}(S, \mathbb{Z}_2)$, the second cohomology group formed by inequivalent classes of cocycles $[\tilde{\omega}]$. As already noted in previous sections, the cocycle $\tilde{\omega}$ taking values in $\{\pm 1\}$ also characterize the projective representation of S formed by $\{\tilde{U}^s \mid s \in S\}$.

To build G_f , next we need to consider the group of spatial symmetries, X . For $g \in X$, \hat{g}_0 preserves the kinetic energy and transforms the fermion operator as

$$\hat{g}_0 \hat{c}_{k\alpha} \hat{g}_0^{-1} = [U_0^g(\mathbf{k})]_{\alpha\beta}^\dagger \hat{c}_{gk\beta} \quad (49)$$

To make this a symmetry of the pairing term, not only do we need to dress it with a compensating phase $e^{-i\phi_g}$ but also with a normal state spin rotation $U_0^{s_0(g)}$ (where $s_0(g) \in S_0$). Since the kinetic energy is invariant under normal state spin rotations and $U(1)$ phase rotations, the resulting transformation \hat{g}' preserves the BdG hamiltonian. Its action on the fermion operator is given by

$$\hat{g}' \hat{c}_{k\alpha} \hat{g}'^{-1} = [\tilde{U}^g(\mathbf{k})]_{\alpha\beta}^\dagger \hat{c}_{gk\beta} \quad (50a)$$

$$\tilde{U}^g(\mathbf{k}) = e^{-i\phi_g} U_0^{s_0(g)} U_0^g(\mathbf{k}) \quad (50b)$$

Although the structure of G_f is in general much more complicated than simply a direct (or even a semi-direct) product of spatial and spin rotation symmetry groups, it is possible to obtain a generic characterization as discussed below. To begin with, let us compare what one obtains by the successive application of \hat{h}' and \hat{g}' on $\hat{c}_{k\alpha}$ and that by applying $\widehat{g\mathbf{h}'}$ on the same. Using (50a) we see that in both these cases we get a fermion operator on the right hand side with the same momentum label $g\mathbf{h}\mathbf{k}$. With both $\hat{g}'\hat{h}'$ and $\widehat{g\mathbf{h}'}$ being symmetries of the BdG hamiltonian, this implies that these are in fact the same upto an internal symmetry transformation $(\eta)^{\hat{F}} \hat{s}'(g, h)$. In other words,

$$\hat{g}' \hat{h}' = (\eta)^{\hat{F}} \hat{s}'(g, h) \widehat{g\mathbf{h}'} \quad (51)$$

Moreover, for any $\hat{s}' \in S_f$, the transformation $\hat{g}' \hat{s}' \hat{g}'^{-1}$ keeps the momentum label of the fermion operator unchanged and hence must belong to S_f . Then again, any element of G_f can be written as a product of a \hat{g}' for some $g \in X$ and an $(\eta)^{\hat{s}'} \in S_f$ such that G_f has the form $G_f = \{(\pm 1)^{\hat{s}'} \hat{s}' \hat{g}' \mid s \in S, g \in X\}$. We thus conclude that S_f is a normal subgroup of G_f and $G_f/S_f = X$. Equivalently S_f , G_f and X satisfy the short exact sequence

$$\mathbf{1} \rightarrow S_f \rightarrow G_f \rightarrow X \rightarrow \mathbf{1} \quad (52)$$

It is hard to find all such extensions in the most general case. However, if S_f is abelian then all such extensions are captured by the second cohomology group $\mathcal{H}_{[\rho]}^{(2)}(X, S_f)$. It is to be noted that the matrices $\{\tilde{U}^s \cdot \tilde{U}^g \mid s \in S, g \in X\}$ form a projective representation of $G = S \times X$ with coefficients in $\{\pm \tilde{U}^s \mid s \in S\}$.

Finally, we discuss the relation between the fermion PSG G_f and the pairing symmetry. In general, the pairing wavefunctions $\Delta_{\alpha, \beta}$ in BdG Hamiltonian (7) form a linear representation \mathcal{R}_{pair} of the bosonic symmetry group $G = S \times X$, where S stands for the global (spin rotational) symmetry group and X stands for the crystalline symmetry group. Meanwhile, in the representation $\{\tilde{U}^s \cdot \tilde{U}^g \mid s \in S, g \in X\}$ introduced above, we can identify a projective representation of group $G = S \times X$:

$$\mathcal{R}_{\Phi}(s, g) = e^{-i(\phi_s + \phi_g)} U_0^{s_0(g)}, \quad \forall s \in S, g \in X. \quad (53)$$

It is evident that the projective representation \mathcal{R}_{Φ} is not 1D in general. Also note that while for the global internal symmetry group S , the transformations that preserve the kinetic energy, preserve the pairing term up to a phase (just as in (14)), that may not be the case for the crystalline group X . Hence \mathcal{R}_{pair} is in general a multi-dimensional linear representation of G . \mathcal{R}_{pair} and \mathcal{R}_{Φ} are related by the following relation:

$$\mathcal{R}_{\Phi} \otimes \mathcal{R}_{\Phi} \otimes \mathcal{R}_{pair} = \mathbb{1} \oplus \dots \quad (54)$$

where $\mathbb{1}$ denotes the trivial one-dimensional (1d) representation of group G . This is because the pairing term (9) must remain invariant under the PSG symmetry transformation $\{\tilde{U}^s \cdot \tilde{U}^g \mid s \in S, g \in X\}$. Notice that in the special case of \mathcal{R}_{Φ} being a 1d irrep, applicable to the situation discussed in Section 3, the general relation (54) reduces to Eq. (22). In order for \mathcal{R}_{pair} to be a multi-dimensional irrep., the tensor product $\mathcal{R}_{\Phi} \otimes \mathcal{R}_{\Phi}$ of two projective representations in Eq. (54) must be a multi-dimensional irrep. of group G . Therefore, a necessary condition for the pairing order parameter to form a multi-dimensional irrep. of symmetry group G (i.e. for \mathcal{R}_{pair} to be multi-dimensional) is that \mathcal{R}_{Φ} is a multi-dimensional projective representation of group G . As we will show below, one such example is the superfluid B phase of Helium 3.

5.2 Examples: superfluid A and B phases in Helium-3

The most famous example of triplet superconductivity (or superfluidity) is perhaps Helium-3 [5]. The normal state preserves continuous spatial rotations and inversion symmetry:

$$X_0 = \mathbf{SO}(3)_{\text{orbital}} \times \mathbb{Z}_2^I \simeq \mathbf{O}(3), \quad (55)$$

along with full spin rotation symmetry, $S_0 = \mathbf{SO}(3)_{\text{spin}}$. Condensation takes place in a spin triplet p -wave state breaking the full spin rotation symmetry down to a proper subgroup. In the basis, $\Psi_{\mathbf{k}} \equiv (c_{\mathbf{k}, \uparrow}, c_{\mathbf{k}, \downarrow}, c_{-\mathbf{k}, \uparrow}^{\dagger}, c_{-\mathbf{k}, \downarrow}^{\dagger})^T$ the BdG Hamiltonian takes the form

$$\hat{H}_{\text{BdG}} = \sum_{\mathbf{k}} \hat{\Psi}_{\mathbf{k}}^{\dagger} \begin{pmatrix} (\frac{k^2}{2m} - \mu) \mathbb{1} & \Delta(\mathbf{k}) \\ \Delta^{\dagger}(\mathbf{k}) & (\mu - \frac{k^2}{2m}) \mathbb{1} \end{pmatrix} \hat{\Psi}_{\mathbf{k}} \quad (56a)$$

$$\Delta(\mathbf{k}) = \mathbf{d}(\mathbf{k}) \cdot \vec{\sigma} (i\sigma_y) \quad (56b)$$

593 To obey Fermi statistics, the three component complex vector $\mathbf{d}(\mathbf{k})$ must satisfy $\mathbf{d}(\mathbf{k}) = -\mathbf{d}(-\mathbf{k})$.
 594 In particular for p -wave ${}^3\text{He}$, the components of $\mathbf{d}(\mathbf{k})$ are linear in \mathbf{k} . The various phases,
 595 characterized by different broken symmetries, are distinguished by the form of the $\mathbf{d}(\mathbf{k})$ vector.
 596 We apply the general framework described above to the two phases: (1) B phase, also known
 597 as the Balian-Werthamer (BW) phase [42], (2) A phase, also known as Anderson-Brinkman-
 598 Morel (ABM) phase [43, 44], discussing the residual symmetry group in the SC state and the
 599 SC state fermion PSG in each case.

600 The transformations that preserve the kinetic energy act on the fermion operators as

$$\text{Spin rot.} \quad \hat{S}_0(\vec{\theta}) \hat{c}_{\mathbf{k}s} \hat{S}_0(\vec{\theta})^{-1} = \left[U_0^{S(\vec{\theta})} \right]_{s s'}^\dagger \hat{c}_{\mathbf{k}s'}, \quad U_0^{S(\vec{\theta})} = e^{i\vec{\theta} \cdot \vec{\sigma} / 2} \quad (57a)$$

$$\text{Space rot.} \quad \hat{\mathcal{R}}_0(\vec{\theta}) \hat{c}_{\mathbf{k}s} \hat{\mathcal{R}}_0(\vec{\theta})^{-1} = \left[U_0^{\mathcal{R}(\vec{\theta})} \right]_{s s'}^\dagger \hat{c}_{\mathcal{R}(\vec{\theta})\mathbf{k}s'}, \quad U_0^{\mathcal{R}(\vec{\theta})} = \mathbb{1} \quad (57b)$$

$$\text{Inversion} \quad \hat{\mathcal{I}}_0 \hat{c}_{\mathbf{k}s} \hat{\mathcal{I}}_0^{-1} = \left[U_0^{\mathcal{I}} \right]_{s s'}^\dagger \hat{c}_{-\mathbf{k}s'}, \quad U_0^{\mathcal{I}} = \mathbb{1} \quad (57c)$$

601 With $[\hat{\mathcal{R}}_0(\pi\hat{\mathbf{n}})]^2 = \hat{\mathcal{I}}_0^2 = \mathbb{1}$ and $[\hat{S}_0(\pi\hat{\mathbf{n}})]^2 = (-1)^{\hat{F}}$, the normal state fermion PSG is of the
 602 form $X_0 \times SU(2)$.

603 5.2.1 Superfluid B phase of Helium-3

604 In the B phase, $\mathbf{d}(\mathbf{k}) = \Delta_0(k_x \hat{\mathbf{x}} + k_y \hat{\mathbf{y}} + k_z \hat{\mathbf{z}})$ [22]. The spin rotation group is broken down from
 605 $S_0 = SO(3)_{\text{spin}}$ to its trivial subgroup $S = \{\mathbb{1}\}$ in the SC state. Using (47), the fermion onsite
 606 symmetry group is simply the fermion parity group $S_f = \mathbb{Z}_2^F$. The system remains isotropic
 607 in the SC state and $X = SO(3)_{\text{orb.}+\text{spin}} \times \mathbb{Z}_2^I \simeq O(3)$. As suggested by the label, the normal
 608 state spatial rotation in (57b) has to be modified by including a normal state spin rotation
 609 and since $\mathbf{d}(\mathbf{k})$ is inversion odd, the normal state inversion in (57c) has to be modified by a
 610 compensating phase rotation by i . The transformations that preserve (56a) are

$$\text{Space rot.} \quad \hat{\mathcal{R}}'(\vec{\theta}) \hat{c}_{\mathbf{k}s} \hat{\mathcal{R}}'(\vec{\theta})^{-1} = \left[\tilde{U}^{\mathcal{R}(\vec{\theta})} \right]_{s s'}^\dagger \hat{c}_{\mathcal{R}(\vec{\theta})\mathbf{k}s'}, \quad \tilde{U}^{\mathcal{R}(\vec{\theta})} = e^{i\vec{\theta} \cdot \vec{\sigma} / 2} \cdot \mathbb{1} \quad (58a)$$

$$\text{Inversion} \quad \hat{\mathcal{I}}' \hat{c}_{\mathbf{k}s} \hat{\mathcal{I}}'^{-1} = \left[\tilde{U}^{\mathcal{I}} \right]_{s s'}^\dagger \hat{c}_{-\mathbf{k}s'}, \quad \tilde{U}^{\mathcal{I}} = i \mathbb{1} \quad (58b)$$

611 With $S_f = \mathbb{Z}_2^F$, the full fermion symmetry group G_f , which by (52) is a group extension of X by
 612 S_f , reduces to a central extension of X by \mathbb{Z}_2^F . Eqns. (58a) and (58b) give $\hat{\mathcal{R}}'(\pi\hat{\mathbf{n}})^2 = \hat{\mathcal{I}}'^2 = (-1)^{\hat{F}}$,
 613 showing that G_f involves non-trivial central extensions of both $SO(3)_{\text{orb.}+\text{spin}}$ and \mathbb{Z}_2^I and is
 614 given by $G_f = (SU(2) \times \mathbb{Z}_4) / \mathbb{Z}_2$.

615 Conversely, one can learn about the pair wavefunction from the SC state PSG in the su-
 616 perfluid B phase. From (58a), we see that $\mathcal{R}_\Phi(\mathcal{R}(\vec{\theta})) = U_0^{S(\vec{\theta})} = e^{i\vec{\theta} \cdot \vec{\sigma} / 2}$ which is a $j = 1/2$
 617 projective representation of $G(\simeq SO(3))$. According to relation (54) and the angular momen-
 618 tum addition rules, $\mathcal{R}_{\text{pair}}$ is either a $j = 0$ or $j = 1$ linear irrep of $G(\simeq SO(3))$. However,
 619 because the projective representation $\mathcal{R}_\Phi(\mathcal{R}(\vec{\theta}))$ coincides with the normal-state spin rotation
 620 in (57a), the $j = 0$ irrep will preserve spin rotation and hence does not apply to the superfluid
 621 B phase. As a result, the pairing term must transform like a $j = 1$ representation under (57b).
 622 This is consistent with $\mathbf{d}(\mathbf{k}) \propto \mathbf{k}$ in this case.

623 5.2.2 Superfluid A phase of Helium-3

624 In the A-phase, without loss of generality, $\mathbf{d}(\mathbf{k}) = \Delta_0(k_x + ik_y)\hat{\mathbf{z}}$ [22]. The spin rotational
 625 symmetry is broken down from $S_0 = SO(3)_{\text{spin}}$ to $S = U(1)^z \times \mathbb{Z}_2^x \simeq O(2)$, which is the
 626 subgroup generated by continuous spin rotations around the $\hat{\mathbf{z}}$ axis, $\mathcal{S}(\theta\hat{\mathbf{z}})$ and π spin ro-
 627 tations about the x -axis. All possible fermion onsite symmetry groups S_f are classified by

628 $\mathcal{H}^2(\mathcal{S}, \mathbb{Z}_2^F) = \mathcal{H}^2(\mathcal{O}(2), \mathbb{Z}_2) = \mathbb{Z}_2^3$. Since under π spin rotation about x -axis $d_x(\mathbf{k}) \rightarrow -d_x(\mathbf{k})$,
 629 the corresponding normal state transformation has to be modified by a phase rotation of i .
 630 No such compensating phase is thus required for spin rotation about z -axis. The SC state spin
 631 rotations are implemented as

$$\text{Spin rot.} \quad \hat{S}'(\theta \hat{z}) \hat{c}_{ks} \hat{S}'(\theta \hat{z})^{-1} = [\tilde{U}^{S(\theta \hat{z})}]_{s s'}^\dagger \hat{c}_{ks'} \quad \tilde{U}^{S(\theta \hat{z})} = e^{i\theta \sigma_z/2} \quad (59a)$$

$$\text{Spin rot.} \quad \hat{S}'(\pi \hat{x}) \hat{c}_{ks} \hat{S}'(\pi \hat{x})^{-1} = [\tilde{U}^{S(\pi \hat{x})}]_{s s'}^\dagger \hat{c}_{ks'} \quad \tilde{U}^{S(\pi \hat{x})} = \sigma_x \quad (59b)$$

632 The central extension is characterized by $\hat{S}'(\pi \hat{x})^2 = 1$ and $\hat{S}'(\pi \hat{z})^2 = \hat{S}'(\pi \hat{y})^2 = (-1)^{\hat{F}}$ and
 633 correspondingly $\mathcal{S}_f \simeq \{\pm \sigma_x^n e^{i\theta \sigma_z/2} \mid 0 \leq \theta < 2\pi, n = 0, 1\} = \{\sigma_x^n e^{i\xi \sigma_z} \mid 0 \leq \xi < 2\pi, n = 0, 1\}$
 634 $\simeq \mathcal{O}(2)$.

635 The spatial $\mathcal{O}(3)$ symmetry is broken down to a subgroup of $X = U(1)^z \times \mathbb{Z}_2^I$, generated
 636 by continuous spatial rotations about z -axis, $\mathcal{R}(\theta \hat{z})$ and inversion \mathcal{I} . In this case, the normal
 637 state transformations need to be modified only by compensating phase rotations. The SC state
 638 transformations are given by

$$\text{Space rot.} \quad \hat{\mathcal{R}}'(\theta \hat{z}) \hat{c}_{ks} \hat{\mathcal{R}}'(\theta \hat{z})^{-1} = [\tilde{U}^{\mathcal{R}(\theta \hat{z})}]_{s s'}^\dagger \hat{c}_{\mathcal{R}(\theta \hat{z})ks'} \quad \tilde{U}^{\mathcal{R}(\theta \hat{z})} = e^{-i\theta/2} \cdot \mathbb{1} \quad (60a)$$

$$\text{Inversion} \quad \hat{\mathcal{I}}' \hat{c}_{ks} \hat{\mathcal{I}}'^{-1} = [\tilde{U}^{\mathcal{I}}]_{s s'}^\dagger \hat{c}_{-ks'} \quad \tilde{U}^{\mathcal{I}} = i \mathbb{1} \quad (60b)$$

639 In this case the fermion symmetry group $G_f \simeq (\mathcal{O}(2) \times U(1) \times \mathbb{Z}_4)/\mathbb{Z}_2$ is a nontrivial extension
 640 of X by \mathcal{S}_f satisfying $[\hat{\mathcal{R}}'(\pi \hat{z})]^2 = \hat{\mathcal{I}}'^2 = (-1)^{\hat{F}}$.

641 6 Conclusion

642 Traditionally, the broken and unbroken symmetries of a superconductor (SC) is described by
 643 the Ginzburg-Landau theory, which characterizes the symmetry properties of all bosonic ex-
 644 citations therein, such as Cooper pairs. In this paper we investigate the same problem of
 645 broken and unbroken symmetries in a SC state from a viewpoint of fermionic excitations.
 646 We showed that the projective symmetry group (PSG) of fermions in a superconductor is the
 647 proper language to capture symmetry-related properties of fermionic excitations in a SC, and
 648 systematically studied the relationship between the pairing symmetry and the fermion PSGs in
 649 a superconductor. We provided a general framework in Section 5 to characterize the fermion
 650 symmetry group after the Cooper pair formation with the concept of PSG, which is a group
 651 extension of the crystalline space group X by the fermion global symmetry group \mathcal{S}_f in the
 652 superconducting phase. Examples of fermion global symmetry groups include the fermion
 653 parity group \mathbb{Z}_2^F in a generic SC without spontaneous breaking of spin rotational symmetries,
 654 and $\mathcal{O}(2)$ as in the case of superfluid A phase of Helium-3. In the case of the fermion global
 655 symmetry group \mathcal{S}_f being an Abelian group, the group extension problem can be classified by
 656 the second group cohomology, which is both conceptually clear and practically easy to com-
 657 pute.

658 When the SC and normal state share the same fermion global symmetries, i.e. in the
 659 absence of spontaneously broken spin rotational symmetries, the fermion PSG of the SC state is
 660 particularly simple: it is a central extension of the crystalline symmetry group X by the fermion
 661 parity group \mathbb{Z}_2^F . In this case, we can classify all fermion PSGs using elements of the 2nd
 662 cohomology group $\mathcal{H}^2(X, \mathbb{Z}_2^F)$. Using the connection between pairing symmetry and fermion
 663 PSG discussed in section 2, we can systematically obtain all the possible pairing symmetries
 664 compatible with the PSGs as delineated in Sec. 3. A distinction was made between the case
 665 of SCs with and without spin-orbital couplings (SOCs), where in the presence of a strong

666 SOC, crystalline symmetries of fermions in the normal state are described by a non-trivial 2-
 667 cocycle $\omega_0 \in \mathcal{H}^2(X, \mathbb{Z}_2^F)$, and the correspondence between PSG and pairing symmetry should
 668 be shifted accordingly. Within this general framework, we calculated all the possible PSGs
 669 for all 3-dimensional point group symmetries both with and without SOCs, and establish the
 670 correspondence between PSGs and pairing symmetries of the SCs. As a demonstration of
 671 the framework, we studied in detail the PSGs and pairing symmetries of several physically
 672 relevant systems in section 4, and hope our work would shed new lights on understandings of
 673 superconductivity in these systems. Considering the crystalline symmetry group X , although
 674 we have restricted our attention to point groups in this work, the case of magnetic point groups
 675 and space groups can be naturally incorporated in our general framework.

676 It is useful to compare the fermion PSGs in this work to PSGs initially introduced in the
 677 context of quantum spin liquids (QSLs) [21,23]. In QSLs, due to the presence of fractionalized
 678 excitations, like spinons, and emergent gauge fields, each element of the PSG is a combination
 679 of physical symmetry operation, such as a crystal symmetry $\mathbf{g} \in X$, and local gauge rotations.
 680 In contrast, in a superconductor each element of the fermion PSG is a combination of an
 681 unbroken crystal symmetry operation $\mathbf{g} \in X$ and a spontaneously-broken *global* symmetry
 682 operation such as a $U(1)$ charge rotation. We emphasize that our analysis does *not* involve the
 683 effects of dynamical *local* gauge fields, which have been proposed to lead to a description of
 684 superconductors as symmetry protected topological states [45] or states with \mathbb{Z}_2 topological
 685 ordered states [46]. We thus treat charged superconductors and neutral paired superfluids on
 686 the same footing as systems with a broken global $U(1)$ possibly in addition to other broken
 687 symmetries.

688 PSGs have important implications on physical properties of a superconductor. As the PSG
 689 G_f is the symmetry group of fermions in a SC, it dictates the symmetry and topological proper-
 690 ties of all the fermionic excitations of the system and its validity extends beyond the mean-field
 691 BdG equations. Therefore, PSG can be used to classify topological superconductors in both
 692 non-interacting (i.e., admitting a mean-field description) and interacting cases. As an illustra-
 693 tion, we discussed systems with two different kinds of symmetry groups where G_f determines
 694 classifications of non-interacting topological superconductors. Moreover, as PSG establishes
 695 a link between pairing symmetry and topological properties of a system, we can utilize topo-
 696 logical properties of the electronic excitations as a diagnosis for the pairing symmetry of a
 697 superconductor. We leave these interesting ideas for future works.

698 Acknowledgements

699 **Funding information** XY, SB, SL, MR and YML acknowledge support by Center for Emergent
 700 Materials at The Ohio State University, a National Science Foundation (NSF) MRSEC through
 701 NSF Award No. DMR-2011876. YML acknowledges support by grant NSF PHY-1748958 to
 702 the Kavli Institute for Theoretical Physics (KITP), and NSF PHY-2210452 to the Aspen Center
 703 for Physics.

704 A A short introduction to projective representation and 2-cocycle

705 In this appendix we want to elucidate the connection between the projective representation of
 706 the crystalline symmetry group as described by the mathematical object called 2-cocycle and
 707 the fermion projective symmetry group G_f .

708 The concept of PSG was first introduced in the study of quantum spin liquids [21]. In the
 709 context of quantum spin liquids, electrons can be thought of as being composed of chargons

710 and spinons which are glued together by an $SU(2)$ gauge field. Due to the emergent gauge
 711 structures, symmetries that are represented linearly on the physical degrees of freedom are
 712 now represented only projectively on the spinons. More specifically, spin operators at site i
 713 can be written as fermionic spinons: $S_i = \frac{1}{2} f_{i,\alpha}^\dagger \vec{\sigma}_{\alpha,\beta} f_{i,\beta}$. A spin Hamiltonian can be described
 714 by a mean-field theory of spinons plus gauge fluctuations. Consider the following mean-field
 715 Hamiltonian:

$$H = \sum_{ij} [\psi_i^\dagger u_{ij} \psi_j + h.c.] + \sum_i a_0^l \psi_i^\dagger \tau^l \psi_i, \quad (\text{A.1})$$

716 where u_{ij} 's are 2×2 matrices encoding pairing and hoppings of fermionic spinons, $\psi_i = (f_\uparrow, f_\downarrow)^T$
 717 are Nambu spinors.

718 The Hamiltonian has a local $SU(2)$ gauge redundancy: a site-dependent $SU(2)$ transfor-
 719 mation $\psi_i \rightarrow W_i \psi_i, u_{ij} \rightarrow W_i u_{ij} W_j^\dagger$ with $W_i \in SU(2)$ which leaves both physical observables
 720 and the Hamiltonian invariant. Due to this gauge redundancy, the symmetry of the spin liquids
 721 are described by the projective symmetry group, which is defined as the collection of all com-
 722 binations of symmetry elements and gauge transformations that leave the mean-field ansatz
 723 $\{u_{ij}\}$ invariant:

$$G_U U(\{u_{ij}\}) = \{u_{ij}\}, \quad (\text{A.2})$$

$$U(\{u_{ij}\}) \equiv \{\tilde{u}_{ij} = u_{U^{-1}(i), U^{-1}(j)}\}, \quad (\text{A.3})$$

$$G_U(u_{ij}) \equiv \{\tilde{u}_{ij} = G_U(i) u_{ij} G_U^\dagger(j)\}, \quad (\text{A.4})$$

$$G_U(i) \in SU(2), \quad (\text{A.5})$$

724 where U is an element of the symmetry group SG of the microscopic system and G_U is the
 725 $SU(2)$ gauge transformation accompanying U that leaves the mean-field ansatz invariant.

726 To encode the emergent gauge fields at low energy for spin liquid states, we introduce
 727 the important concept of invariant gauge group (IGG) which are pure gauge group elements
 728 that leave the mean-field ansatz invariant: $W_i u_{ij} W_j^\dagger = u_{ij}$. It is clear that IGG corresponds
 729 to elements $G_U U$ in PSG where U is the identity. With the concept of IGG it is now easy to
 730 describe the structure of PSG. In fact, IGG is a normal subgroup of PSG, and with the group
 731 homomorphism $\rho(G_U U) = U$ between PSG and SG, we have the following exact sequence:

$$1 \rightarrow \text{IGG} \xrightarrow{\iota} \text{PSG} \xrightarrow{\rho} \text{SG} \rightarrow 1, \quad (\text{A.6})$$

732 where ι is the embedding mapping, and the exactness is ensured by the fact that $\rho(w) \equiv 1 \in$
 733 SG for $w \in \text{IGG}$. The structure of the PSG is now quite clear: it is the group extension of the
 734 SG by the IGG, or alternatively, $SG = \text{PSG}/\text{IGG}$.

735 Equipped with the knowledge of PSG, it is also easy to see that the problem of unbroken
 736 symmetries of the superconductor naturally fits into the general framework of PSG if we notice
 737 that the BdG Hamiltonian takes the same form as the spin liquid mean-field Hamiltonian. More
 738 precisely, as discussed in the main text, fermions in the superconductor has the symmetry
 739 group G_f , which is an extension of the space group X by the fermion global symmetry group
 740 S_f described by the short exact sequence:

$$1 \rightarrow S_f \rightarrow G_f \rightarrow X \rightarrow 1. \quad (\text{A.7})$$

741 The resemblance to Eq. A.6 is immediately seen if we identify the unbroken global symmetry
 742 group S_f as IGG and the fermion symmetry group G_f as PSG. However, there's an important
 743 difference we need to keep in mind: in our study of superconductor, the global symmetry

744 group \mathcal{S}_f should not be regarded as the gauge group corresponding to a fluctuating gauge
745 field, as was in the context of spin liquids.

746 In general \mathcal{S}_f can be non-Abelian, and we refer to Ref. [21] for a general computation
747 scheme to solve the extension problem by obtaining all the inequivalent projective symmetry
748 groups \mathcal{G}_f . Below let's discuss the special case of \mathcal{S}_f being Abelian, which covers most of the
749 practical situations and is mathematically much simpler to deal with. And we will comment
750 briefly on the case of \mathcal{S}_f being non-Abelian in the end.

751 In the case of \mathcal{S}_f being Abelian, the group extension of X by \mathcal{S}_f can be described as an
752 element in the second cohomology group $\mathcal{H}_{[\rho]}^2(X, \mathcal{S}_f)$ with group actions $[\rho] : X \rightarrow \text{Aut}(\mathcal{S}_f)$
753 (note that when the group action is trivial, the group extension is simply a central extension).
754 To see this more clearly, let's label group elements in \mathcal{G}_f as (s, g) with $s \in \mathcal{S}_f, g \in X$. Now,
755 since \mathcal{S}_f is in the center of \mathcal{G}_f , we can represent the group multiplication rule in the following
756 way:

$$(s_g, g) \times (s_h, h) = (s(g, h)s_g s_h, gh), \quad (\text{A.8})$$

757 where $s(g, h)$ is a function $X \times X \rightarrow \mathcal{S}$. The above procedure has an ambiguity since we can
758 alternatively define $g' = \gamma_g g \in \mathcal{G}_f$ ($\gamma_g \in \mathcal{S}_f$) as our canonical choice of g . This then modifies
759 $s(g, h)$ as:

$$s(g, h) \rightarrow s(g, h) \cdot \gamma_g^g \cdot \gamma_h^{-1} \cdot \gamma_{gh}^{-1}, \quad (\text{A.9})$$

760 where the superscript g indicates group actions g on elements in \mathcal{S}_f as described by $[\rho]$.

761 The $s(g, h)$'s satisfy the associativity condition if we apply three group elements in \mathcal{G}_f in
762 two equivalent ways, which yields

$$s(g_1, g_2)s(g_1g_2, g_3) = s(g_1, g_2g_3)s^{g_1}(g_2, g_3). \quad (\text{A.10})$$

763 The coboundary condition A.9 and the cocycle condition A.10 then define an element in
764 $\mathcal{H}^2(X, \mathcal{S}_f)$. Therefore we have found out that in the case of central extension, \mathcal{G}_f is uniquely
765 determined by the 2-cocycle $s(g, h)$, which is further classified by the second cohomology
766 group $\mathcal{H}^2(X, \mathcal{S}_f)$.

767 Before proceeding, let me emphasize an important point: fermions fulfill a 1d representa-
768 tion of \mathcal{S}_f , which we denote as $\rho_{\mathcal{S}} : \mathcal{S}_f \rightarrow U(1)$. Note that $\rho_{\mathcal{S}}$ is determined by the microscopic
769 electrons and can be viewed as a group homomorphism from \mathcal{S}_f to $\text{Image}(\rho_{\mathcal{S}})$.

770 Because elements in \mathcal{G}_f act on fermions in a linear way, let's consider a linear represen-
771 tation \hat{U} of the group \mathcal{G}_f . Since \mathcal{S}_f lies at the center of the group, $\hat{U}((s, 1))$ should be of the
772 form $\rho_{\mathcal{S}}(s) \times \mathbb{1}$ according to Schur's lemma and the fact that the symmetry action of $s \in \mathcal{S}$ on
773 fermions is given by $\rho_{\mathcal{S}}$.

774 If we identify $U(g)$ as $\hat{U}((1, g))$, $U(g)$ would fulfill a projective representation of X :

$$\begin{aligned} U(g)U(h) &= \hat{U}((1, g))\hat{U}((1, h)) = \hat{U}((s(g, h), gh)) \\ &= \hat{U}((s(g, h), 1))\hat{U}((1, gh)) = \omega(g, h)U(gh), \end{aligned} \quad (\text{A.11})$$

775 where $\omega(g, h) \equiv \rho_{\mathcal{S}}(s(g, h))$ is a function $X \times X \rightarrow \text{Image}(\rho_{\mathcal{S}})$. The ω satisfies the following
776 associativity condition if we act three consecutive symmetry operations in two equivalent ways:
777 $g_1g_2g_3 = (g_1g_2)g_3 = g_1(g_2g_3)$, which translates to

$$\omega(g_1, g_2)\omega(g_1g_2, g_3) = \omega(g_1, g_2g_3)\omega^{g_1}(g_2, g_3), \quad (\text{A.12})$$

778 where the superscript g on ω indicates group actions on the $U(1)$ phase induced by the group
779 action $[\rho]$ on elements in \mathcal{S}_f .

780 We can also multiply symmetry actions $U(\mathbf{g})$ by some $U(1)$ phase $\gamma_{\mathbf{g}} \in \text{Image}(\rho_S)$, which
781 then modifies ω in the following way:

$$\omega(\mathbf{g}, \mathbf{h}) \rightarrow \omega(\mathbf{g}, \mathbf{h}) \frac{\gamma_{\mathbf{g}} \gamma_{\mathbf{h}}^{\mathbf{g}}}{\gamma_{\mathbf{gh}}}. \quad (\text{A.13})$$

782 The associativity condition (A.12) and the ambiguity (A.13) thus define a 2-cocycle in
783 the second cohomology group $\mathcal{H}^2(X, \text{Image}(\rho_S))$. And the equation Eq.(A.11) establishes
784 an explicit homomorphism between the projective representation of X (an element in the
785 cohomology group $\mathcal{H}^2(X, \text{Image}(\rho_S))$) and the fermion projective symmetry group G_f (an
786 element in $\mathcal{H}^2(X, S)$).

787 In summary, a linear representation of the fermion projective symmetry group G_f can
788 alternatively be viewed as a projective representation of the group X with cocycle $\omega(\mathbf{g}, \mathbf{h})$
789 which is an element in the cohomology group $\mathcal{H}^2(X, \text{Image}(\rho_S))$, as elucidated by Eq.(A.11).

790 Several remarks are in order:

- 791 1. When S_f is Abelian and ρ_S is injective, the two cohomology groups $\mathcal{H}^2(X, \text{Image}(\rho_S))$
792 and $\mathcal{H}^2(X, S_f)$ are isomorphic to each other, therefore we have sometimes used these
793 terms interchangeably in the main text.
- 794 2. When S_f is non-Abelian, G_f can no longer be described by an element in the second
795 cohomology group. If we restrict our attention to the case where the representation ρ_S
796 of S_f on fermions are one dimensional, then the correspondence Eq.(A.11) still holds,
797 enabling us to carry out calculations within this general framework.
- 798 3. When S_f is non-Abelian, there are cases where the representation of S_f on fermions
799 are at least 2-dimensional, such as spin-1/2 fermions in the superfluid A phase with
800 $S_f = \mathbf{O}(2)$. Such cases are beyond the scope of cohomological description, and we need
801 to solve the projective symmetry groups up to gauge equivalence on a case-by-case basis
802 following the general procedures as described in Ref. [21].

803 B How PSG constrains the pairing symmetry for all crystalline 804 point groups

805 Since G_f is the extension of G by Z_2^F , we can view 1d projective representations $\mathcal{R}_{\Phi}(\mathbf{g})$ of G as
806 regular representations $\bar{\mathcal{R}}_{\Phi}(\hat{\mathbf{g}}')$ for $\hat{\mathbf{g}}' \in G_f$ with $\bar{\mathcal{R}}_{\Phi}(\mathbf{d}) = -1$ ($\mathbf{d} \equiv (-1)^{\hat{\mathbf{d}}}$) when restricted
807 to the subgroup $X = G_f/Z_2^F$. This is confirmed by the following relation:

$$\bar{\mathcal{R}}_{\Phi}((\eta_{\mathbf{g}}, \hat{\mathbf{g}}')) \bar{\mathcal{R}}_{\Phi}((\eta_{\mathbf{h}}, \hat{\mathbf{h}}')) = \bar{\mathcal{R}}_{\Phi}((\eta_{\mathbf{g}} \eta_{\mathbf{h}} \tilde{\omega}(\mathbf{g}, \mathbf{h}), \hat{\mathbf{g}}' \hat{\mathbf{h}}')) = \omega(\mathbf{g}, \mathbf{h}) \bar{\mathcal{R}}_{\Phi}((\eta_{\mathbf{g}} \eta_{\mathbf{h}}, \hat{\mathbf{g}}' \hat{\mathbf{h}}')), \quad (\text{B.1})$$

808 where we have used the fact that Z_2^F is the center of G_f and $\eta_{\mathbf{g}}, \eta_{\mathbf{h}} = \pm 1$.

809 Our strategy then is to first obtain the group extension $G_f \in \mathcal{H}^2(X, Z_2^F)$ and then compute
810 the 1d irreducible representations $\bar{\mathcal{R}}_{\Phi}(\mathbf{g})$ of G_f with $Z_2^F = -1$, from which we can readily
811 obtain $\mathcal{R}_{\text{pair}}$. We used GAP computer algebra program [24] in all these calculations, which is
812 ideally suited for the task. The results are displayed in Table.7.

813 C GAP program for PSG calculation

814 Groups, Algorithms and Programming (GAP) is a software system designed for algebraic com-
815 putations. In this section, we provide further details on how GAP is applied to the theories
816 discussed in Sections 2, 3.

817 To calculate the group cohomology of point groups, we use the HAP and Cryst pack-
 818 ages in GAP. Starting from a point group G , we calculate its second cohomology with coeffi-
 819 cients in \mathbb{Z}_2 using the TwoCohomologyGeneric function, and we determine the representa-
 820 tions of G using the Irr function. For each cohomology class, we obtain the unique (up to
 821 coboundary equivalence) group extension \tilde{X}_f of G via the FpGroupCocycle function. From
 822 the TwoCohomologyGeneric function and the FpGroupCocycle function we can easily con-
 823 struct the quotient map $\tilde{X}_f \rightarrow G$. Next, we calculate the representations and characters of \tilde{X}_f .
 824 For each one-dimensional irrep of \tilde{X}_f , we simply square it and use the quotient map $\tilde{X}_f \rightarrow G$
 825 to obtain the corresponding irrep of G . We then look up the group representation tables in
 826 Ref. [47] to find the corresponding \mathcal{R}_{pair} as listed in the last column of Table. 7. The pseu-
 827 docode is presented in Algorithm 1.

828 One complication in the algorithm is to obtain a complete and linearly-independent list of
 829 gauge-invariant cocycles to label the cohomology classes $\mathcal{H}^2(G, \mathbb{Z}_2)$. This is done manually
 830 by first listing all the gauge-invariant cocycles of the form $\zeta_g \equiv \omega(g, g)$ for group element
 831 g satisfying $g^2 = \mathbb{1}$, and $\eta_{g,h} \equiv \frac{\omega(g,h)}{\omega(h,g)}$ for group elements g, h satisfying $gh = hg$. We
 832 have tested that this list completely characterize all the cohomology classes. We then find the
 833 linearly-independent ones among these gauge-invariant cocycles to label all the cohomology
 834 classes as shown in column 3 of Table. 7.

Algorithm 1 Pseudocode for calculating PSGs and corresponding pairing symmetries

Input: Point group G
Output: the (spinless) PSG ω_Φ
Output: the corresponding pairing symmetry \mathcal{R}_{pair}

- 1: Coh = $\mathcal{H}^2(G, \mathbb{Z}_2)$ //Obtain all information on the second cohomology
- 2: Char = Characters of irreps of G
- 3: **for** $\omega_\Phi \in$ Coh **do**
- 4: \tilde{X}_f = The extended group of G corresponding to ω_Φ
- 5: CharExt = Characters of irreps of \tilde{X}_f
- 6: Identify the group elements of \tilde{X}_f within conjugacy classes of G
- 7: **for** $\mathcal{R}_\Phi \in$ CharExt **do**
- 8: **if** \mathcal{R}_Φ is a 1D irrep **then**
- 9: $\mathcal{R}_{pair} = \mathcal{R}_\Phi^2$
- 10: Identify $\mathcal{R}_{pair} \in$ Char
- 11: **end if**
- 12: **end for**
- 13: **end for**
- 14: **Return** All matching pairs ($\omega_\Phi \in$ Coh, $\mathcal{R}_{pair} \in$ Char)

Table 7: Correspondence between the fermion PSG and the representation of the pairing order parameter for all the crystalline point group. We list gauge-invariant cocycles to label different projective symmetry groups G_f for superconductors both without and with spin-orbital couplings. We follow the convention in Ref. [47] to label irreducible representations $\mathcal{R}_{pair}(\mathbf{g})$ of the pairing order parameter. Some G_f does not admit a 1d projective representation and hence the corresponding \mathcal{R}_{pair} is marked as N/A. For gauge invariant cocycles, we use the following short-hand notations: $\zeta_{\mathbf{g}} \equiv \omega(\mathbf{g}, \mathbf{g})$, and $\eta_{\mathbf{g},h} \equiv \frac{\omega(\mathbf{g},h)}{\omega(h,\mathbf{g})}$.

X	$\mathcal{H}^2(X, \mathbb{Z}_2^F)$	Gauge-invariant 2-cocycles	No SOC (spinless)	w/ SOC (spinful)	$\mathcal{R}_{pair}(\mathbf{g})$
C_1	\mathbb{Z}_1	—	—	—	A
C_i	\mathbb{Z}_2	ζ_i	1	1	A_g
			-1	-1	A_u
C_2	\mathbb{Z}_2	ζ_{C_2}	1	-1	A
			-1	1	B
C_s	\mathbb{Z}_2	ζ_{σ_h}	1	-1	A'
			-1	1	A''
C_{2h}	\mathbb{Z}_2^3	$(\zeta_{C_2}, \zeta_i, \zeta_{\sigma_h})$	(1, 1, 1)	(-1, 1, -1)	A_g
			(1, -1, -1)	(-1, -1, 1)	A_u
			(-1, 1, -1)	(1, 1, 1)	B_g
			(-1, -1, 1)	(1, -1, -1)	B_u
			other cases	other cases	N/A
D_2	\mathbb{Z}_2^3	$(\zeta_{C_{2x}}, \zeta_{C_{2y}}, \zeta_{C_{2z}})$	(1, 1, 1)	(-1, -1, -1)	A
			(-1, -1, 1)	(1, 1, -1)	B_1
			(-1, 1, -1)	(1, -1, 1)	B_2
			(1, -1, -1)	(-1, 1, 1)	B_3
			other cases	other cases	N/A
C_{2v}	\mathbb{Z}_2^3	$(\zeta_{C_2}, \zeta_{\sigma_v}, \zeta_{\sigma'_v})$	(1, 1, 1)	(-1, -1, -1)	A_1
			(1, -1, -1)	(-1, 1, 1)	A_2
			(-1, 1, -1)	(1, -1, 1)	B_1
			(-1, -1, 1)	(1, 1, -1)	B_2
			other cases	other cases	N/A
D_{2h}	\mathbb{Z}_2^6	$(\zeta_{C_{2x}}, \zeta_{C_{2y}}, \zeta_i, \eta_{C_{2x},i}, \eta_{C_{2y},i}, \eta_{C_{2x},C_{2y}})$	(1, 1, 1, 1, 1, 1)	(-1, -1, 1, 1, 1, -1)	A_g
			(1, -1, 1, 1, 1, 1)	(-1, 1, 1, 1, 1, -1)	B_{3g}
			(-1, 1, 1, 1, 1, 1)	(1, -1, 1, 1, 1, -1)	B_{2g}
			(-1, -1, 1, 1, 1, 1)	(1, 1, 1, 1, 1, -1)	B_{1g}
			(1, 1, -1, 1, 1, 1)	(-1, -1, -1, 1, 1, -1)	A_u
			(1, -1, -1, 1, 1, 1)	(-1, 1, -1, 1, 1, -1)	B_{3u}
			(-1, 1, -1, 1, 1, 1)	(1, -1, -1, 1, 1, -1)	B_{2u}
			(-1, -1, -1, 1, 1, 1)	(1, 1, -1, 1, 1, -1)	B_{1u}
			other cases	other cases	N/A
C_4	\mathbb{Z}_2	ζ_{C_2}	1	-1	A, B
			-1	1	E
S_4	\mathbb{Z}_2	ζ_{C_2}	1	-1	A, B
			-1	1	E

Table 7 Continued.

X	$\mathcal{H}^2(X, \mathbb{Z}_2^F)$	Gauge-invariant cocycles	No SOC	With SOC	$\mathcal{R}_{pair}(\mathfrak{g})$
C_{4h}	\mathbb{Z}_2^3	$(\zeta_{C_2}, \zeta_i, \eta_{C_{4,i}})$	(1, 1, 1)	(-1, 1, 1)	A_g, B_g
			(-1, -1, 1)	(1, -1, 1)	E_u
			(1, -1, 1)	(-1, -1, 1)	A_u, B_u
			(-1, 1, 1)	(1, 1, 1)	E_g
			other cases	other cases	N/A
D_4	\mathbb{Z}_2^3	$(\zeta_{C_2}, \zeta_{C'_2}, \zeta_{C''_2})$	(1, 1, 1)	(-1, -1, -1)	A_1
			(1, -1, -1)	(-1, 1, 1)	A_2
			(1, 1, -1)	(-1, -1, 1)	B_1
			(1, -1, 1)	(-1, 1, -1)	B_2
			other cases	other cases	N/A
C_{4v}	\mathbb{Z}_2^3	$(\zeta_{C_2}, \zeta_{\sigma_v}, \zeta_{\sigma_d})$	(1, 1, 1)	(-1, -1, -1)	A_1
			(1, 1, -1)	(-1, -1, 1)	B_1
			(1, -1, 1)	(-1, 1, -1)	B_2
			(1, -1, -1)	(-1, 1, 1)	A_2
			other cases	other cases	N/A
D_{2d}	\mathbb{Z}_2^3	$(\zeta_{C_2}, \zeta_{C'_2}, \zeta_{\sigma_d})$	(1, 1, 1)	(-1, -1, -1)	A_1
			(1, 1, -1)	(-1, -1, 1)	B_1
			(1, -1, 1)	(-1, 1, -1)	B_2
			(1, -1, -1)	(-1, 1, 1)	A_2
			other cases	other cases	N/A
D_{4h}	\mathbb{Z}_2^6	$(\zeta_{C'_2}, \zeta_{C''_2}, \zeta_i, \zeta_{C_2}, \eta_{C'_2,i}, \eta_{C''_2,i})$	(1, 1, 1, 1, 1, 1)	(-1, -1, 1, -1, 1, 1)	A_{1g}
			(1, 1, -1, 1, 1, 1)	(-1, -1, -1, -1, 1, 1)	A_{1u}
			(1, -1, 1, 1, 1, 1)	(-1, 1, 1, -1, 1, 1)	B_{1g}
			(1, -1, -1, 1, 1, 1)	(-1, 1, -1, -1, 1, 1)	B_{1u}
			(-1, -1, 1, -1, 1, 1)	(1, 1, 1, 1, 1, 1)	A_{2g}
			(-1, -1, -1, 1, 1, 1)	(1, 1, -1, -1, 1, 1)	A_{2u}
			(-1, 1, 1, 1, 1, 1)	(1, -1, 1, -1, 1, 1)	B_{2g}
			(-1, 1, -1, 1, 1, 1)	(1, -1, -1, -1, 1, 1)	B_{2u}
			other cases	other cases	N/A
C_3	\mathbb{Z}_1	—	—	—	A_1, E
C_{3i}	\mathbb{Z}_2	ζ_i	+1	+1	A_g, E_g
			-1	-1	A_u, E_u
D_3	\mathbb{Z}_2	ζ_{C_2}	+1	-1	A_1
			-1	+1	A_2
C_{3v}	\mathbb{Z}_2	ζ_{σ_v}	+1	-1	A_1
			-1	+1	A_2
D_{3d}	\mathbb{Z}_2^3	$(\zeta_{C'_2}, \zeta_i, \eta_{C'_2,i})$	(1, 1, 1)	(-1, 1, 1)	A_{1g}
			(1, -1, 1)	(-1, -1, 1)	A_{1u}
			(-1, 1, 1)	(1, 1, 1)	A_{2g}
			(-1, -1, 1)	(1, -1, 1)	A_{2u}
			other cases	other cases	N/A
C_6	\mathbb{Z}_2	ζ_{C_2}	+1	-1	A, E_1
			-1	+1	B, E_2
C_{3h}	\mathbb{Z}_2	ζ_{σ_h}	+1	-1	A', E'
			-1	1	A'', E''

Table 7 Continued.

X	$\mathcal{H}^2(X, Z_2^F)$	Gauge-invariant cocycles	No SOC	With SOC	$\mathcal{R}_{pair}(g)$
C_{6h}	Z_2^3	$(\zeta_{C_2}, \zeta_i, \eta_{C_2, i})$	(1, 1, 1)	(-1, 1, 1)	A_g, E_{1g}
			(1, -1, 1)	(-1, -1, 1)	A_u, E_{1u}
			(-1, 1, 1)	(1, 1, 1)	B_g, E_{2g}
			(-1, -1, 1)	(1, -1, 1)	B_u, E_{2u}
			other cases	other cases	N/A
D_6	Z_2^3	$(\zeta_{C_2}, \zeta_{C'_2}, \eta_{C_2, C'_2})$	(1, 1, 1)	(-1, -1, -1)	A_1
			(1, -1, 1)	(-1, 1, -1)	A_2
			(-1, 1, 1)	(1, -1, -1)	B_2
			(-1, -1, 1)	(1, 1, -1)	B_1
			other cases	other cases	N/A
C_{6v}	Z_2^3	$(\zeta_{C_2}, \zeta_{\sigma_v}, \eta_{C_2, \sigma_v})$	(1, 1, 1)	(-1, -1, -1)	A_1
			(1, -1, 1)	(-1, 1, -1)	A_2
			(-1, 1, 1)	(1, -1, -1)	B_2
			(-1, -1, 1)	(1, 1, -1)	B_1
			other cases	other cases	N/A
D_{3h}	Z_2^3	$(\zeta_{\sigma_v}, \zeta_{\sigma_h}, \eta_{\sigma_h, \sigma_v})$	(1, 1, 1)	(-1, -1, -1)	A'_1
			(1, -1, 1)	(-1, 1, -1)	A''_2
			(-1, 1, 1)	(1, -1, -1)	A''_2
			(-1, -1, 1)	(1, 1, -1)	A'_1
			other cases	other cases	N/A
D_{6h}	Z_2^6	$(\zeta_{C_2}, \zeta_{C'_2}, \zeta_i, \eta_{C_2, C'_2}, \eta_{C_2, i}, \eta_{C'_2, i})$	(1, 1, 1, 1, 1, 1)	(-1, -1, 1, -1, 1, 1)	A_{1g}
			(1, 1, -1, 1, 1, 1)	(-1, -1, -1, -1, 1, 1)	A_{1u}
			(1, -1, 1, 1, 1, 1)	(-1, 1, 1, -1, 1, 1)	A_{2g}
			(1, -1, -1, 1, 1, 1)	(-1, 1, -1, -1, 1, 1)	A_{2u}
			(-1, 1, 1, 1, 1, 1)	(1, -1, 1, -1, 1, 1)	B_{2g}
			(-1, 1, -1, 1, 1, 1)	(1, -1, -1, -1, 1, 1)	B_{2u}
			(-1, -1, 1, 1, 1, 1)	(1, 1, 1, -1, 1, 1)	B_{1g}
			(-1, -1, -1, 1, 1, 1)	(1, 1, -1, -1, 1, 1)	B_{1u}
other cases	other cases	N/A			
T	Z_2	ζ_{C_2}	1	-1	A, E
			-1	1	N/A
T_h	Z_2^2	(ζ_{C_2}, ζ_i)	(1, 1)	(-1, 1)	A_g, E_g
			(1, -1)	(-1, -1)	A_u, E_u
			other cases	other cases	N/A
O	Z_2^2	$(\zeta_{C_2}, \zeta_{C'_2})$	(1, 1)	(-1, -1)	A_1
			(1, -1)	(-1, 1)	A_2
			other cases	other cases	N/A
T_d	Z_2^2	$(\zeta_{C_2}, \zeta_{\sigma_d})$	(1, 1)	(-1, -1)	A_1
			(1, -1)	(-1, 1)	A_2
			other cases	other cases	N/A
O_h	Z_2^4	$(\zeta_{C_2}, \zeta_{C'_2}, \zeta_i, \eta_{i, C'_2})$	(1, 1, 1, 1)	(-1, -1, 1, 1)	A_{1g}
			(1, 1, -1, 1)	(-1, -1, -1, 1)	A_{1u}
			(1, -1, 1, 1)	(-1, 1, 1, 1)	A_{2g}
			(1, -1, -1, 1)	(-1, 1, -1, 1)	A_{2u}
			other cases	other cases	N/A

References

- 835
- 836 [1] G. E. Volovik and L. P. Gorkov, *Superconducting classes in heavy-fermion systems*, Zh. Eksp.
837 Teor. Fiz. **88**, 1412 (1985).
- 838 [2] M. Sigrist and K. Ueda, *Phenomenological theory of unconventional superconductivity*, Rev.
839 Mod. Phys. **63**, 239 (1991), doi:[10.1103/RevModPhys.63.239](https://doi.org/10.1103/RevModPhys.63.239).
- 840 [3] J. F. Annett, *Symmetry of the order parameter for high-temperature superconductivity*, Adv.
841 Phys. **39**(2), 83 (1990), doi:[10.1080/00018739000101481](https://doi.org/10.1080/00018739000101481).
- 842 [4] M. Sigrist, *Introduction to unconventional superconductivity*, AIP Conf. Proc. **789**(1), 165
843 (2005), doi:[10.1063/1.2080350](https://doi.org/10.1063/1.2080350).
- 844 [5] A. J. Leggett, *A theoretical description of the new phases of liquid ^3He* , Rev. Mod. Phys.
845 **47**, 331 (1975), doi:[10.1103/RevModPhys.47.331](https://doi.org/10.1103/RevModPhys.47.331).
- 846 [6] C. Pfleiderer, *Superconducting phases of f -electron compounds*, Rev. Mod. Phys. **81**, 1551
847 (2009), doi:[10.1103/RevModPhys.81.1551](https://doi.org/10.1103/RevModPhys.81.1551).
- 848 [7] C. C. Tsuei and J. R. Kirtley, *Pairing symmetry in cuprate superconductors*, Rev. Mod. Phys.
849 **72**, 969 (2000), doi:[10.1103/RevModPhys.72.969](https://doi.org/10.1103/RevModPhys.72.969).
- 850 [8] A. P. Mackenzie and Y. Maeno, *The superconductivity of Sr_2RuO_4 and the physics of spin-*
851 *triplet pairing*, Rev. Mod. Phys. **75**, 657 (2003), doi:[10.1103/RevModPhys.75.657](https://doi.org/10.1103/RevModPhys.75.657).
- 852 [9] S. Ryu, A. P. Schnyder, A. Furusaki and A. W. W. Ludwig, *Topological insulators and*
853 *superconductors: tenfold way and dimensional hierarchy*, New J. Phys. **12**, 065010 (2010),
854 doi:[10.1088/1367-2630/12/6/065010](https://doi.org/10.1088/1367-2630/12/6/065010).
- 855 [10] A. Kitaev, *Periodic table for topological insulators and superconductors*, AIP Conf. Proc.
856 **1134**, 22 (2009), doi:[10.1063/1.3149495](https://doi.org/10.1063/1.3149495).
- 857 [11] C.-K. Chiu, J. C. Y. Teo, A. P. Schnyder and S. Ryu, *Classification of topo-*
858 *logical quantum matter with symmetries*, Rev. Mod. Phys. **88**, 035005 (2016),
859 doi:[10.1103/RevModPhys.88.035005](https://doi.org/10.1103/RevModPhys.88.035005).
- 860 [12] L. Fidkowski and A. Kitaev, *Effects of interactions on the topological classification of free*
861 *fermion systems*, Phys. Rev. B **81**, 134509 (2010), doi:[10.1103/PhysRevB.81.134509](https://doi.org/10.1103/PhysRevB.81.134509).
- 862 [13] C. Wang and T. Senthil, *Interacting fermionic topological insulators/superconductors in*
863 *three dimensions*, Phys. Rev. B **89**, 195124 (2014), doi:[10.1103/PhysRevB.89.195124](https://doi.org/10.1103/PhysRevB.89.195124).
- 864 [14] Q.-R. Wang and Z.-C. Gu, *Construction and classification of symmetry-protected topo-*
865 *logical phases in interacting fermion systems*, Phys. Rev. X **10**, 031055 (2020),
866 doi:[10.1103/PhysRevX.10.031055](https://doi.org/10.1103/PhysRevX.10.031055).
- 867 [15] N. Read and D. Green, *Paired states of fermions in two dimensions with breaking of parity*
868 *and time-reversal symmetries and the fractional quantum hall effect*, Phys. Rev. B **61**, 10267
869 (2000), doi:[10.1103/PhysRevB.61.10267](https://doi.org/10.1103/PhysRevB.61.10267).
- 870 [16] J. C. Y. Teo and C. L. Kane, *Topological defects and gapless modes in insulators and super-*
871 *conductors*, Phys. Rev. B **82**, 115120 (2010), doi:[10.1103/PhysRevB.82.115120](https://doi.org/10.1103/PhysRevB.82.115120).

- 872 [17] J. C. Teo and T. L. Hughes, *Topological defects in symmetry-protected topological phases*, Annual Review of Condensed Matter Physics **8**(1), 211 (2017),
873 doi:[10.1146/annurev-conmatphys-031016-025154](https://doi.org/10.1146/annurev-conmatphys-031016-025154), <https://doi.org/10.1146/annurev-conmatphys-031016-025154>,
874 <https://doi.org/10.1146/annurev-conmatphys-031016-025154>.
875
- 876 [18] S. Kobayashi, Y. Yamazaki, A. Yamakage and M. Sato, *Majorana multipole response: General theory and application to wallpaper groups*, Phys. Rev. B **103**, 224504 (2021),
877 doi:[10.1103/PhysRevB.103.224504](https://doi.org/10.1103/PhysRevB.103.224504).
878
- 879 [19] X. Yang, S. Ono, S. Biswas, M. Randeria and Y.-M. Lu, *Manuscript under preparation*
880 (2024).
- 881 [20] S. Lu, X. Yang and Y.-M. Lu, *Optical and raman selection rules for odd-parity clean superconductors*, Phys. Rev. B **109**, 245119 (2024), doi:[10.1103/PhysRevB.109.245119](https://doi.org/10.1103/PhysRevB.109.245119).
882
- 883 [21] X.-G. Wen, *Quantum orders and symmetric spin liquids*, Phys. Rev. B **65**, 165113 (2002),
884 doi:[10.1103/PhysRevB.65.165113](https://doi.org/10.1103/PhysRevB.65.165113).
- 885 [22] D. Vollhardt and P. Woelfle, *The Superfluid Phases of Helium 3*, CRC Press, London, 1st
886 edition edn., doi:[10.1201/b12808](https://doi.org/10.1201/b12808) (1990).
- 887 [23] F. Wang and A. Vishwanath, *Spin-liquid states on the triangular and kagomé lattices: A projective-symmetry-group analysis of schwinger boson states*, Phys. Rev. B **74**, 174423
888 (2006), doi:[10.1103/PhysRevB.74.174423](https://doi.org/10.1103/PhysRevB.74.174423).
889
- 890 [24] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.12.2* (2022).
- 891 [25] E. Lake, A. S. Patri and T. Senthil, *Pairing symmetry of twisted bilayer graphene: A phenomenological synthesis*, Phys. Rev. B **106**, 104506 (2022),
892 doi:[10.1103/PhysRevB.106.104506](https://doi.org/10.1103/PhysRevB.106.104506).
893
- 894 [26] Y. Cao, D. Rodan-Legrain, J. M. Park, N. F. Q. Yuan, K. Watanabe, T. Taniguchi, R. M. Fernandes, L. Fu and P. Jarillo-Herrero, *Nematicity and competing orders in superconducting magic-angle graphene*, Science **372**(6539), 264 (2021), doi:[10.1126/science.abc2836](https://doi.org/10.1126/science.abc2836),
895 <https://www.science.org/doi/pdf/10.1126/science.abc2836>.
896
897
- 898 [27] X. Xi, Z. Wang, W. Zhao, J.-H. Park, K. T. Law, H. Berger, L. Forró, J. Shan and K. F. Mak, *Ising pairing in superconducting nbse2 atomic layers*, Nature Physics **12**, 139 (2016),
899 doi:[10.1038/nphys3538](https://doi.org/10.1038/nphys3538).
900
- 901 [28] K. Shiozaki, *The classification of surface states of topological insulators and superconductors with magnetic point group symmetry*, Progress of Theoretical and Experimental Physics **2022**(4), 04A104 (2022), doi:[10.1093/ptep/ptep026](https://doi.org/10.1093/ptep/ptep026).
902
903
- 904 [29] A. Altland and M. R. Zirnbauer, *Nonstandard symmetry classes in mesoscopic normal-superconducting hybrid structures*, Phys. Rev. B **55**, 1142 (1997),
905 doi:[10.1103/PhysRevB.55.1142](https://doi.org/10.1103/PhysRevB.55.1142).
906
- 907 [30] A. P. Schnyder, S. Ryu, A. Furusaki and A. W. W. Ludwig, *Classification of topological insulators and superconductors in three spatial dimensions*, Phys. Rev. B **78**, 195125 (2008),
908 doi:[10.1103/PhysRevB.78.195125](https://doi.org/10.1103/PhysRevB.78.195125).
909
- 910 [31] A. Kapustin, R. Thorngren, A. Turzillo and Z. Wang, *Fermionic symmetry protected topological phases and cobordisms*, Journal of High Energy Physics **2015**(12), 1 (2015),
911 doi:[10.1007/JHEP12\(2015\)052](https://doi.org/10.1007/JHEP12(2015)052).
912

- 913 [32] Y. Ando and L. Fu, *Topological crystalline insulators and topological superconductors:*
914 *From concepts to materials*, Annual Review of Condensed Matter Physics **6**(1), 361
915 (2015), doi:[10.1146/annurev-conmatphys-031214-014501](https://doi.org/10.1146/annurev-conmatphys-031214-014501), <https://doi.org/10.1146/annurev-conmatphys-031214-014501>.
916
- 917 [33] N. Okuma, M. Sato and K. Shiozaki, *Topological classification under nonmag-*
918 *netic and magnetic point group symmetry: Application of real-space atiyah-hirzebruch*
919 *spectral sequence to higher-order topology*, Phys. Rev. B **99**, 085127 (2019),
920 doi:[10.1103/PhysRevB.99.085127](https://doi.org/10.1103/PhysRevB.99.085127).
- 921 [34] S. Ono, K. Shiozaki and H. Watanabe, *Classification of time-reversal symmetric topo-*
922 *logical superconducting phases for conventional pairing symmetries*, arXiv e-prints
923 arXiv:2206.02489 (2022), doi:[10.48550/arXiv.2206.02489](https://doi.org/10.48550/arXiv.2206.02489), [2206.02489](https://arxiv.org/abs/2206.02489).
- 924 [35] Z. Zhang, J. Ren, Y. Qi and C. Fang, *Topological classification of intrinsic three-dimensional*
925 *superconductors using anomalous surface construction*, Phys. Rev. B **106**, L121108 (2022),
926 doi:[10.1103/PhysRevB.106.L121108](https://doi.org/10.1103/PhysRevB.106.L121108).
- 927 [36] K. Shiozaki and S. Ono, *Atiyah-Hirzebruch spectral sequence for topological insula-*
928 *tors and superconductors: E_2 pages for 1651 magnetic space groups*, arXiv e-prints
929 arXiv:2304.01827 (2023), doi:[10.48550/arXiv.2304.01827](https://doi.org/10.48550/arXiv.2304.01827), [2304.01827](https://arxiv.org/abs/2304.01827).
- 930 [37] S. Ono and K. Shiozaki, *Towards complete characterization of topological insula-*
931 *tors and superconductors: A systematic construction of topological invariants based*
932 *on Atiyah-Hirzebruch spectral sequence*, arXiv e-prints arXiv:2311.15814 (2023),
933 doi:[10.48550/arXiv.2311.15814](https://doi.org/10.48550/arXiv.2311.15814), [2311.15814](https://arxiv.org/abs/2311.15814).
- 934 [38] C.-K. Chiu, H. Yao and S. Ryu, *Classification of topological insulators and super-*
935 *conductors in the presence of reflection symmetry*, Phys. Rev. B **88**, 075142 (2013),
936 doi:[10.1103/PhysRevB.88.075142](https://doi.org/10.1103/PhysRevB.88.075142).
- 937 [39] T. Morimoto and A. Furusaki, *Topological classification with additional symmetries from*
938 *clifford algebras*, Phys. Rev. B **88**, 125129 (2013), doi:[10.1103/PhysRevB.88.125129](https://doi.org/10.1103/PhysRevB.88.125129).
- 939 [40] K. Shiozaki and M. Sato, *Topology of crystalline insulators and superconductors*, Phys. Rev.
940 B **90**, 165114 (2014), doi:[10.1103/PhysRevB.90.165114](https://doi.org/10.1103/PhysRevB.90.165114).
- 941 [41] Y.-M. Lu and D.-H. Lee, *Inversion symmetry protected topological insulators and super-*
942 *conductors*, arXiv e-prints arXiv:1403.5558 (2014), doi:[10.48550/arXiv.1403.5558](https://doi.org/10.48550/arXiv.1403.5558),
943 [1403.5558](https://arxiv.org/abs/1403.5558).
- 944 [42] R. Balian and N. R. Werthamer, *Superconductivity with pairs in a relative \mathbf{p} wave*, Phys.
945 Rev. **131**, 1553 (1963), doi:[10.1103/PhysRev.131.1553](https://doi.org/10.1103/PhysRev.131.1553).
- 946 [43] P. W. Anderson and P. Morel, *Generalized bardeen-cooper-schrieffer states and the*
947 *proposed low-temperature phase of liquid He^3* , Phys. Rev. **123**, 1911 (1961),
948 doi:[10.1103/PhysRev.123.1911](https://doi.org/10.1103/PhysRev.123.1911).
- 949 [44] P. W. Anderson and W. F. Brinkman, *Anisotropic superfluidity in ^3He : A possible inter-*
950 *pretation of its stability as a spin-fluctuation effect*, Phys. Rev. Lett. **30**, 1108 (1973),
951 doi:[10.1103/PhysRevLett.30.1108](https://doi.org/10.1103/PhysRevLett.30.1108).
- 952 [45] R. Verresen, U. Borla, A. Vishwanath, S. Moroz and R. Thorngren, *Higgs Conden-*
953 *sates are Symmetry-Protected Topological Phases: I. Discrete Symmetries*, arXiv e-prints
954 arXiv:2211.01376 (2022), doi:[10.48550/arXiv.2211.01376](https://doi.org/10.48550/arXiv.2211.01376), [2211.01376](https://arxiv.org/abs/2211.01376).

- 955 [46] T. Hansson, V. Oganesyan and S. Sondhi, *Superconductors are topologically ordered*, *Annals of Physics* **313**(2), 497 (2004), doi:<https://doi.org/10.1016/j.aop.2004.05.006>.
956
- 957 [47] C. J. Bradley and A. P. Cracknell, *The Mathematical Theory Of Symmetry In Solids: Representation theory for point groups and space groups*, Oxford University Press, ISBN
958 9780199582587, doi:[10.1093/oso/9780199582587.001.0001](https://doi.org/10.1093/oso/9780199582587.001.0001) (2009).
959