

Replica symmetry breaking in spin glasses in the replica-free Keldysh formalism

Johannes Lang^{1*}, Subir Sachdev², Sebastian Diehl¹

¹ Institut für Theoretische Physik, Universität zu Köln, Zùlpicher Straße 77, 50937
Cologne, Germany

² Department of Physics, Harvard University, Cambridge MA 02138, USA

* j.lang@uni-koeln.de

June 19, 2024

1 Abstract

At asymptotically late times ultrametricity can emerge from the persistent slow aging dynamics of the glass phase. We show that this suffices to recover the breaking of replica symmetry in mean-field spin glasses from the late time limit of the time evolution using the Keldysh path integral. This provides an alternative approach to replica symmetry breaking by connecting it rigorously to the dynamic formulation. Stationary spin glasses are thereby understood to spontaneously break thermal symmetry, or the Kubo-Martin-Schwinger relation of a state in global thermal equilibrium. We demonstrate our general statements for the spherical quantum p -spin model and the quantum Sherrington-Kirkpatrick model in the presence of transverse and longitudinal fields. In doing so, we also derive their dynamical Ginzburg-Landau effective Keldysh actions starting from microscopic quantum models..

14

15 Contents

16	1 Introduction	2
17	2 Equivalence of ultrametric Keldysh dynamics and replica symmetry breaking	3
18	2.1 Replica formulation	3
19	2.2 Dynamic theory	5
20		
21	3 Application: The Quantum Sherrington-Kirkpatrick model	11
22	3.1 Effective action	12
23	3.2 Paramagnetic phase	15
24	3.3 Landau action to order u^2	17
25	3.4 Asymptotic solution in the glass phase	18
26	4 Application: The spherical p-spin model	20
27	4.1 Effective action	20
28	4.2 Late-time solution	22
29	5 Discussion	25
30	5.1 Spontaneous breaking of thermal symmetry	25
31	5.2 Zero temperature limit	26

32	6 Outlook	26
33	References	27

34

35

36 1 Introduction

37 The characteristic property of glasses is their slow evolution. As the system approaches
38 the equilibrium state, its evolution becomes increasingly restricted by barriers in the free
39 energy landscape [1–4] that take more and more time to overcome. As the time since the
40 quench increases, relaxation slows down – the system ‘ages’. One finds that accompanying
41 this behavior is an ultrametric structure in the time dependence of correlations [5, 6].
42 Because the dynamic constraints depend on the age of the glass, contrary to most other
43 systems, it develops a sufficiently strong long-term memory for the age of the system
44 to forever remain a relevant time scale [7]. Consequently, aging precludes glasses from
45 reaching thermal equilibrium on accessible time scales [8–13].

46 Simultaneously, the analysis of the putative equilibrium state in systems with quenched
47 disorder has brought forth many surprises, most prominently the breaking of replica sym-
48 metry [14–16], indicating the fragmentation of configuration space into disconnected en-
49 ergetically equivalent regions separated by insurmountable free energy barriers [2, 17, 18].
50 This fragmentation of the phase space breaks ergodicity [19] and gives rise to an ultra-
51 metric structure, observable in correlations [20].

52 Although theoretical research has focused largely on the simplest models exhibiting
53 glassy behavior, namely spin systems with infinite-ranged interactions and quenched dis-
54 order, a connection to fragile glasses exists in mode coupling theory [21–24]. Although
55 lacking a rigorous derivation, numerical [25–28] and experimental evidence [29] support
56 its conclusion that the characteristic properties of mean-field spin glasses carry over to
57 systems with short-ranged interactions and annealed disorder in finite dimensions.

58 From the previous arguments, it is clear that aging dynamics and the absence of
59 ergodicity are related phenomena [17]. In fact, in mean-field spin glasses, several attempts
60 to formalize this relation have been made [30–34]. Nevertheless, despite decades of intense
61 research, it remains unclear how, or even if, the equilibrium solution is connected to the
62 dynamics [13, 35, 36].

63 We aim to make this similarity more concrete by showing that after a quench, at
64 infinitely late times, the dynamic description eventually reproduces the equilibrium results.
65 It is important to point out that the infinite-time limit is taken at the beginning and the
66 equilibrium result is not necessarily smoothly connected to any results obtained at finite
67 times. In particular, we do not claim that the evolution ever reaches the equilibrium state.

68 We present our main result in Sec. 2. There, we show that the algebraic properties
69 of Parisi matrices, characterizing the fragmentation of configuration space, are recovered
70 in the Keldysh formalism under the assumption of a strong hierarchy of time scales. The
71 result then is applied in Sec. 3 to the quantum Sherrington-Kirkpatrick model in a longi-
72 tudinal field and to the spherical quantum p -spin model in Sec. 4. Our approach exposes
73 the spontaneous breaking of thermal symmetry (or the Kubo-Martin-Schwinger relation of
74 a state in thermodynamic equilibrium) as the origin of replica symmetry breaking. This,
75 however, is independent of the breaking of time-translation invariance as emphasized in
76 Sec. 5. There we also apply constraints to the potential quantum critical scaling at zero
77 temperature. In Sec. 6, we conclude with an outlook discussing the connection to glasses

78 of a finite age and to the zero-temperature limit.

79 2 Equivalence of ultrametric Keldysh dynamics and replica 80 symmetry breaking

81 The proposal of a connection between replicas and the classical Langevin theory of spin
82 glasses goes back to the classic early work of Sompolinsky and Zippelius [7, 30, 37]. They
83 proposed that replica symmetry breaking was associated with multiple exponentially long
84 time scales which diverged as the thermodynamic limit was taken. Later, Cugliandolo and
85 Kurchan [5, 6, 38, 39] showed that the classical equations exhibited ‘aging’ dynamics [40]
86 in which the time scales remained finite, although exponentially long, even in the ther-
87 modynamic limit: they established an explicit connection between the aging equations
88 and the replica symmetry breaking of the static problem. The aging dynamics was ex-
89 tended to quantum p -spin spherical models [36, 41, 42] and large M_s $SU(M_s)$ quantum
90 Heisenberg spin models [43] with one-step replica symmetry breaking using the Keldysh
91 formalism: in the slow dynamics regime, the Keldysh equations became identical to the
92 classical Langevin equations. The important case of the quantum Sherrington-Kirkpatrick
93 Ising model, *i.e.* the Ising spin glass in a transverse field, was briefly discussed by Kennett
94 *et al.* [44, 45].

95 This section will present a general and model-independent discussion of the connection
96 between replicas and glassy dynamics. The results apply to quantum and classical spin
97 glasses with possibly full replica symmetry breaking, including the recently studied quan-
98 tum Ising spin glass in both transverse and longitudinal fields with an Almeida-Thouless
99 transition [46], and the $SU(2)$ quantum Heisenberg spin glass [47]. A related connection
100 between supersymmetry and thermal symmetry in the paramagnetic phase of spin glasses
101 was previously found by Kurchan [48].

102 We connect the Parisi spin-glass order parameter characterized by the function $p(u)$,
103 $0 \leq u \leq 1$, in Eq. (1) to the effective time-dependent (half) inverse temperature $X(t)$,
104 defined by Eq. (10). The deviation of $X(t)$ from its equilibrium value $\beta/2$ measures
105 the breaking of the fluctuation-dissipation relation by the glassy dynamics. For each
106 $u < 1$, there is a t which is determined by the solution of Eq. (15), where β is the inverse
107 temperature; smaller u corresponds to larger t , with $u = 1$ mapping to $t = 0$ ($X(0) = \beta/2$),
108 and $u = 0$ mapping to $t = \infty$ ($X(\infty) = 0$).

109 The analogy between the two approaches is complete in the sense that the algebra of
110 ultrametric matrices in Eqs. (3), (4) and (5) is recovered by real-time the Dyson-Keldysh
111 equations in the glassy limit under the assumption of ultrametricity in Eqs. (13), (18) and
112 (20).

113 2.1 Replica formulation

114 On the replica side of the correspondence, we need to recall the algebraic relations satisfied
115 by Parisi matrices, which we then aim to recover in the late-time limit of the dynamical
116 equations.

117 For completeness, we begin by introducing the Parisi matrix P_{ab} and the equivalent
118 Parisi function $p(u)$ $u \in [0, 1]$. Consider some model with N replicas. Its equilibrium
119 correlation functions are N -dimensional square matrices in replica space with an intriguing
120 structure in the physical limit $N \rightarrow 0$. To capture this structure, we define that the
121 N -dimensional symmetric matrix P is called a Parisi matrix if a sequence of integers
122 $\mathcal{N} = \{n_1, n_2, \dots, n_{L-1}\}$ with $n_1 = 1$ exists such that $P_{ab} = p_i$ for $\left[\frac{a-1}{m_i}\right] \neq \left[\frac{b-1}{m_i}\right]$, but

123 $\left[\frac{a-1}{m_{i+1}}\right] = \left[\frac{b-1}{m_{i+1}}\right]$, where $m_i = \prod_{j=1}^i n_j$ and $m_L = N$. Furthermore, we fix the diagonal
 124 to $P_{aa} = p_0$. Simply put, a Parisi matrix consists of a hierarchy of block matrices placed
 125 along the diagonal such that each block itself is again a Parisi matrix (see Fig. 1(g)).
 126 If P is identified with the correlation function, it describes the formation of clusters in
 127 replica space. If the p_i form a decreasing sequence, different realizations of the system
 128 within a cluster are more strongly correlated with each other, than with replicas from
 129 other clusters. Thus, unless the sequence \mathcal{N} contains only one element, the Parisi matrix
 130 describes the breaking of ergodicity.

131 The simple structure of P allows it to be rewritten in terms of the equivalent Parisi
 132 function

$$p(u) = p_i \quad \text{if } m_i < u < m_{i+1} \quad (1)$$

133 with $p(1) = p_0$. Since $m_1 = 1$ and $m_L = N$, with all other values in between, in the
 134 replica limit $N \rightarrow 0$ one has $u \in [0, 1]$, see Fig. 1(f). Note, that due to the limit $N \rightarrow 0$
 135 smaller values of u correspond to terms farther from the diagonal of P . Thus, inverting
 136 (1), $(dp(u)/du)^{-1}$ gives the probability of finding the value p in the Parisi matrix. Again,
 137 if P is interpreted as a correlation function, this determines the probability distribution
 138 of correlations between different realizations.

139 We define an ultrametric space as a metric space M in which the triangle inequality
 140 is replaced by the strong triangle inequality

$$d_{ab} \leq \max\{d_{ac}, d_{cb}\} \quad \forall c, a, b \in M. \quad (2)$$

141 This implies that there are no points between a and b , meaning that all points closer
 142 to a than b are at least a distance d_{ab} from b . The space M thus appears fractured
 143 into a hierarchy of clusters, such that on each level every point is a member of only one
 144 cluster [17]. If the p_i are a decreasing sequence, replica space with the Parisi matrix P as
 145 a measure of the inverse distance is ultrametric, meaning P satisfies (2) with $d_{ab} = 1/P_{ab}$.
 146 The hierarchical structure and the ultrametric condition can then both be read off in
 147 Fig. 1(g).

148 With these definitions, it immediately follows that the Hadamard (or component-wise)
 149 product of two Parisi matrices A and B is again a Parisi matrix C with

$$c(u) = a(u)b(u). \quad (3)$$

150 Following some algebra (for a detailed derivation, see for example [49]), one finds that the
 151 same is true for matrix multiplication, for which one finds in the limit $N \rightarrow 0$

$$\begin{aligned}
 c(u) = & a(u)b(1) + a(1)b(u) - ua(u)b(u) \\
 & - \int_u^1 dv (a(u)b(v) + a(v)b(u)) - \int_0^u dv a(v)b(v).
 \end{aligned} \quad (4)$$

152 Specifically, for the diagonal in replica space, the result simplifies to

$$c(1) = a(1)b(1) - \int_0^1 dv a(v)b(v). \quad (5)$$

153 Some intuition for the interpretation of the Parisi function can be gained by considering
 154 the unmagnetized ergodic case without replica symmetry breaking. In this case, the Parisi
 155 matrix P is diagonal and therefore $p(u) \sim \delta_{1u}$, with δ_{ij} the Kronecker delta. In Fig. 1,
 156 this corresponds to the case with $p_1 = p_2 = p_3 = 0$. This is to be compared with a
 157 ferromagnetic or magnetized phase for which all $p_{n>0}$ are identical but non-zero. We

158 point out that this ergodic solution preserves replica symmetry as P is invariant under
 159 permutations of its indices. Hence, although indistinguishable in terms of the Edwards-
 160 Anderson order parameter [50] $p_{EA} \equiv p(1^-) \equiv p_1$ alone, this gives a clear differentiation
 161 between the ferromagnetic and the spin-glass phase.

162 In general, Parisi functions are not continuous, and for practical purposes, it is often
 163 useful to write $p(u) = p_s(u) + p_f \delta_{1u}$, where $p_s(u)$ is continuous in the limit of $u \rightarrow 1$. In
 164 particular, $p_0 = p_1 + p_f$, $p_s(1) = p_1$, i.e., both p_0 and $p_s(1)$ involve the order parameter
 165 p_1 . Due to the absence of aging in the ergodic phase, it is natural to expect at late
 166 times a relation between the off-diagonal terms $p_s(u)$ in replica space with the slow aging
 167 component of the evolution. Simultaneously, there should be a connection between the
 168 replica diagonal p_f and the fast evolution that at late times becomes independent of the
 169 age of the system. In the following, we will show under which conditions these relations
 170 can be made rigorous.

171 2.2 Dynamic theory

172 We now show that the same rules of computing the Hadamard (or component-wise) prod-
 173 uct and the matrix multiplication of two Parisi matrices are also obtained from a dynamical
 174 approach under the assumption of ultrametricity.

175 Let us therefore turn our attention to the dynamics following a quench, which is de-
 176 scribed in terms of Green's functions and self-energies in Keldysh space (for an introduction
 177 see [51])

$$G = \begin{pmatrix} G^K & G^R \\ G^A & 0 \end{pmatrix} \quad \text{and} \quad \Sigma = \begin{pmatrix} 0 & \Sigma^A \\ \Sigma^R & \Sigma^K \end{pmatrix}. \quad (6)$$

178 In general, these are two-point functions that depend on two times. However, as we send
 179 the time passed since the quench to infinity, the center-of-mass time becomes but an overall
 180 scale that drops out. This will become clear in Sec. 3. For example, for the correlation
 181 function of Ising spins, we write

$$\langle s(t_1)s(t_2) \rangle \equiv G^K(t_1, t_2) = G^K(T, t) \xrightarrow{T \rightarrow \infty} G^K(t), \quad (7)$$

182 with a coarse-grained real scalar variable s . In the second step we have transformed to
 183 Wigner coordinates, i.e., to center-of-mass and relative time $T = (t_1 + t_2)/2$ and $t = t_1 - t_2$
 184 (see Fig. 1(a)). We then split the the Green's functions into a fast part that equilibrates
 185 at late times, and a slow part that describes aging

$$G(t) = G_s(t) + G_f(t) \equiv \int_{-\Lambda/b}^{\Lambda/b} d\omega e^{-i\omega t} G(\omega) \\ + \underbrace{\int_{\Lambda/b}^{\Lambda} d\omega e^{-i\omega t} G(\omega) + \int_{-\Lambda}^{-\Lambda/b} d\omega e^{-i\omega t} G(\omega)}_{=G_f(t)}. \quad (8)$$

186 Here Λ is a high-frequency cutoff. From the view of the scales of the aging variables (index
 187 s) $b \ll 1$, whereas b can be sent to 1 from the view of the fast variables (this makes sense
 188 particularly when the scale separation between aging and stationary field diverges with
 189 $T \rightarrow \infty$). By this construction, the fast field has support in the time domain on scales
 190 $1/\Lambda \leq |t| \leq b/\Lambda$, and the slow one varies with time for $|t| \geq b/\Lambda$, while it is constant
 191 for $|t| \leq b/\Lambda$. We therefore identify $b/\Lambda = \tau_{\text{erg}}$ as the time scale up to which correlations
 192 are ergodic. Furthermore, we anticipate G_f and G_s for $t \sim b/\Lambda$ to correspond to p_f and

193 p_1 for the appropriate Parisi function $p(u)$. The emergence of the scale τ_{erg} for finite T
 194 implying imperfect separation between G_f and G_s is shown in Fig. 1(b). The boundary
 195 value $G^K(t = 2T)$ vanishes as $T \rightarrow \infty$.

196 Next, we address the response to an external (longitudinal) field h which is given by
 197 the retarded Green's function

$$G^R(t_1, t_2) = \frac{\delta \langle s(t_2) \rangle}{\delta h(t_1)}. \quad (9)$$

198 Since the advanced Green's function for real scalar theories can be expressed as $G^A(t_1, t_2) =$
 199 $G^R(t_2, t_1)$, the dynamic theory can be formulated in terms of the two real functions
 200 $G^R(t_1, t_2)$ and $G^K(t_1, t_2)$. Due to causality, the former vanishes for negative relative
 201 times $t < 0$. In the limit $T \rightarrow \infty$, it can therefore be written in the form of a generalized
 202 thermal ansatz [6]

$$G_s^R(t) = -X(t)\theta(t)\partial_t G_s^K(t) = G_s^A(-t), \quad (10)$$

203 where $\theta(t)$ is the Heaviside function and $X(t)$ plays the role of a time-dependent inverse
 204 temperature in the high-temperature expansion of the fluctuation-dissipation relation

$$\begin{aligned} G^R(t) &= -\theta(t) \tan\left(\frac{\beta}{2}\partial_t\right) G^K(t) \\ &= -\theta(t) \frac{\beta}{2} \partial_t G^K(t) + \mathcal{O}(\beta^2). \end{aligned} \quad (11)$$

205 We emphasize that this interpretation becomes particularly meaningful at late times when
 206 the characteristic time scales of the evolution satisfy $\Delta t \gg \beta$, which justifies the expansion
 207 in powers of the inverse temperature β .

208 From its definition (7), it is clear that $G^K(t_1, t_2)$ is symmetric under exchange of t_1 and
 209 t_2 . Without loss of generality, the dynamic theory can therefore be restricted to $t > 0$. At
 210 short relative times ($t < \tau_{\text{erg}}$), all correlations have equilibrated, which fixes the boundary
 211 condition $X(0) = \beta/2$. But at large values of t that diverge as the inverse infrared cutoff
 212 T is sent to infinity, the system becomes increasingly fragmented and thus unresponsive.
 213 Hence, we expect the correlations to decay slowly in t and X to be a decreasing function
 214 of t , see Fig. 1(b). The ansatz (10) is consistent with that of Ref. [6] and a generalization
 215 of the one used in Ref. [35]. From the expansion (11) it also follows that the ansatz (10)
 216 corresponds to a restriction of the Parisi function to the first Matsubara frequency in the
 217 equilibrium approach. Due to the exceedingly slow dynamics in the aging regime, following
 218 the same argument as in the expansion in Eq. (10) this ansatz becomes exact in the limit
 219 $T \rightarrow \infty$ at any finite temperature.

220 Finally, we make the assumption of strong hierarchy [30], which is to say that correla-
 221 tions vary so slowly in time that $G_s^K(t) < G_s^K(t')$ requires $\lim_{T \rightarrow \infty} t/t' = 0$. This implies
 222 that correlations are ultrametric since

$$G_s^K(t + t') = G_s^K(\max(t, t')) \quad (12)$$

223 satisfies the strong triangle inequality $G_s^K(t) \geq \min\{G_s^K(t - \tau), G_s^K(\tau)\} \forall \tau \in \mathbb{R}$. Each
 224 value of G_s^K can be assigned to a characteristic time scale. In case of infinitely many time
 225 scales Eq. (12) is also the only dependence on relative time t in the limit of $T \rightarrow \infty$ that
 226 is consistent with aging dynamics [6]. Conversely, if only a finite number of time scales
 227 emerges, this is not expected to hold true for the late-time dynamics [5]. We will see below
 228 in Sec. 4, that this implies that in the thermodynamic limit a quench in the spin-glass
 229 phase of the spherical p -spin model never reaches the equilibrium.




Figure 1: Illustration of the structural similarity between the dynamical theory at asymptotically late times and replica theory. (a) For the time domain, we work in Wigner coordinates T, t . (b) Correlation function G^K and dynamical temperature X along a cut with fixed T . We parametrize the fields into fast G_f (gray shaded area) and slow fields G_s , for short and long relative times t . (c) The monotonic function $X(t)$, defined in Eq. (10), is used to map time to a compact domain. Since $X(t)$ is decreasing, small values of u correspond to large t . (d) As the number of time scales τ_n are sent to infinity, $G^K(u)$ becomes a smooth function. (e),(f) In the limit $T \rightarrow \infty$, $G^K(u)$ is structurally identical to a Parisi function obtained in the limit $N \rightarrow 0$ from a Parisi matrix. (g) Parisi matrix for $N = 12$ in the specific case with $m_2 = 3$, $m_3 = 6$ and $m_4 = N$. We show the case of 2-step replica symmetry breaking, except for (d) and (e), which demonstrate the extension to full replica symmetry breaking corresponding to an infinite number of time scales or equivalently layers in the Parisi matrix. In this case, the asymptotic correlation functions are ultrametric, such that the equilibrium result is indeed eventually reached by the dynamics. This is in general not the case for n -step replica symmetry breaking with n finite. Although $G^K(u)$ in (c) reproduces the Parisi function $p(u)$ in (f), the absence of ultrametricity means that they do not satisfy the same algebra. This is indicated by the gray arrow between (c) and (f).

replica theory	ultrametric Keldysh theory
A_{ab}	$A^K(t)$
$A_{aa} = a(1) = a_s(1) + a_f$	$A^K(t=0) = A_s^K(0) + A_f^K(0)$
$A_{ab}B_{ab}$	$A^K(t)B^K(t)$
$(A \cdot B)_{ab}$	$(A^K \circ B^A + A^R \circ B^K)(t)$

Table 1: Translation table between replica formalism and ultrametric Keldysh theory. Here, A and B are Parisi matrices evaluated in the limit $N \rightarrow 0$. The corresponding Parisi functions are $a(u)$ and $b(u)$ with $u \in [0, 1]$. $A^{K/R}(t)$ etc., are the associated ultrametric correlation/response functions in relative time t and \circ denotes their convolution.

230 With these preparations, we can now consider the product of two Green's functions in
 231 the time domain and focus on the Keldysh component

$$C^K(t) = A^K(t)B^K(t) = A_s^K(t)B_s^K(t) + A_f^K(t)B_f^K(t). \quad (13)$$

232 In the examples below, we will show how these products enter the equation of motion
 233 for the Keldysh Green's function as a result of memory effects. We identify the equal-
 234 time expression $C^K(t=0)$ with the replica diagonal, i.e. equilibrated, part of the Parisi
 235 function $c(1)$, and the slow component $C_s^K(t)$ with the off-diagonal parts describing replica
 236 symmetry breaking $c(u)$ with $u \in [0, 1[$. The product of two Keldysh Green's functions in
 237 the time domain is therefore equivalent to the Hadamard product of two Parisi matrices
 238 (see Table 1, which summarizes our key results).

239 We next show that this correspondence also extends to convolutions. To do so, we
 240 consider the convolution between a Green's function and a self-energy $C = \Sigma \circ G =$
 241 $\int_{t'} (\Sigma_f + \Sigma_s)(t-t')(G_f + G_s)(t')$ ¹. The non-zero off-diagonal of C in Keldysh space
 242 describes the evolution of the correlation function. It reads $\Sigma^K \circ G^A + \Sigma^R \circ G^K$. This is
 243 also the form of the collision integral that accounts for memory effects in the time evolution
 244 of G^K [51].

245 We first consider the product of the slow components

¹The same structure also emerges from the product of two Green's functions as it appears for example in the action $C = G_1 \sigma^1 G_2 \sigma^1$, with σ^1 the first Pauli matrix.

$$\begin{aligned}
 \Sigma_s^K \circ G_s^A + \Sigma_s^R \circ G_s^K &= - \int_0^t dt' \Sigma_s^K(t+t')X(t')\partial_{t'}G_s^K(t') - \int_t^\infty dt' \Sigma_s^K(t+t')X(t')\partial_{t'}G_s^K(t') \\
 &\quad - \int_0^t dt' G_s^K(t-t')X(t')\partial_{t'}\Sigma_s^K(t') - \int_t^\infty dt' G_s^K(t-t')X(t')\partial_{t'}\Sigma_s^K(t') \\
 &= \int_{X(0)}^{X(t)} dX' \Sigma_s^K(X(t))G_s^K(X') - \Sigma_s^K(X(t))X(t)G_s^K(X(t)) \\
 &\quad + \int_{X(0)}^{X(t)} dX' G_s^K(X(t))\Sigma_s^K(X') - G_s^K(X(t))X(t)\Sigma_s^K(X(t)) \\
 &\quad + \Sigma_s^K(X(t))X(0)G_s^K(X(0)) + G_s^K(X(t))X(0)\Sigma_s^K(X(0)) \\
 &\quad + \Sigma_s^K(X(t))X(t)G_s^K(X(t)) + \int_{X(t)}^0 dX' \Sigma_s^K(X')G_s^K(X') \\
 &= \frac{\beta}{2} \left[- \int_u^1 dv \Sigma_s^K(u)G_s^K(v) - u\Sigma_s^K(u)G_s^K(u) + \Sigma_s^K(u)G_s^K(1) \right. \\
 &\quad \left. - \int_u^1 dv G_s^K(u)\Sigma_s^K(v) + G_s^K(u)\Sigma_s^K(1) - \int_0^u dv \Sigma_s^K(v)G_s^K(v) \right]. \tag{14}
 \end{aligned}$$

246 In the first equality, we have used the generalized thermal ansatz. It implies that the
 247 glass phase becomes stiff as $T \rightarrow \infty$: Since $G_s^K(t)$ decays on a time scale $t \sim T$, the
 248 derivative scales as $\partial_t \sim 1/T$ and compensates the divergence of the integration domain
 249 $\sim T$. This behavior is illustrated in Fig. 2. We point out that this stiffness property
 250 of the classical ansatz (10) ensures a weak long-term memory and thus convergence of
 251 the convolutions even in the aging regime. A more responsive, i.e. more slowly decaying
 252 G^R , would imply a stronger memory and divergent convolutions in Eq. (14) while for
 253 a less responsive ansatz the integrals vanish thereby precluding glassy behavior. The
 254 second equality in Eq. (14), which compactifies time, follows from ultrametricity and
 255 partial integration. It is important to point out that due to this change of variables, the
 256 information on time scales is lost. In the last step, we have introduced the dimensionless
 257 variable $u \in [0, 1]$ as

$$X(t) = \beta u/2 \tag{15}$$

258 with the boundary conditions $X(0) = \beta/2$ and $X(\infty) = 0$. The same relation holds be-
 259 tween v and X' . $X(t)$ is a decreasing function. Consequently, small values of u correspond
 260 to late times t and while the system equilibrates at short relative times, $X(\infty) = 0$ implies
 261 a maximally unresponsive infinite temperature state at large relative times.

262 As a consequence of the stiffness implied by the generalized fluctuation-dissipation
 263 relation (10) with $X(t) \leq \beta/2$ the slow retarded Green's function decays faster than the
 264 slow Keldysh component and can therefore be neglected at sufficiently late times t (see
 265 also Fig. 2). Consequently, one finds

$$\Sigma_f^K \circ G_s^A + \Sigma_s^R \circ G_f^K = 0, \tag{16}$$

266 while the other term mixing fast and slow parts is finite

$$\begin{aligned}
 \Sigma_s^K \circ G_f^A + \Sigma_f^R \circ G_s^K &= \int_t^\infty [G_f^R(t)\Sigma_s^K(u) + \Sigma_f^R(t)G_s^K(u)] \\
 &= \frac{\beta}{2} [G_f^K(1)\Sigma_s^K(u) + \Sigma_f^K(1)G_s^K(u)]. \tag{17}
 \end{aligned}$$




Figure 2: Stiffness of the glass phase. We show a logarithmic plot of typical correlation and response functions $G^K(t)$ and $G^R(t)$ at intermediate center-of-mass times T . In the aging regime $t > \tau_{\text{erg}}$ the correlation function $G^K(t)$ varies slowly, such that the generalized thermal ansatz Eq. (10) implies that $G^R(t \gtrsim \tau_{\text{erg}}) \lesssim 1/T$ vanishes as $T \rightarrow \infty$. The arrows indicate this behavior of the response function as T is increased. For times $t < \tau_{\text{erg}}$ the system is in thermal equilibrium. The boundary effects for $t \approx T$ that cause G^R to rise quickly become irrelevant as $T \rightarrow \infty$.

267 The first line follows from the condition of strong hierarchy: The slow parts are constant
 268 on the scale on which the fast functions decay. To obtain the simplified expression in
 269 the second line, we have used the high-temperature expansion of the standard fluctuation-
 270 dissipation relation Eq. (11) to linear order in β for the fast field, which makes the analogy
 271 to the replica formalism more apparent. It will therefore be used throughout this article.
 272 We emphasize, however, that it is not essential to the argument. Combining all terms, we
 273 find

$$\begin{aligned} & \Sigma^K \circ G^A + \Sigma^R \circ G^K \\ &= \frac{\beta}{2} \left[\Sigma_s^K(u) G^K(1) + G_s^K(u) \Sigma^K(1) - u \Sigma_s^K(u) G_s^K(u) \right. \\ & \quad \left. - \int_u^1 dv (\Sigma_s^K(u) G_s^K(v) + G_s^K(u) \Sigma_s^K(v)) - \int_0^u dv \Sigma_s^K(v) G_s^K(v) \right], \end{aligned} \quad (18)$$

274 which is to be compared with Eq. (4). In Eq. (18) that fast fields G_f and Σ_f enter only
 275 via $G^K(u=1)$ and $\Sigma^K(u=1)$ in the first two terms as is the case for the replica diagonal
 276 in Eq. (4).

277 We are left with the task of evaluating the time diagonal in the Keldysh formulation.

278 Sending $t \rightarrow 0$ and using the same arguments as above, we find

$$\begin{aligned}
 \Sigma_f^K \circ G_s^A + G_f^K \circ \Sigma_s^R \Big|_{t=0} &= 0, \\
 \Sigma_f^K \circ G_f^A + G_f^K \circ \Sigma_f^R \Big|_{t=0} &= \frac{\beta}{2} \Sigma_f^K(1) G_f^K(1), \\
 \Sigma_s^K \circ G_s^A + G_s^K \circ \Sigma_s^R \Big|_{t=0} &= \frac{\beta}{2} \left[\Sigma_s^K(1) G_s^K(1) - \int_0^1 dv \Sigma_s^K(v) G_s^K(v) \right], \\
 \Sigma_s^K \circ G_f^A + G_s^K \circ \Sigma_f^R \Big|_{t=0} &= \frac{\beta}{2} \left[G_f^K(1) \Sigma_s^K(1) + \Sigma_f^K(1) G_s^K(1) \right].
 \end{aligned} \tag{19}$$

279 Putting everything together, this gives for the time-diagonal

$$\Sigma^K \circ G^A + \Sigma^R \circ G^K \Big|_{t=0} = \frac{\beta}{2} \left[\Sigma^K(1) G^K(1) - \int_0^1 dv \Sigma_s^K(v) G_s^K(v) \right]. \tag{20}$$

280 Comparison with Eq. (5) shows that the matrix multiplication in Keldysh formalism at
 281 asymptotically late times using ultrametricity and a generalized thermal ansatz in the
 282 classical limit is identical to the matrix multiplication in the replica formalism.

283 As has previously been reported by Cugliandolo and Kurchan [5], this approach also
 284 gives an interpretation of the replica average in equilibrium theory. For example, the
 285 averaged correlation function

$$Q_\infty = \int_0^1 du q(u), \tag{21}$$

286 with $q(u)$ the Parisi function of the replica matrix $Q_{ab} = \langle s_i^a s_i^b \rangle$, is related to the integrated
 287 response function

$$Q_\infty = 1 + \int_0^\infty dt X(t) \partial_t Q^K(t) = 1 - \int_0^\infty dt Q^R(t). \tag{22}$$

288 In summary, we have shown that, under the assumption of ultrametricity, the Keldysh
 289 component of convolution integrals in time $\Sigma \circ G$ reproduces the algebra of replica matrices
 290 in the limit $N \rightarrow 0$.

291 3 Application: The Quantum Sherrington-Kirkpatrick model

292 We now turn our attention to the most general case of replica symmetry breaking, known as
 293 full RSB and realized by the Sherrington-Kirkpatrick model. We begin with the derivation
 294 of the Landau action valid near the critical point. The procedure can be understood as
 295 the out-of-equilibrium version of the Landau theory presented in Ref. [46]. Our approach
 296 is similar in spirit to that of Sompolinsky and Zippelius [7, 30, 37] that culminated in the
 297 analytical solution of the late-time relaxation obtained by Cugliandolo and Kurchan [6].
 298 Despite several attempts at recovering the results of replica theory from the dynamical
 299 equations at late times [30–33], discrepancies remain [48]. The problem arises from the
 300 order of limits. If one considers a finite system at infinitely late times and eventually
 301 sends the system size to infinity [30], the system is time-translation invariant, but must
 302 also obey the fluctuation-dissipation relation [52], as at infinite times, any finite system
 303 is fully equilibrated [13]. Consequently, a violation of the fluctuation-dissipation relation
 304 in this limit contradicts the underlying assumptions. In the opposite limit considering an

305 infinite system before sending the time to infinity [6] time-translation invariance is always
 306 broken because the equilibration time is determined by the system size and therefore the
 307 largest time scale in the problem. Our approach avoids these problems by measuring time
 308 in terms of the inverse temperature of the generalized fluctuation-dissipation relation.
 309 Therefore, by construction, we have access to all relevant time scales even though the
 310 system size is taken as infinite from the outset.

311 Recent experimental development have resulted in renewed interest in spin glasses.
 312 In particular, the precise positioning of large numbers of Rydberg atoms with tweezers
 313 provides an avenue towards the realization of spin glasses with long-ranged interactions
 314 [53–58]. The idea is as follows, lasers are used to drive the Rabi transition between
 315 ground-state atoms and a highly excited long-lived Rydberg state. As no other states get
 316 occupied, it is sufficient to describe the atoms as two level systems that interact via van-
 317 der Waals interactions only when in the large and therefore highly polarizable Rydberg
 318 state. By positioning the atoms at random but fixed sites using tweezers, the strength of
 319 the interactions are randomized [54]. Finally, the occupation of the Rydberg states can be
 320 controlled by adjusting the detuning δ of the driving laser, which leads to a longitudinal
 321 field $h = \delta$ in the effective spin model [59]

$$H = \sum_{ij} J_{ij} \sigma_i^3 \sigma_j^3 - \sum_i \sigma_i^1 - h \sum_i \sigma_i^3. \quad (23)$$

322 Here the Rabi coupling has been set to one and J_{ij} denotes the van-der Waals interaction
 323 between atoms i and j .

324 We point out that other experimental schemes such as Rydberg dressing which uses
 325 lasers far detuned from the Rabi transition to increase the lifetime at the expense of
 326 weaker interactions or microwave coupling between different Rydberg states leading to
 327 longer-ranged interactions $\sim R^{-3}$ result in the same Hamiltonian [59, 60]. Furthermore,
 328 random long-range interactions can be achieved with quantum simulators based on su-
 329 perconducting qubits [61] or by trapping atoms in a confocal cavity [62, 63]. Although,
 330 in the latter case the driven-dissipative cavity prevents the system from reaching thermal
 331 equilibrium, significant similarities with the classical Sherrington-Kirkpatrick model have
 332 been found in theory [64, 65] and experiment [66].

333 3.1 Effective action

334 An important distinction between the new platforms and classical glasses is the finite
 335 lifetime of the excited states due to spontaneous emission. It is therefore important to
 336 develop a minimal dynamical description applicable to late but finite times. We will
 337 achieve this by deriving the effective Ginzburg-Landau action of the random Ising model
 338 in a longitudinal and a transversal field as defined in (23). The quenched disordered
 339 coupling strengths J_{ij} are drawn from a Gaussian distribution independent of the site
 340 indices i and j . Hence, this model first introduced in Ref. [67], is effectively infinite-
 341 dimensional and described by mean-field theory. Its equilibrium Landau action has been
 342 studied in Ref. [68], with aging dynamics analyzed in Ref. [45] and previously based on
 343 the approach of Sompolinsky and Zippelius [7, 37] in Ref. [6]. Near the phase transition,
 344 we can average the spins over a small domain, such that the discrete spins are replaced by
 345 a real bosonic variable S_i . Integrating out the disordered coupling strengths J_{ij} , the site

346 index can be dropped, and the effective action is given by [45]

$$\begin{aligned}
 s[S] &= s_0[S] + s_u[S] + s_\kappa[S] + s_h[S], \\
 s_0[S] &= -\frac{1}{2} \int_t \sum_\sigma \sigma S_\sigma(t) [\partial_t^2 + m^2] S_\sigma(t), \\
 s_h[S] &= \int_t \sum_\sigma \sigma h_\sigma(t) S_\sigma(t), \\
 s_u[S] &= -\frac{u}{2} \int_t \sum_\sigma \sigma S_\sigma^4(t), \\
 s_\kappa[S] &= i \frac{\kappa}{4} \int_{t_1, t_2} S_\sigma(t_1) \sigma_{\sigma\rho}^3 S_\rho(t_1) S_{\sigma'}(t_2) \sigma_{\sigma'\rho'}^3 S_{\rho'}(t_2).
 \end{aligned} \tag{24}$$

347 Here, the quartic term s_u provides a soft constraint for the spin length and the transverse
 348 field gives rise to the inertial dynamic term in s_0 . The disorder is encoded in the term
 349 s_κ , with $\kappa = J_{ij}^2$ the variance of the Gaussian distribution $\mathcal{P}(J_{ij})$. The dynamical theory
 350 requires a doubling of the time contour. Following the standard procedure of the Keldysh
 351 path-integral (for an introduction see [51, 69]), we therefore introduce Greek indices that
 352 take the values $\{+, -\}$ to denote the branch of the contour.

353 Due to the infinite range of the random couplings J_{ij} , the site index in (24) is irrelevant
 354 and will be suppressed in the following.

355 It is convenient to introduce the spin bilinear $q_{\alpha\beta} \equiv q_{\rho\rho'}(t_1, t_2) = S_\sigma(t_1) \sigma_{\sigma\rho}^3 S_{\rho'}(t_2)$.
 356 Here and in the following α and β denote multi-indices incorporating the Keldysh index
 357 and time. We now decouple the disorder-induced non-linearity by a Hubbard-Stratonovich
 358 transformation with

$$s_{\text{aux}}[S, Q] = \frac{i}{4\kappa} \text{Tr} [(RQ\sigma^1 R - i\kappa q)^2], \tag{25}$$

359 where $R = (\sigma^1 + \sigma^3)/\sqrt{2}$. We have furthermore introduced the trace Tr over the multi
 360 index, i.e. $\text{Tr}[A^2] = \int_{t, t'} A_{\sigma\rho}(t, t') A_{\rho\sigma}(t', t)$. Rewriting the soft constraint s_u in terms of
 361 a functional derivative with respect to a source field K , we can perform the remaining
 362 Gaussian integral over S_σ . Rotating the result to the $R/A/K$ basis, we define classical
 363 and quantum fields as $h_c = \sum_\sigma h_\sigma/\sqrt{2}$ and $h_q = \sum_\sigma \sigma h_\sigma/\sqrt{2}$. Although external fields
 364 are classical, we keep the notation symmetric and express the Keldysh partition function
 365 as

$$\begin{aligned}
 Z &= \int \mathcal{D}Q e^{i s_u[\frac{\delta}{\delta K}]} e^{-\frac{i}{2} (K^\top + h^\top) \sigma^1 [G_0^{-1} + Q]^{-1} \sigma^1 (K + h)} \\
 &\times e^{-\frac{1}{4\kappa} \text{Tr}[Q\sigma^1 Q\sigma^1] - \frac{1}{2} \text{Tr} \log(1 + G_0 Q) + \text{const.}} \Big|_{K=0}.
 \end{aligned} \tag{26}$$

366 Here, \top denotes the transpose in Keldysh space. Finally, the bare spin propagator can be
 367 expanded in powers of m^{-2}

$$\begin{aligned}
 G_0(t_1, t_2) &= -\delta(t_1 - t_2) \sigma^1 (\partial_{t_2}^2 + m^2)^{-1} \\
 &\approx \delta(t_1 - t_2) \sigma^1 \left(-\frac{1}{m^2} + \frac{1}{m^4} \partial_{t_2}^2 \right).
 \end{aligned} \tag{27}$$

368 Due to the saddle point condition

$$0 = -2i\kappa \frac{\delta s_{\text{aux}}}{\delta Q_{\alpha\beta}} = (\sigma^1 Q \sigma^1 - i\kappa \sigma^1 R q R)_{\beta\alpha} \tag{28}$$

369 the average of Q can be given a physical interpretation in terms of the full spin propagator
 370 $G = (G_0^{-1} + Q)^{-1}$

$$\frac{1}{\kappa} \langle Q_{\alpha\beta} \rangle = i \langle RqR\sigma^1 \rangle_{\alpha\beta} = -(\sigma^1 G \sigma^1)_{\alpha\beta}. \quad (29)$$

371 It is however not a valid order parameter as it does not vanish at the critical point.
 372 We arrange for this property by a shift operation, which can also be viewed as a UV
 373 renormalization operating on short time distances. Thereby, we fix the value of the critical
 374 coupling strength via the requirement that the renormalized order parameter vanishes
 375 at the transition, which will be verified explicitly below. To this end, we decompose
 376 $Q(t, t') = ((m^2 - \tilde{m}^2)\sigma^1 + Q_{\text{EA}}) \delta(t - t') + \tilde{Q}(t, t') \equiv Q_0(t, t') + \tilde{Q}(t, t')$ into a UV shift
 377 $Q_0(t, t')$ with $Q_{\text{EA}} = iq_{\text{EA}}(\mathbb{1} - \sigma^3)/2$ and a small field \tilde{Q} . We also define $\tilde{G}_0^{-1} = G_0^{-1} + Q_0$,
 378 use again the exact relation Eq. (29), and expand in powers of \tilde{Q}

$$\begin{aligned} & \frac{1}{\kappa} \left[\sigma^1 (Q_0 + \tilde{Q}) \sigma^1 \right]_{\alpha\beta} - i \left[\tilde{G}_0 \sigma^1 (hh^\top) \sigma^1 \tilde{G}_0 \right]_{\alpha\beta} \\ &= - \left[(\tilde{G}_0^{-1} + \tilde{Q})^{-1} \right]_{\alpha\beta} \\ &\approx \left[-\tilde{G}_0 + \tilde{G}_0 \tilde{Q} \tilde{G}_0 - \tilde{G}_0 \tilde{Q} \tilde{G}_0 \tilde{Q} \tilde{G}_0 + \dots \right]_{\alpha\beta}. \end{aligned} \quad (30)$$

379 We approximate $\tilde{G}_0 \approx -\tilde{m}^{-2} \sigma^1 \delta(t - t')$ everywhere except for the zero-order term in \tilde{Q} ,
 380 where we also expand in the time derivative to leading order. Furthermore, we make use of
 381 the fact that the magnetic field is classical and time-independent. With this, the term due
 382 to the magnetic field simplifies to $i[\tilde{G}_0 \sigma^1 (hh^\top) \sigma^1 \tilde{G}_0]_{\alpha\beta} = ih^2 \tilde{G}_0^R \tilde{G}_0^A \delta_{\alpha 1} \delta_{\beta 1} \approx \frac{ih^2}{\tilde{m}^4} \delta_{\alpha 1} \delta_{\beta 1}$.
 383 Most importantly, we notice, that both the magnetic field and the order parameter q_{EA}
 384 only affect the Keldysh component of the matrix equation (30). This is an exact statement,
 385 that follows from the causal structure of the Green's function. It implies that the magnetic
 386 field can be absorbed into q_{EA} .

387 Explicitly, upon Fourier transformation, the Keldysh and retarded components of

$$\tilde{Q} = \begin{pmatrix} \tilde{Q}^V & \tilde{Q}^A \\ \tilde{Q}^R & \tilde{Q}^K \end{pmatrix} \quad (31)$$

388 satisfy the equations

$$\begin{aligned} (m^2 - \tilde{m}^2) + \tilde{Q}^R &= -\frac{\kappa}{\omega^2 - \tilde{m}^2 + \tilde{Q}^R} \\ &\approx \frac{\kappa}{\tilde{m}^2} + \frac{\kappa}{\tilde{m}^4} (\omega^2 + \tilde{Q}^R) + \frac{\kappa}{\tilde{m}^6} \left[\tilde{Q}^R \right]^2 \\ 2\pi i q_{\text{EA}} \delta(\omega) + \tilde{Q}^K &= \kappa \frac{2\pi i (q_{\text{EA}} + h^2) \delta(\omega) + \tilde{Q}^K}{|\omega^2 - \tilde{m}^2 + \tilde{Q}^R|^2}. \end{aligned} \quad (32)$$

389 Causality of the spin response function requires $Q(t, t') \sim \theta(t - t')$. Furthermore, we have
 390 used that $\tilde{Q}^A(\omega)$ is the complex conjugate of $\tilde{Q}^R(\omega)$ and that \tilde{Q}^V vanishes due to the
 391 normalization of the partition function $Z = 1$. Clearly, in the first equation, the linear
 392 term in \tilde{Q}^R disappears for $\tilde{m}^4 = \kappa$, independent of h (because $u = 0$ here). We conclude

$$\tilde{Q}^R(\omega) = -\sqrt{-\sqrt{\kappa} [(\omega + i0^+)^2 - r]}, \quad (33)$$

393 which is causal in the paramagnetic phase and gives rise to a phase transition when
 394 $r = m^2 - 2\sqrt{\kappa}$ vanishes at $\kappa = m^4/4$.

395 The second equation evaluated at $\omega = 0$ fixes the order parameter (we demand that
 396 \tilde{Q}^K is a continuous function). Multiplying both sides with $G^R G^A$, inserting the retarded
 397 and advanced $\tilde{Q}^{R/A}$ and keeping only the leading term in r one finds

$$q_{\text{EA}} = \frac{h^2 \kappa^{1/4}}{2\sqrt{r}}. \quad (34)$$

398 As expected, in the paramagnetic phase, the magnetization is linear in the longitudinal
 399 field $m \sim S \sim h$ and the Hubbard-Stratonovich field $Q \sim S^2$ is proportional to h^2 .
 400 Furthermore, at the critical point, the system is gapless and has a divergent response to
 401 the external field, signified by $q_{\text{EA}} \sim r^{-1/2}$.

402 Below we obtain the Landau action by expanding in the small field \tilde{Q} near the critical
 403 point. The above ensures that there will be no contribution $\sim Q^2 = \int_{t'} Q(t_1, t') Q(t', t_2)$
 404 to the Landau action.

405 In the following, we will exclusively work with \tilde{Q} , \tilde{G}_0 , and \tilde{m} and therefore drop the
 406 tilde from here on.

407 3.2 Paramagnetic phase

408 Having established the proper order parameter field, we can now expand the action in the
 409 soft constraint s_u . For the discussion of the paramagnetic phase, an expansion to first
 410 order in u is enough to obtain stable results known from equilibrium theory. On the other
 411 hand, an expansion to second order in u is necessary to recover the spin glass phase [68].

412 Following the discussion above, we expand in small fields Q to find the unconstrained
 413 action

$$\begin{aligned}
 i s_0[Q] = & -\frac{1}{2\kappa} \int_{t,t'} (\partial_t^2 + r) \text{tr}(\sigma^1 Q(t, t')|_{t=t'}) \\
 & + \frac{i}{2\kappa} \int_{t,t'} h^\top(t) Q(t, t') h(t') - \frac{1}{6\kappa^{3/2}} \text{Tr}[(\sigma^1 Q)^3].
 \end{aligned} \quad (35)$$

414 To first order in u , the constraint on the spin length contributes a Hartree term

$$\begin{aligned}
 i \Delta s_u[Q] = & \frac{3iu}{2} \int_t (G^K + G^V)(t, t) (G^R + G^A)(t, t) \\
 \approx & -\frac{3iu}{2\kappa^2} \int_t (Q^K + Q^V)(t, t) (Q^R + Q^A)(t, t).
 \end{aligned} \quad (36)$$

415 to the Landau action, which then reads

$$s[Q] = s_0[Q] + \Delta s_u[Q]. \quad (37)$$

416 To highlight the temporal structure of this action, it is useful to consider its diagrammatic
 417 representation shown in Fig. 3. We observe that the disorder gives rise to a term $\sim Q^3$ that
 418 relates the order-parameter fields at different times. As we will see below, it corresponds to
 419 a memory that for sufficiently large κ causes the order parameter to become stiff, thereby
 420 excluding its relaxation at large relative times that is characteristic of the paramagnetic
 421 phase.

422 To find the critical disorder strength where the paramagnet freezes, we consider the
 423 equations of motion for Q obtained from the saddle point condition

$$\begin{aligned}
 0 \stackrel{!}{=} \frac{\delta i s[Q]}{\delta Q(t_1, t_2)} \approx & -\frac{1}{2\kappa} \sigma^1 \delta(t_1 - t_2) (r + \partial_{t_2}) + \frac{1}{2\kappa^{3/2}} \int_t \sigma^1 Q(t_1, t) \sigma^1 Q(t, t_2) \sigma^1 \\
 & + \frac{i}{2\kappa} h(t_1) h^\top(t_2) + \frac{\delta i \Delta s_u[Q]}{\delta Q(t_1, t_2)}
 \end{aligned} \quad (38)$$

424 with

$$\frac{\delta i \Delta s_u[Q]}{\delta Q_{\bar{\sigma}\rho}(t_1, t_2)} = -\frac{3iu}{2\kappa} Q^K(t_1, t_1) \delta(t_1 - t_2) \delta_{\bar{\sigma}\rho}, \quad (39)$$

425 where $\bar{\sigma}$ denotes the opposite of σ .




Figure 3: Diagrammatic representation of the Landau action (37) at linear order in the soft-spin constraint u . For simplicity, we have suppressed the Keldysh structure. The inverse bare spin propagator for $Q = 0$ reading $1/(2\kappa)\sigma^1\delta(t_1 - t_2)(r + \partial_{t_2}^2)$ is depicted as open rectangle. Q is shown as a straight line, h is represented by a cross, the vertex $\frac{iu}{2\kappa^2}$ as a dot, and G_0 as open circle.

426 Following the general procedure outline in Sec. 2.2, we split the order parameter field
 427 into fast and slow components $Q = Q_f + Q_s$, where the evolution of Q_s slows down
 428 indefinitely for $T \rightarrow \infty$. Looking for a paramagnetic solution, we require $Q_s = 0$ and
 429 make a time-translation invariant ansatz $Q(t, t') = Q(t - t')$. The equations of motion
 430 (38) therefore become diagonal in frequency space. Due to the absence of scattering
 431 at the current level of approximation, there is only one non-trivial equation of motion.
 432 Expanding for $h = 0$ in small frequencies ω , one finds

$$\omega^2 + \frac{1}{\kappa^{1/2}} [Q^R]^2(\omega) = r - \frac{3iu}{\kappa} \int_{\nu} Q^K(\nu), \quad (40)$$

433 which has the thermal paramagnetic solution

$$\begin{aligned} Q^R(\omega) &= -\kappa^{1/4} \sqrt{\Delta^2 - (\omega + i0^+)^2}, \\ Q^K(\omega) &= 2\kappa^{1/4} \coth \frac{\beta|\omega|}{2} \sqrt{\Delta^2 - \omega^2} \theta(|\omega| - |\Delta|) \end{aligned} \quad (41)$$

434 with the shifted mass $\Delta^2 = r - \frac{3iu}{\kappa} \int_{\nu} Q^K(\nu)$.

435 This reproduces the form of results from the analytically continued replica theory
 436 for $h = 0$ (up to relabelling of coefficients) developed in [68] and in [45] in the Keldysh
 437 framework. In particular, for small u we reach a critical point $\Delta(r_c0) = 0$, with

$$r_{c0} = -\frac{6u}{\kappa^{3/4}} \int_{\omega} \omega \coth \frac{\beta\omega}{2} \xrightarrow{\beta \rightarrow \infty} \frac{6u}{\kappa^{3/4}} \int_{\omega} |\omega| \quad (42)$$

438 as well as $Q(\omega = 0) = 0$, which verifies the assumption that the shifted Q is an order
 439 parameter for the Landau theory. After crossing the phase transition, we expect that Δ
 440 remains pinned to zero and that this can be achieved by introducing an Edwards-Anderson
 441 order parameter into the occupation function component, $Q_{\text{EA}}^K(\omega) = Q^K(\omega) + 2\pi i q^{\text{EA}} \delta(\omega)$,
 442 $q^{\text{EA}} > 0$. Indeed, inserting this ansatz into the equation of motion (40) we reproduce the
 443 known results [45, 46]

$$\begin{aligned} Q^R(\omega) &= i\kappa^{1/4} \omega, \\ q^{\text{EA}} &= i \int_{\nu} Q^K(\nu) \Big|_{\Delta=0} - \frac{\kappa}{3u} r = \frac{\kappa}{3u} (r_{c0} - r), \\ Q^K(\omega) &= 2i\kappa^{1/4} \omega \coth \frac{\beta\omega}{2}. \end{aligned} \quad (43)$$

444 In particular, there is a gapless, damped mode.

445 3.3 Landau action to order u^2

446 The discussion of the spin glass phase requires a more careful discussion of the memory
 447 terms. In particular, beyond the critical point the disorder term $\sim Q^3$ renders the Landau
 448 action in Eq. (37) unstable. It is therefore necessary to continue the perturbative expansion
 449 in the soft-spin constraint u to second order. As is shown in Fig. 4, there is only one term
 450 in the effective action that is of order u^2 and two-particle irreducible. It involves time-non-
 451 local fields and thus gives rise to memory effects that are essential for the stability of a spin
 452 glass. All other diagrams $\sim u^2$ are either disconnected or not two-particle irreducible and
 453 thus constitute at most a quantitative correction to the Hartree shift already discussed in
 454 the previous section. We therefore exclusively focus on the memory term at this order

$$\begin{aligned}
 i\Delta s_{u^2}[Q] &= -\frac{3u^2}{4\kappa^4} \int_{t,t'} [(\text{tr}(Q(t,t')Q(t',t))\text{tr}(Q(t,t')\sigma^1 Q(t',t)\sigma^1) \\
 &\quad + \text{tr}(Q(t,t')Q(t',t)\sigma^1)\text{tr}(Q(t,t')\sigma^1 Q(t',t))]. \tag{44}
 \end{aligned}$$

455 Terms of this form are known as the primary cause of relaxation and thermalization in
 456 quench dynamics, see for example [70]. For the stability of the spin-glass it is therefore
 457 important to investigate the competition between the terms $\sim u^2$ that favor ergodicity
 458 and the disorder term $\sim Q^3$ that favors freezing.

$$\begin{aligned}
 i\Delta s_{u^2}[Q] &= \frac{1}{2} \left(90 \times \text{Diagram 1} \right. \\
 &\quad \left. + 3 \times \text{Diagram 2} + 12 \times \text{Diagram 3} \right)
 \end{aligned}$$

Figure 4: Diagrammatic representation of the second order contribution of the soft-spin constraint u to the effective action. The first diagram is not one-particle irreducible and corresponds to quantitative corrections to the tadpole diagram in Fig. 3. The second diagram is disconnected and therefore cancels against the normalization of the partition function. Consequently, we only retain the last contribution, which involves time-non-local fields and thus introduces a memory to the equation of motion that competes with the disorder term in the spin glass phase.

459 Expanding the trace-log as before, we find the Landau action of the Sherrington-
 460 Kirkpatrick model with longitudinal and transversal fields

$$s[Q] = s_0[Q] + \Delta s_u[Q] + \Delta s_{u^2}[Q]. \tag{45}$$

461 This action is the dynamical equivalent of the result recently reported in Ref. [46].

462 3.4 Asymptotic solution in the glass phase

463 In the previous section, we have found the Keldysh action corresponding to the equilibrium
 464 Landau action in replica theory. We will now consider the limit of late times and apply
 465 the general results of Sec. 2 to show how full replica symmetry breaking is recovered in
 466 the time domain.

467 In the limit of late times $T = (t_1 + t_2)/2 \rightarrow \infty$ the forward evolution scale drops out.
 468 This does exclude the spontaneous breaking of time translation invariance globally. Time
 469 translation invariance can, however, be broken in a scale-dependent way, as suggested by
 470 the reparametrization invariance of the aging action [6].

471 We thus bring the action into a form in which time translation invariance is used:
 472 $Q(t_1, t_2) = Q(t = t_1 - t_2)$. To this extent, one performs a Wigner expansion of the action
 473 (45) and drops all derivatives ∂_T . In all terms of the action the length of the time domain,
 474 T , factors out

$$\begin{aligned}
 is[Q]/T = & -\frac{1}{2\kappa} (\partial_t^2 + r) \text{tr}[\sigma^1 Q(t)] \Big|_{t=0} + \frac{ih_c^2}{2\kappa} \int_t Q^V(t) \\
 & + \frac{1}{6\kappa^{3/2}} \int_{t,t'} \text{tr} \left[Q(t) \sigma^1 Q(t') Q^\top(t+t') \right] + \frac{3iu}{2\kappa^2} [\text{tr}(Q(t=0)) \text{tr}(\sigma^1 Q(t=0))] \\
 & - \frac{3u^2}{4\kappa^4} \int_t \left[\text{tr}(Q(t) Q^\top(t)) \text{tr}(Q(t) \sigma^1 Q^\top(t) \sigma^1) + \text{tr}(Q(t) Q^\top(t) \sigma^1) \text{tr}(Q(t) \sigma^1 Q^\top(t)) \right].
 \end{aligned} \tag{46}$$

475 Following the procedure of Sec. 2, we split the field Q in a slow and a fast component
 476 and similarly divide the action into a 'spin glass' part s_{sg} that involves the slow field and
 477 a quantum part s_q that describes the equilibration at short relative times. Since $Q_f(t)$
 478 approaches zero for large arguments, we require $q_{EA} = -iQ_s^K(t=0)$. In analogy to the
 479 paramagnetic phase, in s_q the terms $\sim u^2$ are not important at small frequencies, so we
 480 will neglect these by writing $s_q = s_{q,0} + \mathcal{O}(u^2)$. Since in the following, we will mostly
 481 concern ourselves with the slow field, we will drop its index from now on and simply refer
 482 to it as Q (i.e. $Q \equiv Q_s$). One then has

$$\begin{aligned}
 s[Q] & \approx s_{sg}[Q] + s_{q,0}[Q], \\
 is_{sg}[Q]/T = & -\int_t \text{tr}[R_1 Q(t) Q^\top(t) \sigma^1] + \frac{ih_c^2}{2\kappa} \int_t Q^V(t) + \frac{R_2}{3} \int_{t,t'} \text{tr}[Q(t) \sigma^1 Q(t') Q^\top(t+t')] \\
 & - \frac{R_3}{3} \int_t \left\{ \text{tr}[Q(t) Q^\top(t)] \text{tr}[\sigma^1 Q(t) \sigma^1 Q^\top(t)] + \text{tr}[Q(t) Q^\top(t) \sigma^1] \text{tr}[Q(t) \sigma^1 Q^\top(t)] \right\}, \\
 is_{q,0}[Q]/T = & \frac{1}{2\kappa} \int_\omega (\omega^2 - r) \text{tr}[\sigma^1 Q_f(\omega)] - \frac{ih_c^2}{2\kappa} \text{tr}[Q_f(\omega=0)] \\
 & + \frac{R_2}{3} \int_\omega \text{tr}[\sigma^1 Q_f(\omega) \sigma^1 Q_f(\omega) \sigma^1 Q_f(\omega)] + iR_2 q_{EA} Q_{f11}(\omega=0) \text{tr}[\sigma^1 Q_f(\omega=0)] \\
 & + \frac{3iu}{2\kappa^2} \int_{\omega,\omega'} \text{tr}[Q_f(\omega) + 2\pi\delta(\omega) Q_{EA}] \text{tr}[\sigma^1 (Q_f(\omega') + 2\pi\delta(\omega') Q_{EA})],
 \end{aligned} \tag{47}$$

483 with

$$R_1 = -\frac{1}{2\kappa^{3/2}} \sigma^1 Q_f(\omega=0) \sigma^1 \equiv \begin{pmatrix} R_1^K & R_1^R \\ R_1^A & R_1^V \end{pmatrix}, \quad R_2 = \frac{1}{2\kappa^{3/2}}, \quad R_3 = \frac{9u^2}{4\kappa^4}, \tag{48}$$

484 where $R_1^A = R_1^R$. The saddlepoint of s_{sg} and $s_{q,0}$ with respect to Q and Q_f respectively
 485 gives the coupled dynamical equations of the aging and ergodic components. Since we

486 are looking for the qualitative form of the slow component Q , it is enough to consider
 487 $\delta s_{sg}[Q]/\delta Q$, which gives

$$\begin{aligned}
 0 &= 2R_1^R Q^R(t) - R_2 \int_{t'} Q^R(t') Q^R(t-t') + \frac{2R_3}{3} [Q^{R^2}(t) + 3Q^{K^2}(t)] Q^R(t), \\
 0 &= R_1^K [Q^R(t) + Q^A(t)] + 2R_1^R Q^K(t) - i \frac{h^2}{2\kappa} \\
 &\quad - R_2 \left(\int_0^{T+t/2} dt' Q^K(t-t') Q^R(t') + \int_{t/2-T}^0 dt' Q^K(t-t') Q^A(t') \right) \\
 &\quad + \frac{2R_3}{3} [Q^{K^2}(t) + 3(Q^{R^2}(t) + Q^{A^2}(t))] Q^K(t),
 \end{aligned} \tag{49}$$

488 where we have kept the integration boundaries explicit, even though we have not done
 489 so before. The reason is, that, although we expect that the integration boundaries are
 490 irrelevant when we send $T \rightarrow \infty$, we want to show so explicitly in the following.

491 As is the case for the replica formulation, Eq. (49) has a ferromagnetic solution. In
 492 equilibrium, it can be shown that this solution is thermodynamically unstable [71], with
 493 the exact solution instead given by the Parisi function with full replica symmetry breaking
 494 [72, 73]. When considering quench dynamics on the other hand, one has to fix a boundary
 495 condition for Q at large values of $|t|$. The difference between a system that exhibits aging
 496 and more conventional spontaneous symmetry breaking has to be encoded in the time
 497 scale on which the order parameter q_{EA} recovers from the perturbation at the boundary.
 498 Here, we will only discuss the equivalent of the spin glass solution, not the (im)possibility
 499 of a ferromagnetic phase.

500 Following the discussion of Kurchan [74], we expect $Q(t)$ to vary increasingly slowly
 501 as t grows. In fact, each value of $Q^K(t)$ corresponds to a time scale on which the system
 502 thermalizes to an effective inverse temperature $X(t)$. This time scale is much longer than
 503 those of all previous (larger) values of $Q^K(t' < t)$. We can exploit this to simplify e.g.
 504 integrals of the form $\int_0^t dt' Q(t') Q(t-t')$. Specifically, for all values of t' on the scale of t
 505 one has $Q^K(t') = Q^K(t)$, while for $t-t' \sim t$ one has $Q^K(t-t') = Q^K(t)$. In other terms,
 506 the correlation Q^K is ultrametric. At the same time, the generalized thermal response
 507 function $Q^R(t) = iX(t)\partial_t Q^K(t)$ vanishes much more quickly than $Q^K(t)$. This justifies
 508 the classical approximation of the general scheme in Sec. 2. In Eqs. (49) we therefore
 509 only keep the memory terms $\sim R_3$ with the highest power in Q^K and drop the term
 510 proportional to R_1^K .

511 Following these preparations, the second equation in (49) only involves time-local or
 512 Hadamard products of Keldysh Green's functions as well as the Keldysh component of
 513 causal convolutions. Both have been discussed in Sec. 2. Applying the partial integration
 514 Eq. (14), we find

$$\begin{aligned}
 0 &= -2R_1^R q(u) + \frac{h^2}{2\kappa} + \frac{2R_3}{3} q(u)^3 \\
 &\quad - R_2 \beta \left(2q(u) \int_u^1 dv q(v) + \int_0^u dv q(v)^2 + uq(u)^2 - 2q_{EA} q(u) \right).
 \end{aligned} \tag{50}$$

515 Since $X(t=0) = \beta/2$ is fixed by the temperature of the equilibrated part, we have
 516 parametrized $X(t) = \beta u/2$ with $u \in [0, 1]$ and $Q^K(X(t)) = -iq(u)$. We thereby exactly
 517 recover the replica result [46]². Consequently, the Keldysh structure for $T \rightarrow \infty$ derives

²The only difference between their result and ours is the addition of $2\beta q_{EA} q(u)$ in Eq. (50). This is a consequence of the replica diagonal of the Parisi matrix being removed. It exactly compensates for the difference in the definition of R_1

518 the rules of the replica limit. Physically, we can say that replica symmetry breaking
 519 corresponds to the inability of the system to fully thermalize to a single global inverse
 520 temperature β even at arbitrarily late times/the steady state. The assumption of the
 521 replica off-diagonal being independent of Matsubara frequency (hence including only the
 522 zeroth Matsubara frequency) is equivalent to the classical limit involving only the time-
 523 local generalized FDR (10).

524 Exploiting the analogy to the known solution from replica theory, it is easy to show
 525 that

$$q(u) = \begin{cases} q_h = \frac{1}{2}(\frac{3}{\kappa R_3} h^2)^{1/3} & u \leq \frac{q_h}{q_{EA}} x \\ q_{EA} \frac{u}{x} & \frac{q_h}{q_{EA}} x < u < x \\ q_{EA} & x \leq u \end{cases} \quad (51)$$

526 with $q_{EA}^2 = R_1^R/R_3$ and $x = \frac{2R_3 q_{EA}}{R_2 \beta} > 0$. Consequently,

$$X(q) = \begin{cases} 0 & q < q_h \\ \frac{R_3}{R_2} q & q_h < q < q_{EA} \\ \frac{\beta}{2} & q = q_{EA} \end{cases}, \quad (52)$$

527 which is consistent with the solution of Ref. [6]. We point out that $x > 0$ requires
 528 $Q^R(\omega = 0) > 0$, which, as we saw in Sec. 3.2, requires the disorder strength to exceed the
 529 critical value $\kappa > \kappa_c = m^4/4$.

530 What is left is to show that this is also a solution to the classical limit of the first
 531 equation in (49). This can be seen by integrating that equation with respect to t and
 532 exploiting that with $Q^K(t)$ also $X(t)$ is an ultrametric function. One then finds

$$0 = 2R_1 \int_{q_{EA}}^q dq' X(q') + R_2 \left(\int_{q_{EA}}^q dq' X(q') \right)^2 - 2R_3 \int_{q_{EA}}^q dq' q'^2 X(q'), \quad (53)$$

533 which is indeed solved by (52).

534 4 Application: The spherical p -spin model

535 Our second application is the spherical p -spin model. We begin with a brief derivation
 536 of its effective action in the Keldysh formalism. The procedure is analogous to that
 537 of Ref. [46] with minor modifications owed to the doubling of the time contour in the
 538 Keldysh approach. We then apply the generalized thermal ansatz to the ultrametric aging
 539 component of the spin correlations. This procedure is then shown to reproduce the results
 540 known from replica formalism.

541 4.1 Effective action

542 The spherical p -spin model was first introduced in Ref [75] with the Ising counterpart
 543 discussed in Ref [76]. It is known to exhibit 1-step replica symmetry breaking in thermal
 544 equilibrium [77]. Additionally, the transition between the paramagnetic and glass phase
 545 changes from second to first order at low temperatures. There, the dynamical equations of
 546 motion predict a higher critical field strength than the equilibrium theory [35]. It therefore
 547 poses a critical test to the general arguments of Sec. 2.

548 The following discussion follows closely that of Ref. [46]. In fact, it can be seen as a
 549 translation of their discussion to the Keldysh formalism.

550 The spherical p -spin model is given by the Hamiltonian

$$H_{\text{int}} = \sum_{1 \leq i_1 < i_2 < \dots < i_p \leq N} J_{i_1 i_2 \dots i_p} \sigma_{i_1} \sigma_{i_2} \dots \sigma_{i_p} \quad (54)$$

551 with Ising spins $\sigma_i = \pm 1$, $p \geq 3$ and the global spherical constraint $\sum_{i=1}^N \sigma_i^2 = N$. As
 552 for the Sherrington-Kirkpatrick model, we allow for longitudinal and transverse fields to
 553 couple to the spins (but neglect all commutators). The coupling constants $J_{i_1 i_2 \dots i_p}$ are
 554 chosen randomly with a Gaussian distribution

$$\mathcal{P}(J_{i_1 \dots i_p}) \propto \exp\left(-\frac{N^{p-1} J_{i_1 \dots i_p}^2}{p! J^2}\right). \quad (55)$$

555 Averaging the spins over some small regions, the Keldysh partition function

$$Z = \int \mathcal{D}J_{i_1 \dots i_p} \mathcal{P}(J_{i_1 \dots i_p}) \int \mathcal{D}S e^{is[S]} \quad (56)$$

556 can be written in terms of the continuous bosonic variable $S_{\sigma,i}$, where the Latin index
 557 indicates the lattice site and the Greek index $\sigma \in \{+, -\}$ denotes the branch of the
 558 Keldysh contour (see for example [69]). Due to the transverse field, the averaged spins
 559 obtain a massive dispersion. Hence, we can write the action as

$$\begin{aligned} s[S] &= s_0[S] + s_h[S] + s_\kappa[S], \\ s_0[S] &= -\frac{1}{2} \int_t \sum_{\sigma,i} \sigma S_{\sigma,i} (\partial_t^2 + m^2) S_{\sigma,i}(t), \\ s_h[S] &= \int_t \sum_{\sigma,i} \sigma h_{\sigma,i}(t) S_{\sigma,i}(t), \\ s_\kappa[S] &= -i \int dt \sum_{\sigma} \sum_{1 \leq i_1 < \dots < i_p \leq N} \sigma J_{i_1 \dots i_p} S_{\sigma,i_1} \dots S_{\sigma,i_p}, \end{aligned} \quad (57)$$

560 where the second term describes the coupling to the longitudinal external field and s_κ
 561 accounts for the effect of the disorder Hamiltonian H_{int} .

562 Averaging over the Gaussian distribution of the coupling constants $J_{i_1 \dots i_p}$ the disorder
 563 term is simplified to

$$\begin{aligned} s_\kappa[S] &= \int_{t,t'} \frac{iJ^2}{p! N^{p-1}} \sum_{i_1 < \dots < i_p} \left(\sum_{\sigma} \sigma S_{\sigma,i_1} \dots S_{\sigma,i_p} \right)^2 \\ &= \frac{i\kappa}{4} \int_{t,t'} \frac{1}{N^{p-1}} \sum_{\sigma\mu} \sigma\mu \left(\sum_{i=1}^N S_{\sigma,i}(t) S_{\mu,i}(t') \right)^p \end{aligned} \quad (58)$$

564 with $\kappa = J^2$. The global spherical constraint can be included using an auxiliary field $z_\sigma(t)$
 565 as

$$Z = \int \mathcal{D}S \mathcal{D}z e^{is[S,z]} \quad (59)$$

566 with

$$s[S, z] = s[S] + \int_t \sum_{\sigma} \sigma z_{\sigma} (S_{\sigma,i}^2 - N). \quad (60)$$

567 At this point, the action has become purely local in the site index i . Without loss of
 568 generality, we may thus focus only on a single site, dropping the irrelevant site index.

569 Next, we introduce the bilocal field $\tilde{Q}_{\sigma\mu}(t, t')$ as

$$\begin{aligned} 1 &= \int \mathcal{D}\tilde{Q} \delta \left[\tilde{Q}_{\sigma\mu}(t, t') - S_\sigma(t) S_\mu(t') \right] \\ &= \int \mathcal{D}\tilde{Q} \mathcal{D}\lambda \exp \left(\frac{i}{2} \int_{t, t'} \sum_{\sigma\mu} \lambda_{\sigma\mu}(t, t') \left[\tilde{Q}_{\sigma\mu}(t, t') - S_\sigma(t) S_\mu(t') \right] \right), \end{aligned} \quad (61)$$

570 such that the disorder term becomes

$$s_\kappa[\tilde{Q}] = \frac{i\kappa}{4} \int_{t, t'} \sum_{\sigma\mu} \sigma_\mu \tilde{Q}_{\sigma\mu}^p(t, t'). \quad (62)$$

571 We can then perform the Gaussian integral over the averaged spin fields S , which gives

$$\begin{aligned} Z &= \int \mathcal{D}\tilde{Q} \mathcal{D}\lambda \mathcal{D}z e^{is[\tilde{Q}, \lambda, z]}, \\ s[\tilde{Q}, \lambda, z] &= \frac{1}{2} \int_{t, t'} \sum_{\sigma\mu} \sigma_\mu h_\sigma(t) G_{\sigma\mu}(t, t') h_\mu(t') - \int_t \sum_\sigma \sigma z_\sigma(t) + \frac{1}{2} \text{Tr} \left(\lambda \tilde{Q} \right) \\ &\quad + \frac{i\kappa}{4} \int_{t, t'} \sum_{\sigma\mu} \sigma_\mu \tilde{Q}_{\sigma\mu}^p(t, t') - \frac{i}{2} \text{Tr} \ln(G), \end{aligned} \quad (63)$$

572 where the trace is performed over time and the contour index alike, and we have introduced
 573 the inverse spin propagator

$$G^{-1}(t, t') = \delta(t - t') \left[-(\partial_t^2 + m^2) \sigma^3 + 2 \text{diag}(z_+, -z_-) \right] - \begin{pmatrix} \lambda_{11} & \lambda_{12} \\ \lambda_{21} & \lambda_{22} \end{pmatrix} (t, t'). \quad (64)$$

574 We now turn our attention to the saddle point equations of the action $s[\tilde{Q}, \lambda, z]$. As
 575 these are most conveniently written in the $R/A/K$ basis, we introduce $z_{c/q} = z_+ \pm z_-$ such
 576 that in the new basis

$$G^{-1}(t, t') = \delta(t - t') \left[(-\partial_t^2 - m^2 + z_c(t)) \sigma^1 + z_q(t) \mathbb{1} \right] - \begin{pmatrix} \lambda^V & \lambda^A \\ \lambda^R & \lambda^K \end{pmatrix} (t, t'). \quad (65)$$

577 4.2 Late-time solution

578 We assume a constant longitudinal field $h_c = h = (h_+ + h_-)/2$, use that at the saddle
 579 point quantum fields vanish, and remember that $G^R(t, t) + G^A(t, t) = 0$ to write the saddle
 580 point equations

$$\begin{aligned} 0 &\stackrel{!}{=} \frac{\delta s}{\delta z_q(t)} = -1 + \frac{i}{2} G^K(t, t) + \frac{h^2}{2} \int_{t', t''} G^R(t', t) G^A(t, t''), \\ 0 &\stackrel{!}{=} \frac{\delta s}{\delta z_c(t)} = 0, \\ 0 &\stackrel{!}{=} \frac{\delta s}{\delta \tilde{Q}^{R/K}(t, t')} = \frac{1}{2} \lambda^{R/K}(t, t') + \frac{i\kappa}{4} p \left[\tilde{Q}^{p-1} \right]^{R/K} (t, t'), \\ 0 &\stackrel{!}{=} \frac{\delta s}{\delta \lambda^A(t, t')} = \frac{1}{2} \tilde{Q}^R(t, t') - \frac{i}{2} G^R(t, t'), \\ 0 &\stackrel{!}{=} \frac{\delta s}{\delta \lambda^V(t, t')} = \frac{1}{2} \tilde{Q}^K(t, t') - \frac{i}{2} G^K(t, t') - \frac{h^2}{2} \int_{t'', t'''} G^R(t'', t) G^A(t', t'''). \end{aligned} \quad (66)$$

581 Here $[\tilde{Q}^p]^{R/K}$ refers to the retarded/Keldysh component of the p -th power of the matrix
 582 \tilde{Q} . These equations are to be compared with Eq. (3.17) in Ref. [46].

583 To simplify these equations even further, we specify $p = 3$. Furthermore, we introduce
 584 the real fields $Q^R(t, t') = i\tilde{Q}^R(t, t')$ and $Q^K(t, t') = \tilde{Q}^K(t, t')$, which then satisfy

$$\begin{aligned} Q^K(t, t) &= 2 \\ Q^R(t, t') &= [\delta(t - t') (\partial_t^2 + m^2 - z_c(t)) - \Sigma^R(t, t')]^{-1} \\ Q^K(t, t') &= \int_{t'', t'''} Q^R(t, t'') \Sigma^K(t'', t''') Q^A(t''', t') \end{aligned} \quad (67)$$

585 with the self-energies

$$\begin{aligned} \Sigma^R(t, t') &= 3\kappa Q^R(t, t') Q^K(t, t') \\ \Sigma^K(t, t') &= \frac{3\kappa}{2} \left([Q^K]^2(t, t') - [Q^R]^2(t, t') - [Q^A]^2(t, t') \right) + h(t)h(t'), \end{aligned} \quad (68)$$

586 which are both real and in the case of Σ^K non-negative.

587 Following the arguments of Sec. 2, we distinguish between fast and slow fields $Q_{f/s}(t)$
 588 in the time-translation invariant ansatz $Q(t) = Q_f(t) + Q_s(t)$. We then once again make
 589 a generalized thermal ansatz $Q_s^R(t) = -X(t)\theta(t)\partial_t Q_s^K(t)$. Since the slow field varies on
 590 a time scale that diverges as $T \rightarrow \infty$ this implies that the retarded Green's function
 591 decays more quickly than the Keldysh component. Consequently, the Keldysh self-energy
 592 simplifies as follows

$$\begin{aligned} \Sigma_s^K(t) &= \frac{3\kappa}{2} \left([Q_s^K]^2(t) - [Q_s^R]^2(t) - [Q_s^A]^2(t) \right) \\ &= \frac{3\kappa}{2} [Q_s^K]^2(t). \end{aligned} \quad (69)$$

593 From this, it follows immediately that the self-energy satisfies the generalized fluctuation-
 594 dissipation relation $\Sigma_s^R(t) = -X(t)\partial_t \Sigma_s^K(t)$. Similarly, the most slowly decaying contribu-
 595 tion to the Keldysh component Q_s^K must involve Σ_s^K such that we can write

$$Q_s^K = Q^R \circ \Sigma_s^K \circ Q^A. \quad (70)$$

596 It is now more convenient to rewrite the equations of motion of the slow field in the more
 597 conventional form

$$[Q_f^R]^{-1} \circ Q_s^R = \Sigma_s^R \circ Q^R, \quad (71)$$

598

$$[Q_f^R]^{-1} \circ Q_s^K = \Sigma_s^K \circ Q^A + \Sigma_s^R \circ Q_s^K. \quad (72)$$

599 In the case of dissipative dynamics, these equations coincide with those derived by Som-
 600 polinsky and Zippelius [7, 37] and solved by Cugliandolo and Kurchan [5]. As has been
 601 noted before [21], we find that these equations of motion satisfied by the p -spin model are
 602 surprisingly similar to those derived from mode coupling theory in the context of structural
 603 glasses [22].

604 Assuming ultrametricity, we satisfy all conditions required for the general argument
 605 of Sec. 2, where we showed that the matrix multiplication in replica space is identical to
 606 the Keldysh component of the product of functions in Keldysh space. From the general

607 matrix multiplication follows the same statement also for matrix inversion. Hence, we
 608 conclude that the equation for the Keldysh component of the expression

$$Q(t) = [\delta(t)\sigma^1(\partial_t^2 + m^2 - z_c) - \Sigma(t)]^{-1} \quad (73)$$

609 or equivalently the solution to (72) is similar to that obtained in replica formalism (see
 610 for example Eq. (3.17) in Ref. [46], which differs in the conventions for mass and coupling
 611 strength).

612 In summary, we find

$$Q_s^K(u) = \begin{cases} q_0 = -\frac{q_f\sigma_0(\sigma_f-2z)}{(\sigma_f+x(\sigma_1-\sigma_0)-2z)^2} & u < x \\ q_1 = q_0 - \frac{q_f(\sigma_1-\sigma_0)}{\sigma_f+x(\sigma_1-\sigma_0)-2z} & u > x \end{cases} . \quad (74)$$

613 with the shorthand notation

$$\Sigma_s^K(u) = \frac{3\kappa}{2} [Q_s^K]^2(u) + h^2 = \begin{cases} \sigma_0 & u < x \\ \sigma_1 & u > x \end{cases} . \quad (75)$$

614 Furthermore, the fast field satisfies

$$\begin{aligned} G_f^K(t) &= G_f^R \circ \Sigma_f^K \circ G_f^A, \\ \Sigma_f^K(t) &= \frac{3\kappa}{2} (Q_f^K(t) + 2q_1) Q_f^K(t), \end{aligned} \quad (76)$$

615 which we abbreviated above as $q_f = Q_f^K(t=0)$ and $\sigma_f = \Sigma_f^K(t=0)$. Finally, the
 616 Lagrange parameter $z = (z_c - m^2)/\beta$ is fixed by the additional constraint

$$Q^K(u=1) \equiv q_1 + q_f \equiv Q_s^K(t=0) + Q_f^K(t=0) \stackrel{!}{=} 2. \quad (77)$$

617 Conversely to Eq. (74), the effective inverse temperature mirrors the structure of 1-step
 618 RSB

$$X(q) = \begin{cases} 0 & q < q_0 \\ \frac{\beta x}{2} & q_0 < q < q_1 \\ \frac{\beta}{2} & q = q_1 \end{cases} . \quad (78)$$

619 Note, that once again, it is not possible to reconstruct $Q_s^K(t)$ because the information
 620 on the time-dependence was lost during the change of variables $t \rightarrow X(t)$ in Eq. (14).
 621 Furthermore, the breakpoint x has to be determined by an additional criterion, requiring
 622 either marginal stability or minimization of the free energy [35, 46].

623 Due to the equivalence between the ultrametric Keldysh and the replica formalism, we
 624 conclude that our approach finds the same critical point and one-step RSB as reported
 625 in Ref. [46], provided the same condition for x is used. On the other hand, at any finite
 626 time T , ultrametric relations must be violated and an analysis similar to that of Ref. [22]
 627 shows that on a finite time interval in the one-time formulation, the spin glass phase is
 628 indistinguishable from a ferromagnet.

629 The comparison between ultrametric Keldysh and replica formalism for the p -spin
 630 model has already been addressed by Crisanti et al. some 31 years ago [35]. Although
 631 they use a slightly different ansatz for the generalized fluctuation-dissipation relation in
 632 the aging regime

$$Q^R(t) = -x\theta(t)\partial_t Q^K(t), \quad (79)$$

633 where $x \in [0, 1]$ corresponds to the position of the discontinuity in the replica formalism.
 634 In the case of 1-step RSB without a longitudinal field, this ansatz also reproduces the
 635 replica equations. The reported difference between the dynamical and equilibrium critical
 636 temperature is related in part to the different conditions used to fix x . In the dynamical
 637 case, matching with the fast dynamics implies a marginal stability condition as opposed
 638 to a minimization of the free energy in equilibrium. Furthermore, as we had anticipated
 639 below Eq.(12) for models with a finite number of replica symmetry breaking steps, the
 640 Keldysh Green's function of the spherical p -spin model does not become ultrametric at late
 641 times [5]. Consequently, the aging dynamics of the spherical p -spin model never reaches
 642 equilibrium.

643 An intuitive explanation of this observation can be given using the Thouless-Anderson-
 644 Palmer free energy [78]. One finds that the dynamics of the spherical p -spin model gets
 645 stuck in local minima that are separated from the equilibrium solution by energy barri-
 646 ers that diverge in the thermodynamic limit. For comparison, the slow evolution of the
 647 Sherrington-Kirkpatrick model is explained by an entropic effect: As the system relaxes,
 648 it evolves through a series of saddle points with an ever decreasing number of unstable
 649 directions resulting in long, but finite escape times [17].

650 5 Discussion

651 The results presented in this article rely on the existence of a finite temperature to which
 652 the system equilibrates on short relative times $t < \tau_{\text{erg}}$, see Fig. 1(b). Specifically, as we
 653 send the center-of-mass time $T \rightarrow \infty$, the ultrametric solutions (52)(78) are parametrized
 654 by the inverse temperature β . However, in a spin glass, no global equilibrium is reached.
 655 We identify the absence of a global temperature as the characteristic property of the
 656 ultrametric spin glass. This is independent of the breaking of time translation invariance
 657 at finite center-of-mass times T . We also address to which extent these conclusions apply
 658 to quantum critical quenches at zero temperature.

659 5.1 Spontaneous breaking of thermal symmetry

660 The non-analytic behavior of the ultrametric solution at x emerges in the temporal ther-
 661 modynamic limit $T \rightarrow \infty$ (in space, the mean-field system is assumed to be in the thermo-
 662 dynamic limit by construction). The ultrametric solution corresponds to a spontaneous
 663 breaking of the thermal (or Kubo-Martin-Schwinger, KMS) symmetry [79–82]

$$S_{\sigma,i}(t) \rightarrow S_{\sigma,i}(-t + i\sigma\beta/2), \quad i \rightarrow -i, \quad h \rightarrow -h, \quad (80)$$

664 which is present in the stationary state of an ergodic system with a time-independent
 665 Hamiltonian generator of dynamics characterized by an inverse temperature β . Via our
 666 construction, replica symmetry breaking thus gets stringently tied to the spontaneous
 667 breaking of thermal symmetry – or more physically speaking, of ergodicity.

668 We emphasize that, since T drops out of the equations of motion at asymptotically
 669 late times, which can be seen explicitly in Eq. (46), all microscopic details of the quench
 670 protocol disappear from the problem. The time-translation invariant discussion presented
 671 here is, therefore, independent of the details of the aging process. It instead extracts
 672 solely the universal property common to all classical glasses: The spontaneous breaking of
 673 thermal symmetry. The emergence of this broken symmetry at finite times was previously
 674 anticipated by Kurchan [74].

675 For glasses, it is found that a weak long-term memory is necessary to preclude ther-
 676 malization on all scales. Although this implies that time translation symmetry remains

677 broken at any finite time T following a quench, our time translation invariant approach
 678 clarifies that the persistence of broken time translation invariance, and thus aging, should
 679 not be equated to ergodicity breaking in the stationary state. Instead, the emergence of
 680 reparametrization invariance lifts this connection at asymptotically late times [5, 6].

681 5.2 Zero temperature limit

682 The finite temperature spin glasses discussed here are solved by the classical ansatz
 683 $G^R(t) \sim \beta \partial_t G^K(t)$ with different scaling dimensions for response and correlation func-
 684 tions. The classical scaling, therefore, requires the existence of a time scale that enters the
 685 asymptotic solution as inverse temperature. For a quench through the quantum critical
 686 point at zero temperature, one, therefore, expects one of two options: Either β emerges
 687 as a result of the finite energy density imposed upon the system during the quench, or the
 688 absence of a fixed time scale suggests quantum scaling

$$G_s^R \sim G_s^K. \quad (81)$$

689 In the following, we will address the implications of quantum scaling. With Eq. (81), it is
 690 not possible to expand the equations of motion in powers of G^R . Furthermore, the failure
 691 of the generalized thermal ansatz indicates the necessity of a dynamic Parisi function.

692 The characteristic observable feature of a glass is aging, which implies that correlations
 693 $G_s^K(t)$ decay infinitely slowly as $T \rightarrow \infty$. In the quantum regime, assuming the above
 694 scaling, the same must apply to the response function G_s^R , and thus the self-energy Σ_s^R .
 695 Hence, as the infrared cutoff T^{-1} is sent to zero, memory integrals of the form $\Sigma_s^R \circ G_s^K$
 696 diverge. We emphasize the similarity of this argument to the Mermin-Wagner theorem
 697 that prevents spontaneous symmetry breaking due to infrared fluctuations – here, these
 698 fluctuations prevent the ergodicity breaking identified in the classical case above, upon
 699 removing the infrared cutoff $T \rightarrow \infty$. Consequently, the quantum regime characterized
 700 by Eq. (81) is always transient and bounded by the energy density imparted upon the
 701 system by the initial quench. According to this argument, at asymptotically late times,
 702 spin glasses are necessarily classical (see also [41, 83]) with a temperature determined by
 703 the energy density after the quench.

704 We re-emphasize, however, that the argument here relies on the assumption of a com-
 705 mon scaling of retarded and Keldysh Green’s functions. This raises the question of whether
 706 more general forms of ergodicity breaking could be realized at zero temperature.

707 Recent experiments are performed at very low temperatures and finite times [53–58].
 708 In addition to possible asymptotic symmetry-breaking phenomena, weak quenches at zero
 709 temperature could also display interesting intermediate-time dynamical phenomena related
 710 to their quantum mechanical microscopic physics.

711 At the current level of the analysis presented here, it is not possible to recover the
 712 time scales associated with the effective temperature X , which hinders the investigation
 713 of transient regimes. However, by continuing the Wigner expansion, it is possible to
 714 systematically restore corrections due to the boundary at $t = 2T$ and derivatives with
 715 respect to the center-of-mass time. It is then possible to work backward from the latest
 716 times to recover the explicit time dependence of the aging solution, including a potential
 717 transient quantum critical regime.

718 6 Outlook

719 Recent realizations of spin glasses with Rydberg atoms are affected by decoherence due
 720 to dephasing caused by fluctuations in the external fields (i.e. lasers) and spontaneous

721 emission from the Rydberg state [54, 59]. A more realistic description of the system will
 722 take these into account. This necessitates the treatment of an open system with a time
 723 evolution governed by the Lindblad equation

$$\partial_t \rho(t) = -i[H, \rho] + \kappa \sum_i \left(L_i \rho L_i^\dagger - \frac{1}{2} \{L_i L_i^\dagger, \rho\} \right). \quad (82)$$

724 Here, $\rho(t)$ denotes the density matrix, the Hamiltonian H is that of Eq. (23), and the
 725 Hermitian Lindblad operators $L_i = \sigma_i^3$ describe dephasing noise that acts incoherently on
 726 all atoms. The decoherence introduced by the Lindblad operators causes heating. Specifi-
 727 cally, for Hermitian L_i , the stationary state has infinite temperature. Dephasing, therefore,
 728 introduces a time scale beyond which the system becomes paramagnetic, independent of
 729 the initial quench. At late times, dephasing needs to be taken into account by simulations
 730 of the experimental systems.

731 It is a strength of the Keldysh field theory that the inclusion of decoherence is very nat-
 732 ural and requires little additional effort [69]. This is in contrast to microscopic approaches
 733 like exact diagonalization or matrix product states, particularly in quantum systems at
 734 low temperatures, when the system becomes highly entangled [66]. Despite this advan-
 735 tage, simulations of the glass phase, even in mean-field models, remain challenging. The
 736 reason is the weak long-term memory, which precludes using a finite cutoff time for mem-
 737 ory integrals. The numerical effort therefore scales with time to the third power, which
 738 currently limits this method to short times. However, these limitations can be lifted [84]
 739 and long-time simulations of the quench dynamics will be addressed in the future [85].

740 Finally, we mention the connection to Sachdev-Ye-Kitaev (SYK) models, which have
 741 quantum ‘spin liquid’ ground states [86]. These states are quite distinct from the spin glass
 742 ground states considered in the present paper, as they do not have any aging behavior,
 743 and are described by a replica diagonal saddle point. The low energy theory of SYK
 744 models exhibits an emergent time reparameterization symmetry while preserving thermal
 745 symmetry. This has enabled a detailed understanding of their quantum dynamics at a
 746 finite number of spins N_s , well beyond the $N_s = \infty$ saddle point. The quantum spin glass
 747 states considered in the present paper also have an emergent time reparameterization
 748 symmetry, but the glassy dynamics break thermal symmetry [87]. All our analysis here
 749 has been in the $N_s = \infty$ saddle point theory, and it would be interesting to adapt the
 750 SYK technology to understand the structure of the finite N_s theory. However, the broken
 751 thermal symmetry makes this task considerably more difficult.

752 *Acknowledgements* – We thank Alexander Altland, Giulio Biroli, Leticia Cugliandolo,
 753 Antoine Georges, Nikita Kavokine, Jorge Kurchan, Olivier Parcollet, Rhine Samajdar,
 754 Marco Schiró, and Maria Tikhanovskaya for valuable discussions. S.S. thanks Rhine Sama-
 755 jdar and Maria Tikhanovskaya for an earlier collaboration [46]. The work of J.L. and
 756 S.D. was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research
 757 Foundation) CRC 1238 project number 277146847. S.S. has been supported by the U.S.
 758 Department of Energy under Grant DE-SC0019030.

759 References

- 760 [1] N. D. Mackenzie and A. P. Young, *Lack of Ergodicity in the Infinite-Range Ising*
 761 *Spin-Glass*, Phys. Rev. Lett. **49**, 301 (1982), doi:10.1103/PhysRevLett.49.301.
- 762 [2] F. Ritort, *Glassiness in a Model without Energy Barriers*, Phys. Rev. Lett. **75**, 1190
 763 (1995), doi:10.1103/PhysRevLett.75.1190.

- 764 [3] S. Franz and F. Ritort, *Dynamical Solution of a Model without Energy Barriers*,
765 Europhysics Letters **31**(9), 507 (1995), doi:10.1209/0295-5075/31/9/001.
- 766 [4] C. Godreche, J. P. Bouchaud and M. Mézard, *Entropy barriers and slow relaxation in*
767 *some random walk models*, Journal of Physics A: Mathematical and General **28**(23),
768 L603 (1995), doi:10.1088/0305-4470/28/23/002.
- 769 [5] L. F. Cugliandolo and J. Kurchan, *Analytical solution of the off-equilibrium*
770 *dynamics of a long-range spin-glass model*, Phys. Rev. Lett. **71**, 173 (1993),
771 doi:10.1103/PhysRevLett.71.173.
- 772 [6] L. F. Cugliandolo and J. Kurchan, *On the out-of-equilibrium relaxation of the*
773 *Sherrington-Kirkpatrick model*, Journal of Physics A: Mathematical and General
774 **27**(17), 5749 (1994), doi:10.1088/0305-4470/27/17/011.
- 775 [7] H. Sompolinsky and A. Zippelius, *Dynamic Theory of the Spin-Glass Phase*, Phys.
776 Rev. Lett. **47**, 359 (1981), doi:10.1103/PhysRevLett.47.359.
- 777 [8] L. C. E. Struik, *Physical aging in amorphous polymers and other materials*, Elsevier
778 Scientific Pub. Co. ; Distributors for the U.S. and Canada, Elsevier North-Holland,
779 Amsterdam, New York, ISBN 0444416552; 9780444416551 (1978).
- 780 [9] M. Alba, J. Hammann and M. Nogues, *Phase diagrams of two dilute insulat-*
781 *ing systems with competing interactions: CdCr_{2x}In_{2-2x}S₄ and ZnCr_{2x}Al_{2-2x}S₄*,
782 Journal of Physics C: Solid State Physics **15**(26), 5441 (1982), doi:10.1088/0022-
783 3719/15/26/022.
- 784 [10] L. Lundgren, P. Svedlindh, P. Nordblad and O. Beckman, *Dynamics of the*
785 *Relaxation-Time Spectrum in a CuMn Spin-Glass*, Phys. Rev. Lett. **51**, 911 (1983),
786 doi:10.1103/PhysRevLett.51.911.
- 787 [11] P. Nordblad, P. Svedlindh, L. Lundgren and L. Sandlund, *Time decay of the*
788 *remanent magnetization in a CuMn spin glass*, Phys. Rev. B **33**, 645 (1986),
789 doi:10.1103/PhysRevB.33.645.
- 790 [12] M. Alba, M. Ocio and J. Hammann, *Ageing Process and Response Function in Spin*
791 *Glasses: An Analysis of the Thermoremanent Magnetization Decay in Ag:Mn (2.6 %)*,
792 Europhysics Letters **2**(1), 45 (1986), doi:10.1209/0295-5075/2/1/007.
- 793 [13] J.-P. Bouchard, L. F. Cugliandolo, J. Kurchan and M. Mézard, *Out of equilibrium*
794 *dynamics in spin-glasses and other glassy systems*, pp. 161–223, World Scientific,
795 doi:10.1142/9789812819437-0006 (1998).
- 796 [14] G. Parisi, *Toward a mean field theory for spin glasses*, Physics Letters A **73**(3), 203
797 (1979), doi:https://doi.org/10.1016/0375-9601(79)90708-4.
- 798 [15] G. Parisi, *Infinite Number of Order Parameters for Spin-Glasses*, Phys. Rev. Lett.
799 **43**, 1754 (1979), doi:10.1103/PhysRevLett.43.1754.
- 800 [16] G. Parisi, *A sequence of approximated solutions to the S-K model for spin*
801 *glasses*, Journal of Physics A: Mathematical and General **13**(4), L115 (1980),
802 doi:10.1088/0305-4470/13/4/009.
- 803 [17] M. Mézard, G. Parisi and M. Virasoro, *Spin Glass Theory and Beyond*, WORLD
804 SCIENTIFIC, doi:10.1142/0271 (1986), <https://www.worldscientific.com/doi/pdf/10.1142/0271>.
- 805

- 806 [18] T. Castellani and A. Cavagna, *Spin-glass theory for pedestrians*, Journal of Statistical
807 Mechanics: Theory and Experiment **2005**(05), P05012 (2005), doi:10.1088/1742-
808 5468/2005/05/P05012.
- 809 [19] G. Parisi, *Order Parameter for Spin-Glasses*, Phys. Rev. Lett. **50**, 1946 (1983),
810 doi:10.1103/PhysRevLett.50.1946.
- 811 [20] Mézard, M. and Virasoro, M. A., *The microstructure of ultrametricity*, J. Phys.
812 France **46**(8), 1293 (1985), doi:10.1051/jphys:019850046080129300.
- 813 [21] T. R. Kirkpatrick and D. Thirumalai, *p-spin-interaction spin-glass models: Con-*
814 *nections with the structural glass problem*, Phys. Rev. B **36**, 5388 (1987),
815 doi:10.1103/PhysRevB.36.5388.
- 816 [22] E. Leutheusser, *Dynamical model of the liquid-glass transition*, Phys. Rev. A **29**,
817 2765 (1984), doi:10.1103/PhysRevA.29.2765.
- 818 [23] U. Bengtzelius, W. Götze and A. Sjolander, *Dynamics of supercooled liquids and*
819 *the glass transition*, Journal of Physics C: Solid State Physics **17**(33), 5915 (1984),
820 doi:10.1088/0022-3719/17/33/005.
- 821 [24] W. Götze and L. Sjogren, *Relaxation processes in supercooled liquids*, Reports on
822 Progress in Physics **55**(3), 241 (1992), doi:10.1088/0034-4885/55/3/001.
- 823 [25] W. Kob and H. C. Andersen, *Scaling Behavior in the β -Relaxation Regime*
824 *of a Supercooled Lennard-Jones Mixture*, Phys. Rev. Lett. **73**, 1376 (1994),
825 doi:10.1103/PhysRevLett.73.1376.
- 826 [26] W. Kob and H. C. Andersen, *Testing mode-coupling theory for a supercooled binary*
827 *Lennard-Jones mixture I: The van Hove correlation function*, Phys. Rev. E **51**, 4626
828 (1995), doi:10.1103/PhysRevE.51.4626.
- 829 [27] W. Kob and H. C. Andersen, *Testing mode-coupling theory for a supercooled binary*
830 *Lennard-Jones mixture. II. Intermediate scattering function and dynamic susceptibil-*
831 *ity*, Phys. Rev. E **52**, 4134 (1995), doi:10.1103/PhysRevE.52.4134.
- 832 [28] W. Götze, *Recent tests of the mode-coupling theory for glassy dynamics*, Journal of
833 Physics: Condensed Matter **11**(10A), A1 (1999), doi:10.1088/0953-8984/11/10A/002.
- 834 [29] C. Angell, *Perspective on the glass transition*, Journal of Physics and Chemistry of
835 Solids **49**(8), 863 (1988), doi:https://doi.org/10.1016/0022-3697(88)90002-9.
- 836 [30] H. Sompolinsky, *Time-Dependent Order Parameters in Spin-Glasses*, Phys. Rev.
837 Lett. **47**, 935 (1981), doi:10.1103/PhysRevLett.47.935.
- 838 [31] H. Horner, *Dynamic mean field theory of the SK-spin glass*, Zeitschrift für Physik B
839 Condensed Matter **57**(1), 29 (1984), doi:10.1007/BF01679923.
- 840 [32] H. Horner, *Dynamic mean field theory of the SK-spin glass*, Zeitschrift für Physik B
841 Condensed Matter **57**(1), 39 (1984), doi:10.1007/BF01679924.
- 842 [33] L. B. Ioffe, *Quasiequilibrium states of spin glasses*, Phys. Rev. B **38**, 5181 (1988),
843 doi:10.1103/PhysRevB.38.5181.
- 844 [34] R. G. Palmer, D. L. Stein, E. Abrahams and P. W. Anderson, *Models of Hierarchi-*
845 *cally Constrained Dynamics for Glassy Relaxation*, Phys. Rev. Lett. **53**, 958 (1984),
846 doi:10.1103/PhysRevLett.53.958.

- 847 [35] A. Crisanti, H. Horner and H.-J. Sommers, *The spherical p -spin interaction spin-glass*
848 *model*, Zeitschrift für Physik B Condensed Matter **92**(2), 257 (1993).
- 849 [36] G. Biroli and L. F. Cugliandolo, *Quantum Thouless-Anderson-Palmer equations for*
850 *glassy systems*, Phys. Rev. B **64**, 014206 (2001), doi:10.1103/PhysRevB.64.014206.
- 851 [37] H. Sompolinsky and A. Zippelius, *Relaxational dynamics of the Edwards-Anderson*
852 *model and the mean-field theory of spin-glasses*, Phys. Rev. B **25**, 6860 (1982),
853 doi:10.1103/PhysRevB.25.6860.
- 854 [38] L. F. Cugliandolo and J. Kurchan, *Weak ergodicity breaking in mean-*
855 *field spin-glass models*, Philosophical Magazine B **71**(4), 501 (1995),
856 doi:10.1080/01418639508238541.
- 857 [39] J. Kurchan, *Supersymmetry in spin glass dynamics*, J. Phys. I France **2**(7), 1333
858 (1992), doi:10.1051/jp1:1992214.
- 859 [40] J. P. Bouchaud, *Weak ergodicity breaking and aging in disordered systems*, J. Phys.
860 I France **2**(9), 1705 (1992), doi:10.1051/jp1:1992238.
- 861 [41] L. F. Cugliandolo and G. Lozano, *Real-time nonequilibrium dynamics of quantum*
862 *glassy systems*, Phys. Rev. B **59**, 915 (1999), doi:10.1103/PhysRevB.59.915.
- 863 [42] S. J. Thomson, P. Urbani and M. Schiró, *Quantum Quenches in Isolated Quan-*
864 *tum Glasses out of Equilibrium*, Phys. Rev. Lett. **125**(12), 120602 (2020),
865 doi:10.1103/PhysRevLett.125.120602, 1904.03147.
- 866 [43] G. Biroli and O. Parcollet, *Out-of-equilibrium dynamics of a quantum Heisenberg*
867 *spin glass*, Phys. Rev. B **65**, 094414 (2002), doi:10.1103/PhysRevB.65.094414.
- 868 [44] M. P. Kennett and C. Chamon, *Time Reparametrization Group and the Long*
869 *Time Behavior in Quantum Glassy Systems*, Phys. Rev. Lett. **86**, 1622 (2001),
870 doi:10.1103/PhysRevLett.86.1622.
- 871 [45] M. P. Kennett, C. Chamon and J. Ye, *Aging dynamics of quantum spin glasses of*
872 *rotors*, Phys. Rev. B **64**, 224408 (2001), doi:10.1103/PhysRevB.64.224408.
- 873 [46] M. Tikhanovskaya, S. Sachdev and R. Samajdar, *Equilibrium Dynamics of Infinite-*
874 *Range Quantum Spin Glasses in a Field*, PRX Quantum **5**(2), 020313 (2024),
875 doi:10.1103/PRXQuantum.5.020313, 2309.03255.
- 876 [47] N. Kavokine, M. Müller, A. Georges and O. Parcollet, *Exact numerical solution of the*
877 *classical and quantum Heisenberg spin glass*, arXiv e-prints arXiv:2312.14598 (2023),
878 doi:10.48550/arXiv.2312.14598, 2312.14598.
- 879 [48] J. Kurchan, *Supersymmetry in spin glass dynamics*, J. Phys. I France **2**(7), 1333
880 (1992), doi:10.1051/jp1:1992214.
- 881 [49] I. Aref'eva, M. Khramtsov, M. Tikhanovskaya and I. Volovich, *Replica-nondiagonal*
882 *solutions in the SYK model*, Journal of High Energy Physics **2019**(7), 113 (2019),
883 doi:10.1007/JHEP07(2019)113.
- 884 [50] S. F. Edwards and P. W. Anderson, *Theory of spin glasses*, Journal of Physics F:
885 Metal Physics **5**(5), 965 (1975), doi:10.1088/0305-4608/5/5/017.
- 886 [51] A. Kamenev, *Field Theory of Non-Equilibrium Systems*, Cambridge University Press,
887 ISBN 9781139500296 (2011).

- 888 [52] A. Houghton, S. Jain and A. P. Young, *Role of initial conditions in the*
889 *mean-field theory of spin-glass dynamics*, Phys. Rev. B **28**, 2630 (1983),
890 doi:10.1103/PhysRevB.28.2630.
- 891 [53] H. Labuhn, D. Barredo, S. Ravets, S. de Léséleuc, T. Macrì, T. Lahaye and
892 A. Browaeys, *Tunable two-dimensional arrays of single Rydberg atoms for realizing*
893 *quantum Ising models*, Nature **534**(7609), 667 (2016), doi:10.1038/nature18274.
- 894 [54] A. Signoles, T. Franz, R. Ferracini Alves, M. Gärttner, S. Whitlock, G. Zürn and
895 M. Weidemüller, *Glassy Dynamics in a Disordered Heisenberg Quantum Spin System*,
896 Phys. Rev. X **11**, 011011 (2021), doi:10.1103/PhysRevX.11.011011.
- 897 [55] M. Kim, K. Kim, J. Hwang, E.-G. Moon and J. Ahn, *Rydberg quantum*
898 *wires for maximum independent set problems*, Nature Physics **18**(7), 755 (2022),
899 doi:10.1038/s41567-022-01629-5.
- 900 [56] A. Byun, M. Kim and J. Ahn, *Finding the Maximum Independent Sets*
901 *of Platonic Graphs Using Rydberg Atoms*, PRX Quantum **3**, 030305 (2022),
902 doi:10.1103/PRXQuantum.3.030305.
- 903 [57] M.-T. Nguyen, J.-G. Liu, J. Wurtz, M. D. Lukin, S.-T. Wang and H. Pichler, *Quantum*
904 *Optimization with Arbitrary Connectivity Using Rydberg Atom Arrays*, PRX
905 Quantum **4**, 010316 (2023), doi:10.1103/PRXQuantum.4.010316.
- 906 [58] S. Jeong, M. Kim, M. Hhan, J. Park and J. Ahn, *Quantum programming of the*
907 *satisfiability problem with Rydberg atom graphs*, Phys. Rev. Res. **5**, 043037 (2023),
908 doi:10.1103/PhysRevResearch.5.043037.
- 909 [59] A. Browaeys and T. Lahaye, *Many-body physics with individually controlled Rydberg*
910 *atoms*, Nature Physics **16**(2), 132 (2020), doi:10.1038/s41567-019-0733-z.
- 911 [60] P. Scholl, H. J. Williams, G. Bornet, F. Wallner, D. Barredo, L. Henriët, A. Signoles,
912 C. Hainaut, T. Franz, S. Geier, A. Tebben, A. Salzinger *et al.*, *Microwave Engineering*
913 *of Programmable XXZ Hamiltonians in Arrays of Rydberg Atoms*, PRX Quantum
914 **3**, 020303 (2022), doi:10.1103/PRXQuantum.3.020303.
- 915 [61] A. D. King, J. Raymond, T. Lanting, R. Harris, A. Zucca, F. Altomare, A. J. Berkley,
916 K. Boothby, S. Ejtemaee, C. Enderud, E. Hoskinson, S. Huang *et al.*, *Quantum*
917 *critical dynamics in a 5,000-qubit programmable spin glass*, Nature **617**(7959), 61
918 (2023), doi:10.1038/s41586-023-05867-2.
- 919 [62] V. D. Vaidya, Y. Guo, R. M. Kroeze, K. E. Ballantine, A. J. Kollár, J. Keeling
920 and B. L. Lev, *Tunable-Range, Photon-Mediated Atomic Interactions in Multimode*
921 *Cavity QED*, Phys. Rev. X **8**, 011002 (2018), doi:10.1103/PhysRevX.8.011002.
- 922 [63] Y. Guo, R. M. Kroeze, V. D. Vaidya, J. Keeling and B. L. Lev, *Sign-Changing*
923 *Photon-Mediated Atom Interactions in Multimode Cavity Quantum Electrodynamics*,
924 Phys. Rev. Lett. **122**, 193601 (2019), doi:10.1103/PhysRevLett.122.193601.
- 925 [64] H. Hosseinabadi, D. E. Chang and J. Marino, *Dynamics of spin glass formation under*
926 *tunable fluctuations in frustrated cavity QED experiments* (2023), 2311.05682.
- 927 [65] H. Hosseinabadi, D. E. Chang and J. Marino, *Non-equilibrium Dyson equations for*
928 *strongly coupled light and matter: spin glass formation in multi-mode cavity QED*
929 (2023), 2312.11624.

- 930 [66] B. P. Marsh, R. M. Kroeze, S. Ganguli, S. Gopalakrishnan, J. Keeling and B. L. Lev,
931 *Entanglement and Replica Symmetry Breaking in a Driven-Dissipative Quantum Spin*
932 *Glass*, Phys. Rev. X **14**, 011026 (2024), doi:10.1103/PhysRevX.14.011026.
- 933 [67] J. Ye, S. Sachdev and N. Read, *Solvable spin glass of quantum rotors*, Phys. Rev.
934 Lett. **70**, 4011 (1993), doi:10.1103/PhysRevLett.70.4011.
- 935 [68] N. Read, S. Sachdev and J. Ye, *Landau theory of quantum spin glasses of rotors and*
936 *Ising spins*, Phys. Rev. B **52**, 384 (1995), doi:10.1103/PhysRevB.52.384.
- 937 [69] L. M. Sieberer, M. Buchhold and S. Diehl, *Keldysh field theory for driven open quan-*
938 *tum systems*, Reports on Progress in Physics **79**(9), 096001 (2016), doi:10.1088/0034-
939 4885/79/9/096001.
- 940 [70] J. Berges, *Non-equilibrium quantum fields: from cold atoms to cosmology*, Strongly
941 Interacting Quantum Systems out of Equilibrium. Oxford University Press, Oxford,
942 ISBN 9780198768166, doi:10.1093/acprof:oso/9780198768166.003.0002 (2016).
- 943 [71] J. R. L. de Almeida and D. J. Thouless, *Stability of the Sherrington-Kirkpatrick*
944 *solution of a spin glass model*, Journal of Physics A: Mathematical and General
945 **11**(5), 983 (1978), doi:10.1088/0305-4470/11/5/028.
- 946 [72] M. Talagrand, *The Parisi Formula*, Annals of Mathematics **163**(1), 221 (2006).
- 947 [73] D. Panchenko, *The Parisi ultrametricity conjecture*, Annals of Mathematics **177**(1),
948 383 (2013).
- 949 [74] J. Kurchan, *Time-reparametrization invariances, multithermalization and the Parisi*
950 *scheme*, SciPost Phys. Core **6**, 001 (2023), doi:10.21468/SciPostPhysCore.6.1.001.
- 951 [75] B. Derrida, *Random-energy model: An exactly solvable model of disordered systems*,
952 Phys. Rev. B **24**, 2613 (1981), doi:10.1103/PhysRevB.24.2613.
- 953 [76] Mézard, M., Parisi, G., Sourlas, N., Toulouse, G. and Virasoro, M., *Replica symmetry*
954 *breaking and the nature of the spin glass phase*, J. Phys. France **45**(5), 843 (1984),
955 doi:10.1051/jphys:01984004505084300.
- 956 [77] A. Crisanti and H.-J. Sommers, *The spherical p-spin interaction spin glass*
957 *model: the statics*, Zeitschrift für Physik B Condensed Matter **87**(3), 341 (1992),
958 doi:10.1007/BF01309287.
- 959 [78] D. J. Thouless, P. W. Anderson and R. G. Palmer, *Solution of 'Solvable model of*
960 *a spin glass'*, The Philosophical Magazine: A Journal of Theoretical Experimental
961 and Applied Physics **35**(3), 593 (1977), doi:10.1080/14786437708235992, <https://doi.org/10.1080/14786437708235992>.
962
- 963 [79] L. M. Sieberer, A. Chiocchetta, A. Gambassi, U. C. Täuber and S. Diehl, *Thermo-*
964 *dynamic equilibrium as a symmetry of the Schwinger-Keldysh action*, Phys. Rev. B
965 **92**, 134307 (2015), doi:10.1103/PhysRevB.92.134307.
- 966 [80] M. Crossley, P. Glorioso and H. Liu, *Effective field theory of dissipative fluids*, J.
967 High Energy Phys. **2017**(9), 95 (2017), doi:10.1007/JHEP09(2017)095, 1511.03646.
- 968 [81] F. M. Haehl, R. Loganayagam and M. Rangamani, *The fluid manifesto: emergent*
969 *symmetries, hydrodynamics, and black holes*, J. High Energy Phys. **2016**(1), 184
970 (2016), doi:10.1007/JHEP01(2016)184, 1510.02494.

- 971 [82] C. Aron, G. Biroli and L. Cugliandolo, *(Non) equilibrium dynamics: a (bro-*
972 *ken) symmetry of the Keldysh generating functional*, SciPost Physics **4**(1) (2018),
973 doi:10.21468/scipostphys.4.1.008.
- 974 [83] A. Andreanov and M. Müller, *Long-Range Quantum Ising Spin Glasses at $T=0$:*
975 *Gapless Collective Excitations and Universality*, Phys. Rev. Lett. **109**, 177201 (2012),
976 doi:10.1103/PhysRevLett.109.177201.
- 977 [84] J. Kaye and D. Golež, *Low rank compression in the numerical solu-*
978 *tion of the nonequilibrium Dyson equation*, SciPost Phys. **10**, 091 (2021),
979 doi:10.21468/SciPostPhys.10.4.091.
- 980 [85] J. Lang, S. Sachdev and S. Diehl, *in preparation* (2024).
- 981 [86] D. Chowdhury, A. Georges, O. Parcollet and S. Sachdev, *Sachdev-Ye-Kitaev models*
982 *and beyond: Window into non-Fermi liquids*, Rev. Mod. Phys. **94**(3), 035004 (2022),
983 doi:10.1103/RevModPhys.94.035004, 2109.05037.
- 984 [87] S. Sachdev, *Quantum glasses, reparameterization invariance, Sachdev-Ye-Kitaev*
985 *models* (2024), 2402.17824.