

Necklace Ansatz for strongly repulsive spin mixtures on a ring

Gianni Aupetit-Diallo ^{1*}, Giovanni Pecci ^{1†}, Artem Volosniev ^{2‡}, Mathias Albert ^{3,4◦},
Anna Minguzzi ^{5§} and Patrizia Vignolo ^{3,4¶}

¹ SISSA, Via Bonomea 265, I-34136 Trieste, Italy

² Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, DK-8000
Aarhus C, Denmark

³ Université Côte d'Azur, CNRS, Institut de Physique de Nice, 06200 Nice, France

⁴ Institut Universitaire de France

⁵ Université Grenoble Alpes, CNRS, LPMMC, 38000 Grenoble, France

* gaupetit@sissa.it, † gpecci@sissa.it, ‡ artem@phys.au.dk, ◦
mathias.albert@univ-cotedazur.fr, § anna.minguzzi@lpmmc.cnrs.fr, ¶
patrizia.vignolo@univ-cotedazur.fr

Abstract

We propose an alternative to the Bethe Ansatz method for **repulsive** strongly-interacting fermionic (or bosonic) mixtures on a ring. Starting from the knowledge of the solution for single-component non-interacting fermions (or strongly-interacting bosons), we explicitly impose periodic condition on the amplitudes of the spin configurations. This reduces drastically the number of independent complex amplitudes that we determine by constrained diagonalization of an effective Hamiltonian. This procedure allows us to obtain a complete basis for the exact low-energy many-body solutions for mixtures with a large number of particles, both for $SU(\kappa)$ and symmetry-breaking systems.

Copyright attribution to authors.

This work is a submission to SciPost Physics.

License information to appear upon publication.

Publication information to appear upon publication.

Received Date

Accepted Date

Published Date

1

2 Contents

3	1 Introduction	2
4	2 Strongly-interacting quantum gases on a ring	3
5	2.1 Single-component particles on a ring	3
6	2.2 Quantum mixtures on a ring	4
7	2.3 The Bethe Ansatz solution at strong interaction	4
8	3 The necklace Ansatz	5
9	4 Illustration of the method	8
10	4.1 Spectrum and eigenstates of a 2+2 $SU(2)$ bosonic mixture	8
11	4.2 Spectrum and eigenstates of a 4+2 $SU(2)$ fermionic mixture	10
12	4.3 3+3 mixtures: from $SU(2)$ to symmetry breaking mixtures	10
13	4.4 Persistent current in a $SU(3)$ fermionic mixture	12
14	5 Concluding remarks	17

15	A Alternative derivation of the necklace Ansatz	18
16	B Derivation of the effective Hamiltonian	19
17	C Contact matrices	20
18	D Energy correction to order $\mathcal{O}\left(\frac{1}{g}\right)$: Bethe Ansatz derivation	21
19	E Analysis of the strongly interacting limit	24
20	References	24

21
22

23 1 Introduction

24 Exactly solvable quantum many-body systems are rare in physics, and generally exist only
25 in one-dimensional (1D) spatial geometries. For a long time these systems have been seen
26 more as toy models rather than models that can describe real physical set-ups. However,
27 in the last decades they gained also this status thanks to their implementation in cold-atom
28 laboratories that enabled quantum simulations of many-body [1–3] as well as of few-body
29 physics in 1D [4–6].

30 Ultracold gases are extremely rich and versatile. They can be realized with bosonic or
31 fermionic atoms, which can be non-interacting or with tunable interactions up-to very strong
32 repulsive or attractive interactions, and eventually with a spin or color degree of freedom
33 that can be very large [7]. The stronger the interaction strength, the more correlated the
34 atoms are, and the more difficult it is to get numerically an accurate description of the system,
35 especially for long-time dynamics. For these reasons, exact solutions for quantum systems are
36 gaining more and more interest and are becoming essential both for deep understanding of
37 fundamental physics and for benchmarking classical and quantum simulators.

38 Exact solutions for 1D homogeneous systems are well-known in the literature. Celebrated
39 examples are 1D bosons or fermions with contact interactions that are solvable by the Bethe
40 Ansatz both in the case of repulsions [8–12] and attractions [3, 13–18], the latter giving rise
41 to many-body bound states or pairing. In the presence of an inhomogeneous confinement
42 there are generally no exact solutions for interacting systems, except for systems with infinite
43 repulsive interactions such as impenetrable bosons, the Tonks-Girardeau (TG) gas [19], or
44 impenetrable bosonic or fermionic mixtures [2, 4, 5, 20–23]. The key point for this category of
45 exact solutions is that impenetrable particles behave like non-interacting fermions as long as
46 the correct symmetry exchange is taken into account. The mixture many-body wavefunction is
47 thus mapped on that for non-interacting fermions, and the exchange properties are determined
48 by the diagonalization of an effective Hamiltonian related to the contact matrix [21, 23–26].

49 Until now, this method was essentially applied only to mixtures under external confine-
50 ment, such as harmonic or box traps, and not to ring systems, with the sole exception of the
51 ground-state of non-degenerate symmetric mixtures with vanishing momentum [27]. Ring
52 trapping potentials have become available in the experiments with ultracold atoms, and are
53 nowadays realized with unprecedented precision and smoothness (see e.g. Ref. [28] and ref-
54 erences therein). The ring geometry, corresponding to imposing periodic boundary conditions,
55 is the most suitable geometry to study the thermodynamic limit, due to absence of boundaries
56 or inhomogeneities. Also, finite-size rings are important for studying mesoscopic effects, such

57 the response to applied gauge fields [29, 30] as well as for applications to atomtronics [31].

58 The exact solutions for mixtures on a ring are generally provided by Bethe Ansatz [10, 11,
59 13, 32], whose resolution becomes increasingly complex as the number of atoms and compo-
60 nents increases [33–37]. Inspired by the contact matrix method introduced in [21, 23], we
61 propose here an alternative to the Bethe Ansatz, the necklace Ansatz, for calculating the spec-
62 trum and the many-body eigenstates for fermionic or bosonic mixtures in the strongly repulsive
63 limit.

64 The first building block of our procedure is the solution for a single-component quantum
65 gas on a ring. It allows us to write the many-body wavefunction of a fermionic/bosonic mixture
66 for a given sector, i.e., a given order of particles. The second step consists in regrouping the
67 sectors that are equivalent up to a permutation of identical particles in snippets. The number
68 of snippets fixes the number of independent solutions for the spin components. The third
69 step, which is the crucial point of the procedure, is to regroup the snippets that are the same
70 on the ring up to a rotation, i.e. that belong to the same *necklace*. The fact that the many-
71 body wavefunction has to be the same on snippets belonging at the same necklace fixes a
72 phase relation between the snippets' amplitudes. This reduces further the complexity of the
73 problem: one needs to determine a number of complex coefficients that is equal to the number
74 of different necklaces minus one, because of the normalization condition. The last step is to
75 determine these complex amplitudes. This is done by solving a constrained diagonalization of
76 an effective Hamiltonian for each value of the quantized total momentum.

77 The article is organized as following. In Sec. 2 we remind the solution for a spinless Fermi
78 gas and for a single-component TG gas on a ring and we remind the main steps of the Bethe
79 Ansatz in order to find the solution for a fermionic/bosonic mixture in the strongly repulsive
80 limit. The necklace Ansatz is detailed in Sec. 3 and many examples are given in Sec. 4. Our
81 concluding remarks are given in Sec. 5.

82 2 Strongly-interacting quantum gases on a ring

83 2.1 Single-component particles on a ring

84 Let us consider the ground state for N spinless fermions or N TG bosons on a ring of length L ,
85 with coordinates $X = \{x_1, \dots, x_N\}$ [38, 39]:

$$\Psi_{GS}(X) = \begin{cases} \Psi_{SD}(k_\ell^F x_j) \\ \mathcal{A}\Psi_{SD}(k_\ell^B x_j) \end{cases}, \quad (1)$$

86 with $\Psi_{SD}(k_\ell x_j)$ being the Slater determinant built with the ring single-particle orbital solutions
87 $\sim e^{ik_\ell x_j}$ and \mathcal{A} is the symmetrization operator.

88 For the case of N even fermions, $k_\ell^F = \{-\frac{N}{2}\frac{2\pi}{L}, (-\frac{N}{2} + 1)\frac{2\pi}{L}, \dots, 0, \dots, (\frac{N}{2} - 1)\frac{2\pi}{L}\}$; for
89 the case of N odd fermions, $k_\ell^F = \{-\frac{N-1}{2}\frac{2\pi}{L}, \dots, 0, \dots, \frac{N-1}{2}\frac{2\pi}{L}\}$; for the case of N even TG
90 bosons, $k_\ell^B = \{-\frac{N-1}{2}\frac{2\pi}{L}, \dots, -\frac{\pi}{L}, +\frac{\pi}{L}, \dots, \frac{N-1}{2}\frac{2\pi}{L}\}$; and for the case of N odd TG bosons,
91 $k_\ell^B = \{-\frac{N-1}{2}\frac{2\pi}{L}, \dots, 0, \dots, \frac{N-1}{2}\frac{2\pi}{L}\}$.

92 Remark that, for all $\mathcal{K} = \frac{2\pi n}{L}$,

$$\Psi_{\mathcal{K}}(X) = e^{i\mathcal{K}(\sum_j x_j)/N} \Psi_{GS}(X) = \begin{cases} \Psi_{SD}((k_\ell^F + \mathcal{K}/N)x_j) \\ \mathcal{A}\Psi_{SD}((k_\ell^B + \mathcal{K}/N)x_j) \end{cases}, \quad (2)$$

93 is a solution with total momentum

$$\mathcal{P}^{N,n} = \sum_{j=1}^N \hbar \left(\mathbf{k}_j^{F,B} + \frac{\mathcal{K}}{N} \right) = \sum_{j=1}^N \hbar \mathbf{k}_j^{F,B} + \frac{2n\pi\hbar}{L} = \mathcal{P}_{F,B}^{N,n} + \hbar \mathcal{K}, \quad (3)$$

94 and energy

$$E_\infty^{N,n} = \frac{\hbar^2}{2m} \sum_{j=1}^N \left(\mathbf{k}_j^{F,B} + \frac{\mathcal{K}}{N} \right)^2 = \frac{\hbar^2}{2m} \sum_{j=1}^N \left(\mathbf{k}_j^{F,B} + \frac{2n\pi}{LN} \right)^2. \quad (4)$$

95 2.2 Quantum mixtures on a ring

96 Let us now consider a fermionic or a bosonic spin mixture with κ components, obeying the
97 Hamiltonian

$$\hat{H} = \sum_{\sigma=1}^{\kappa} \sum_{i=1}^{N_\sigma} -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial \mathbf{x}_{i,\sigma}^2} + \sum_{\sigma} g_{\sigma\sigma} \sum_{i=1}^{N_\sigma} \sum_{j>i}^{N_\sigma} \delta(\mathbf{x}_{i,\sigma} - \mathbf{x}_{j,\sigma}) + \frac{1}{2} \sum_{\sigma \neq \sigma'=1}^{\kappa} g_{\sigma\sigma'} \sum_{i=1}^{N_\sigma} \sum_{j=1}^{N_{\sigma'}} \delta(\mathbf{x}_{i,\sigma} - \mathbf{x}_{j,\sigma'}), \quad (5)$$

98 where the i, j 's are the particle's indices that go from 1 to N_σ ($N_{\sigma'}$), the σ, σ' 's are the spin
99 indices that go from 1 to κ , the number of spin species, and the $g_{\sigma\sigma'}$'s are the inter- and intra-
100 species interaction strengths. This latter concerns only identical bosons since for identical
101 fermions s -waves contact interactions are not allowed. Here and in the following, we are
102 interested in the limit $g_{\sigma\sigma'} \rightarrow +\infty$, for any σ, σ' . In this strongly interacting regime, the
103 many-body wavefunction vanishes whenever $\mathbf{x}_i = \mathbf{x}_j$.

104 2.3 The Bethe Ansatz solution at strong interaction

105 In this section, we briefly summarize the Bethe Ansatz solution for strongly-interacting quan-
106 tum mixtures on a ring. In $SU(\kappa)$ mixtures, when $g_{\sigma\sigma'} = g$ for any σ, σ' , the systems de-
107 scribed by Hamiltonian (5) can be solved exactly at any interaction strength and for generic κ
108 using Bethe Ansatz [11, 40–44].

109 In the following, we only treat the strongly interacting regime $g \rightarrow \infty$, as it constitutes
110 the main focus of this work. In this regime, in each coordinate sector $\theta_Q(X) = \theta(\mathbf{x}_{Q(1)} < \mathbf{x}_{Q(2)}$
111 $\dots < \mathbf{x}_{Q(N)})$, Q being the permutation operator, the Bethe Ansatz wavefunction reads:

$$\Psi_{BA,Q}(X) = a_Q(\Lambda_1^\sigma, \dots, \Lambda_{N_\sigma}^\sigma) \sum_P (-1)^{(1-\eta_B)|P|} \exp\left\{i \sum_j k_{P(j)} \mathbf{x}_{Q(j)}\right\}, \quad (6)$$

112 where the wavevectors k_j , $j = 1 \dots N$ are the charge rapidities and Λ_m^σ are the spin rapidities
113 rescaled by the interaction constant [45, 46]. Here, $\sigma = 2 \dots \kappa$ labels the spin species and
114 the index m runs from 1 to the number of particles of the σ -th spin species N_σ . The sum
115 is performed over all the possible permutations P in the symmetric group S_N ; $\eta_B = 0, 1$ for
116 fermionic and bosonic mixtures, respectively. The notation $|P|$ indicates the number of trans-
117 positions linking the permutation P with the identical sector defined by $k_1 \leq k_2 \leq \dots \leq k_N$.
118 We remark that, at intermediate or weak interactions, the amplitudes a_Q depend also on the
119 charge rapidities. As a consequence, in the general case, the amplitudes also depend on the
120 permutation P . However, in the strongly interacting limit we consider, such dependence drops
121 out due to the decoupling of spin and charge degrees of freedom, and the amplitudes of the
122 Bethe wavefunction only depend on the permutation Q indicating the coordinate sector.

123 The charge and spin rapidities are specified by a set of charge and a set of spin quantum
124 numbers, respectively $\{\mathcal{L}_j\}$, $j = 1 \dots N$ and $\{\mathcal{J}_m^{(\sigma)}\}$, $m = 1 \dots N_\sigma$, and they are determined

125 by imposing the periodic boundary conditions for the charge and for the spin part of the wave-
126 function [47, 48].

127 In the strongly interacting regime, the charge rapidities can be calculated explicitly using:

$$Lk_j = 2\pi \left(\mathcal{I}_j \pm \frac{1}{N} \sum_{\sigma=2}^{\kappa} \sum_{n=1}^{N_{\sigma}} \mathcal{J}_m^{\sigma} \right), \quad (7)$$

128 where the + sign is for fermions and the − is for bosons. The possible values of the quantum
129 numbers depend on the statistics of the particles. In the next paragraph, we outline the Bethe
130 Ansatz solution for two-component mixtures at infinite interaction strength.

131 **SU(2) mixtures** For the case $\kappa = 2$, the quantum numbers obey the following rules. For
132 bosonic mixtures, $\{\mathcal{I}_j\}$ and $\{\mathcal{J}_m^{(2)}\}$ are integers if N and N_2 have the same parity, and half-
133 integers otherwise [44]. For Fermi gases, the nature of the quantum numbers is more com-
134 plicated. For odd N , both $\{\mathcal{I}_j\}$ and $\{\mathcal{J}_m^{(2)}\}$ are integers or half-integers depending on N_2
135 being even or odd respectively. For even number of particles, $\{\mathcal{I}_j\}$ are integers and $\{\mathcal{J}_m^{(2)}\}$
136 are half-integers for even N_2 , while for N_2 odd the quantum numbers are $\{\mathcal{I}_j\}$ and $\{\mathcal{J}_m^{(2)}\}$ are
137 half-integers and integers respectively.

138 Bethe Ansatz provides an analytical expression for the amplitudes $\mathbf{a}_Q(\Lambda_1^{(2)}, \dots, \Lambda_{N_2}^{(2)})$:

$$\mathbf{a}_Q(\Lambda_1^{(2)}, \dots, \Lambda_{N_2}^{(2)}) \propto (-1)^{|\mathcal{Q}|} \sum_R \prod_{1 \leq m < n \leq N_2} \frac{\Lambda_{R(m)}^{(2)} - \Lambda_{R(n)}^{(2)} - 2i}{\Lambda_{R(m)}^{(2)} - \Lambda_{R(n)}^{(2)}} \prod_{l=1}^{N_2} \left(\frac{\Lambda_{R(l)}^{(2)} - i}{\Lambda_{R(l)}^{(2)} + i} \right)^{y_{Q(l)}}, \quad (8)$$

139 where the integer $y_{Q(l)}$ labels the position of the l -th spin down in the coordinate sector $\theta_Q(X)$
140 and $|\mathcal{Q}|$ indicates the number of transpositions mapping \mathcal{Q} into the identical coordinate sector
141 $\mathbf{x}_1 \leq \mathbf{x}_2 \leq \dots \leq \mathbf{x}_N$. The summation runs over all the possible permutations R of the N_2 spin
142 rapidities.

143 For each set of $\{\mathbf{k}_j\}$ and of spin quantum numbers $\{\mathcal{J}_m^{(2)}\}$, the spin rapidities can be ob-
144 tained by imposing periodic boundary conditions on the amplitudes \mathbf{a}_Q . This yields the N_2
145 *non-linear* coupled Bethe equations:

$$2N \arctan(\Lambda_m^{(2)}) = 2\pi \mathcal{J}_m^{(2)} + \sum_{n=1}^{N_2} 2 \arctan(\Lambda_m^{(2)} - \Lambda_n^{(2)}) \quad m = 1 \dots N_2. \quad (9)$$

146 In this limit, amplitudes (8) coincide, up to a normalization constant, with the ones of the
147 Bethe wavefunction for the isotropic Heisenberg spin chain [49–53]. The Bethe equations (9)
148 coincide with the ones of the Heisenberg model. This implies that the $1/g$ -correction to the
149 energy spectrum of the quantum mixture can be calculated from the Heisenberg spin chain
150 with a suitable definition of an effective exchange coupling [46]. Remarkably, this equivalence
151 holds for a generic value of κ [54].

152 Finding all the solutions to the Bethe equations is a challenging task, already for moderate
153 values of N [33–35]. In order to obtain a complete set of complex roots [55] one has to include
154 exceptional (or "singular") solutions and introduce regularizations of the equations [36, 56, 57].
155 Furthermore, the Bethe equations become significantly more intricate as κ increases [37].

156 3 The necklace Ansatz

157 In this section we will outline an alternative procedure to the Bethe Ansatz at strong interaction
158 for deriving the amplitudes \mathbf{a}_Q that we will call the necklace Ansatz. The word ‘necklace’




Figure 1: Schematic representation of the necklaces, their link to the snippets and the corresponding wavefunction amplitudes for the case of a 2+2 bosonic (fermionic) mixture.

159 appears in combinatorics to describe a string of N coloured beads, which can have up to κ
 160 different colours, assuming that all rotations are equivalent [58].

161 Analogously to the trapped case [21, 23], in the fermionized regime, we can write the
 162 many-body wavefunction for a mixture starting from the single-component many-body wave-
 163 function $\Psi_{\mathcal{K}}(\mathbf{X})$ (see Eq. (2)) on the basis of particle sectors $\theta_{\mathbf{Q}}(\mathbf{X})$ yielding

$$\Psi(\mathbf{X}) = \sum_{\mathbf{Q} \in \mathcal{S}_N} a_{\mathbf{Q}} \theta_{\mathbf{Q}}(\mathbf{X}) \Psi_{\mathcal{K}}(\mathbf{X}). \quad (10)$$

164 The $N!$ sectors can be regrouped in $N_s = N! / (\prod_{\nu=1}^{\kappa} N_{\nu}!)$ snippets, with N_s being the dimen-
 165 sion of the Hilbert space, where each snippet is an ensemble of sectors that are equivalent
 166 under permutation of identical particles [5]. Notice that, for each snippet \mathbf{s} , $a_{\mathbf{Q}} = a_{\mathbf{s}}$ for all
 167 $\mathbf{Q} \in \mathbf{s}$, using the symmetry under exchange of identical particles within the mixture. We can
 168 then rewrite Eq. (10) as

$$\Psi(\mathbf{X}) = \sum_{\mathbf{s}=1}^{N_s} a_{\mathbf{s}} \Psi_{\mathcal{K},\mathbf{s}}(\mathbf{X}). \quad (11)$$

169 where $\Psi_{\mathcal{K},\mathbf{s}}(\mathbf{X}) = \sum_{\mathbf{Q} \in \mathbf{s}} \theta_{\mathbf{Q}}(\mathbf{X}) \Psi_{\mathcal{K}}(\mathbf{X})$ is the wavefunction that includes all coordinate sectors
 170 corresponding to the same snippet. We outline below the procedure for finding the $a_{\mathbf{s}}$ coeffi-
 171 cients for the N_s independent spin-configurations corresponding to the ground- and excited
 172 states of the spin Hamiltonian.

173 In order to build the necklace Ansatz, the key observation is that, because of the periodic
 174 boundary conditions, there are families of equivalent snippets. This implies that the $a_{\mathbf{s}}$ coeffi-
 175 cients are not all independent. Let us clarify this point by making an example on two sectors
 176 for the sake of clarity. We consider the sector $\mathbf{x}_1 < \mathbf{x}_2 < \mathbf{x}_3 < \dots < \mathbf{x}_N$ corresponding to the
 177 identity permutation $\mathbf{Q} = \text{Id}$, and the sector $\mathbf{x}_2 < \mathbf{x}_3 < \dots < \mathbf{x}_N < \mathbf{x}_1$ obtained by the cyclic
 178 permutation $\mathbf{Q}'(j) = j + 1$, where the cyclic condition implies $\mathbf{Q}'(N) = 1$. On a ring these two
 179 sectors correspond – up to a rotation – to the same necklace, hence the many-body wavefunc-
 180 tion has to be the same on these two sectors [49]. This means that, observing that the sector
 181 $\theta_{\mathbf{Q}'}(\mathbf{x})$ can be obtained from $\theta_{\text{Id}}(\mathbf{X})$ by applying the transformation $\mathbf{x}_1 \rightarrow \mathbf{x}_1 + L$, and taking
 182 into account that $\Psi_{\mathcal{K}}(\mathbf{x}_1 + L, \mathbf{x}_2, \dots, \mathbf{x}_N) = e^{i\kappa L/N} \Psi_{\mathcal{K}}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N)$, the amplitudes in the
 183 sector $\theta_{\mathbf{Q}'}(\mathbf{x})$ have to satisfy $a_{\mathbf{Q}'} = e^{-i\kappa L/N} a_{\text{Id}}$.

Using the fact that all the sectors that contribute to a given snippet have the same weight $a_{\mathbf{Q}}$, we can extend the reasoning above to the snippets and regroup them in families when they correspond to the same necklace up-to a rotation (see Fig. 1). The \mathcal{N}_{ℓ} snippets, belonging to




Figure 2: Necklaces for a 3+3 bosonic (fermionic) mixture. There is one high-symmetry necklace with $p_\ell = 2$ and three 6-period necklaces.

the same ℓ -th necklace family, are connected by a q -cycle permutation $\tilde{Q}_q(i) = Q(i + q)$ with $q = 1, \dots, \mathcal{N}_\ell - 1$. Setting c_ℓ the complex amplitude for a given snippet of the ℓ -th necklace, the periodicity of the wavefunction along the ring imposes that in the many-body wavefunction the amplitudes of the snippets obtained by the \tilde{Q}_q permutation coincide with c_ℓ times the phase factor $e^{-i\frac{qKL}{N}}$, such that all the amplitudes of the snippets belonging to the ℓ -th necklace read

$$c_\ell, e^{-i\frac{KL}{N}}c_\ell, e^{-i\frac{2KL}{N}}c_\ell, \dots, e^{-i\frac{(\mathcal{N}_\ell-1)KL}{N}}c_\ell.$$

184 The number of snippets \mathcal{N}_ℓ belonging to the same necklace is generally N , except for high-
 185 symmetry necklaces that have a pattern that repeats itself with a period $p_\ell < N$, in such a case
 186 $\mathcal{N}_\ell = p_\ell$.

187 The condition for the ℓ th rotated necklace to come to its initial position is that $\mathcal{N}_\ell KL/N =$
 188 $2\pi n$, with n a relative integer. This recovers the condition $\mathcal{K} = 2\pi n/L$ for most families where
 189 $\mathcal{N}_\ell = N$, but gives a more restrictive condition for high-symmetry configurations. The allowed
 190 values of the quantum number n depend on the type of necklace, see Sec. 4 for examples.

191 Notice that this ansatz reduces drastically the number of coefficients required to define the
 192 wavefunction, from the number of snippets N_s to the number of independent necklaces N_{neck}
 193 (minus one, because of the normalization condition). The number of necklaces is given by

$$N_{\text{neck}} = (N_s - \sum_{j=1}^M p_j)/N + M, \quad (12)$$

194 M being the number of high-symmetry configurations with period $p_\ell < N$. Notice that our
 195 ansatz provides by construction a complete set of solutions at strong coupling. This can be a
 196 noticeably harder task if the Bethe Ansatz methods are used instead [34,36,55,57,59]. As an il-
 197 lustrative example, let us consider a two-component balanced mixture. For $N = 4$, $N_s = 6$ and
 198 $M = 1$ with period p_1 of length 2 ($\uparrow\downarrow\uparrow\downarrow$). This gives $N_{\text{neck}} = (6-2)/4+1 = 2$. Indeed we have
 199 two families of snippets that correspond to two necklaces (see Fig. 1): $\{\uparrow\uparrow\downarrow\downarrow, \uparrow\downarrow\uparrow\downarrow, \downarrow\downarrow\uparrow\uparrow, \downarrow\uparrow\downarrow\uparrow\}$
 200 and $\{\uparrow\downarrow\uparrow\downarrow, \downarrow\uparrow\downarrow\uparrow\}$. For $N = 6$ we have $N_s = 20$ and again $M = 1$ with period of length $p_1 = 2$
 201 ($\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow$). This gives $N_{\text{neck}} = (6-2)/6+1 = 4$, see Fig. 2. For the case $N = 8$, $N_s = 70$. There
 202 are 2 possible periodic configurations with period smaller than N : one with period of length
 203 $p_1 = 2$, and the other with period of length $p_2 = 4$. This gives $N_{\text{neck}} = (70-2-4)/8+2 = 10$.

204 The coefficients in the wavefunction (10) are determined by solving the Schrödinger equa-
 205 tion in the limit of large repulsive interactions. In this regime, spin and orbital part decouple
 206 and we are left to solve the eigenvalue problem for the contact matrix V whose form depends

207 on the type of mixture under consideration (see Appendices B and C for details and examples).
 208 In the fermionic $SU(2)$ case the contact matrix in the snippets' basis reads

$$[V^{SU^F}]_{i,j} = \frac{\hbar^4}{m^2} \begin{cases} \sum_{\ell_i} \delta_d \alpha_{\ell_i} & j = i \\ -v_{i,j} & j \neq i \end{cases}, \quad (13)$$

209 while in the bosonic $SU(2)$ case we have

$$[V^{SU^B}]_{i,j} = \frac{\hbar^4}{m^2} \begin{cases} \sum_{\ell_i} (\delta_d + 2\delta_b) \alpha_{\ell_i} & j = i \\ v_{i,j} & j \neq i \end{cases}, \quad (14)$$

210 where δ_d is equal to one if ℓ_i and $\ell_i + 1$ correspond to distinguishable particles (two different
 211 spins) and zero otherwise, while δ_b is equal to one only for next-neighbor identical bosons.
 212 We indicate α_{ℓ_i} as the nearest-neighbor exchange constant, given by

$$\alpha_{\ell_i} = N! \int \mathbf{d}\mathbf{x}_1, \dots, \mathbf{d}\mathbf{x}_N \theta_{\text{Id}}(\mathbf{x}_1, \dots, \mathbf{x}_N) \delta(\mathbf{x}_{\ell_i} - \mathbf{x}_{\ell_i+1}) \left[\frac{\partial \Psi_{\mathcal{K}}}{\partial \mathbf{x}_{\ell_i}} \right]^2. \quad (15)$$

213 For the off-diagonal terms we have $v_{i,j} = \alpha_{\ell_i}$ if the snippets i and j differ from the exchange
 214 of two nearest-neighbor particles with different spins, set at positions ℓ_i ($\ell_j + 1$) and $\ell_i + 1$
 215 (ℓ_j), and zero otherwise.

216 In the homogeneous system, translation symmetry implies that all α_{ℓ_i} 's are equal and
 217 depend solely on the number of particles. From now on we will set $\alpha_{\ell_i} = \alpha^{(N)}$. We notice that
 218 the matrix $-V/g$ corresponds to the Hamiltonian of a Heisenberg spin chain with a hopping
 219 amplitude $\alpha^{(N)}/g$, that depends on the number of particles and interaction strength g [21,
 220 23–25, 27, 60].

221 4 Illustration of the method

222 In the next subsections, we will illustrate our method using a few representative cases. We
 223 show that the necklace Ansatz provides the same solutions as the Bethe Ansatz in the strongly
 224 interacting limit (see also results in Appendix D, whose range of validity is discussed in Ap-
 225 pendix E). The illustrative examples below herald analysis of more complicated problems
 226 where the direct solution to the Bethe ansatz equations is very difficult to access.

227 4.1 Spectrum and eigenstates of a 2+2 $SU(2)$ bosonic mixture

228 Let us consider the case of a bosonic $SU(2)$ mixture with $N_{\uparrow} = 2$ $N_{\downarrow} = 2$ on a ring of length
 229 L . For such a system, the number of snippets is $N_s = 6$ and we have two necklaces, i.e. two
 230 families of snippets. To the snippets of the first necklace, given by $\{\uparrow\uparrow\downarrow\downarrow, \uparrow\downarrow\downarrow\uparrow, \downarrow\downarrow\uparrow\uparrow, \downarrow\uparrow\uparrow\downarrow\}$,
 231 we assign the coefficients $\{c_1, c_1 e^{-in\pi/2}, c_1 e^{-in\pi}, c_1 e^{-i3n\pi/2}\}$, while to the second necklace
 232 $\{\uparrow\downarrow\uparrow\downarrow, \downarrow\uparrow\downarrow\uparrow\}$ we assign the coefficients $\{c_2, c_2 e^{-in\pi/2}\}$, see Fig. 1. Remark that we have set
 233 $\mathcal{K}/N = 2n\pi/4$, n being a relative integer, and that solutions with c_2 different from zero must
 234 have even values of n ($c_2 e^{-in\pi} = c_2$). Thus, we expect two spin states if n is even and only
 235 1 when n is odd and that Néel spin-configurations, namely configurations with alternating
 236 spin-up and spin-down, are forbidden in this case.

237 In order to find \mathbf{c}_1 and \mathbf{c}_2 as functions of \mathbf{n} , we solve the conditioned eigenvalue problem

$$\alpha^{(4)} \begin{pmatrix} 6 & 0 & 0 & 0 & 1 & 1 \\ 0 & 6 & 0 & 0 & 1 & 1 \\ 0 & 0 & 6 & 0 & 1 & 1 \\ 0 & 0 & 0 & 6 & 1 & 1 \\ 1 & 1 & 1 & 1 & 4 & 0 \\ 1 & 1 & 1 & 1 & 0 & 4 \end{pmatrix} \begin{pmatrix} c_1 \\ c_1 e^{-in\pi/2} \\ c_1 e^{-in\pi} \\ c_1 e^{-i3n\pi/2} \\ c_2 \\ c_2 e^{-in\pi/2} \end{pmatrix} = \xi_n \begin{pmatrix} c_1 \\ c_1 e^{-in\pi/2} \\ c_1 e^{-in\pi} \\ c_1 e^{-i3n\pi/2} \\ c_2 \\ c_2 e^{-in\pi/2} \end{pmatrix}. \quad (16)$$

238 Setting $\tilde{\xi}_n = \xi_n / \alpha^{(4)}$, we obtain the following system of equations

$$\begin{cases} (6 - \tilde{\xi}_n) c_1 & + (1 + e^{-in\pi/2}) c_2 & = 0 \\ (6 - \tilde{\xi}_n) c_1 e^{-in\pi/2} & + (1 + e^{-in\pi/2}) c_2 & = 0 \\ (6 - \tilde{\xi}_n) c_1 e^{-in\pi} & + (1 + e^{-in\pi/2}) c_2 & = 0 \\ (6 - \tilde{\xi}_n) c_1 e^{-i3n\pi/2} & + (1 + e^{-in\pi/2}) c_2 & = 0 \\ (1 + e^{-in\pi/2} + e^{-in\pi} + e^{-i3n\pi/2}) c_1 & + (4 - \tilde{\xi}_n) c_2 & = 0 \\ (1 + e^{-in\pi/2} + e^{-in\pi} + e^{-i3n\pi/2}) c_1 & + (4 - \tilde{\xi}_n) c_2 e^{-in\pi/2} & = 0 \end{cases}, \quad (17)$$

239 that has six solutions summarized in Table 1. Note that there are two families of equations
 240 in Eq. (17): the first 4 differ only by the phase factor of the first member, and the second 2
 241 by the phase factor of the second member. As Eq. (17) is overdetermined, we conclude that
 242 for $\mathbf{n} \neq \mathbf{0}$, the solution has either $\mathbf{c}_1 = \mathbf{0}$ or $\mathbf{c}_2 = \mathbf{0}$. Only if $\mathbf{n} = \mathbf{0}$ (modulo 4), we can have
 243 a situation when both coefficients are non-vanishing. We notice that the solution of Eq.(17)
 244 for $\mathbf{c}_1 = \mathbf{0}$ shows that \mathbf{c}_2 is different from zero only if $(1 + e^{-in\pi/2}) = 0$, implying that \mathbf{n} is
 245 even, as was anticipated above. Similarly, the solution of Eq. (17) for $\mathbf{c}_2 = \mathbf{0}$ yields as possible
 246 values $\mathbf{n} = 1, 2, 3$ (modulo 4), showing that the allowed values of \mathcal{K} are all the multiples of
 247 $2\pi\hbar/L$. This is at the origin of the fractionalization of the period of persistent currents, see
 248 Sec. 4.4. It follows from the above structure of the solution that the \mathbf{c}_j 's solution with a given
 249 \mathbf{n} is a solution also for $\mathbf{n}' = \mathbf{n} + 4$.

250 One could think that there are too many equations for determining \mathbf{c}_1 and \mathbf{c}_2 , but the
 251 imposition of this sort of gauge invariance condition for the two coefficients is necessary in
 252 order to find and select all (and only) the physical solutions. Indeed if one tries to get \mathbf{c}_1 and
 253 \mathbf{c}_2 directly from the minimization of the energy one obtains the equations

$$\begin{cases} (12 - 4\tilde{\xi}_n) c_1 & + (1 + 2 \cos(n\pi/2) + \cos(n\pi)) c_2 & = 0 \\ (1 + 2 \cos(n\pi/2) + \cos(n\pi)) c_1 & + (4 - 2\tilde{\xi}_n) c_2 & = 0 \end{cases}, \quad (18)$$

254 that allows solutions that are not physical, such as $\mathbf{c}_2 \neq \mathbf{0}$ for $\mathbf{n} = 1$.

255 To ascertain the validity of these solutions, one can verify that they have the correct sym-
 256 metry. This could be done using the natural representations of $SU(\kappa)$ (or equivalently S_N),
 257 the Young diagrams. These diagrams are collections of boxes that schematically represent the
 258 particle-exchange symmetry of a physical state. Precisely, boxes in line (resp. column) refer to
 259 symmetric (resp. anti-symmetric) exchanges so that line diagram $\square\square$ corresponds to a three-
 260 particles fully symmetric state and a column $\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}$ to a three-particles fully anti-symmetric state.

261 All other configurations represent more exotic states with mixed symmetries, each correspond-
 262 ing to a different representation of S_N . The usual procedure to connect our physical states to
 263 these diagrams is through the use of class sum operators, verifying that they are eigenstates of
 264 the 2-cycle class-sum operator $\Gamma^{(2)} = \frac{1}{2} \sum_{i < j} P_{i,j}$, [61,62] where $P_{i,j}$ is the operator which per-
 265 mutes the i -th and j -th elements. $\Gamma^{(2)}$'s eigenvalues are directly connected to the irreducible
 266 representations of $SU(\kappa)$, and thus to the Young diagrams. Indeed the relation between the

n	$\tilde{\xi}_n$	c_1	c_2	$\gamma^{(2)}$	YD
-1	6	1/2	0	2	$\square\square$
0	8	$1/\sqrt{6}$	$1/\sqrt{6}$	6	$\square\square\square$
	2	$1/(2\sqrt{3})$	$-1/\sqrt{3}$	0	\square
1	6	1/2	0	2	$\square\square$
2	6	1/2	0	0	\square
	4	0	$1/\sqrt{2}$	2	$\square\square$

Table 1: Solution for the 2+2 SU(2) bosonic mixture. $\tilde{\xi}_n$ are the rescaled eigenvalues $\xi_n/\alpha^{(4)}$. The last column indicates the symmetry of the solution with the associated Young diagram (YD).

267 eigenvalues $\gamma^{(2)}$'s and a Young diagrams with a number of boxes μ_i at row i is [63]

$$\gamma^{(2)} = \frac{1}{2} \sum_i [\mu_i(\mu_i - 2i + 1)]. \quad (19)$$

268 For $n = 0$ the total momentum \mathcal{P} is zero and we find (i) the fully symmetric state $\square\square\square$ with
 269 $c_1 = c_2 = 1/\sqrt{6}$ with eigenvalue $\xi_0/\alpha^{(4)} = 8$, and (ii) the solution $c_1 = 1/(2\sqrt{3})$, $c_2 = -1/\sqrt{3}$
 270 with $\xi_0/\alpha^{(4)} = 2$ that corresponds to the symmetry represented by the Young diagram \square . For
 271 $n = \pm 1$ ($\mathcal{P} = \pm 2\pi\hbar/L$), we find the solutions (iii) and (iv) with amplitudes $c_1 = 1/\sqrt{2}$, $c_2 = 0$
 272 and $\xi_{\pm 1}/\alpha^{(4)} = 6$. Both have symmetry $\square\square$. For $n = 2$, we find two solutions: (v) one with
 273 $c_1 = 1/2$, $c_2 = 0$ and $\xi_2/\alpha^{(4)} = 6$ ($\square\square$), and the last one (vi) with $c_1 = 0$, $c_2 = 1/\sqrt{2}$ and
 274 $\xi_2/\alpha^{(4)} = 4$ ($\square\square$). The values for the spin-states coefficients are summarized in Table 1. These
 275 states constitute a complete and orthogonal basis for the spin configurations. One can readily
 276 check that the obtained values for the c_j yield the same wavefunctions as those obtained from
 277 the Bethe Ansatz solution in the strongly interacting limit (see e.g. [49, 64]).

278 4.2 Spectrum and eigenstates of a 4+2 SU(2) fermionic mixture

279 Let us now consider a 4+2 SU(2) fermionic mixture. For such a system $N_s = 6!/(4!2!) = 15$
 280 and $N_{\text{neck}} = (15 - 3)/6 + 1 = 3$. To the snippets of the first necklace $\{\uparrow\uparrow\uparrow\uparrow\downarrow, \uparrow\uparrow\downarrow\downarrow\uparrow,$
 281 $\uparrow\uparrow\downarrow\uparrow\uparrow, \uparrow\downarrow\uparrow\uparrow\uparrow, \downarrow\uparrow\uparrow\uparrow\uparrow, \downarrow\uparrow\uparrow\uparrow\downarrow\}$ we assign the coefficients $\{c_1, c_1e^{-in\pi/3}, c_1e^{-i2n\pi/3}, c_1e^{-in\pi},$
 282 $c_1e^{-i4n\pi/3}, c_1e^{-i5n\pi/3}\}$, to the second $\{\uparrow\uparrow\uparrow\downarrow\downarrow, \uparrow\uparrow\downarrow\uparrow\uparrow, \uparrow\downarrow\uparrow\uparrow\uparrow, \downarrow\downarrow\uparrow\uparrow\uparrow, \uparrow\downarrow\uparrow\uparrow\downarrow, \downarrow\uparrow\uparrow\downarrow\uparrow\}$, we
 283 assign the coefficients $\{c_2, c_2e^{-in\pi/3}, c_2e^{-i2n\pi/3}, c_2e^{-in\pi}, c_2e^{-i4n\pi/3}, c_2e^{-i5n\pi/3}\}$, and to the
 284 third $\{\uparrow\uparrow\downarrow\uparrow\uparrow\downarrow, \uparrow\downarrow\uparrow\uparrow\downarrow, \downarrow\uparrow\uparrow\downarrow\uparrow\}$ we assign the coefficients $\{c_3, c_3e^{-in\pi/3}, c_3e^{-i2n\pi/3}\}$. Remark
 285 that we have set $\mathcal{K}/N = 2n\pi/6$, n being a relative integer, and that solutions with c_3 different
 286 from zero must have even values of n ($c_3e^{-in\pi} = c_3$). Thus, we expect 3 possible spin con-
 287 figurations if n is even and only 2 if n is odd. Indeed, by solving the conditioned eigenvalue
 288 system (40), we find that every even value of n allows for 3 solutions and every odd n allows
 289 for 2 solutions. Therefore, there are 15 independent solutions for n from $n = -2$ to $n = 3$,
 290 that are given in Table 2.

291 4.3 3+3 mixtures: from SU(2) to symmetry breaking mixtures

292 The necklace Ansatz can also be used to analyze systems with $g_{\sigma\sigma'} \neq g_{\sigma\sigma}$ in Eq. (5) as long
 293 as the system is strongly interacting. In this subsection, we consider arguably the simplest
 294 scenario. The generalization to other cases is straightforward just as in the trapped cases [24,
 295 60] or for the problem in a ring with vanishing total momentum [27].

n	$\tilde{\xi}_n$	c_1	c_2	c_3	$\gamma^{(2)}$	YD
-2	3	$e^{2i\pi/3}$	1	$2e^{i\pi/3}$	-9	
	$(7 - \sqrt{17})/2$	$-(3 + \sqrt{17})/4 e^{2i\pi/3}$	1	$(\sqrt{17} - 1)/4 e^{i\pi/3}$	-5	
	$(7 + \sqrt{17})/2$	$(-3 + \sqrt{17})/4 e^{2i\pi/3}$	1	$(\sqrt{17} + 1)/4 e^{-i2\pi/3}$	-5	
-1	1	$\sqrt{3}e^{-i\pi/6}$	1	0	-9	
	5	$e^{i5\pi/6}/\sqrt{3}$	1	0	-5	
0	0	1	1	1	-15	
	$5 - \sqrt{5}$	$-(1 + \sqrt{5})/4$	$(-1 + \sqrt{5})/4$	1	-5	
	$5 + \sqrt{5}$	$(-1 + \sqrt{5})/4$	$-(1 + \sqrt{5})/4$	1	-5	
1	1	$\sqrt{3}e^{i\pi/6}$	1	0	-9	
	5	$e^{-i5\pi/6}/\sqrt{3}$	1	0	-5	
2	3	$e^{-2i\pi/3}$	1	$2e^{-i\pi/3}$	-9	
	$(7 - \sqrt{17})/2$	$-(3 + \sqrt{17})/4 e^{-2i\pi/3}$	1	$(\sqrt{17} - 1)/4 e^{-i\pi/3}$	-5	
	$(7 + \sqrt{17})/2$	$(-3 + \sqrt{17})/4 e^{-2i\pi/3}$	1	$(\sqrt{17} + 1)/4 e^{i2\pi/3}$	-5	
3	2	1	0	0	-5	
	4	0	1	0	-9	

Table 2: Solution for the 4+2 SU(2) fermionic mixture. $\tilde{\xi}_n$ are the rescaled eigenvalues $\xi_n/\alpha^{(6)}$. The coefficients c_j are not normalized. The last column indicates the symmetry of the solution with the associated Young diagram (YD).

296 Here we will consider a $3 + 3$ mixture and we will compare the cases of a $SU(2)$ fermionic
 297 mixture and of a $SU(2)$ bosonic mixture $\mathbf{g}_{\uparrow\uparrow} = \mathbf{g}_{\downarrow\downarrow} = \mathbf{g}_{\uparrow\downarrow}$, with a symmetry breaking (SB) case of
 298 a bosonic mixture where the $SU(2)$ symmetry is explicitly broken ($1/\mathbf{g}_{\uparrow\uparrow} = 1/\mathbf{g}_{\downarrow\downarrow} = \mathbf{0}$ and $\mathbf{g}_{\uparrow\downarrow}$
 299 is large but finite) [27]. For a $3 + 3$ mixture there are 20 snippets and $N_{\text{neck}} = 4$ independent
 300 necklaces, one of which has a period of length 2. We assign to the snippets of the first necklace
 301 $\{\uparrow\uparrow\uparrow\downarrow\downarrow, \uparrow\uparrow\downarrow\downarrow\uparrow, \uparrow\downarrow\downarrow\uparrow\uparrow, \downarrow\downarrow\downarrow\uparrow\uparrow, \downarrow\uparrow\uparrow\uparrow\downarrow, \downarrow\uparrow\uparrow\downarrow\downarrow\}$ the amplitudes $\{c_1, c_1 e^{-i\phi}, c_1 e^{-2i\phi}, c_1 e^{-3i\phi},$
 302 $c_1 e^{-4i\phi}, c_1 e^{-5i\phi}\}$; to those of the second necklace $\{\uparrow\uparrow\downarrow\uparrow\downarrow, \uparrow\uparrow\downarrow\downarrow\uparrow, \uparrow\downarrow\downarrow\uparrow\uparrow, \uparrow\downarrow\uparrow\uparrow\downarrow, \downarrow\downarrow\uparrow\uparrow\downarrow,$
 303 $\downarrow\uparrow\uparrow\downarrow\downarrow\}$ the amplitudes $\{c_2, c_2 e^{-i\phi}, c_2 e^{-2i\phi}, c_2 e^{-3i\phi}, c_2 e^{-4i\phi}, c_2 e^{-5i\phi}\}$; to those of the third
 304 necklace $\{\uparrow\downarrow\uparrow\uparrow\downarrow, \uparrow\downarrow\downarrow\uparrow\uparrow, \uparrow\uparrow\downarrow\uparrow\downarrow, \uparrow\downarrow\uparrow\uparrow\downarrow, \downarrow\downarrow\uparrow\uparrow\downarrow, \downarrow\uparrow\uparrow\downarrow\downarrow\}$ the amplitudes $\{c_3, c_3 e^{-i\phi}, c_3 e^{-2i\phi},$
 305 $c_3 e^{-3i\phi}, c_3 e^{-4i\phi}, c_3 e^{-5i\phi}\}$; and finally to the necklace of period 2 $\{\uparrow\downarrow\uparrow\uparrow\downarrow, \uparrow\downarrow\uparrow\downarrow\uparrow\}$, $\{c_4, c_4 e^{-i\phi}\}$
 306 where $\phi = 2n\pi/6$ with n being a relative integer. From the condition that $c_4 e^{-2i\phi} = c_4$, we
 307 expect that c_4 has to be zero if n is not zero or a multiple of 3. So we expect 4 solutions if
 308 $n = 0$ and if $n = 3$, and 3 solutions for the cases $n = -2, -1, 1$ and 2 , namely 20 independent
 309 solutions.

310 The contact V matrices corresponding to the three different cases are given in Appendix
 311 C. The resulting spin states are presented in Table 3 for the $SU(2)$ fermionic mixture, in Table
 312 4 for the $SU(2)$ bosonic mixture, and in Table 5 for the SB bosons. Remark that the SB states
 313 for a given n are very similar to the $SU(2)$ fermionic ones for $n \neq 3$. This is due to the fact
 314 that this SB bosonic mixture can be seen as a $SU(2)$ fermionic mixture up-to a symmetrization
 315 operation of the particles in each component [27].

316 4.4 Persistent current in a $SU(3)$ fermionic mixture

317 We will consider here the case of three $SU(3)$ fermions rotating in a ring of length L with
 318 rotation frequency Ω [54, 64]. The Hamiltonian of the system reads

$$\mathcal{H} = \sum_{j=1}^3 \frac{1}{2m} \left(p_j - \frac{m\Omega L}{2\pi} \right)^2 + g \sum_{j<\ell} \delta(x_j - x_\ell). \quad (20)$$

319 The effect of the rotation is to produce an artificial gauge field yielding an effective flux
 320 $\Phi = \Omega L^2 / (2\pi)$ [28–31], which in turn induces a persistent current of particles in the ring.
 321 Such currents can be used for characterizing the different phases of the system [65–69]. In
 322 analogy with superconducting rings, the ground state energy and the current are periodic
 323 functions of the gauge flux, with a period that is defined as the quantum of flux of the parti-
 324 cles [70].

325 Strong repulsive [54, 64] and attractive [18, 71, 72] interactions induce a fractionalization
 326 of the period of the persistent current. In the attractive case, this phenomenon is related to the
 327 formation of two-body and many-body bound states, respectively for fermionic and bosonic
 328 mixtures. The period is reduced by a factor equal to the number of particles forming the bound
 329 state. In the repulsive case, the period of the persistent current is reduced, both for bosons
 330 and fermions, by a factor equal to the number of particles in the mixture. In this interaction
 331 regime, the fractionalization is due to the formation of spin excitations in the ground state of
 332 the gas, occurring as one applies the gauge flux [45, 46, 54].

333 Here, we use the necklace Ansatz to compute the energy levels of the $SU(3)$ fermionic mix-
 334 ture, at infinite and at large but finite repulsive interactions, as a function of an effective gauge
 335 flux Φ . In this case the snippets can be divided into two necklaces, one $\{\bullet \circ \triangleright, \circ \triangleright \bullet, \triangleright \bullet \circ\}$ with
 336 the amplitudes $\{c_1, c_1 e^{-i2\pi n/3}, c_1 e^{-i4\pi n/3}\}$, and another $\{\circ \bullet \triangleright, \bullet \triangleright \circ, \triangleright \circ \bullet\}$ with the amplitudes
 337 $\{c_2, c_2 e^{-i2\pi n/3}, c_2 e^{-i4\pi n/3}\}$. The computed spin states and the corresponding eigenvalues are
 338 given in Table 6.

339 The effect of the rotation enters in the energy landscape. Indeed the energy of the spin

n	$\tilde{\xi}_n$	c_1	c_2	c_3	c_4	$\gamma^{(2)}$	YD
-2	$\frac{1}{2}(7 - \sqrt{17})$	$\frac{e^{2i\pi/3}}{2}(3 + \sqrt{17})$	$e^{2i\pi/3}$	1	0	-5	
	$\frac{1}{2}(7 + \sqrt{17})$	$\frac{e^{-i\pi/3}}{2}(-3 + \sqrt{17})$	$e^{2i\pi/3}$	1	0	-5	
	3	0	$e^{-i\pi/3}$	1	0	-9	
-1	1	$2e^{i\pi/3}$	$e^{i\pi/3}$	1	0	-9	
	4	$e^{i2\pi/3}$	$e^{i\pi/3}$	1	0	-3	
	5	0	$e^{-i2\pi/3}$	1	0	-5	
0	0	1	1	1	1	-15	
	6	0	1	-1	0	-3	
	$5 - \sqrt{5}$	$-\frac{1}{2}(3 + \sqrt{5})$	1	1	$\frac{3}{2}(-1 + \sqrt{5})$	-5	
	$5 + \sqrt{5}$	$\frac{1}{2}(-3 + \sqrt{5})$	1	1	$-\frac{3}{2}(1 + \sqrt{5})$	-5	
1	1	$2e^{-i\pi/3}$	$e^{-i\pi/3}$	1	0	-9	
	4	$e^{-i2\pi/3}$	$e^{-i\pi/3}$	1	0	-3	
	5	0	$e^{i2\pi/3}$	1	0	-5	
2	$\frac{1}{2}(7 - \sqrt{17})$	$\frac{e^{-2i\pi/3}}{2}(3 + \sqrt{17})$	$-2i\pi/3$	1	0	-5	
	$\frac{1}{2}(7 + \sqrt{17})$	$\frac{e^{i\pi/3}}{2}(-3 + \sqrt{17})$	$-2i\pi/3$	1	0	-5	
	3	0	$e^{i\pi/3}$	1	0	-9	
3	2	0	1	1	0	-5	
	4	-1	1	-1	-3	-9	
	$5 - \sqrt{13}$	$\frac{1}{2}(3 + \sqrt{13})$	1	-1	$\frac{1}{2}(1 - \sqrt{13})$	-3	
	$5 + \sqrt{13}$	$\frac{1}{2}(3 - \sqrt{13})$	1	-1	$\frac{1}{2}(1 + \sqrt{13})$	-3	

Table 3: Solution for the 3+3 SU(2) fermionic mixture. $\tilde{\xi}_n$ are the rescaled eigenvalues $\xi_n/\alpha^{(6)}$. The coefficients c_j are not normalized. The last column indicates the symmetry of the solution with the associated Young diagram (YD).

n	$\tilde{\xi}_n$	c_1	c_2	c_3	c_4	$\gamma^{(2)}$	YD
-2	$\frac{1}{2}(17 - \sqrt{17})$	$\frac{e^{-i\pi/3}}{2}(-3 + \sqrt{17})$	$e^{2i\pi/3}$	1	0	5	
	$\frac{1}{2}(17 + \sqrt{17})$	$\frac{e^{i\pi/3}}{2}(3 + \sqrt{17})$	$e^{2i\pi/3}$	1	0	5	
	9	0	$e^{-i\pi/3}$	1	0	9	
-1	11	$2e^{i\pi/3}$	$e^{i\pi/3}$	1	0	9	
	8	$e^{-i2\pi/3}$	$e^{i\pi/3}$	1	0	3	
	7	0	$-e^{i\pi/3}$	1	0	5	
0	12	1	1	1	1	15	
	6	0	1	-1	0	3	
	$7 - \sqrt{5}$	$\frac{1}{2}(-3 + \sqrt{5})$	1	1	$-\frac{3}{2}(1 + \sqrt{5})$	5	
	$7 + \sqrt{5}$	$-\frac{1}{2}(3 + \sqrt{5})$	1	1	$\frac{3}{2}(1 - \sqrt{5})$	5	
1	11	$2e^{-i\pi/3}$	$e^{-i\pi/3}$	1	0	9	
	8	$e^{i2\pi/3}$	$e^{-i\pi/3}$	1	0	3	
	7	0	$-e^{-i\pi/3}$	1	0	5	
2	$\frac{1}{2}(17 - \sqrt{17})$	$\frac{e^{i\pi/3}}{2}(-3 + \sqrt{17})$	$e^{-2i\pi/3}$	1	0	5	
	$\frac{1}{2}(17 + \sqrt{17})$	$\frac{e^{-i\pi/3}}{2}(3 + \sqrt{17})$	$e^{-2i\pi/3}$	1	0	5	
	9	0	$e^{i\pi/3}$	1	0	9	
3	10	0	1	1	0	5	
	8	-1	1	-1	-3	9	
	$7 - \sqrt{13}$	$\frac{1}{2}(3 - \sqrt{13})$	1	-1	$\frac{1}{2}(1 + \sqrt{13})$	3	
	$7 + \sqrt{13}$	$\frac{1}{2}(3 + \sqrt{13})$	1	-1	$\frac{1}{2}(1 - \sqrt{13})$	3	

Table 4: Solution for the 3+3 SU(2) bosonic mixture. $\tilde{\xi}_n$ are the rescaled eigenvalues $\xi_n/\alpha^{(6)}$. The coefficients c_j are not normalized. The last column indicates the symmetry of the solution with the associated Young diagram (YD).

n	$\tilde{\xi}_n$	c_1	c_2	c_3	c_4	$\langle \Gamma^{(2)} \rangle$
-2	1	$2e^{-i\pi/3}$	$e^{i2\pi/3}$	1	0	5
	4	$e^{i2\pi/3}$	$e^{i2\pi/3}$	1	0	5
	5	0	$e^{-i2\pi/3}$	1	0	8
-1	$\frac{1}{2}(7 - \sqrt{17})$	$\frac{e^{-2i\pi/3}}{2}(3 + \sqrt{17})$	$e^{i\pi/3}$	1	0	4.78
	$\frac{1}{2}(7 + \sqrt{17})$	$\frac{e^{i\pi/3}}{2}(-3 + \sqrt{17})$	$e^{i\pi/3}$	1	0	7.21
	3	0	$e^{-i2\pi/3}$	1	0	5
0	2	0	1	-1	0	3
	4	1	1	1	-3	7
	$5 - \sqrt{13}$	$-\frac{1}{2}(3 + \sqrt{13})$	1	1	$\frac{1}{2}(1 - \sqrt{13})$	5.67
	$5 + \sqrt{13}$	$\frac{1}{2}(-3 + \sqrt{13})$	1	1	$\frac{1}{2}(1 + \sqrt{13})$	9.72
1	$\frac{1}{2}(7 - \sqrt{17})$	$\frac{e^{2i\pi/3}}{2}(3 + \sqrt{17})$	$e^{-i\pi/3}$	1	0	4.78
	$\frac{1}{2}(7 + \sqrt{17})$	$\frac{e^{-i\pi/3}}{2}(-3 + \sqrt{17})$	$e^{-i\pi/3}$	1	0	7.21
	3	0	$e^{i2\pi/3}$	1	0	5
2	1	$2e^{i\pi/3}$	$e^{-i2\pi/3}$	1	0	5
	4	$e^{-i2\pi/3}$	$e^{-i2\pi/3}$	1	0	5
	5	0	$e^{i2\pi/3}$	1	0	8
3	0	-1	1	-1	1	4.2
	6	0	1	1	0	5
	$5 - \sqrt{5}$	$\frac{1}{2}(3 + \sqrt{5})$	1	-1	$\frac{3}{2}(-1 + \sqrt{5})$	3.611
	$5 + \sqrt{5}$	$-\frac{1}{2}(-3 + \sqrt{5})$	1	-1	$-\frac{3}{2}(1 + \sqrt{5})$	7.18

Table 5: Solution for the 3+3 symmetry breaking bosonic mixture. $\tilde{\xi}_n$ are the rescaled eigenvalues $\xi_n/\alpha^{(6)}$. The coefficients c_j are not normalized. The last column indicates the expectation value of the 2-cycle class-sum operator $\Gamma^{(2)}$

n	$\tilde{\xi}_n$	c_1	c_2	$\gamma^{(2)}$	YD
-1	3	$1/\sqrt{2}$	0	0	$\begin{array}{ c } \hline \square \\ \hline \end{array}$
	3	0	$1/\sqrt{2}$	0	$\begin{array}{ c } \hline \square \\ \hline \end{array}$
0	6	1/2	-1/2	3	$\begin{array}{ c c } \hline \square & \square \\ \hline \end{array}$
	0	1/2	1/2	-3	$\begin{array}{ c } \hline \square \\ \hline \end{array}$
1	3	$1/\sqrt{2}$	0	0	$\begin{array}{ c } \hline \square \\ \hline \end{array}$
	3	0	$1/\sqrt{2}$	0	$\begin{array}{ c } \hline \square \\ \hline \end{array}$

Table 6: Solution for 3 SU(3) particles. $\tilde{\xi}_n$ are the rescaled eigenvalues $\xi_n/\tilde{\alpha}^{(3)}$. The last column indicates the symmetry of the solution with the associated Young diagram (YD).




Figure 3: Energy landscape $\tilde{E}_j = E_j mL^2/(\hbar^2 \pi^2)$ for a SU(3) three-fermions system, as a function of $\tilde{\Phi}$ at infinite interactions (thin lines) and for $g = 100mL/\hbar^2$ (thick lines). From left to right the different parabolas correspond to $n_j = -1$ (violet curves), 0 (green curves), 1 (blue curves), 2 (orange curves) and 3 (yellow curves). The black dotted horizontal line corresponds to the ground state energy of the three particles at $\tilde{\Phi} = 0$.

340 state η with the eigenvalue $\tilde{\xi}_{n_\eta}$ and the quantum number n_η

$$E_\eta = \varepsilon^{N, n_\eta} - \frac{\tilde{\xi}_{n_\eta}}{g} \alpha^{(N)}, \quad (21)$$

341 with

$$\varepsilon^{N, n_\eta} = \frac{\hbar^2}{2m} \sum_{j=1}^N \left(k_j^{F,B} + \frac{2n_\eta \pi}{LN} - \frac{2\pi}{L} \tilde{\Phi} \right)^2, \quad (22)$$

342 where $\tilde{\Phi} = \Phi/\Phi_0$, $\Phi_0 = h/m$.

343 We plot in Fig. 3 the energy landscape for the cases $g \rightarrow \infty$ (thin lines) and $g = 100mL/\hbar^2$.
 344 At infinite interactions, we observe the expected $1/N$ fractionalization of the periodicity of the
 345 energy landscape [45, 46, 54, 64]. At finite interactions, the parabola branches that display the
 346 most symmetric spin configurations and are centered in $\tilde{\Phi} = 0, 1, \dots$ have the largest energy
 347 corrections and thus correspond to the lowest energy (see Fig. 3), while the intermediate
 348 parabolas have a higher energy, as expected when fractionalization is not yet achieved. This
 349 result displays the same features as those predicted in [46, 54] for the case of mixtures on a
 350 ring lattice: at increasing interactions, the parabolas corresponding to fractional values of the
 351 flux quantum decrease more and more in energy with respect to the ones centered at integers
 352 values of the reduced flux until they become all degenerate. To simplify the discussion, we
 353 have considered odd number of particles for each spin component. Conversely, we would have
 354 to include the parity effect, occurring in fermionic systems with even number of particles per
 355 spin species [70]. However, this does not change qualitatively the outcomes as it only results
 356 in a shift of the whole energy landscape by $\Phi_0/2$. We finally point out that, in the presence of
 357 a lattice, the energy correction depends also on the center-of-mass motion. In this case, not
 358 only the offset of the parabola branches is modified, but also their curvature [46, 54].

359 5 Concluding remarks

360 In this article, we have presented an alternative to the Bethe Ansatz, called the necklace Ansatz,
361 which allows to access the many-body wavefunction of a quantum mixture on a ring in the
362 limit of strong repulsive contact interactions. Our method applies to both $SU(\kappa)$ mixtures
363 and mixtures where the exchange symmetry is broken by taking different values for intra- and
364 inter-component interaction strengths.

365 The necklace Ansatz allows one to obtain all the possible spin configurations very directly
366 and brings a deep insight on their connection with the total momentum. The simplicity of this
367 approach with respect to solving the Bethe's equations relies on the following facts: (i) being
368 in the strongly interacting limit allows us to build the solution for an N -fermion (boson) mix-
369 ture using as a building block the solution for N free fermions (N TG bosons); (ii) we reduce
370 the dimension of the problem by organizing the sectors in snippets – groups of sectors that are
371 equivalent under the permutation of identical particles, and the snippets in necklaces – groups
372 of snippets that are equivalent up-to a rotation and thus that can be represented by the same
373 necklace. This drastically reduces the number of independent amplitudes to be determined in
374 the many-body wavefunction. The remaining amplitudes, which are as many as the number
375 of necklaces, are finally obtained by the constrained diagonalization of an effective Hamilto-
376 nian represented on the snippet basis. This approach yields by construction the complete basis
377 of solutions in the given energy subspace. It also complements the method previously devel-
378 oped for mixtures under external confinement [21], solving the open issue on how to impose
379 periodic boundary conditions within this formalism.

380 The necklace Ansatz tames the factorial increase of the Hilbert space with increasing the
381 particle number. In addition, the connection between our Ansatz and combinatorial neck-
382 laces can mitigate the problem of generating and enumerating [58, 73, 74] possible orderings
383 of strongly interacting atoms on a ring as well as spin configurations of the dual Heisen-
384 berg chain¹. This will further simplify numerical calculations and allow one to obtain the
385 many-body wavefunction of strongly correlated mixtures of relatively large systems. Such a
386 wavefunction will be used as a starting point in the future for accurate calculations of both
387 equilibrium and dynamical properties of 1D mixtures in ring geometries.

388 Acknowledgments

389 We thank Luigi Amico, Wayne Chetcuti, Nathan Harshman and Véronique Terras for enlight-
390 ening discussions. We acknowledge funding from the ANR-21-CE47-0009 Quantum-SOPHA
391 project, and the support of the Institut Henri Poincaré (UAR 839 CNRS-Sorbonne Université),
392 and LabEx CARMIN (ANR-10-LABX-59-01).

393 GP and GAD have contributed equally to this work.

¹Furthermore, this connection might guide the studies of mathematical symmetries of a few-body system (see, e.g., Refs. [75, 76] for corresponding studies in a trap) using the existing literature, see, e.g., Ref. [77].

394 A Alternative derivation of the necklace Ansatz

395 For completeness, we present an alternative derivation of the ansatz introduced in the main
396 text. To this end, we study the auxiliary Hamiltonian

$$\begin{aligned} \hat{h} = & -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial y^2} + \sum_{\sigma=1}^{\kappa} \sum_{i=1}^{N_{\sigma}} g_{I\sigma} \delta(y - x_{i,\sigma}) - \sum_{\sigma=1}^{\kappa} \sum_{i=1}^{N_{\sigma}} \frac{\hbar^2}{2m} \frac{\partial^2}{\partial x_{i,\sigma}^2} \\ & + \sum_{\sigma=1}^{\kappa} g_{\sigma\sigma} \sum_{i=1}^{N_{\sigma}} \sum_{j>i}^{N_{\sigma}} \delta(x_{i,\sigma} - x_{j,\sigma}) + \frac{1}{2} \sum_{\sigma \neq \sigma'=1}^{\kappa} g_{\sigma\sigma'} \sum_{i=1}^{N_{\sigma}} \sum_{j=1}^{N_{\sigma'}} \delta(x_{i,\sigma} - x_{j,\sigma'}). \end{aligned} \quad (23)$$

397 Here, we consider $N = \sum_{\sigma=1}^{\kappa} N_{\sigma} + 1$ particles, introducing explicitly a single ‘impurity’ atom
398 (y coordinate) in the Hamiltonian \hat{h} in comparison to Eq. (5). Note that with a proper re-
399 definition of the coordinates and the numbers of particles, the Hamiltonian \hat{H} can always be
400 written in the form of Eq. (23). By explicitly introducing the ‘impurity’ in this way, we set a
401 reference frame, which allows us to order the snippets in a natural way.

402 Any solution to Eq. (23) has the form (see, e.g., Ref. [78])

$$\Psi_h(y, X) = e^{iPy/\hbar} \phi_{\mathcal{P}}(\tilde{X}), \quad (24)$$

403 where \tilde{X} is a set of coordinates of the majority particles measured with respect to the position
404 of the impurity y , i.e., $\tilde{x}_{i,\sigma} = L\theta(y - x_{i,\sigma}) + x_{i,\sigma} - y$; \mathcal{P} is the total momentum of the system.
405 In the impenetrable limit, $1/g = 0$ and $1/g_{\sigma\sigma'} = 0$, the function $\phi_{\mathcal{P}}$ is an eigenstate of the
406 Hamiltonian

$$\hat{h}_{\mathcal{P}} = -\frac{\hbar^2}{2m} \sum_{\sigma=1}^{\kappa} \sum_{i=1}^{N_{\sigma}} \frac{\partial^2}{\partial \tilde{x}_{i,\sigma}^2} - \frac{\hbar^2}{2m} \left(\sum_{\sigma=1}^{\kappa} \sum_{i=1}^{N_{\sigma}} \frac{\partial}{\partial \tilde{x}_{i,\sigma}} \right)^2 + i \frac{\hbar \mathcal{P}}{m} \sum_{\sigma=1}^{\kappa} \sum_{i=1}^{N_{\sigma}} \frac{\partial}{\partial \tilde{x}_{i,\sigma}}. \quad (25)$$

407 The last term in $\hat{h}_{\mathcal{P}}$ can be eliminated by the gauge transformation:

$$\phi_{\mathcal{P}}(\tilde{X}) = \exp \left[i \frac{\mathcal{P}}{\hbar} \frac{\sum_{i,\sigma} \tilde{x}_{i,\sigma}}{1 + \sum_{\sigma} N_{\sigma}} \right] f_{\mathcal{P}}(\tilde{X}), \quad (26)$$

408 where $f_{\mathcal{P}}$ solves the following Schrödinger equation

$$\left[-\frac{\hbar^2}{2m} \sum_{\sigma=1}^{\kappa} \sum_{i=1}^{N_{\sigma}} \frac{\partial^2}{\partial \tilde{x}_{i,\sigma}^2} - \frac{\hbar^2}{2m} \left(\sum_{\sigma=1}^{\kappa} \sum_{i=1}^{N_{\sigma}} \frac{\partial}{\partial \tilde{x}_{i,\sigma}} \right)^2 \right] f_{\mathcal{P}} = \epsilon f_{\mathcal{P}}. \quad (27)$$

409 First, we note that for every ordering $Q(\tilde{X})$, $f_{\mathcal{P}}$ can be written as

$$f_{\mathcal{P}}(Q(\tilde{X})) = \tilde{c}_Q \exp \left[-i \frac{\mathcal{P}_F y}{\hbar} - i \frac{\mathcal{P}_F}{\hbar} \frac{\sum_{i,\sigma} \tilde{x}_{i,\sigma}}{1 + \sum_{\sigma} N_{\sigma}} \right] \Psi_F, \quad (28)$$

410 where Ψ_F is a state that describes a system of $1 + \sum_{\sigma} N_{\sigma}$ spin-polarized fermions, and \mathcal{P}_F is the
411 total momentum of that state; \tilde{c}_Q is an arbitrary coefficient. Indeed, the solution in Eq. (28)
412 satisfies the boundary conditions, i.e., it vanishes whenever any two particles meet. To argue
413 that any solution of Eq. (27) has the form of Eq. (28), one can design a proof by contradiction,
414 i.e., show that one can construct a fermionic state Ψ_F from a solution Eq. (27). To this end, it
415 is useful to consider the known solution to the problem of one impurity in a Fermi gas in the
416 frame co-moving with the impurity [79–81].

417 We see that any eigenstate of \hat{h} has the form

$$\Psi_h(y, X) = \sum_Q \tilde{c}_Q \Theta_Q(\tilde{X}) \exp \left[i \frac{(\mathcal{P} - \mathcal{P}_F)y}{\hbar} + i \frac{\mathcal{P} - \mathcal{P}_F}{\hbar} \frac{\sum_{i,\sigma} \tilde{x}_{i,\sigma}}{1 + \sum_{\sigma} N_{\sigma}} \right] \Psi_F(y, X), \quad (29)$$

418 where the sum runs over all orderings of \tilde{X} coordinates. The connection of this expression to
 419 the ansatz in Eq. (10) becomes clear when working with the sector basis and remarking that
 420 $\mathcal{K} = (\mathcal{P} - \mathcal{P}_F)/\hbar$. Indeed, let us consider an example: three SU(3) fermions, i.e., the system
 421 in Sec. 4.4 with $\Phi = \mathbf{0}$. Without loss of generality, we assign the coordinates y, x_1, x_2 to the
 422 particles $\bullet \circ \triangleright$. Let us now write the function $\Psi_h(y, X)$ on the six possible sectors

$$(\bullet \circ \triangleright) = \{y < x_1 < x_2; \tilde{x}_1 < \tilde{x}_2\} \quad \Psi_h(y, X) = \tilde{c}_1 \Psi_{\mathcal{K}}(y, X) \quad (30)$$

$$(\bullet \triangleright \circ) = \{y < x_2 < x_1; \tilde{x}_2 < \tilde{x}_1\} \quad \Psi_h(y, X) = \tilde{c}_2 \Psi_{\mathcal{K}}(y, X) \quad (31)$$

$$(\triangleright \bullet \circ) = \{x_2 < y < x_1; \tilde{x}_1 < \tilde{x}_2\} \quad \Psi_h(y, X) = \tilde{c}_1 e^{\frac{i\mathcal{K}L}{3}} \Psi_{\mathcal{K}}(y, X) \quad (32)$$

$$(\circ \bullet \triangleright) = \{x_1 < y < x_2; \tilde{x}_2 < \tilde{x}_1\} \quad \Psi_h(y, X) = \tilde{c}_2 e^{\frac{i\mathcal{K}L}{3}} \Psi_{\mathcal{K}}(y, X) \quad (33)$$

$$(\circ \triangleright \bullet) = \{x_1 < x_2 < y; \tilde{x}_1 < \tilde{x}_2\} \quad \Psi_h(y, X) = \tilde{c}_1 e^{\frac{2i\mathcal{K}L}{3}} \Psi_{\mathcal{K}}(y, X) \quad (34)$$

$$(\triangleright \circ \bullet) = \{x_2 < x_1 < y; \tilde{x}_2 < \tilde{x}_1\} \quad \Psi_h(y, X) = \tilde{c}_2 e^{\frac{2i\mathcal{K}L}{3}} \Psi_{\mathcal{K}}(y, X) \quad (35)$$

423 where $\Psi_{\mathcal{K}}(y, X) = e^{i\mathcal{K}(y+x_1+x_2)/3} \Psi_F(y, X)$. To write down the wavefunction on sectors in this
 424 way, we used the definition of $\tilde{x}_i = L\theta(y - x_i) + x_i - y$ to connect the two orderings of
 425 $\{\tilde{x}_1, \tilde{x}_2\}$ to the six orderings of $\{y, x_1, x_2\}$. We see that Ψ_h indeed reproduces the necklace
 426 Ansatz reported in Sec. 4.4 when choosing $\mathcal{K} = 2\pi n/L$, $c_1 = \tilde{c}_1$ and $c_2 = \tilde{c}_2 e^{2\pi ni/3}$.

427 Note that if the impurity belongs to the component σ' , by defining $Q' = Q(y \rightarrow x_{1,\sigma'})$, the
 428 values of \tilde{c}_Q must satisfy the property: $\tilde{c}_Q = \tilde{c}_{Q'} \exp \left[i \frac{\mathcal{P} - \mathcal{P}_F}{\hbar(1 + \sum_{\sigma} N_{\sigma})} L j_{y, x_{1,\sigma'}} \right]$, where an integer
 429 $j_{y, x_{1,\sigma'}}$ is the distance between y and $x_{1,\sigma'}$ on a lattice for the ordering Q .

430 Finally, let us consider the system in Eq. (23) coupled to the Aharonov-Bohm flux as in
 431 Sec. 4.4. To this end, we substitute: $-i\hbar\partial/(\partial x_{i,\sigma}) \rightarrow -i\hbar\partial/(\partial x_{i,\sigma}) - m\Omega L/(2\pi)$ and for
 432 the impurity $-i\hbar\partial/(\partial y) \rightarrow -i\hbar\partial/(\partial y) - m\Omega L/(2\pi)$ in Eq. (23). The transformation to the
 433 co-moving with the ‘impurity’ frame of reference according to Eq. (24) separates the flux from
 434 the interaction potential explicitly, see, e.g., Supplementary of Ref. [82]. This means that the
 435 relative dynamics of the system is independent of the value of Φ . In particular, the interaction
 436 between particles is disconnected from Φ in agreement with Sec. 4.4.

437 B Derivation of the effective Hamiltonian

438 In order to obtain the effective low-energy Hamiltonian, we have followed the procedure out-
 439 lined in [23]. We expand in order of $1/g_{\sigma\sigma'}$ the Hamiltonian (5) on the snippet basis $\Psi_{\mathcal{K},s}$:

$$[\hat{H}]_{s,s'} \simeq [\hat{H}_{g_{\sigma\sigma'} \rightarrow \infty}]_{s,s'} + [H_{\text{eff}}]_{s,s'} \quad (36)$$

440 where $[\hat{H}_{g_{\sigma\sigma'} \rightarrow \infty}]_{s,s'} = E_{\infty}^{N, n_s} \delta_{s,s'}$

$$[H_{\text{eff}}]_{s,s'} = - \sum_{\sigma}^{\kappa} \frac{1}{g_{\sigma\sigma}} \left[\int dX \Psi_{\mathcal{K},s}^* g_{\sigma\sigma}^2 \sum_{i=1}^{N_{\sigma}} \sum_{j>i}^{N_{\sigma}} \delta(x_{i,\sigma} - x_{j,\sigma}) \Psi_{\mathcal{K},s'} \right]_{g_{\sigma\sigma} \rightarrow \infty} \\ - \sum_{\sigma \neq \sigma'=1}^{\kappa} \frac{1}{g_{\sigma\sigma'}} \left[\frac{1}{2} \int dX \Psi_{\mathcal{K},s}^* g_{\sigma\sigma'}^2 \sum_{i=1}^{N_{\sigma}} \sum_{j=1}^{N_{\sigma'}} \delta(x_{i,\sigma} - x_{j,\sigma'}) \Psi_{\mathcal{K},s'} \right]_{g_{\sigma\sigma'} \rightarrow \infty}. \quad (37)$$

441 By setting $g_{\sigma\sigma'} = \beta_{\sigma\sigma'} g$, we can write

$$[H_{\text{eff}}]_{s,s'} = -\frac{1}{g}[V]_{s,s'}, \quad (38)$$

442 where V is the contact matrix, whose elements

$$[V]_{s,s'} = \sum_{\sigma}^{\kappa} \frac{1}{\beta_{\sigma\sigma}} \left[\int dX \Psi_{\mathcal{K},s}^* g_{\sigma\sigma}^2 \sum_{i=1}^{N_{\sigma}} \sum_{j>i}^{N_{\sigma}} \delta(x_{i,\sigma} - x_{j,\sigma}) \Psi_{\mathcal{K},s'} \right]_{g_{\sigma\sigma} \rightarrow \infty} \quad (39)$$

$$\sum_{\sigma \neq \sigma'=1}^{\kappa} \frac{1}{\beta_{\sigma\sigma'}} \left[\frac{1}{2} \int dX \Psi_{\mathcal{K},s}^* g_{\sigma\sigma'}^2 \sum_{i=1}^{N_{\sigma}} \sum_{j=1}^{N_{\sigma'}} \delta(x_{i,\sigma} - x_{j,\sigma'}) \Psi_{\mathcal{K},s'} \right]_{g_{\sigma\sigma'} \rightarrow \infty},$$

443 can be evaluated using the cusp condition [83].

444 For SU(2) mixtures $\beta_{\sigma\sigma'}$ is the same for any σ, σ' , so that we can set $\beta_{\sigma\sigma'} = 1$. We thus
445 obtain Eqs. (13) and (14) respectively for fermions and bosons.

446 Remark that the contact matrix elements (15) do not depend on the momentum \mathcal{K} (n) [83].
447 Indeed $\alpha^{(N)}$ is equal, up to the dimensional constant $\hbar^2/(mL)$, to the difference between the
448 total kinetic energy and the center-of-mass kinetic energy [27, 84], thus it does not depend
449 on the ring current [83]. This result is consistent with the Bethe formalism, by doing the
450 expansion of the Bethe equations with respect to the inverse of the interaction strength (see
451 Appendix D).

452 C Contact matrices

453 In this Appendix, we provide the contact matrices V that have been used in order to obtain
454 the results outlined in the main text.

455 The conditioned eigenvalues problem for the 4+2 SU(2) fermionic mixture takes the form

$$\begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 & -1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 & 0 & -1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & -1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & -1 & -1 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & -1 \\ 0 & -1 & -1 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & -1 & -1 & 0 & 0 \\ 0 & 0 & -1 & -1 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & -1 & -1 & -1 \\ 0 & 0 & 0 & -1 & -1 & 0 & 0 & 0 & 4 & 0 & 0 & -1 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 & -1 & 0 & 0 & 0 & 4 & 0 & -1 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 4 & 0 & -1 & -1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & -1 & 0 & -1 & -1 & 0 & 4 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & -1 & 0 & -1 & 0 & 4 & 0 & 4 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & -1 & -1 & 0 & -1 & 0 & 4 & 0 \end{pmatrix} \begin{pmatrix} c_1 \\ c_1 e^{-in\pi/3} \\ c_1 e^{-2in\pi/3} \\ c_1 e^{-in\pi} \\ c_1 e^{-4in\pi/3} \\ c_1 e^{-5in\pi/3} \\ c_2 \\ c_2 e^{-in\pi/3} \\ c_2 e^{-2in\pi/3} \\ c_2 e^{-in\pi} \\ c_2 e^{-4in\pi/3} \\ c_2 e^{-5in\pi/3} \\ c_3 \\ c_3 e^{-in\pi/3} \\ c_3 e^{-2in\pi/3} \end{pmatrix} = \frac{\xi_n}{\alpha^{(6)}} \begin{pmatrix} c_1 \\ c_1 e^{-in\pi/3} \\ c_1 e^{-2in\pi/3} \\ c_1 e^{-in\pi} \\ c_1 e^{-4in\pi/3} \\ c_1 e^{-5in\pi/3} \\ c_2 \\ c_2 e^{-in\pi/3} \\ c_2 e^{-2in\pi/3} \\ c_2 e^{-in\pi} \\ c_2 e^{-4in\pi/3} \\ c_2 e^{-5in\pi/3} \\ c_3 \\ c_3 e^{-in\pi/3} \\ c_3 e^{-2in\pi/3} \end{pmatrix}. \quad (40)$$

456 For the case of 3+3 SU(2) fermions, the V matrix reads

$$\frac{V}{\alpha^{(6)}} = \begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ -1 & -1 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & -1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & -1 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & -1 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & 0 & 0 & 4 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & -1 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 4 & 0 & -1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & -1 & -1 & 0 & 0 & 0 & -1 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & -1 & 0 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & -1 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & -1 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 4 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & -1 & -1 & 0 & -1 & 0 & -1 & 0 & -1 & 0 & 6 & 0 & 6 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & -1 & 0 & -1 & 0 & -1 & 0 & -1 & 0 & -1 & 0 & 6 & 0 & 6 & 0 \end{pmatrix} \quad (41)$$

457 For the case of 3+3 SU(2) bosons, the V matrix reads

$$\frac{V}{\alpha^{(6)}} = \begin{pmatrix} 10 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 10 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 10 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 10 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 10 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 10 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 8 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 8 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 8 & 0 & 0 & 0 & 8 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 8 & 0 & 0 & 0 & 8 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 8 & 0 & 0 & 0 & 8 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 8 & 0 & 0 & 0 & 8 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 8 & 0 & 0 & 0 & 8 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 6 & 0 & 6 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 6 & 0 & 6 \end{pmatrix}. \quad (42)$$

458 For the case of 3+3 SB bosons, the V matrix reads

$$\frac{V}{\alpha^{(6)}} = \begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 4 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 4 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 6 & 0 & 6 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 6 & 0 & 6 \end{pmatrix}. \quad (43)$$

459 The contact matrix for 3 SU(3) fermions reads

$$\frac{V}{\alpha^{(3)}} = \begin{pmatrix} 3 & 0 & 0 & -1 & -1 & -1 \\ 0 & 3 & 0 & -1 & -1 & -1 \\ 0 & 0 & 3 & -1 & -1 & -1 \\ -1 & -1 & -1 & 3 & 0 & 0 \\ -1 & -1 & -1 & 0 & 3 & 0 \\ -1 & -1 & -1 & 0 & 0 & 3 \end{pmatrix}. \quad (44)$$

460 D Energy correction to order $\mathcal{O}\left(\frac{1}{g}\right)$: Bethe Ansatz derivation

461 In this section, we explicitly derive the energy correction at large but finite interaction strength
 462 g for a SU(2) mixture of N particles. To do so, we consider the first-order expansion of the
 463 Bethe equations close to the limit $\frac{1}{g} \rightarrow 0$. In the following, we consider the rescaled interaction
 464 strength $u \doteq \frac{2m}{\hbar^2} g$, so that u has the dimension of a wavevector. We start from the Bethe
 465 equations valid for any interaction strength, which read:

$$\begin{cases} Lk_j = 2\pi\mathcal{I}_j + (-1)^{\eta_B} \sum_{b=1}^{N_b} 2 \arctan\left(\frac{2(\tilde{\lambda}_b - k_j)}{u}\right) + \eta_B \sum_{\ell=1}^N 2 \arctan\left(\frac{k_\ell - k_j}{u}\right) \\ \sum_{j=1}^N 2 \arctan\left(\frac{2(\tilde{\lambda}_a - k_j)}{u}\right) = 2\pi\mathcal{J}_a + \sum_{b=1}^{N_b} 2 \arctan\left(\frac{\tilde{\lambda}_a - \tilde{\lambda}_b}{u}\right) \end{cases}, \quad (45)$$

466 where $\eta_B = 1$ for bosons and $\eta_B = 0$ for fermions. The quantum numbers \mathcal{I}_j and \mathcal{J}_m are de-
 467 fined in Section 2.3. In the limit $u \rightarrow \infty$, one has $\frac{\tilde{\lambda}_a}{u} \gg \frac{k_j}{u}$, therefore we expand the arcotan-
 468 gent function according to the first-order expansions $\arctan(a+x) = \arctan(a) + \frac{x}{1+a^2} + \mathcal{O}(x^2)$

469 and $\arctan(x) = x + \mathcal{O}(x^2)$. We introduce the rescaled spin rapidities $\Lambda_a \doteq \frac{2\tilde{\lambda}}{u}$ and obtain:

$$\begin{cases} Lk_j = 2\pi\mathcal{I}_j + (-1)^{\eta_B} \sum_{b=1}^{N_\downarrow} 2 \arctan(\Lambda_b) - (-1)^{\eta_B} \frac{4}{u} k_j \sum_{b=1}^{N_\downarrow} \frac{1}{1 + \Lambda_b^2} + \frac{2\eta_B}{u} \left(\sum_{\ell} k_\ell - Nk_j \right) \\ 2 \arctan(\Lambda_a) = \frac{4}{uN} \sum_j k_j \frac{1}{1 + \Lambda_a^2} + \frac{2\pi}{N} \mathcal{J}_a + \sum_{b=1}^{N_\downarrow} \frac{2}{N} \arctan\left(\frac{\Lambda_a - \Lambda_b}{2}\right) \end{cases} \quad (46)$$

470 A straightforward substitution of the right-hand-side of the second equation in the first equa-
471 tion yields:

$$\begin{cases} Lk_j = 2\pi\mathcal{I}_j + (-1)^{\eta_B} \frac{2\pi}{N} \sum_{b=1}^{N_\downarrow} \mathcal{J}_b + \frac{1}{u} \left(\frac{1}{N} \sum_{\ell} k_\ell - k_j \right) \left(2N\eta_B + (-1)^{\eta_B} \sum_{b=1}^{N_\downarrow} \frac{4}{1 + \Lambda_b^2} \right) \\ 2 \arctan(\Lambda_a) = \frac{1}{uN} \sum_j k_j \frac{4}{1 + \Lambda_a^2} + \frac{2\pi}{N} \mathcal{J}_a + \sum_{b=1}^{N_\downarrow} \frac{2}{N} \arctan\left(\frac{\Lambda_a - \Lambda_b}{2}\right) \end{cases} \quad (47)$$

472 where we use $\sum_{a,b=1}^{N_\downarrow} \arctan\left(\frac{\Lambda_a - \Lambda_b}{2}\right) = 0$. In the limit $\frac{1}{u} \rightarrow 0$, one recovers Eqs.(7) and (9).

473 We can write the first equation in a more compact form:

$$k_j = \frac{2\pi}{L} \mathcal{I}_j + \frac{\chi}{L} + \frac{1}{uL} \delta k_j, \quad (48)$$

474 where we defined:

$$\chi \doteq (-1)^{\eta_B} \frac{2\pi}{N} \sum_{b=1}^{N_\downarrow} \mathcal{J}_b, \quad (49)$$

$$\delta k_j \doteq \left(\frac{1}{N} \sum_{\ell} k_\ell - k_j \right) \left(2N\eta_B + (-1)^{\eta_B} \sum_{b=1}^{N_\downarrow} \frac{4}{1 + \Lambda_b^2} \right). \quad (50)$$

475 Remarkably, χ can include any shift to the rapidities k_j which does not depend on the index
476 j , as for instance an artificial gauge field.

477 The energy in this strongly interacting limit is:

$$\begin{aligned} \frac{2m}{\hbar^2} E_{1/u} &= \sum_j k_j^2 = \sum_j \left(\frac{2\pi}{L} \mathcal{I}_j + \frac{\chi}{L} + \frac{1}{uL} \delta k_j \right)^2 = \\ &= \sum_j \left(\frac{2\pi}{L} \mathcal{I}_j + \frac{\chi}{L} \right)^2 + \frac{2}{uL} \sum_j \left(\frac{2\pi}{L} \mathcal{I}_j + \frac{\chi}{L} \right) \delta k_j + \mathcal{O}\left(\frac{1}{u^2}\right) \doteq \frac{2m}{\hbar^2} \left(E_\infty + \delta E_{1/u} + \mathcal{O}\left(\frac{1}{u^2}\right) \right), \end{aligned} \quad (51)$$

478 where we introduced the energy correction:

$$\delta E_{1/u} = \frac{\hbar^2}{uLm} \sum_j \left(\frac{2\pi}{L} \mathcal{I}_j + \frac{\chi}{L} \right) \delta k_j, \quad (52)$$

479 Neglecting the $\mathcal{O}\left(\frac{1}{u^2}\right)$ terms, the energy correction reads:

$$\begin{aligned}\delta E_{1/u} &= \frac{\hbar^2}{uLm} \sum_j \left(\frac{2\pi}{L} \mathcal{I}_j + \frac{\chi}{L} \right) \left(\frac{1}{N} \sum_\ell k_\ell - k_j \right) \left(2N\eta_B + (-1)^{\eta_B} \sum_{b=1}^{N_1} \frac{4}{1 + \Lambda_b^2} \right) \\ \delta E_{1/u} &= \frac{\hbar^2}{uLm} \left(\frac{1}{N} \left(\frac{2\pi}{L} \sum_\ell \mathcal{I}_\ell + \frac{N\chi}{L} \right)^2 - \sum_j \left(\frac{2\pi}{L} \mathcal{I}_j + \frac{\chi}{L} \right)^2 \right) \left(2N\eta_B + (-1)^{\eta_B} \sum_{b=1}^{N_1} \frac{4}{1 + \Lambda_b^2} \right) \\ \delta E_{1/u} &= -\frac{\hbar^2}{uLm} \left(\frac{4\pi^2}{L^2} \sum_j \mathcal{I}_j^2 - \frac{4\pi^2}{NL^2} \left(\sum_\ell \mathcal{I}_\ell \right)^2 \right) \left(2N\eta_B + (-1)^{\eta_B} \sum_{b=1}^{N_1} \frac{4}{1 + \Lambda_b^2} \right).\end{aligned}\quad (53)$$

480 In order to have only first-order correction in $\frac{1}{u}$, the spin rapidities Λ_b are obtained solving
481 the corresponding Bethe equation in the limit $\frac{1}{u} \rightarrow 0$, Eq. (9). From the last line of Eq. (53),
482 we see that the center of mass momentum does not contribute to the first-order corrections to
483 the total energy. Moreover, if we introduce an effective coupling

$$J_{\text{eff}} = \frac{\hbar^2}{uLm} \left(\frac{4\pi^2}{L^2} \sum_j \mathcal{I}_j^2 - \frac{4\pi^2}{NL^2} \left(\sum_\ell \mathcal{I}_\ell \right)^2 \right), \quad (54)$$

484 the correction to the energy can be expressed as:

$$\delta E_{1/u} = -J_{\text{eff}} \left(2N\eta_B + (-1)^{\eta_B} \sum_{b=1}^{N_1} \frac{4}{1 + \Lambda_b^2} \right), \quad (55)$$

485 which, up to a constant shift in the bosonic case, coincides with the Bethe Ansatz solution
486 for the energy of an isotropic Heisenberg spin chain. The ferromagnetic or antiferromagnetic
487 nature of the ground state is determined by the value of η_B and therefore by the statistics of
488 the mixture. Remark that $J_{\text{eff}}uL/2$ is equal to the difference between the total kinetic energy
489 and the center-of-mass kinetic energy, and that $J_{\text{eff}}uL/2 = J_{\text{eff}}g mL/\hbar^2 = mL/\hbar^2 \alpha^{(N)}$.

Table 7: Energy corrections to first order in $1/u$ for $N = 4$ and $N_1 = 2$ bosons. The spin rapidities are calculated from Eq. (9). The results are in units of J_{eff} .

Λ_1	Λ_2	$-\frac{\delta E_{1/u}}{J_{\text{eff}}}$
$1/\sqrt{3}$	$-1/\sqrt{3}$	2
0	∞	4
1	∞	6
-1	∞	6
i	$-i$	6
∞	$-\infty$	8

490 We computed explicitly the energy correction in the case of $N = 4$ and $N_1 = 2$ bosons.
491 First, to determine the correct Λ_1 and Λ_2 , we solved the Bethe equations (9). Then, we used
492 Eq. (55) to evaluate the first-order correction to the energy at infinite interactions. Since
493 Eq. (55) diverges for $\Lambda_{1,2} = \pm i$, in order to obtain the corresponding correction we had to
494 introduce the regularization $\Lambda_{1,2} = \epsilon \pm i$ and then compute the limit of Eq. (55) for $\epsilon \rightarrow 0$.

495 We show the results in Table 7. The energy corrections coincide with the results presented
496 in the second column of Table 1 ($-\frac{\delta E_{1/u}}{J_{\text{eff}}} = \tilde{\xi}_n$), where we computed the first-order energy
497 correction using the necklace Ansatz. More details on the Bethe Ansatz solution for $N = 4$ and
498 $N_1 = 2$, including the exact expression of the corresponding eigenstates, are available in [64].

n	$\tilde{\xi}_n$	c_1	$\gamma^{(2)}$	YD	$E_\infty^{3,n}$
0	0	1	-3	$\begin{array}{ c } \hline \square \\ \hline \end{array}$	$4\pi^2$
1	3	1	0	$\begin{array}{ c c } \hline \square & \square \\ \hline \end{array}$	$\frac{14\pi^2}{3}$
2	3	1	0	$\begin{array}{ c c c } \hline \square & \square & \square \\ \hline \end{array}$	$\frac{20\pi^2}{3}$

Table 8: Solution for the fermionic 2+1 system. $\tilde{\xi}_n$ are the rescaled eigenvalues $\xi_n/\tilde{\alpha}^{(3)}$. $\gamma^{(2)}$ indicates the symmetry of the solution, and the associated Young diagram. The last column corresponds to the energies at infinite interactions (in units of $\hbar^2/(mL^2)$).

499 E Analysis of the strongly interacting limit

500 With the aim to specify the validity range of the strong-interaction expansion used in Sec.
 501 D and in the necklace Ansatz, we compare the solution obtained in the strongly interacting
 502 limit for a system of 2+1 fermions with that obtained by the Bethe equations for intermediate
 503 interactions given in Eq. (45).

504 In Fig. 4, we plot the first three energy levels of the Bethe Ansatz solution as a function of
 505 the inverse of the interaction strength. The ground state at infinite interactions corresponds
 506 to the fully antisymmetric state (see Table 8). It does not depend on the interaction strength
 507 (circles) and coincides with the necklace solution with $n = 0$ (horizontal blue line). The
 508 two excited states at $1/g = 0$ (cross and plus symbols) correspond to the necklace solutions
 509 (tangent thick green and violet lines) with $n = 1$ and $n = 2$ respectively.

510 We observe that the solutions obtained in the strong-interaction limit agree with the ener-
 511 gies from the Bethe Ansatz equations for $1/g \lesssim 0.03mL/\hbar^2$. This corresponds to an effective
 512 interaction parameter $\hbar^2mLg/N = 10$, which provides an estimate for the ratio between the
 513 interaction and kinetic energies for which the necklace Ansatz can still produce accurate re-
 514 sults.

515 References

- 516 [1] I. Bloch, J. Dalibard and W. Zwerger, *Many-body physics with ultracold gases*, Rev. Mod.
 517 Phys. **80**, 885 (2008), doi:[10.1103/RevModPhys.80.885](https://doi.org/10.1103/RevModPhys.80.885).
- 518 [2] M. A. Cazalilla, R. Citro, T. Giamarchi, E. Orignac and M. Rigol, *One dimensional bosons:
 519 From condensed matter systems to ultracold gases*, Rev. Mod. Phys. **83**, 1405 (2011),
 520 doi:[10.1103/RevModPhys.83.1405](https://doi.org/10.1103/RevModPhys.83.1405).
- 521 [3] X.-W. Guan, M. T. Batchelor and C. Lee, *Fermi gases in one dimension: From Bethe ansatz
 522 to experiments*, Rev. Mod. Phys. **85**, 1633 (2013), doi:[10.1103/RevModPhys.85.1633](https://doi.org/10.1103/RevModPhys.85.1633).
- 523 [4] T. Sowiński and M. Ángel García-March, *One-dimensional mixtures of several ul-
 524 tracold atoms: a review*, Reports on Progress in Physics **82**(10), 104401 (2019),
 525 doi:[10.1088/1361-6633/ab3a80](https://doi.org/10.1088/1361-6633/ab3a80).
- 526 [5] A. Minguzzi and P. Vignolo, *Strongly interacting trapped one-dimensional quantum gases:
 527 Exact solution*, AVS Quantum Science **4**(2), 027102 (2022), doi:[10.1116/5.0077423](https://doi.org/10.1116/5.0077423).




Figure 4: First three energy levels for a system of 2+1 fermions, in units of $E^{3,0} = 4\pi^2\hbar^2/(mL^2)$ as functions of $1/g$ (in units of mL/\hbar^2). The horizontal blue line corresponds to a fully antisymmetric state, and the necklace solution ($n = 0$ in Table 8) coincides with the Bethe Ansatz one for any interaction strength (blue circles). The cross and the plus symbols (the lines are guides to the eyes) lines are the Bethe solutions that correspond to the first two excited states at infinite interactions. The tangent thick lines correspond to the necklace solution respectively for $n = 1$ (green line) and $n = 2$ (violet line).

- 528 [6] S. Mistakidis, A. Volosniev, R. Barfknecht, T. Fogarty, T. Busch, A. Foerster, P. Schmelcher
529 and N. Zinner, *Few-body bose gases in low dimensions—A laboratory for quantum dynam-*
530 *ics*, Physics Reports **1042**, 1–108 (2023), doi:[10.1016/j.physrep.2023.10.004](https://doi.org/10.1016/j.physrep.2023.10.004).
- 531 [7] G. Pagano, M. Mancini, G. Cappellini, P. Lombardi, F. Schäfer, H. Hu, X.-J. Liu, J. Catani,
532 C. Sias, M. Inguscio and L. Fallani, *A one-dimensional liquid of fermions with tunable spin*,
533 Nature Physics **10**(3), 198–201 (2014), doi:[10.1038/nphys2878](https://doi.org/10.1038/nphys2878).
- 534 [8] E. H. Lieb and W. Liniger, *Exact Analysis of an Interacting Bose Gas. I. The General Solution*
535 *and the Ground State*, Phys. Rev. **130**, 1605 (1963), doi:[10.1103/PhysRev.130.1605](https://doi.org/10.1103/PhysRev.130.1605).
- 536 [9] E. H. Lieb, *Exact Analysis of an Interacting Bose Gas. II. The Excitation Spectrum*, Phys.
537 Rev. **130**, 1616 (1963), doi:[10.1103/PhysRev.130.1616](https://doi.org/10.1103/PhysRev.130.1616).
- 538 [10] C. N. Yang, *Some Exact Results for the Many-Body Problem in one Dimen-*
539 *sion with Repulsive Delta-Function Interaction*, Phys. Rev. Lett. **19**, 1312 (1967),
540 doi:[10.1103/PhysRevLett.19.1312](https://doi.org/10.1103/PhysRevLett.19.1312).
- 541 [11] B. Sutherland, *Further Results for the Many-Body Problem in One Dimension*, Phys. Rev.
542 Lett. **20**, 98 (1968), doi:[10.1103/PhysRevLett.20.98](https://doi.org/10.1103/PhysRevLett.20.98).
- 543 [12] J. B. McGuire, *Interacting Fermions in One Dimension. I. Repulsive Potential*, Journal of
544 Mathematical Physics **6**(3), 432 (1965), doi:[10.1063/1.1704291](https://doi.org/10.1063/1.1704291).
- 545 [13] J. B. McGuire, *Study of Exactly Soluble One-Dimensional N-Body Problems*, Journal of
546 Mathematical Physics **5**(5), 622 (1964), doi:[10.1063/1.1704156](https://doi.org/10.1063/1.1704156).
- 547 [14] P. Calabrese and J.-S. Caux, *Correlation Functions of the One-Dimensional Attractive Bose*
548 *Gas*, Phys. Rev. Lett. **98**, 150403 (2007), doi:[10.1103/PhysRevLett.98.150403](https://doi.org/10.1103/PhysRevLett.98.150403).

- 549 [15] L. Piroli and P. Calabrese, *Local correlations in the attractive one-dimensional Bose gas:*
550 *From Bethe ansatz to the Gross-Pitaevskii equation*, Phys. Rev. A **94**, 053620 (2016),
551 doi:[10.1103/PhysRevA.94.053620](https://doi.org/10.1103/PhysRevA.94.053620).
- 552 [16] M. Gaudin, *Un système a une dimension de fermions en interaction*, Physics Letters A
553 **24**(1), 55 (1967), doi:[10.1016/0375-9601\(67\)90193-4](https://doi.org/10.1016/0375-9601(67)90193-4).
- 554 [17] X.-W. Guan and T.-L. Ho, *Quantum criticality of a one-dimensional attractive Fermi gas*,
555 Phys. Rev. A **84**, 023616 (2011), doi:[10.1103/PhysRevA.84.023616](https://doi.org/10.1103/PhysRevA.84.023616).
- 556 [18] P. Naldesi, J. Polo, V. Dunjko, H. Perrin, M. Olshanii, L. Amico and A. Minguzzi, *Enhancing*
557 *sensitivity to rotations with quantum solitonic currents*, SciPost Phys. **12**, 138 (2022),
558 doi:[10.21468/SciPostPhys.12.4.138](https://doi.org/10.21468/SciPostPhys.12.4.138).
- 559 [19] M. Girardeau, *Relationship between Systems of Impenetrable Bosons and Fermions in One*
560 *Dimension*, Journal of Mathematical Physics **1**(6), 516 (1960), doi:[10.1063/1.1703687](https://doi.org/10.1063/1.1703687).
- 561 [20] M. D. Girardeau and A. Minguzzi, *Soluble Models of Strongly Interacting Ul-*
562 *tracold Gas Mixtures in Tight Waveguides*, Phys. Rev. Lett. **99**, 230402 (2007),
563 doi:[10.1103/PhysRevLett.99.230402](https://doi.org/10.1103/PhysRevLett.99.230402).
- 564 [21] A. G. Volosniev, D. V. Fedorov, A. S. Jensen, M. Valiente and N. T. Zinner, *Strongly inter-*
565 *acting confined quantum systems in one dimension*, Nature Communications **5**(1), 5300
566 (2014), doi:[10.1038/ncomms6300](https://doi.org/10.1038/ncomms6300).
- 567 [22] F. Deuretzbacher, K. Fredenhagen, D. Becker, K. Bongs, K. Sengstock and D. Pfannkuche,
568 *Exact Solution of Strongly Interacting Quasi-One-Dimensional Spinor Bose Gases*, Phys.
569 Rev. Lett. **100**, 160405 (2008), doi:[10.1103/PhysRevLett.100.160405](https://doi.org/10.1103/PhysRevLett.100.160405).
- 570 [23] F. Deuretzbacher, D. Becker, J. Bjerlin, S. M. Reimann and L. Santos, *Quantum magnetism*
571 *without lattices in strongly interacting one-dimensional spinor gases*, Phys. Rev. A **90**,
572 013611 (2014), doi:[10.1103/PhysRevA.90.013611](https://doi.org/10.1103/PhysRevA.90.013611).
- 573 [24] A. G. Volosniev, D. Petrosyan, M. Valiente, D. V. Fedorov, A. S. Jensen and N. T. Zinner, *En-*
574 *gineering the dynamics of effective spin-chain models for strongly interacting atomic gases*,
575 Phys. Rev. A **91**, 023620 (2015), doi:[10.1103/PhysRevA.91.023620](https://doi.org/10.1103/PhysRevA.91.023620).
- 576 [25] J. Decamp, P. Armagnat, B. Fang, M. Albert, A. Minguzzi and P. Vignolo, *Exact density*
577 *profiles and symmetry classification for strongly interacting multi-component Fermi gases*
578 *in tight waveguides*, New Journal of Physics **18**(5), 055011 (2016), doi:[10.1088/1367-](https://doi.org/10.1088/1367-2630/18/5/055011)
579 [2630/18/5/055011](https://doi.org/10.1088/1367-2630/18/5/055011).
- 580 [26] J. Decamp, J. Jünemann, M. Albert, M. Rizzi, A. Minguzzi and P. Vignolo,
581 *High-momentum tails as magnetic-structure probes for strongly correlated $SU(\kappa)$*
582 *fermionic mixtures in one-dimensional traps*, Phys. Rev. A **94**, 053614 (2016),
583 doi:[10.1103/PhysRevA.94.053614](https://doi.org/10.1103/PhysRevA.94.053614).
- 584 [27] G. Aupetit-Diallo, G. Pecci, C. Pignol, F. Hébert, A. Minguzzi, M. Albert and P. Vignolo,
585 *Exact solution for $SU(2)$ -symmetry-breaking bosonic mixtures at strong interactions*, Phys.
586 Rev. A **106**, 033312 (2022), doi:[10.1103/PhysRevA.106.033312](https://doi.org/10.1103/PhysRevA.106.033312).
- 587 [28] L. Amico, M. Boshier, G. Birkl, A. Minguzzi, C. Miniatura, L.-C. Kwek, D. Aghamalyan,
588 V. Ahufinger, D. Anderson, N. Andrei, A. S. Arnold, M. Baker *et al.*, *Roadmap on Atom-*
589 *tronics: State of the art and perspective*, AVS Quantum Science **3**(3), 039201 (2021),
590 doi:[10.1116/5.0026178](https://doi.org/10.1116/5.0026178).

- 591 [29] J. Dalibard, *Introduction to the physics of artificial gauge fields*, IOS Press,
592 doi:[10.3254/978-1-61499-694-1-1](https://doi.org/10.3254/978-1-61499-694-1-1) (2015).
- 593 [30] J. Dalibard, F. Gerbier, G. Juzeliūnas and P. Öhberg, *Colloquium: Artificial gauge potentials*
594 *for neutral atoms*, Rev. Mod. Phys. **83**, 1523 (2011), doi:[10.1103/RevModPhys.83.1523](https://doi.org/10.1103/RevModPhys.83.1523).
- 595 [31] L. Amico, D. Anderson, M. Boshier, J.-P. Brantut, L.-C. Kwek, A. Minguzzi and W. von Kl-
596 itzing, *Colloquium: Atomtronic circuits: From many-body physics to quantum technologies*,
597 Rev. Mod. Phys. **94**, 041001 (2022), doi:[10.1103/RevModPhys.94.041001](https://doi.org/10.1103/RevModPhys.94.041001).
- 598 [32] C. K. Lai and C. N. Yang, *Ground-State Energy of a Mixture of Fermions and Bosons in*
599 *One Dimension with a Repulsive δ -Function Interaction*, Phys. Rev. A **3**, 393 (1971),
600 doi:[10.1103/PhysRevA.3.393](https://doi.org/10.1103/PhysRevA.3.393).
- 601 [33] C. Marboe and D. Volin, *Fast analytic solver of rational Bethe equations*, Journal of
602 Physics A: Mathematical and Theoretical **50**(20), 204002 (2017), doi:[10.1088/1751-](https://doi.org/10.1088/1751-8121/aa6b88)
603 [8121/aa6b88](https://doi.org/10.1088/1751-8121/aa6b88).
- 604 [34] W. Hao, R. I. Nepomechie and A. J. Sommes, *Completeness of solutions of Bethe's equa-*
605 *tions*, Phys. Rev. E **88**, 052113 (2013), doi:[10.1103/PhysRevE.88.052113](https://doi.org/10.1103/PhysRevE.88.052113).
- 606 [35] J. S. Van Dyke, G. S. Barron, N. J. Mayhall, E. Barnes and S. E. Economou, *Preparing*
607 *Bethe Ansatz Eigenstates on a Quantum Computer*, PRX Quantum **2**, 040329 (2021),
608 doi:[10.1103/PRXQuantum.2.040329](https://doi.org/10.1103/PRXQuantum.2.040329).
- 609 [36] A. N. Kirillov and R. Sakamoto, *Singular solutions to the Bethe ansatz equations and rigged*
610 *configurations*, Journal of Physics A: Mathematical and Theoretical **47**(20), 205207
611 (2014), doi:[10.1088/1751-8113/47/20/205207](https://doi.org/10.1088/1751-8113/47/20/205207).
- 612 [37] X.-W. Guan, Z.-Q. Ma and B. Wilson, *One-dimensional multicomponent fermions with δ -*
613 *function interaction in strong- and weak-coupling limits: κ -component Fermi gas*, Phys.
614 Rev. A **85**, 033633 (2012), doi:[10.1103/PhysRevA.85.033633](https://doi.org/10.1103/PhysRevA.85.033633).
- 615 [38] S. Viefers, P. Koskinen, P. Singha Deo and M. Manninen, *Quantum rings for beginners:*
616 *energy spectra and persistent currents*, Physica E: Low-dimensional Systems and Nanos-
617 tructures **21**(1), 1 (2004), doi:[10.1016/j.physe.2003.08.076](https://doi.org/10.1016/j.physe.2003.08.076).
- 618 [39] M. Manninen, S. Viefers and S. Reimann, *Quantum rings for beginners II: Bosons ver-*
619 *sus fermions*, Physica E: Low-dimensional Systems and Nanostructures **46**, 119 (2012),
620 doi:[10.1016/j.physe.2012.09.013](https://doi.org/10.1016/j.physe.2012.09.013).
- 621 [40] A. Imambekov and E. Demler, *Exactly solvable case of a one-dimensional Bose–Fermi mix-*
622 *ture*, Phys. Rev. A **73**, 021602 (2006), doi:[10.1103/PhysRevA.73.021602](https://doi.org/10.1103/PhysRevA.73.021602).
- 623 [41] C. N. Yang, *Some Exact Results for the Many-Body Problem in one Dimen-*
624 *sion with Repulsive Delta-Function Interaction*, Phys. Rev. Lett. **19**, 1312 (1967),
625 doi:[10.1103/PhysRevLett.19.1312](https://doi.org/10.1103/PhysRevLett.19.1312).
- 626 [42] M. Gaudin and J. Caux, *The Bethe Wavefunction*, Cambridge University Press,
627 doi:[10.1017/CBO9781107053885](https://doi.org/10.1017/CBO9781107053885) (2014).
- 628 [43] N. Oelkers, M. T. Batchelor, M. Bortz and X.-W. Guan, *Bethe ansatz study of one-*
629 *dimensional Bose and Fermi gases with periodic and hard wall boundary conditions*, Jour-
630 nal of Physics A: Mathematical and General **39**(5), 1073 (2006), doi:[10.1088/0305-](https://doi.org/10.1088/0305-4470/39/5/005)
631 [4470/39/5/005](https://doi.org/10.1088/0305-4470/39/5/005).

- 632 [44] Y.-Q. Li, S.-J. Gu, Z.-J. Ying and U. Eckern, *Exact results of the ground state and excitation*
633 *properties of a two-component interacting Bose system*, Europhysics Letters (EPL) **61**(3),
634 368 (2003), doi:[10.1209/epl/i2003-00183-2](https://doi.org/10.1209/epl/i2003-00183-2).
- 635 [45] F. V. Kusmartsev, *Magnetic resonance on a ring of aromatic molecules*, Journal of Physics:
636 Condensed Matter **3**(18), 3199 (1991), doi:[10.1088/0953-8984/3/18/014](https://doi.org/10.1088/0953-8984/3/18/014).
- 637 [46] N. Yu and M. Fowler, *Persistent current of a Hubbard ring threaded with a magnetic flux*,
638 Phys. Rev. B **45**, 11795 (1992), doi:[10.1103/PhysRevB.45.11795](https://doi.org/10.1103/PhysRevB.45.11795).
- 639 [47] C. N. Yang, *Some Exact Results for the Many-Body Problem in one Dimension*
640 *with Repulsive Delta-Function Interaction*, Phys. Rev. Lett. **19**, 1312 (1967),
641 doi:[10.1103/PhysRevLett.19.1312](https://doi.org/10.1103/PhysRevLett.19.1312).
- 642 [48] A. Imambekov and E. Demler, *Applications of exact solution for strongly inter-*
643 *acting one-dimensional Bose–Fermi mixture: Low-temperature correlation functions,*
644 *density profiles, and collective modes*, Annals of Physics **321**(10), 2390 (2006),
645 doi:[10.1016/j.aop.2005.11.017](https://doi.org/10.1016/j.aop.2005.11.017).
- 646 [49] H. A. Bethe, *Zur Theorie der Metalle. i. Eigenwerte und Eigenfunktionen der linearen Atom-*
647 *kette*, Zeit. für Physik **71**, 205 (1931), doi:[10.1007/BF01341708](https://doi.org/10.1007/BF01341708).
- 648 [50] F. Franchini, *An Introduction to Integrable Techniques for One-Dimensional Quantum Sys-*
649 *tems*, Lecture Notes in Physics **940** (2017), doi:[10.1007/978-3-319-48487-7](https://doi.org/10.1007/978-3-319-48487-7).
- 650 [51] F. H. L. Essler, H. Frahm, F. Göhmann, A. Klümper and V. E. Korepin, *The One-Dimensional*
651 *Hubbard Model*, Cambridge University Press, doi:[10.1017/CBO9780511534843](https://doi.org/10.1017/CBO9780511534843) (2005).
- 652 [52] B. Sutherland, *Model for a multicomponent quantum system*, Phys. Rev. B **12**, 3795
653 (1975), doi:[10.1103/PhysRevB.12.3795](https://doi.org/10.1103/PhysRevB.12.3795).
- 654 [53] M. Ogata and H. Shiba, *Bethe-ansatz wave function, momentum distribution, and spin*
655 *correlation in the one-dimensional strongly correlated hubbard model*, Phys. Rev. B **41**,
656 2326 (1990), doi:[10.1103/PhysRevB.41.2326](https://doi.org/10.1103/PhysRevB.41.2326).
- 657 [54] W. J. Chetcuti, T. Haug, L.-C. Kwek and L. Amico, *Persistent current of $SU(N)$ fermions*,
658 SciPost Phys. **12**, 033 (2022), doi:[10.21468/SciPostPhys.12.1.033](https://doi.org/10.21468/SciPostPhys.12.1.033).
- 659 [55] V. Tarasov, *Completeness of the Bethe Ansatz for the Periodic Isotropic Heisen-*
660 *berg model*, Reviews in Mathematical Physics **30**(08), 1840018 (2018),
661 doi:[10.1142/S0129055X18400184](https://doi.org/10.1142/S0129055X18400184).
- 662 [56] R. I. Nepomechie and C. Wang, *Algebraic Bethe ansatz for singular solutions*, Journal of
663 Physics A: Mathematical and Theoretical **46**(32), 325002 (2013), doi:[10.1088/1751-](https://doi.org/10.1088/1751-8113/46/32/325002)
664 [8113/46/32/325002](https://doi.org/10.1088/1751-8113/46/32/325002).
- 665 [57] R. I. Nepomechie and C. Wang, *Twisting singular solutions of Bethe’s equations*, Journal
666 of Physics A: Mathematical and Theoretical **47**(50), 505004 (2014), doi:[10.1088/1751-](https://doi.org/10.1088/1751-8113/47/50/505004)
667 [8113/47/50/505004](https://doi.org/10.1088/1751-8113/47/50/505004).
- 668 [58] J. Riordan, *Introduction to Combinatorial Analysis*, Dover Books on Mathematics. Dover
669 Publications, Mineola, NY (2002).
- 670 [59] P. R. Giri and T. Deguchi, *Singular eigenstates in the even(odd) length heisenberg spin*
671 *chain*, Journal of Physics A: Mathematical and Theoretical **48**(17), 175207 (2015),
672 doi:[10.1088/1751-8113/48/17/175207](https://doi.org/10.1088/1751-8113/48/17/175207).

- 673 [60] P. Massignan, J. Levinsen and M. M. Parish, *Magnetism in Strongly Interact-*
674 *ing One-Dimensional Quantum Mixtures*, Phys. Rev. Lett. **115**, 247202 (2015),
675 doi:[10.1103/PhysRevLett.115.247202](https://doi.org/10.1103/PhysRevLett.115.247202).
- 676 [61] A. Novoselsky and J. Katriel, *Hyperspherical functions with arbitrary permutational sym-*
677 *metry*, Phys. Rev. A **49**, 833 (1994), doi:[10.1103/PhysRevA.49.833](https://doi.org/10.1103/PhysRevA.49.833).
- 678 [62] G. James and M. Liebeck, *Conjugacy classes*, In *Representations and Characters of Groups*
679 *(2nd ed.)*, chap. 12. Cambridge University Press, doi:[10.1017/CBO9780511814532](https://doi.org/10.1017/CBO9780511814532)
680 (2001).
- 681 [63] J. Katriel, *Representation-free evaluation of the eigenvalues of the class-sums of the sym-*
682 *metric group*, Journal of Physics A: Mathematical and General vol. **26**(4), page L135
683 (1993), doi:[10.1088/0305-4470/26/4/002](https://doi.org/10.1088/0305-4470/26/4/002).
- 684 [64] G. Pecci, G. Aupetit-Diallo, M. Albert, P. Vignolo and A. Minguzzi, *Persistent currents*
685 *in a strongly interacting multicomponent Bose gas on a ring*, Comptes Rendus. Physique
686 (2023), doi:[10.5802/crphys.157](https://doi.org/10.5802/crphys.157).
- 687 [65] Y. Cai, D. G. Allman, P. Sabharwal and K. C. Wright, *Persistent Currents*
688 *in Rings of Ultracold Fermionic Atoms*, Phys. Rev. Lett. **128**, 150401 (2022),
689 doi:[10.1103/PhysRevLett.128.150401](https://doi.org/10.1103/PhysRevLett.128.150401).
- 690 [66] G. Del Pace, K. Khani, A. Muzi Falconi, M. Fedrizzi, N. Grani, D. Hernandez Rajkov,
691 M. Inguscio, F. Scazza, W. J. Kwon and G. Roati, *Imprinting Persistent Currents in Tunable*
692 *Fermionic Rings*, Phys. Rev. X **12**, 041037 (2022), doi:[10.1103/PhysRevX.12.041037](https://doi.org/10.1103/PhysRevX.12.041037).
- 693 [67] W. J. Chetcuti, A. Osterloh, L. Amico and J. Polo, *Interference dynamics of matter-waves*
694 *of $SU(N)$ fermions*, SciPost Phys. **15**, 181 (2023), doi:[10.21468/SciPostPhys.15.4.181](https://doi.org/10.21468/SciPostPhys.15.4.181).
- 695 [68] G. Pecci, P. Naldesi, A. Minguzzi and L. Amico, *Single-particle versus many-body phase*
696 *coherence in an interacting Fermi gas*, Quantum Science and Technology **8**(1), 01LT03
697 (2022), doi:[10.1088/2058-9565/aca712](https://doi.org/10.1088/2058-9565/aca712).
- 698 [69] W. J. Chetcuti, J. Polo, A. Osterloh, P. Castorina and L. Amico, *Probe for bound states*
699 *of $SU(3)$ fermions and colour deconfinement*, Communications Physics **6**(1), 1 (2023),
700 doi:[10.1038/s42005-023-01256-3](https://doi.org/10.1038/s42005-023-01256-3), Publisher: Nature Publishing Group.
- 701 [70] A. J. Leggett, *Dephasing and Non-Dephasing Collisions in Nanostructures*, pp. 297–311,
702 Springer US, Boston, MA, doi:[10.1007/978-1-4899-3689-9_19](https://doi.org/10.1007/978-1-4899-3689-9_19) (1991).
- 703 [71] X. Waintal, G. Fleury, K. Kazymyrenko, M. Houzet, P. Schmitteckert and D. Weinmann,
704 *Persistent Currents in One Dimension: The Counterpart of Leggett’s Theorem*, Phys. Rev.
705 Lett. **101**, 106804 (2008), doi:[10.1103/PhysRevLett.101.106804](https://doi.org/10.1103/PhysRevLett.101.106804).
- 706 [72] G. Pecci, P. Naldesi, L. Amico and A. Minguzzi, *Probing the BCS-BEC*
707 *crossover with persistent currents*, Phys. Rev. Research **3**, L032064 (2021),
708 doi:[10.1103/PhysRevResearch.3.L032064](https://doi.org/10.1103/PhysRevResearch.3.L032064).
- 709 [73] H. Fredricksen and I. J. Kessler, *An algorithm for generating necklaces of beads in*
710 *two colors*, Discrete Mathematics **61**(2–3), 181–188 (1986), doi:[10.1016/0012-](https://doi.org/10.1016/0012-365x(86)90089-0)
711 [365x\(86\)90089-0](https://doi.org/10.1016/0012-365x(86)90089-0).
- 712 [74] F. Ruskey, C. Savage and T. Min Yih Wang, *Generating necklaces*, Journal of Algorithms
713 **13**(3), 414–430 (1992), doi:[10.1016/0196-6774\(92\)90047-g](https://doi.org/10.1016/0196-6774(92)90047-g).

- 714 [75] N. L. Harshman, *One-dimensional traps, two-body interactions, few-body symme-*
715 *tries: I. One, two, and three particles*, *Few-Body Systems* **57**(1), 11–43 (2015),
716 doi:[10.1007/s00601-015-1024-6](https://doi.org/10.1007/s00601-015-1024-6).
- 717 [76] N. L. Harshman, *One-dimensional traps, two-body interactions, few-body symmetries. II. N*
718 *particles*, *Few-Body Systems* **57**(1), 45–69 (2015), doi:[10.1007/s00601-015-1025-5](https://doi.org/10.1007/s00601-015-1025-5).
- 719 [77] R. L. Miller, *Necklaces, symmetries and self-reciprocal polynomials*, *Discrete Mathematics*
720 **22**(1), 25–33 (1978), doi:[10.1016/0012-365x\(78\)90043-2](https://doi.org/10.1016/0012-365x(78)90043-2).
- 721 [78] S. I. Mistakidis, A. G. Volosniev, N. T. Zinner and P. Schmelcher, *Effective approach to im-*
722 *purity dynamics in one-dimensional trapped Bose gases*, *Physical Review A* **100**(1) (2019),
723 doi:[10.1103/physreva.100.013619](https://doi.org/10.1103/physreva.100.013619).
- 724 [79] D. M. Edwards, *Magnetism in Single-Band Models: Exact One-Dimensional Wave Functions*
725 *Generalised to Higher Dimensions*, *Progress of Theoretical Physics Supplement* **101**, 453
726 (1990), doi:[10.1143/PTP101.453](https://doi.org/10.1143/PTP101.453).
- 727 [80] H. Castella and X. Zotos, *Exact calculation of spectral properties of a particle in-*
728 *teracting with a one-dimensional fermionic system*, *Phys. Rev. B* **47**, 16186 (1993),
729 doi:[10.1103/PhysRevB.47.16186](https://doi.org/10.1103/PhysRevB.47.16186).
- 730 [81] O. Gamayun, A. G. Pronko and M. B. Zvonarev, *Impurity Green’s func-*
731 *tion of a one-dimensional Fermi gas*, *Nuclear Physics B* **892**, 83 (2015),
732 doi:[10.1016/j.nuclphysb.2015.01.004](https://doi.org/10.1016/j.nuclphysb.2015.01.004).
- 733 [82] F. Brauneis, A. Ghazaryan, H.-W. Hammer and A. G. Volosniev, *Emergence of a Bose*
734 *polaron in a small ring threaded by the Aharonov-Bohm flux*, *Communications Physics*
735 **6**(1) (2023), doi:[10.1038/s42005-023-01281-2](https://doi.org/10.1038/s42005-023-01281-2).
- 736 [83] M. D. Girardeau, *Dynamics of Lieb-Liniger Gases*, *Phys. Rev. Lett.* **91**, 040401 (2003),
737 doi:[10.1103/PhysRevLett.91.040401](https://doi.org/10.1103/PhysRevLett.91.040401).
- 738 [84] R. E. Barfknecht, A. Foerster, N. T. Zinner and A. G. Volosniev, *Generation of spin currents*
739 *by a temperature gradient in a two-terminal device*, *Communications Physics* **4**(1) (2021),
740 doi:[10.1038/s42005-021-00753-7](https://doi.org/10.1038/s42005-021-00753-7).