

Stochastic Resetting and Large Deviations

Martin R. Evans^{1*} and John C. Sunil^{1†}

¹ SUPA, School of Physics and Astronomy, University of Edinburgh, Peter Guthrie Tait Road, Edinburgh, EH9 3FD, United Kingdom

* m.evans@ed.ac.uk, † j.chakanal-sunil@sms.ed.ac.uk

Abstract

Stochastic resetting has been a subject of considerable interest within statistical physics, both as means of improving completion times of complex processes such as searches and as a paradigm for generating nonequilibrium stationary states. In these lecture notes we give a self-contained introduction to the toy model of diffusion with stochastic resetting. We also discuss large deviation properties of additive functionals of the process such as the cost of resetting. Finally, we consider the generalisation from Poissonian resetting, where the resetting process occurs with a constant rate, to non-Poissonian resetting.

Copyright attribution to authors.

This work is a submission to SciPost Physics Lecture Notes.

License information to appear upon publication.

Publication information to appear upon publication.

Received Date

Accepted Date

Published Date

1

Contents

3	1 Introduction	2
4	1.1 Motivations	3
5	2 Preliminaries: Diffusion equation and solution	3
6	2.1 Diffusion equation	3
7	2.2 Solution by Laplace transform	5
8	3 Preliminaries: Diffusion with an Absorbing Target	6
9	3.1 Survival probability	6
10	3.2 Mean first passage time (MFPT)	7
11	4 Diffusion with Stochastic Resetting	8
12	4.1 Renewal equation approach	9
13	4.2 Relaxation to stationary state	10
14	5 Diffusion with Resetting and Absorbing Target	11
15	5.1 Mean first passage time	12
16	5.2 Long time behaviour of survival probability	12
17	5.3 Mini-summary	14
18	6 Large Deviations in Stochastic Resetting	14
19	6.1 Additive functional	14
20	7 The Cost of Stochastic Resetting	15

21	7.1 Additive cost function	15
22	7.2 Mean total cost	17
23	7.3 Large deviation function	18
24	8 Non-Poissonian Resetting	19
25	8.1 Survival probability	19
26	8.2 Optimising the waiting time distribution	20
27	9 Conclusion	20
28	A Inverting Laplace transform of Gaussian using the Bromwich integral	21
29	B Proof of the integral identity (15)	22
30	C Proof of the Laplace transform of the error function (24)	23
31	D Laplace's method	23
32	E Calculation of Mean Costs for Resetting	25
33	E.1 Linear cost per reset	25
34	E.2 Quadratic cost per reset	26
35	References	26

37

38 1 Introduction

39 These are the lecture notes from two lectures given at the 2024 summer school on Large Devia-
 40 tions held at Les Houches School of Physics. The lectures give an introduction to diffusion with
 41 stochastic resetting and how this relates to the general theme of the school, large deviations.

42 In recent years there has been considerable interest in stochastic resetting for a number
 43 of reasons. As we shall motivate below, resetting can speed up some complex task such as a
 44 search process by cutting off errant trajectories that go wandering off in the wrong direction.
 45 Resetting also provides a paradigm for creating a nonequilibrium state, by virtue of continually
 46 restarting the dynamics and not letting the system relax to equilibrium. Perhaps the simplest
 47 dynamical process, to which one can add a resetting process, is diffusion. Indeed, diffusion
 48 with stochastic resetting [1] has become an important toy model where an in-depth under-
 49 standing of resetting can be obtained from exact calculations of relevant quantities such as
 50 survival probabilities, mean first passage times etc.

51 The aim of these lecture notes is to provide a self-contained introduction to diffusion
 52 with stochastic resetting that will allow the reader to execute the calculations in full, using
 53 straightforward techniques such as Laplace transforms and renewal equations. These notes
 54 will hopefully be complementary to Reference [2], which is a more detailed review and con-
 55 tains a comprehensive bibliography up to 2020. The present notes also contain some more
 56 recent material. We also note that a historical perspective on resetting can be found in [3] and
 57 a brief review is given in [4].


58 These notes are organised as follows. After motivating the study of stochastic resetting
 59 in Section 1.1, we begin by reviewing the diffusion equation in Section 2 and first passage
 60 times 3 and Laplace transform techniques. We then use these techniques to study diffusion

with Poissonian resetting in Section 4 and diffusion with resetting and an absorbing target in Section 5. We discuss large deviations of additive functionals of diffusion with resetting in Section 6 and the example of the cost of resetting in Section 7. Finally we generalise to non-Poissonian resetting, defined by a waiting time distribution between resets, in Section 8.

1.1 Motivations

The process of stochastic resetting can be motivated by a simple everyday experience of searching for misplaced keys. During the search process, we periodically reset back to the place where the keys usually should be and restart the search. It turns out that such resetting expedites the completion time of the search process [1, 2]. More generally, motivations for studying stochastic resetting are

- (i) **Searching for a target:** In search processes it has been established that mixing local moves with long range moves improves the search process. This is termed as an “Intermittent search strategy” [5]. In the case of searches with stochastic resetting, the local moves are diffusion and the long range moves are the resets.
- (ii) **Expediting completion times:** More generally, it has been observed that restarting a complex task, which has a long time tail of completion times, can speed up performance. For example, consider a complicated chemical reaction [6], such as,



The above equation represents a reaction where E is the enzyme, and S is the substrate which combines to form ES through a binding rate k_{on} and then takes a random time τ (which could have long tails in its distribution) to produce the product P . The rate of unbinding of the enzyme, k_{off} , can be thought of as the resetting rate, and it turns out that having such a rate enhances the completion time of the process by cutting off the long tails in the distribution of τ .

- (iii) **Generating non-equilibrium steady states (NESS):** Resetting the system to its initial state prevents the system from relaxing to its equilibrium state. The resetting process generates a probability current back to the initial condition from all other configurations, thus ensuring detailed balance is not satisfied. This results in a simple, but non-trivial, NESS exhibiting circulation of probability.

2 Preliminaries: Diffusion equation and solution

Diffusion is perhaps the simplest dynamical process in nature. Our aim is to study how introducing resetting fundamentally changes the behaviour of a diffusive particle.

2.1 Diffusion equation

We begin by deriving the diffusion equation. Consider an over-damped diffusing particle starting at $\mathbf{x}(t = 0) = \mathbf{x}_0$, which can be represented as,

$$\mathbf{x}_{t+\Delta t} = \mathbf{x}_t + \boldsymbol{\xi}, \quad (1)$$

97 where ξ is a Gaussian white noise given by the distribution

$$P(\xi) = \frac{1}{\sqrt{4\pi D \Delta t}} e^{-\frac{\xi^2}{4D \Delta t}}.$$

98 Averaging over all possible events between t and $t + \Delta t$

$$P(x, t + \Delta t | x_0) = \langle P(x - \xi, t | x_0) \rangle \quad (2)$$

99 where the angle brackets indicates an average over the noise ξ . Expanding for small ξ

$$P(x, t + \Delta t) = \left[P(x, t) - \frac{\partial P(x, t)}{\partial x} \langle \xi \rangle + \frac{1}{2} \frac{\partial^2 P(x, t)}{\partial x^2} \langle \xi^2 \rangle + \dots \right] \quad (3)$$

$$= P(x, t) + \Delta t \left[D \frac{\partial^2}{\partial x^2} P(x, t) + \dots \right]. \quad (4)$$

100 where in (3),(4) we have suppressed the dependence on x_0 for ease of presentation. In going
101 to equation (4), we have made use of the following properties of white noise

$$\begin{aligned} \langle \xi(t) \rangle &= 0, \\ \langle \xi(t) \xi(t') \rangle &= 2D \Delta t \delta(t - t'). \end{aligned}$$

102 Dividing (4) by Δt and taking $\Delta t \rightarrow 0$, yields the diffusion equation

$$\frac{\partial}{\partial t} P(x, t | x_0) = D \frac{\partial^2}{\partial x^2} P(x, t | x_0) \quad (5)$$

103 In a similar way we can average over events between $t = 0$ and $t = \Delta t$ and write

$$P(x, t + \Delta t | x_0) = \langle P(x, t | x_0 + \xi) \rangle \quad (6)$$

104 which leads to a backward equation. For diffusion this looks the same as the forward equation,
105 where the spatial variable is now x_0

$$\frac{\partial}{\partial t} P(x, t | x_0) = D \frac{\partial^2}{\partial x_0^2} P(x, t | x_0). \quad (7)$$

106 Taking the initial condition

$$P(x, t = 0 | x_0) = \delta(x - x_0),$$

107 the solution of (5) is the familiar Gaussian

$$P(x, t | x_0) = \frac{1}{\sqrt{4\pi D t}} e^{-\frac{(x-x_0)^2}{4D t}}. \quad (8)$$

108 Perhaps the simplest method to obtain the solution (8) is to assume a scaling solution

$$P = (Dt)^{-1/2} f(z) \quad \text{where} \quad z = \frac{(x - x_0)}{(Dt)^{1/2}} \quad (9)$$

so that z is the dimensionless scaling variable and $f(z)$ is the scaling function. The prefactor $(Dt)^{-1/2}$ in P ensures normalisation of probability through

$$\int_{-\infty}^{\infty} dx P(x, t | x_0) = \int_{-\infty}^{\infty} dz f(z) = 1.$$

109 Substituting expression (9) in (5) leads to

$$-\frac{1}{2} \frac{d}{dz} [z f(z)] = \frac{d^2}{dz^2} f(z). \quad (10)$$

110 The solution which tends to zero as $z \rightarrow \infty$ and is normalised is $f(z) = (4\pi)^{-1/2} e^{-z^2/2}$.

2.2 Solution by Laplace transform

Let us also derive the solution of the diffusion equation by Laplace transform and discuss the subsequent inversion using a Bromwich contour to obtain the diffusion propagator. The reasons for choosing this method will become apparent when we move on to more complicated problems. We define the Laplace transform with respect to time for the PDF as,

$$\tilde{P}(x, s|x_0) = \int_0^\infty dt e^{-st} P(x, t|x_0). \quad (11)$$

By making use of integration by parts, we can rewrite the Laplace transform of the time derivative of $P(x, t|x_0)$ as

$$\begin{aligned} \int_0^\infty e^{-st} \frac{\partial P}{\partial t} dt &= [e^{-st} P]_0^\infty + s \int_0^\infty e^{-st} P dt \\ &= -\delta(x - x_0) + s\tilde{P}. \end{aligned}$$

The space derivative goes through unaltered as the Laplace transform only acts on the time variable. Hence we obtain the Laplace transform of the diffusion equation as

$$D \frac{\partial^2}{\partial x^2} \tilde{P}(x, s|x_0) - s\tilde{P}(x, s|x_0) = -\delta(x - x_0) \quad (12)$$

The solution of (12) is

$$\tilde{P}(x, s|x_0) = \frac{1}{\sqrt{4sD}} e^{-\sqrt{\frac{s}{D}}|x-x_0|}, \quad (13)$$

which can be verified by making use of the identity

$$\frac{d^2}{dx^2} [e^{-a|x-b|}] = -2a\delta(x-b) + a^2 e^{-a|x-b|}.$$

Finally, to obtain the solution back in the time domain, we have to invert the Laplace transform by making use of the Bromwich integral

$$P(x, t|x_0) = \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} ds e^{st} \tilde{P}(x, s|x_0), \quad (14)$$

where γ is a real number which lies to the right of all the singularities of the function to be inverted in the complex s plane. The singularity in (13) is a branch point at $s = 0$ due to the presence of $s^{1/2}$. Therefore the inversion integral for (13) has to be evaluated around a branch cut along the negative real axis, which will finally result in (8) (See Appendix A for details). Rather than performing the contour integral, a simpler method to obtain the final result is to make use of a known integral (a particular case of identity 3.174.9 from [7], see Appendix B for an elementary proof)

$$\int_0^\infty dt t^{-1/2} e^{-\frac{\beta}{t} - st} = \frac{\pi^{1/2}}{s^{1/2}} e^{-2(\beta s)^{1/2}} \quad (15)$$

and match $\beta = \frac{(x-x_0)^2}{4D}$. One then deduces that the function whose Laplace transform is (13) is (8).

Exercise: Perform the calculation both ways and verify that you obtain the diffusive propagator (8).

3 Preliminaries: Diffusion with an Absorbing Target

Now consider the situation where in addition to a diffusing particle, there is an absorbing target $[2, 8]$ at $\mathbf{x} = \mathbf{x}_T$. When the particle touches the target situated, it is absorbed, and the process is considered to be completed. We will usually take the absorbing target to be at the origin $\mathbf{x}_T = \mathbf{0}$.

3.1 Survival probability

We are interested in the probability that the particle survive up to a time t . The presence of the absorbing target can be modelled as an additional a boundary condition $P(\mathbf{x}_T, t) = 0$. using the backward Fokker-Planck approach, we obtain the following equation,

$$\frac{\partial}{\partial t} P(\mathbf{x}, t | \mathbf{x}_0) = D \frac{\partial^2}{\partial \mathbf{x}_0^2} P(\mathbf{x}, t | \mathbf{x}_0). \quad (16)$$

The survival probability $q(t | \mathbf{x}_0)$ is obtained by integrating over all final positions at time t

$$q(t | \mathbf{x}_0) = \int_0^\infty d\mathbf{x} P(\mathbf{x}, t | \mathbf{x}_0). \quad (17)$$

Integrating over \mathbf{x} from $\mathbf{0}$ to ∞ , and exchanging the order of differentiation and integration (since integration is with respect to \mathbf{x} while the derivatives are with respect to \mathbf{x}_0), we end up with an equation for the survival probability as

$$\frac{\partial}{\partial t} q(t | \mathbf{x}_0) = D \frac{\partial^2}{\partial \mathbf{x}_0^2} q(t | \mathbf{x}_0), \quad (18)$$

with the boundary condition

$$q(t | 0) = 0,$$

and initial condition

$$q(0 | \mathbf{x}_0) = 1.$$

The solution to (18) with the above mentioned boundary and initial conditions is given by

$$q(t | \mathbf{x}_0) = \operatorname{erf}\left(\frac{\mathbf{x}_0}{\sqrt{4Dt}}\right), \quad (19)$$

where the error function is defined as

$$\operatorname{erf}(z) = \frac{2}{\sqrt{\pi}} \int_0^z du e^{-u^2}. \quad (20)$$

The simplest way to obtain (19) is to assume a scaling form $q(t | \mathbf{x}_0) = g(z)$ where the scaling variable is now $z = \mathbf{x}_0 / \sqrt{4Dt}$. Substituting in (18) one obtains

$$-\frac{z}{2} \frac{dg}{dz} = \frac{d^2g}{dz^2}. \quad (21)$$

Integrating twice and using the boundary conditions $g(0) = 0$, $g(\infty) = 1$ yields (19).

Again, one can also use the method of Laplace transform to obtain this solution. The Laplace transform of (18) is given by

$$D \frac{\partial^2 \tilde{q}}{\partial x_0^2} - s \tilde{q} = -1, \quad (22)$$

with boundary conditions $\tilde{q}(s|0) = 0$ and $\tilde{q}(s|\infty) = 1/s$, which can be solved to obtain

$$\tilde{q}(s|x_0) = \frac{1 - e^{-x_0 \sqrt{\frac{s}{D}}}}{s}. \quad (23)$$

Again, (23) can be inverted using the Bromwich integral by integrating around the branch cut at $s = 0$ to obtain (19), but is easier to use the known integral

$$\int_0^\infty dt e^{-st} \operatorname{erf}\left(\frac{\beta}{t^{1/2}}\right) = \frac{1 - e^{-2\beta s^{1/2}}}{s}, \quad (24)$$

which can be derived using (15) (see Appendix C).

The great advantage of using the Laplace transform is that we can obtain the large t behaviour quite easily, without having to invert the full Laplace transform. For this we expand $\tilde{q}(s|x_0)$ for small values of s and invert it term by term using the inverse Laplace transform identity

$$\mathcal{L}_{s \rightarrow t}^{-1} \left\{ \frac{1}{s^\alpha} \right\} = \frac{1}{\Gamma(\alpha)} t^{\alpha-1}, \quad (25)$$

where the r.h.s. is zero when $\alpha = 0, -1, -2, \dots$. Following this approach we get, expanding (23)

$$\tilde{q}(s|x_0) \simeq \frac{x_0}{\sqrt{sD}} - \frac{x_0^2}{2D} + \mathcal{O}(\sqrt{s}), \quad (26)$$

$$q(t|x_0) \simeq \frac{x_0}{(\pi Dt)^{1/2}} + \mathcal{O}(t^{-3/2}). \quad (27)$$

where we have used $\Gamma(1/2) = \pi^{1/2}$.

3.2 Mean first passage time (MFPT)

The mean first passage time to the absorbing target is obtained from the survival probability as follows

$$\langle T(x_0) \rangle = \int_0^\infty dt t F(t, x_0), \quad (28)$$

where

$$F(t, x_0) = -\frac{\partial q(t|x_0)}{\partial t} \quad (29)$$

is the first passage time distribution i.e. minus the rate of change of the survival probability is the rate at which the particle is absorbed. But due to the $t^{-1/2}$ tails for the survival probability for diffusive processes as obtained in (27), $F(t, x_0) \sim t^{-3/2}$ and the integral (28) for the mean first passage time diverges. That is, the mean first passage time for a diffusive process is infinite even though the particle is absorbed with probability one!

178 **Note:** When the survival probability goes to zero faster than t^{-1} , the mean first passage time
 179 (28) can be simplified using integration by parts as

$$\begin{aligned}\langle T(x_0) \rangle &= -[q(t|x_0)t]_0^\infty + \int_0^\infty dt q(t|x_0) \\ &= \int_0^\infty dt q(t|x_0) \\ &= \tilde{q}(s=0|x_0).\end{aligned}\tag{30}$$

180 We will use this expression for $\langle T(x_0) \rangle$ later on.

181 4 Diffusion with Stochastic Resetting

182 We now consider adding Poissonian resetting to the diffusive process discussed in the previous
 183 sections [1, 2]. This is defined by a constant resetting rate (per unit time) r to a resetting site
 184 x_r . This leads to the following process:

$$x_{t+\Delta t} = \begin{cases} x_t + \xi & \text{with prob. } 1 - r\Delta t, \\ x_r & \text{with prob. } r\Delta t \end{cases}\tag{31}$$

185 As for diffusion, we first write the forward master equation for the process with resetting. We
 186 denote $P_r(x, t)$ as the probability density at time t under resetting at rate r , where we have
 187 suppressed the dependence on initial condition x_0 . Upon averaging over all possible events in
 188 between the time t to $t + \Delta t$, we get

$$\begin{aligned}P_r(x, t + \Delta t) &= r\Delta t \delta(x - x_r) + (1 - r\Delta t) \langle P_r(x - \xi, t) \rangle \\ &= r\Delta t \delta(x - x_r) + (1 - r\Delta t) \\ &\quad \times \left[P_r(x, t) - \frac{\partial P_r(x, t)}{\partial x} \langle \xi \rangle + \frac{1}{2} \frac{\partial^2 P_r(x, t)}{\partial x^2} \langle \xi^2 \rangle + \dots \right] \\ &= P_r(x, t) + \Delta t \left[r\delta(x - x_r) - rP_r(x, t) + D \frac{\partial^2}{\partial x^2} P_r(x, t) \right] + \dots,\end{aligned}$$

189 where as before the angle brackets $\langle \cdot \rangle$ indicates an average of the the noise ξ .

190 Then in the limit $\Delta t \rightarrow 0$, we obtain the equation for the evolution of distribution with
 191 resetting as

$$\frac{\partial}{\partial t} P_r(x, t) = D \frac{\partial^2}{\partial x^2} P_r(x, t) - rP_r(x, t) + r\delta(x - x_r).\tag{32}$$

192 We thus obtain a diffusion equation with two additional terms proportional to r on the r.h.s..
 193 The first represents the loss of probability with rate r from any position x , the second rep-
 194 represents the gain of probability at resetting position x_r with rate r from all other positions
 195 x .

196 As $t \rightarrow \infty$ we reach a stationary state $P_r^*(x)$ with $\frac{\partial P_r^*}{\partial t} = 0$. The forward equation becomes

$$D \frac{\partial^2}{\partial x^2} P_r^*(x, t) - rP_r^*(x, t) = -r\delta(x - x_r),\tag{33}$$

197 which is the same as equation (12) with s replaced by r and a factor of r in the r.h.s.. Thus
 198 the solution is

$$P_r^*(x) = \frac{\alpha_0}{2} e^{-\alpha_0 |x - x_r|},\tag{34}$$




Figure 1: A schematic of trajectory for a particle undergoing stochastic resetting. The particle starts at x_0 and resets to x_r . t' represents the time of last reset and τ represents the time of first reset.

199 where

$$\alpha_0 = \sqrt{\frac{r}{D}}. \quad (35)$$

200 Due to the existence of a non-zero probability current in the system with resetting, the resulting
 201 steady state is a Non-Equilibrium Steady State (NESS). We note that there is a cusp in the
 202 distribution at the resetting site x_r .

203 Similarly, to obtain the backward master equation, we consider the evolution of the particle
 204 in the first time step from 0 to Δt . We then obtain (restoring the dependence on the initial
 205 condition),

$$P_r(x, t + \Delta t | x_0) = r \Delta t P_r(x, t | x_r) + (1 - r \Delta t) \langle P_r(x, t | x_0 + \xi) \rangle, \quad (36)$$

206 which upon expanding and taking the $\Delta t \rightarrow 0$ limit leads to

$$\frac{\partial}{\partial t} P_r(x, t | x_0) = D \frac{\partial^2}{\partial x_0^2} P_r(x, t | x_0) + r P_r(x, t | x_r) - r P_r(x, t | x_0). \quad (37)$$

207 4.1 Renewal equation approach

208 Another equivalent, but simpler, method to obtain the propagator for the system with resetting
 209 is to use the renewal approach [1, 2, 9]. This approach is based on the fact that between resets,
 210 the process behaves exactly like a normal diffusion process with the resetting position as the
 211 initial condition (See Fig:1).

212 To show this more concretely, the last renewal equation can be written as:

$$P_r(x, t | x_0) = e^{-rt} P_0(x, t | x_0) + \int_0^t dt' r e^{-r(t-t')} \int_{-\infty}^{\infty} dx' P_r(x', t' | x_0) P_0(x, t - t' | x_r). \quad (38)$$

213 The first term is the product of the probability of no resets in time t , e^{-rt} , and reaching x at
 214 time t without resetting, $P_0(x, t | x_0)$. The subscript 0 just denotes usual diffusion. The second
 215 term represents the probability of reaching x at time t when there has been at least one reset.
 216 We integrate over the time of the *last* reset, t' , and $dt' r e^{-r(t-t')}$ is the probability of a reset
 217 in time $t' \rightarrow t' + dt'$ and no further resets in time $t - t'$. We also integrate over the position

218 \mathbf{x}' at time t' . However, as we have conservation of probability $\int_{-\infty}^{\infty} d\mathbf{x}' P_r(\mathbf{x}', t' | \mathbf{x}_0) = 1$.
 219 Finally, changing integration variable to $u = t - t'$ the last renewal equation becomes

$$P_r(\mathbf{x}, t | \mathbf{x}_0) = \underbrace{e^{-rt} P_0(\mathbf{x}, t | \mathbf{x}_0)}_{\text{Probability of no reset up to time } t \text{ and reaching } \mathbf{x} \text{ by usual diffusion.}} + \int_0^t \underbrace{du r e^{-ru}}_{\text{Probability of reset between } t' \text{ and } t' + dt' \text{ and no resets for the remaining time } t - t'.} \underbrace{P_0(\mathbf{x}, u | \mathbf{x}_r)}_{\text{After the last reset, the system evolves like usual diffusion starting from } \mathbf{x}_r.} . \quad (39)$$

220 To obtain the steady-state solution from the (39), we look at the $t \rightarrow \infty$ limit, which makes
 221 the first term zero, and the remaining term gives the steady state solution as

$$P^*(\mathbf{x}) = r \int_0^{\infty} du e^{-ru} P_0(\mathbf{x}, u | \mathbf{x}_r), \quad (40)$$

222 which is the Laplace transform of a Gaussian, which we derived in Section 2 Eq. (13), and
 223 yields (34).

224 4.2 Relaxation to stationary state

225 The last renewal equation may also be used to study the relaxation to the stationary state.
 226 Consider a large time t . The important term to consider in (39) is the second term. We
 227 introduce variables $w = u/t$ and $y = |\mathbf{x} - \mathbf{x}_r|/t$ so that the second term may be written as

$$r \int_0^t du e^{-ru} P_0(\mathbf{x}, u | \mathbf{x}_r) = \frac{r t^{1/2}}{\sqrt{4\pi D}} \int_0^1 \frac{dw}{w^{1/2}} \exp\left[-t \left[r w + \frac{y^2}{4Dw} \right]\right]. \quad (41)$$

228 Then the integral is in a form that may be evaluated by Laplace's method (which is a particular
 229 case of the saddle point method). Since t is large, the integrand will be sharply peaked at the
 230 maximum of the argument of the exponential, which is given by

$$w^* = \frac{y}{\sqrt{4Dr}}. \quad (42)$$

231 Then, if w^* is within the limits of integration, i.e. $w^* \leq 1$, we obtain

$$r \int_0^t du e^{-ru} P_0(\mathbf{x}, u | \mathbf{x}_r) \sim \exp\left(-t[r/D]^{1/2} y\right) \quad (43)$$

232 where we have ignored the prefactors to the exponential. But if w^* is outwith the limits of
 233 integration, i.e. $w^* > 1$, the integral is dominated by $w = 1$ and the contribution of the
 234 integral is of the same order as the first term in (39), $e^{-rt} P_0(\mathbf{x}, t | \mathbf{x}_0)$. See Appendix D for
 235 more details).

236 The condition $w^* < 1$ corresponds to $y = |\mathbf{x} - \mathbf{x}_r|/t < \sqrt{4Dr}$. Thus there is an equilibra-
 237 tion front travelling with speed $v = \sqrt{4Dr}$. When $|\mathbf{x} - \mathbf{x}_r| < vt$, $P_r(\mathbf{x}, t | \mathbf{x}_0)$ has relaxed to
 238 its stationary form, but if $|\mathbf{x} - \mathbf{x}_r| > vt$, $P_r(\mathbf{x}, t | \mathbf{x}_0)$ is still time dependent and is dominated
 239 by trajectories in which no resets have occurred or the last reset was near $t' = 0$. See [9] for
 240 a more detailed discussion.

241 We also note that, ignoring prefactors, may write for large t and $|\mathbf{x} - \mathbf{x}_r|$ with $y = |\mathbf{x} - \mathbf{x}_r|/t$

$$P_r(\mathbf{x}, t | \mathbf{x}_0) \sim \exp[-t I(|\mathbf{x} - \mathbf{x}_r|/t)] \quad (44)$$

242 where

$$I(y) = \begin{cases} (r/D)^{1/2} y & \text{when } y < \sqrt{4Dr}, \\ r + y^2/4D & \text{when } y \geq \sqrt{4Dr}. \end{cases} \quad (45)$$

243 Here $I(y)$ is our first example of a large deviation function (see [9] for more details).




Figure 2: A schematic of a trajectory that does not cross $x = 0$ and hence survives till time t . t' represents the time of last reset and τ represents the time of first reset.

244 5 Diffusion with Resetting and Absorbing Target

245 We now consider our diffusing particle under resetting as a searcher searching for the target.
 246 As in Section 3, we model the target with an absorbing site at $x_T = 0$ [1, 2, 11]. Using a
 247 last renewal approach (See Fig. 2) we can write the survival probability for the system with
 248 resetting in terms of the system without resetting as

$$q_r(t|x_0) = \underbrace{e^{-rt}q_0(t|x_0)}_{\text{Probability of surviving with no reset up to time } t} + \int_0^t \underbrace{dt' r e^{-r(t-t')}}_{\text{Probability of reset between } t' \text{ and } t' + dt' \text{ and no resets for the remaining time.}} \underbrace{q_r(t'|x_0)}_{\text{Probability of survival till } t' \text{ with resetting.}} \underbrace{q_0(t-t'|x_r)}_{\text{Probability of survival without resetting for the remaining time.}}. \quad (46)$$

249 Again there are two terms: the first is the probability of no resets and survival, the second
 250 integrates over the time t' of the last reset. Recognizing that in (46), the term inside the
 251 integral is a convolution, we can make use of the convolution theorem to obtain the Laplace
 252 transform of $q_r(t|x_0)$ as,

$$\tilde{q}_r(s|x_0) = \tilde{q}_0(s+r|x_0) + r\tilde{q}_r(s|x_0)\tilde{q}_0(s+r|x_r). \quad (47)$$

253 Then we obtain

$$\tilde{q}_r(s|x_0) = \frac{\tilde{q}_0(s+r|x_0)}{1-r\tilde{q}_0(s+r|x_r)}, \quad (48)$$

254 which upon taking the initial and reset position to be the same ie, $x_r = x_0$, becomes

$$\begin{aligned} \tilde{q}_r(s|x_0) &= \frac{\tilde{q}_0(s+r|x_0)}{1-r\tilde{q}_0(s+r|x_0)} \\ &= \frac{1-e^{-\alpha(s)x_0}}{s+r e^{-\alpha(s)x_0}}, \end{aligned} \quad (49) \quad (50)$$

255 where $\alpha(s) = \sqrt{\frac{r+s}{D}}$, and we have used (13) to obtain (50).

256

257 **Note:** Instead of a last renewal equation for the survival probability as given in (46), one could
 258 also write an equivalent first renewal equation with τ representing the time of first reset as

(See Fig. 2),

$$q_r(t|x_0) = \underbrace{e^{-rt} q_0(t|x_0)}_{\text{Probability of surviving with no reset up to time } t} + \int_0^t \underbrace{d\tau r e^{-r\tau}}_{\text{Probability of a single reset between } \tau \text{ and } \tau + d\tau} \underbrace{q_0(\tau|x_0)}_{\text{Probability of survival till } \tau \text{ without resetting.}} \underbrace{q_r(t-\tau|x_r)}_{\text{Probability of survival with resetting for the remaining time.}} . \quad (51)$$

Taking the Laplace transform of (51) then yields an alternative equation for the Laplace transforms

$$\tilde{q}_r(s|x_0) = \tilde{q}_0(s+r|x_0) + r \tilde{q}_0(r+s|x_0) \tilde{q}_r(s|x_r), \quad (52)$$

which reduces to (47) when $x_0 = x_r$.

5.1 Mean first passage time

Using (30), the MFPT, calculated from (50), reduces to the simple expression

$$\begin{aligned} \langle T_r(x_0) \rangle &= \tilde{q}_r(s=0|x_0) \\ &= \frac{e^{\alpha_0 x_0} - 1}{r}. \end{aligned} \quad (53)$$

Expression (54) has diverging behaviour for both the small r and large r limits,

- For $r \ll 1$: $\langle T_r(x_0) \rangle \rightarrow \frac{x_0}{\sqrt{Dr}}$.
- For $r \gg 1$: $\langle T_r(x_0) \rangle \rightarrow \frac{e^{\sqrt{\frac{r}{D}} x_0}}{r}$.

The limit $r \rightarrow 0$ recovers the diverging diffusive mean first passage time discussed in Section 3. The limit $r \rightarrow \infty$ implies resetting to x_r with infinite rate so that the particle is unable to diffuse away and find the target. In between these two limits the MFPT has an optimal value of resetting which minimizes the MFPT. This optimal value, r^* , can be obtained by evaluating $\frac{d\langle T_r(x_0) \rangle}{dr} = 0$, which results in

$$\frac{y}{2} = 1 - e^{-y}, \quad (55)$$

where $y = \frac{x_0}{(D/r)^{1/2}}$. y is a dimensionless parameter, which is the ratio of the two length scales in the system: x_0 , the distance to the target and $(D/r)^{1/2}$, the typical distance diffused between resets. Equation (55) has a unique non-zero solution given by $y^* = 1.5936\dots$, that minimizes the MFPT for searches with resetting.

Note that here the optimisation of the MFPT assumes that we know the distance to the target x_0 . Obviously, this requires significant information. Nevertheless, the calculation demonstrates the principle that the MFPT can be optimised by tuning the resetting rate. Also we see that any (finite) resetting rate performs better than the diffusive process without resetting.

5.2 Long time behaviour of survival probability

It is difficult to invert the Laplace transform of the survival probability for all time t . For an integral expression see [12]. However, as in the diffusive case discussed in Section 3, the asymptotic behaviour can be easily obtained.

The asymptotic behaviour of the survival probability (50) can be obtained from the singularities of the Laplace transform [1, 2]. The singularity furthest to the right (with largest real part) in the complex s plane dominates the long term behaviour. The denominator of (50) has two singularities:

289 (i) Branch point at $s = -r$.

(ii) Pole at

$$s_0 + r \exp\left(-\sqrt{\frac{r+s}{D}}x_0\right) = 0.$$

290 Since $s_0 > -r$, the pole is the furthest singularity to the right and the contribution from the
291 pole determines the leading order behaviour,

$$q(t|x_0) \underset{t \gg 1}{\simeq} A e^{s_0 t}, \quad (56)$$

292 where the constant A is the residue from the pole, which may be calculated to be

$$A = \frac{1 + s_0/r}{1 + s_0 x_r / (4(r + s_0)D)^{1/2}}. \quad (57)$$

293 If $|s_0| \ll 1$ then,

$$s_0 \simeq -r e^{-y}, \quad (58)$$

294 where $y = \sqrt{\frac{r}{D}}x_0$, which in turn implies $y \gg 1$. Then we have

$$q(t|x_0) \simeq e^{-r t e^{-y}}, \quad (59)$$

295 which is of the form of the Gumbel distribution.

296 Aside on Gumbel distribution

297 Consider a situation where N iid random variables are drawn from a PDF, $f(x)$. We are
298 interested in finding the probability that the maximum of these N random numbers is less
299 than a given number M . That is,

$$\mathbb{P}(\max_{1 \leq i \leq N} X_i \leq M) = [\mathbb{P}(X_1 \leq M)]^N = \left[\int_{-\infty}^M f(x) dx \right]^N = \exp\left(N \ln \left[1 - \int_M^{\infty} f(x) dx \right]\right). \quad (60)$$

300 Now if $f(x)$ has an exponential tail, $f(x) \sim A e^{-ax}$ for $x \gg 1$, we obtain the Gumbel distri-
301 bution from (60),

$$\mathbb{P}(\max_{1 \leq i \leq N} X_i \leq M) \approx \exp\left[-\frac{NA}{a} e^{-aM}\right]. \quad (61)$$

302 In the case of the survival of a diffusive particle under stochastic resetting, we have a similar
303 situation [2, 11]. Between each reset the particle must survive, that is, its maximum excursion
304 to the left must be less than x_0 . The maximum excursion between resets $i-1$ and i , is a random
305 variable that we'll call X_i , and due to renewal the X_i are independent random variables. For
306 survival through N resets we must have $X_i < x_0$ for $i = 1, \dots, N$. That is, the maximum of
307 the X_i must be less than x_0 which is the problem described above. We obtain the probability
308 that $X_i < x_0$ by averaging the survival probability for a diffusive particle $q_0(t|x_0)$ over the
309 duration of each excursion. Then, with the expected number of resets rt in the long time
310 limit, we obtain the Gumbel distribution through

$$\begin{aligned} q_r(t|x_0) &\approx \left[\int_0^{\infty} dt r e^{-rt} q_0(t|x_0) \right]^{rt} && \text{(Using (23))} \\ &= \left[1 - e^{-x_0 \sqrt{\frac{r}{D}}} \right]^{rt} \\ &\simeq e^{-r t e^{-y}}. \end{aligned} \quad (62)$$

5.3 Mini-summary

So far we have seen how stochastic resetting changes the behaviour of diffusion with an absorbing target. First, the mean first passage time is rendered finite with the introduction of resetting whereas it is infinite without resetting. Second, the survival probability decays as an exponential for large time rather than as a power law, as is the case without resetting. The form of the exponential decay can be understood from extreme value statistics. As an aside, we note that in some circumstances resetting can also generate a non-trivial power decay of the survival probability [13]. We now turn to the subject of this summer school: large deviations.

6 Large Deviations in Stochastic Resetting

6.1 Additive functional

We are interested in calculating the large deviations of an additive functional for a process with resetting [14–16]. Consider a path with N resets. Let $f(x_t)$ be an additive functional of the trajectory, using which we can define

$$\begin{aligned} A_t &= \int_0^t dt f(x_t) \\ &= \sum_{i=1}^{N+1} A_{(t_n - t_{n-1})}, \end{aligned} \quad (63)$$

where t_n is the time of the n^{th} reset and we have defined $t_0 = 0$ and $t_{N+1} = t$. For simplicity we will assume $f \geq 0$ so that $A_t \geq 0$ (although more general cases can be considered [14–16]).

We expect a large deviation principle to hold with the form

$$P_r \left(\frac{A_t}{t} = a, t \right) \sim \exp(-t I(a)), \quad (64)$$

where $I(a)$ is the large deviation function, sometimes referred to as the rate function. Using the first renewal formalism, we can write an equation for $P_r(A_t, t)$ as

$$P_r(A_t, t) = e^{-rt} P_0(A_t, t) + \int_0^t d\tau e^{-r\tau} r \left(\int_0^{A_t} dA_\tau P_0(A_\tau, \tau) P_r(A - A_\tau, t - \tau) \right), \quad (65)$$

where we have used the additive property of the functional to write a convolution over A_τ . Define a generating function (which in the case of $A_t \geq 0$ is simply a Laplace transform over A_t)

$$G_r(k, t) = \langle e^{-kA_t} \rangle = \int_0^\infty dA_t e^{-kA_t} P_r(A_t, t), \quad (66)$$

using which we can rewrite (65), making use of the convolution theorem as

$$G_r(k, t) = e^{-rt} G_0(k, t) + \int_0^t d\tau r e^{-r\tau} G_0(k, \tau) G_r(k, t - \tau). \quad (67)$$

Now define the Laplace transform of $G_r(A_t, t)$ as

$$\tilde{G}_r(k, s) = \int_0^\infty dt e^{-st} G_r(k, t), \quad (68)$$

334 from which we obtain again, making use of the convolution theorem,

$$\tilde{G}_r(k, s) = \tilde{G}_0(k, s + r) + r \tilde{G}_0(k, s + r) \tilde{G}_r(k, s), \quad (69)$$

335 which can be rearranged to obtain,

$$\tilde{G}_r(k, s) = \frac{\tilde{G}_0(k, s + r)}{1 - r \tilde{G}_0(k, s + r)}. \quad (70)$$

336 In (70), we have obtained the generating function for the system with resetting in terms of the
 337 generating function for the system without resetting. (70) has a pole at $1 - r \tilde{G}_0(k, s + r) = 0$,
 338 which we denote by $s_0(k, r)$. So upon inverting, we obtain the leading behaviour of $G_r(k, t)$
 339 as

$$G_r(k, t) \sim e^{s_0(k, r)t}. \quad (71)$$

340 Then upon a second inversion with respect to k , we obtain (ignoring sub-exponential terms),

$$P_r(A_t, t) \sim \int dk \exp\left(t \left[\frac{A_t}{t} k + s_0(k, r)\right]\right), \quad (72)$$

341 which is dominated by the saddle point ,

$$\frac{A_t}{t} + \frac{d}{dk} s_0(k, r) = 0. \quad (73)$$

342 So we obtain the large deviation form for $P_r(A_t, t)$ as

$$P_r(A, t) \sim e^{tI(a)} \quad (74)$$

343 where $I(a)$ is the Legendre-Fenchel transform of s_0

$$I(a) = -\sup_k (ka + s_0(k, r)), \quad (75)$$

344 with $a = \frac{A_t}{t}$.

345 7 The Cost of Stochastic Resetting

346 In the model so far, we have assumed that the resets are instantaneous and cost free, but this
 347 assumption is unrealistic as resets must come at a price. To rectify this and make the model
 348 more applicable, we introduce a cost to each of the resets in the model [17, 18], which may
 349 also be thought of as time penalties.

350 We consider here the simple case of a diffusing particle under resetting with cost but no
 351 absorbing target [18]. A different ensemble of trajectories is to consider the cost of resetting
 352 up to absorption by a target at the origin [17].

353 7.1 Additive cost function

354 We consider additive costs that occur at the resets (See Fig. 3) of the following form,

$$C = \sum_{i=1}^N c(|x_i|), \quad (76)$$

355 where we take x_i as the position just before the reset and the resetting position as the origin.
 356 The resetting costs are functionals of resetting jumps. Example costs include:




Figure 3: For simplicity, we set the particle to start and reset to $x = 0$. Cost is incurred at the jumps denoted by x_i . τ represents the time of first reset.

- A constant cost $c(x) = c$. This can be thought of as a fixed charge or time penalty such as refractory period for each reset [19].
- Linear cost per reset: $c(x) = \frac{|x|}{V}$. This can be interpreted as the time to return to the origin at a constant speed V . [20]
- Quadratic cost per reset $c(x) = x^2$. This is related to “thermodynamics of resetting” [21].
- Exponential cost per reset: $c(x) = e^{|x|}$. This could be used to model situation where resets from a large distance are highly unfavourable. [17]

We are interested in calculating the statistics of the total cost, such as the mean $\langle C \rangle$ for the process up to a fixed time t and also the large deviation function.

To begin with the calculation for the mean total cost, we first write down a first renewal equation for $P_r(C, t)$, which is the joint PDF of having incurred a total cost C at time t . We obtain the first renewal equation as

$$P_r(C, t) = \delta(C) e^{-rt} + \int_0^t d\tau r e^{-r\tau} \int_{-\infty}^{\infty} dx P_0(x, \tau) P_r(C - c(x), t - \tau) \Theta(C - c(|x_i|)), \quad (77)$$

where $\Theta(C - c(|x_i|))$ ensures that the total cost is kept positive. The first term accounts for trajectories with no resets and the second term integrates over trajectories where the first reset occurs at time τ .

We then define a generating function

$$G_r(k, t) = \int_0^{\infty} dC e^{-kC} P_r(C, t). \quad (78)$$

Using the definition of the generating function on (77), we obtain after a few steps of calculation,

$$G_r(k, t) = e^{-rt} + \int_0^t d\tau r e^{-r\tau} \int_{-\infty}^{\infty} dx P_0(x, \tau) e^{-kc(|x|)} G_r(k, t - \tau). \quad (79)$$

376 Finally taking a Laplace transform of (79), we get

$$\tilde{G}_r(k, s) = \frac{1}{s+r} + r \int_{-\infty}^{\infty} dx \tilde{P}_0(x, r+s) e^{-kc(|x|)} \tilde{G}_r(k, s), \quad (80)$$

377 which can be rewritten as

$$\tilde{G}_r(k, s) = \frac{1}{s+r} \frac{1}{\left[1 - r \int_{-\infty}^{\infty} dx \tilde{P}_0(x, r+s) e^{-kc(|x|)}\right]}. \quad (81)$$

378 Equation (81) gives an exact expression for the Laplace transform of the generating function
379 of the cost, from which exact expressions for the moments of the cost, can be computed.

380 7.2 Mean total cost

381 For example, the mean total cost can then be obtained as

$$\langle C \rangle = \mathcal{L}_{s \rightarrow t}^{-1} \left[-\partial_k \tilde{G}_r(k, s) \Big|_{k \rightarrow 0^+} \right] \quad (82)$$

$$= \mathcal{L}_{s \rightarrow t}^{-1} \left[\frac{2r(r+s)}{s^2} \int_0^{\infty} dx \tilde{P}_0(x, r+s) c(|x|) \right]. \quad (83)$$

382 Performing these calculations for the linear and quadratic cost per reset (see Appendix E), leads
383 to the following mean total costs, which is written in terms of the dimensionless resetting rate
384 $R = r t$.

- 385 • Linear cost per resetting: $c(x) = |x|$

$$\langle C \rangle_{\text{lin}} = \sqrt{Dt} \left(\frac{e^{-R}}{\sqrt{\pi}} + \frac{(2R-1) \text{erf}(\sqrt{R})}{2\sqrt{R}} \right). \quad (84)$$

386 For large values of t (and therefore R), this increases as, $\langle C \rangle_{\text{lin}} \simeq \sqrt{D r t}$.

- 387 • Quadratic cost per resetting: $c(x) = |x|^2$

$$\langle C \rangle_{\text{quad}} = \frac{2Dt(R + e^{-R} - 1)}{R}. \quad (85)$$

388 For large values of t , the mean total quadratic cost increases as $\langle C \rangle_{\text{quad}} \simeq 2Dt$. In-
389 terestingly, this does not depend on r and therefore is finite in the limit $r \searrow 0$. The
390 explanation is that in this limit the probability of a reset is $O(rt)$ but the typical cost of
391 a reset is $O(1/r)$, which results in an $O(t)$ contribution to the mean cost as $r \searrow 0$.

392 The large time behaviours can be explained by the following scaling argument. The distri-
393 bution of the position x at a reset is given by integrating the Gaussian distribution over the
394 waiting time distribution between resets

$$P(x) \simeq \int_0^{\infty} dt \frac{r e^{-rt}}{\sqrt{4\pi Dt}} \exp(-x^2/4Dt) \quad (86)$$

$$= \frac{\alpha_0}{2} e^{-\alpha_0 |x|}, \quad (87)$$

where we have used our trusty formula (13), and as before $\alpha_0 = (r/D)^{1/2}$. Then, for large times, the mean cost will be the mean number of resets rt multiplied by the mean cost per reset. For the case of $c(|x|) = |x|^\beta$ this gives

$$\langle C \rangle = rt \int_0^\infty dx \alpha_0 e^{-\alpha_0 x} x^\beta \quad (88)$$

$$= \frac{\Gamma(\beta + 1)rt}{\alpha_0^\beta}, \quad (89)$$

which precisely recovers the long time limits of the exact results above for the case $\beta = 1, 2$.

7.3 Large deviation function

The large deviation function of the cost can be obtained from (81) following the approach of Section 6. Here, we work out the case of linear cost per reset $c = |x|$.

The integral in (81) becomes

$$\begin{aligned} r \int_{-\infty}^\infty dx \tilde{P}_0(x, r+s) e^{-kc(|x|)} &= r \int_{-\infty}^\infty dx \frac{1}{\sqrt{4(r+s)D}} e^{-\sqrt{\frac{r+s}{D}}|x|} e^{-k|x|} \\ &= \frac{r}{r+s+k\sqrt{D(r+s)}} \end{aligned} \quad (90)$$

and we obtain

$$\tilde{G}_r(k, s) = \frac{r+s+kD^{1/2}(r+s)^{1/2}}{(r+s)(s+kD^{1/2}(r+s)^{1/2})}. \quad (91)$$

It is easiest to invert the Laplace transform over k first as the only singularity is a simple pole at $k = -s/(D(r+s))^{1/2}$. Therefore we obtain

$$\tilde{P}_r(C, s) = \frac{r}{(r+s)^{3/2}D^{1/2}} \exp\left(-\frac{sC}{D^{1/2}(r+s)^{1/2}}\right). \quad (92)$$

We now invert the Laplace transform over s by using the Bromwich contour

$$P_r(C, t) = \frac{1}{2\pi i} \int_{-i\infty+\gamma}^{+i\infty+\gamma} ds e^{st} \tilde{P}_r(C, s) \quad (93)$$

where γ should be to the right of the singularity at $s = -r$. Setting $C = at$, with t large, we evaluate the inversion integral over s by the saddle point method,

$$P_r\left(\frac{C}{t} = a, t\right) \sim \int_{-i\infty+s_0}^{+i\infty+s_0} ds \exp\left(-t \left[\frac{as}{D^{1/2}(r+s)^{1/2}} - s \right]\right), \quad (94)$$

where we have chosen $\gamma = s_0$ so that the contour goes through the saddle point s_0 , in a direction parallel to the imaginary axis, and for simplicity we have ignored prefactors to the exponential. The saddle point equation for s_0 reduces to

$$r + s_0 = -r + \frac{2D^{1/2}}{a}(r + s_0)^{3/2}, \quad (95)$$

which has a unique real solution with $(r + s_0) > 0$.

Therefore $P_r(C/t = a, t) \sim \exp(-tI(a))$ where the large deviation function is

$$I(a) = \frac{as_0}{D^{1/2}(r + s_0)^{1/2}} - s_0, \quad (96)$$

where $s_0(a, r)$ is the solution of (95). Note that for $a = (Dr)^{1/2}$ the solution of (95) is $s_0 = 0$ and $I(a = (Dr)^{1/2}) = 0$, which is consistent with the mean cost $\langle C \rangle \simeq (Dr)^{1/2}t$, as we saw in Section 7.2.

417 8 Non-Poissonian Resetting

418 So far we have assumed that the resetting is Poissonian, i.e., we have assumed that the waiting
 419 time between resets is distributed exponentially. We now relax this assumption and assume
 420 a general waiting time distribution $\psi(t)$ [22, 23]. This means that a reset occurs, in time
 421 $t \rightarrow t + dt$ after the last reset, with probability $\psi(t)dt$. We also define the probability of no
 422 resets up to time t as

$$\Psi(t) = \int_t^\infty d\tau \psi(\tau). \quad (97)$$

423 Note that we recover all the results for Poissonian resetting if we set $\psi(t) = r e^{-rt}$. Another
 424 situation that is also of interest is the case of deterministic reset (or sharp restart) which is given
 425 by $\psi(t) = \delta(t - t_r)$.

426 Similarly to Section 4.1, in the absence of an absorbing target and with $x_0 = x_r$, we can
 427 write the propagator for system with resetting using a first renewal equation as

$$P_r(x, t|x_0) = \Psi(t)P_0(x, t|x_0) + \int_0^t d\tau \psi(\tau)P_r(x, t - \tau|x_0). \quad (98)$$

428 Then using Laplace transform we get,

$$\begin{aligned} \tilde{P}_r(x, s|x_0) &= \frac{\int_0^\infty dt e^{-st} \Psi(t)P_0(x, t|x_0)}{1 - \int_0^\infty dt e^{-st} \psi(t)} \\ &= \frac{\int_0^\infty dt e^{-st} \Psi(t)P_0(x, t|x_0)}{s \int_0^\infty dt e^{-st} \Psi(t)}, \end{aligned} \quad (99)$$

429 where the last equality is obtained by performing integration by parts on the denominator. The
 430 stationary state will then be given by the coefficient of the $\frac{1}{s}$ term in the small s expansion.
 431 So we obtain the stationary state to be

$$P^*(x) = \lim_{s \rightarrow 0} \frac{\int_0^\infty dt e^{-st} \Psi(t)P_0(x, t|x_0)}{\int_0^\infty dt e^{-st} \Psi(t)}. \quad (100)$$

432 Thus, for the NESS to exist, this limit should not vanish. Therefore, we require

$$\int_0^\infty \Psi(t)dt = \int_0^\infty \tau \psi(\tau)d\tau = \mathbb{E}(\tau) < \infty. \quad (101)$$

433 That is, for a stationary state to exist for diffusion under resetting we require the mean time
 434 between resets to be finite.

435 8.1 Survival probability

436 Let us also consider the survival probability with an arbitrary waiting time distribution. As we
 437 did in Section 5, we can write down a first renewal equation for the survival probability as
 438 follows:

$$q_r(t|x_0) = \Psi(t)q_0(t|x_0) + \int_0^t d\tau \psi(\tau)q_0(\tau|x_0)q_r(t - \tau|x_r). \quad (102)$$

Then, as usual, taking the Laplace transform and setting $\mathbf{x}_0 = \mathbf{x}_r$ yields

$$\tilde{q}_r(s|\mathbf{x}_0) = \frac{\int_0^\infty dt e^{-st} \Psi(t) q_0(t|\mathbf{x}_0)}{1 - \int_0^\infty d\tau e^{-s\tau} \psi(\tau) q_0(\tau|\mathbf{x}_0)}. \quad (103)$$

Finally, we obtain the MFPT by setting $s = 0$

$$\begin{aligned} \langle T_r(\mathbf{x}_0) \rangle &= \frac{\int_0^\infty dt \psi(t) \int_0^t d\tau q_0(\tau|\mathbf{x}_0)}{1 - \int_0^\infty d\tau \psi(\tau) q_0(\tau|\mathbf{x}_0)} \\ &= \frac{\int_0^\infty dt \psi(t) \int_0^t d\tau q_0(\tau|\mathbf{x}_0)}{\int_0^\infty d\tau \psi(\tau) (1 - q_0(\tau|\mathbf{x}_0))}. \end{aligned} \quad (104)$$

8.2 Optimising the waiting time distribution

In Section 5.1, for the case of Poissonian resetting, we optimised the MFPT to an absorbing target, with respect to the resetting rate r . More generally, we can ask, what is the best choice of waiting time distribution to minimize the mean time for first passage?

It turns out that when the target is at a known distance from the resetting site (in the case of resetting to \mathbf{x}_0 and a target at the origin this distance is \mathbf{x}_0) the answer to this question is that a deterministic resetting, $\psi(t) = \delta(t - t_r)$ is the best [23–25]. But the resetting period, t_r , has to be chosen appropriately and relies on us knowing the distance to the target.

A more realistic problem is when we know where the target ought to be (at the origin say) but the actual distance from this position is a random variable, \mathbf{x}_T , drawn from a target distribution, $\mathcal{P}_T(\mathbf{x}_T)$ centred at the origin [26]. Thus we do not know the position of the target exactly. A sensible strategy for searching for the target would be to reset to the origin and select an optimal waiting time distribution for resetting. The relevant quantity to optimise is then the MFPT averaged over the target position, \mathbf{x}_T ,

$$\overline{\langle T_r(\mathbf{x}_0) \rangle} = \int d\mathbf{x}_T \mathcal{P}_T(\mathbf{x}_T) \langle T_r(\mathbf{x}_T) \rangle \quad (105)$$

where $\langle T_r(\mathbf{x}_T) \rangle$ is given by (104) with \mathbf{x}_0 replaced by \mathbf{x}_T .

Although this optimisation problem is beyond the scope of these lectures, we mention that the optimal waiting time distribution depends on the target distribution. For example, in the case of an exponential target distribution $\mathcal{P}_T(\mathbf{x}_T) = \beta/2 \exp(-\beta|\mathbf{x}_T|)$ the optimal waiting time distribution is exponential, i.e. Poissonian resetting with rate $r = \beta^2 D/4$ [27].

9 Conclusion

In these lectures we have presented a detailed account of diffusion with stochastic resetting and highlighted some large deviation principles that apply. We have endeavoured to make the lecture notes self-contained as far as possible so that an interested reader can learn the tricks of the trade. The field of stochastic resetting is constantly expanding and evolving and these notes, whilst not aiming to provide comprehensive coverage of all recent developments, will hopefully provide an entry point to this rich and complex topic.

Acknowledgements

The authors thank the organisers of the Les Houches School on *Theory of Large Deviations and Applications*. M.R.E. thanks Satya Majumdar, Baruch Meerson and Grégory Schehr for helpful feedback on the lectures. J.C.S. thanks the University of Edinburgh for the award of Edinburgh Doctoral College Scholarship. For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

A Inverting Laplace transform of Gaussian using the Bromwich integral

The task is to evaluate the inverse Laplace transform using the integral obtained from (13) and (14),

$$P(x, t|x_0) = \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} ds e^{st} \frac{1}{\sqrt{4sD}} e^{-\sqrt{\frac{s}{D}}|x-x_0|}. \quad (\text{A.1})$$

To evaluate the Bromwich integral (A.1), we integrate along the contour Γ , shown in Fig. 4,




Figure 4: We define the keyhole contour Γ composed of $C_\gamma, C_1, C_a, C_\epsilon, C_b$ and C_2 . C_γ is a straight segment that vertically intersects the real axis with all the singularities lying to its left. Here $\gamma > 0$ and s runs from $s = \gamma - iR$ to $s = \gamma + iR$. C_ϵ is a small circular contour around the branch point singularity at $s = 0$. In the limit $R \rightarrow \infty$ and $\epsilon \rightarrow 0$, we see that the only contributions to the integral are from C_γ, C_a and C_b , using which we can calculate the required Bromwich integral.

where we have a keyhole construction around the branch point at $s = 0$. From Cauchy's theorem, since there are no singularities inside the contour, we obtain

$$\begin{aligned} \frac{1}{2\pi i} \oint_{\Gamma} ds e^{st} \frac{1}{\sqrt{4sD}} e^{-\sqrt{\frac{s}{D}}|x-x_0|} \\ = \frac{1}{2\pi i} \left(\int_{C_\gamma} + \int_{C_1} + \int_{C_a} + \int_{C_\epsilon} + \int_{C_b} + \int_{C_2} \right) dz e^{zt} \frac{1}{\sqrt{4zD}} e^{-\sqrt{\frac{z}{D}}|x-x_0|} = 0. \end{aligned} \quad (\text{A.2})$$

Since we eventually take the limits $R \rightarrow \infty$ and $\epsilon \rightarrow 0$, the contribution from the integral from C_1, C_2 and C_ϵ goes to zero, as a consequence of Jordan's lemma. So we obtain the required

483 integral as

$$\frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} ds e^{st} \frac{1}{\sqrt{4sD}} e^{-\sqrt{\frac{s}{D}}|x-x_0|} = -\frac{1}{2\pi i} \left(\int_{C_a} + \int_{C_b} \right) dz e^{st} \frac{1}{\sqrt{4sD}} e^{-\sqrt{\frac{s}{D}}|x-x_0|}. \quad (\text{A.3})$$

484 To evaluate the integral along C_a , we choose $s = re^{i\theta}$ with $\theta = \pi$, with the limits of the
485 integral running from $r = \infty$ to $r = 0$, which gives us

$$\begin{aligned} \int_{C_a} ds e^{st} \frac{1}{\sqrt{4sD}} e^{-\sqrt{\frac{s}{D}}|x-x_0|} &= \int_{\infty}^0 dr e^{i\pi} \frac{\exp(re^{i\pi}t)}{\sqrt{4rDe^{i\pi/2}}} \exp\left(-\sqrt{\frac{r}{D}}e^{i\pi/2}|x-x_0|\right) \\ &= -i \int_0^{\infty} dr \frac{\exp(-rt)}{\sqrt{4rD}} \exp\left(-i\sqrt{\frac{r}{D}}|x-x_0|\right) \\ &= -\frac{2i}{\sqrt{4Dt}} \int_0^{\infty} du e^{-u^2} e^{\frac{iu}{\sqrt{Dt}}|x-x_0|} \\ &= -\frac{2i}{\sqrt{4Dt}} \int_0^{\infty} du e^{-\left(u - \frac{i|x-x_0|}{\sqrt{4Dt}}\right)^2} e^{-\frac{|x-x_0|^2}{4Dt}} \\ &= -i \sqrt{\frac{\pi}{4Dt}} e^{-\frac{|x-x_0|^2}{4Dt}}. \end{aligned} \quad (\text{A.4})$$

486 To perform the calculations we have used a substitution $u^2 = rt$, followed by completion of
487 the squares and Gaussian integral.

488 Similarly, to evaluate the integral over C_b , we choose $s = re^{i\theta}$ with $\theta = -\pi$, with the
489 limits from $r = 0$ to $r = \infty$. Performing exactly the same steps used to obtain (A.4), we get
490 the same result

$$\int_{C_b} ds e^{st} \frac{1}{\sqrt{4sD}} e^{-\sqrt{\frac{s}{D}}|x-x_0|} = -i \sqrt{\frac{\pi}{4Dt}} e^{-\frac{|x-x_0|^2}{4Dt}}. \quad (\text{A.5})$$

491 Plugging in (A.4) and (A.5) into (A.3), we obtain the inverse Laplace transform

$$P(x, t|x_0) = \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} ds e^{st} \frac{1}{\sqrt{4sD}} e^{-\sqrt{\frac{s}{D}}|x-x_0|} = \frac{1}{\sqrt{4\pi Dt}} e^{-\frac{|x-x_0|^2}{4Dt}}. \quad (\text{A.6})$$

492 B Proof of the integral identity (15)

493 In this appendix we show how to obtain

$$\int_0^{\infty} dt t^{-1/2} e^{-\frac{\beta}{t}-st} = \frac{\pi^{1/2}}{s^{1/2}} e^{-2(\beta s)^{1/2}}. \quad (\text{B.1})$$

494 Note that the above integral converges only when $s > 0$ and $\beta > 0$. We start by defining

$$I = \int_0^{\infty} dt t^{-1/2} e^{-\frac{\beta}{t}-st}, \quad (\text{B.2})$$

495 Using the properties of a Gaussian integral, we can rewrite the integrand in (B.2) as

$$t^{-1/2} e^{-\frac{\beta}{t}} = \pi^{-1/2} \int_{-\infty}^{\infty} du \exp\{(-tu^2 + 2i\sqrt{\beta}u)\}, \quad (\text{B.3})$$

496 using which we obtain,

$$\begin{aligned} I &= \pi^{-1/2} \int_{-\infty}^{\infty} du \exp(2i\sqrt{\beta}u) \int_0^{\infty} dt \exp(-[u^2 + s]t) \\ &= \pi^{-1/2} \int_{-\infty}^{\infty} du \frac{\exp(2i\sqrt{\beta}u)}{u^2 + s}. \end{aligned} \quad (\text{B.4})$$

497 The last integral can be evaluated using a contour integral along a contour C that runs along
498 the real axis, and is closed in the upper half plane since $\beta > 0$. (This is a consequence of
499 Jordan's lemma.) Thus we obtain,

$$\begin{aligned} I &= \pi^{-1/2} \oint_C dz \frac{\exp(2i\sqrt{\beta}z)}{(z - i\sqrt{s})(z + i\sqrt{s})} \\ &= 2\pi^{1/2} i \operatorname{Res}_{z=i\sqrt{s}} \frac{\exp(\pm 2i\sqrt{\beta}z)}{(z - i\sqrt{s})(z + i\sqrt{s})}, \end{aligned} \quad (\text{B.5})$$

500 Upon evaluating the residue, we obtain

$$I = \frac{2\sqrt{\pi}i \exp(-2\sqrt{\beta}s)}{2i\sqrt{s}} = \frac{\pi^{1/2}}{s^{1/2}} e^{-2(\beta s)^{1/2}}. \quad (\text{B.6})$$

501 C Proof of the Laplace transform of the error function (24)

502 Using the definition of error function (20), we can re-write (24) as,

$$\begin{aligned} \int_0^{\infty} dt e^{-st} \operatorname{erf}\left(\frac{\beta}{t^{1/2}}\right) &= \frac{2}{\sqrt{\pi}} \int_0^{\infty} dt e^{-st} \int_0^{\frac{\beta}{\sqrt{t}}} du e^{-u^2} \\ &= \frac{2}{\sqrt{\pi}} \int_0^1 dw \int_0^{\infty} dt t^{-1/2} e^{-st} e^{-\frac{\beta^2 w^2}{t}}, \end{aligned} \quad (\text{C.1})$$

503 where for the second equality we have performed the substitution, $w = \frac{\sqrt{t}}{\beta}u$ and changed the
504 order of the integration. Now we observe that the second integral in (C.1) is of the form (15)
505 once we make the identification $\beta \rightarrow (\beta w)^2$, after which we obtain

$$\begin{aligned} \int_0^{\infty} dt e^{-st} \operatorname{erf}\left(\frac{\beta}{t^{1/2}}\right) &= \frac{2\beta}{s^{1/2}} \int_0^1 dw e^{-2\beta w s^{1/2}} \\ &= \frac{1 - e^{-2\beta s^{1/2}}}{s}. \end{aligned} \quad (\text{C.2})$$

506 D Laplace's method

507 In this appendix we give the details of evaluating (39) by Laplace's method. Using $w = u/t$
508 and $y = |x - x_r|/t$, we can rewrite the second term of (39) as

$$P_r(x, t|x_0) = \frac{e^{-t\phi(1,y)}}{\sqrt{4\pi Dt}} + \frac{r t^{1/2}}{\sqrt{4\pi D}} \int_0^1 \frac{dw}{w^{1/2}} e^{-t\phi(w,y)} \quad (\text{D.1})$$

509 where

$$\phi(w, y) = rw + \frac{y^2}{4Dw} \quad (D.2)$$

510 Since we are interested in the large time behaviour, we can evaluate (D.1) using Laplace
511 method, where we expand the integrand around the maxima of the argument of the expo-
512 nential and integrate. In (D.2) it occurs at the minimum of $\phi(w, y)$, which we obtain as

$$w^* = \frac{y}{\sqrt{4Dr}}. \quad (D.3)$$

513 We get two different large time behaviours for (D.1) depending on whether $w^* < 1$ or $w^* \geq 1$.
514 For the first case, choose $w^* < 1$, which is the case where the minimum of $\phi(w, y)$ is
515 within the interval $[0, 1]$. Then, the dominant contribution will be from the integral in (D.1).
516 We can evaluate the term by expanding $\phi(w, y)$ to the second order in w

$$\phi(w, y) = \phi(w^*, y) + \frac{\phi''(w^*, y)}{2}(w - w^*)^2 + \mathcal{O}(w^3). \quad (D.4)$$

517 Since w^* is a minimum of $\phi(w, y)$, we have $\phi'(w^*, y) = 0$ and $\phi''(w^*, y) > 0$. Using the
518 substitution

$$s = \sqrt{\frac{t\phi''(w^*, y)}{2}}(w - w^*), \quad (D.5)$$

519 we obtain

$$\frac{rt^{1/2}}{\sqrt{4\pi D}} \int_0^1 \frac{dw}{w^{1/2}} e^{-t\phi(w, y)} \simeq \frac{re^{-t\phi(w^*, y)}}{\sqrt{2\pi D w^* \phi''(w^*, y)}} \int_{-\infty}^{\infty} dw e^{-s^2}. \quad (D.6)$$

520 Note that we can safely take the upper and lower limits to $\pm\infty$ as we are interested in the
521 large t behaviour and the contributions from the values different from w^* will be exponentially
522 small. Evaluating all the terms, we obtain

$$P_r(x, t|x_0) \simeq \frac{1}{2} \sqrt{\frac{r}{D}} e^{-\sqrt{\frac{r}{D}} yt}, \quad (D.7)$$

523 which is the same as the steady-state distribution obtained for diffusion with resetting in (34).
524 Note that the contribution from the first term in (D.1) is sub-leading in t as it will be of the
525 order of $e^{-t\phi(1, y)}/\sqrt{t}$.

526 In the second case, when $w^* \geq 1$, which occurs when the minimum of $\phi(w, y)$ is outside
527 the interval $(0, 1)$, the major contribution of the integral will be from the boundary close to 1.
528 So the relevant expansion of $\phi(w, y)$ is

$$\phi(w, y) = \phi(1, y) + \phi'(1, y)(w - 1) + \mathcal{O}((w - 1)^2). \quad (D.8)$$

529 Note that in this case $\phi'(1, y) < 0$. We can approximate the second term in (D.1), by evalu-
530 ating the integral in the interval $[1 - \epsilon, 1]$. We then obtain

$$\begin{aligned} \frac{rt^{1/2}}{\sqrt{4\pi D}} \int_0^1 \frac{dw}{w^{1/2}} e^{-t\phi(w, y)} &\simeq -\frac{re^{-\sqrt{4\pi D} t \phi(1, y)}}{t\phi'(1, y)} \int_0^{-t\phi'(1, y)\epsilon} ds e^{-s} \\ &\simeq -\frac{re^{-t\phi(1, y)}}{\sqrt{4\pi D} t \phi'(1, y)} \int_0^{\infty} ds e^{-s} \\ &= -\frac{e^{-t\phi(1, y)}}{\sqrt{4\pi D} t \phi'(1, y)}. \end{aligned} \quad (D.9)$$

To obtain the above expression we have used the substitution $s = t\phi'(1, y)(w - 1)$, and changed the upper limit to ∞ as we are interested in the large t behaviour. We see that unlike the previous case, the contribution from the integral and the first term in (D.1) are of the same order. Thus for $w^* \geq 1$, we obtain

$$P_r(x, t|x_0) \simeq \left(\frac{y^2}{y^2 - 4Dr} \right) \frac{e^{-t\left(r + \frac{y^2}{4D}\right)}}{\sqrt{4\pi Dt}}. \quad (\text{D.10})$$

E Calculation of Mean Costs for Resetting

In the main text we derived

$$\langle C \rangle = \mathcal{L}_{s \rightarrow t}^{-1} \left[\frac{2r(r+s)}{s^2} \int_0^\infty dx \tilde{P}_0(x, r+s) c(|x|) \right], \quad (\text{E.1})$$

where

$$\tilde{P}_0(x, r+s) = \frac{1}{\sqrt{4D(r+s)}} e^{-(r+s)/D)^{1/2} x}. \quad (\text{E.2})$$

E.1 Linear cost per reset

For the case of linear cost per resetting, plugging in $c(x) = |x|$ we obtain

$$\int_0^\infty dx \tilde{P}_0(x, r+s) |x| = \frac{D^{1/2}}{2(r+s)^{3/2}}. \quad (\text{E.3})$$

Therefore we obtain from (E.1),

$$\langle C \rangle_{\text{lin}} = r D^{1/2} \mathcal{L}_{s \rightarrow t}^{-1} \left[\frac{1}{s^2 \sqrt{r+s}} \right]. \quad (\text{E.4})$$

Noting that

$$\mathcal{L}_{s \rightarrow t}^{-1} [s^{-2}] = t, \quad (\text{E.5})$$

and

$$\mathcal{L}_{s \rightarrow t}^{-1} [(r+s)^{-1/2}] = \frac{e^{-rt}}{\sqrt{\pi t}}, \quad (\text{E.6})$$

we can make use of the convolution theorem to obtain the product of the two Laplace transforms as a convolution, which gives

$$\begin{aligned} \mathcal{L}_{s \rightarrow t}^{-1} \left[\frac{1}{s^2 \sqrt{r+s}} \right] &= \int_0^t d\tau (t-\tau) \frac{e^{-r\tau}}{\sqrt{\pi\tau}} \\ &= \frac{e^{-rt} \sqrt{t}}{\sqrt{\pi r}} + \frac{(2rt-1) \text{erf}(\sqrt{rt})}{2r^{3/2}}. \end{aligned} \quad (\text{E.7})$$

To obtain the above integral, we have made the substitution $\tau = u^2$ and made use of the definition of the error function (20). Upon plugging (E.7) into (E.4) and defining $R = rt$, we obtain

$$\langle C \rangle_{\text{lin}} = \sqrt{Dt} \left(\frac{e^{-R}}{\sqrt{\pi}} + \frac{(2R-1) \text{erf}(\sqrt{R})}{2\sqrt{R}} \right). \quad (\text{E.8})$$

E.2 Quadratic cost per reset

For the case of quadratic cost per reset, $c(\mathbf{x}) = |\mathbf{x}|^2$, following exactly the same procedure, we obtain

$$\int_0^\infty d\mathbf{x} \tilde{P}_0(\mathbf{x}, \mathbf{r} + \mathbf{s}) |\mathbf{x}|^2 = \frac{D}{(\mathbf{r} + \mathbf{s})^2}. \quad (\text{E.9})$$

Therefore we obtain from (E.1),

$$\langle C \rangle_{\text{quad}} = 2Dr \mathcal{L}_{s \rightarrow t}^{-1} \left[\frac{1}{s^2(\mathbf{r} + \mathbf{s})} \right]. \quad (\text{E.10})$$

Splitting (E.10) into partial fractions, and inverting term by term and again defining $\mathbf{R} = \mathbf{r} \mathbf{t}$, we obtain,

$$\begin{aligned} \langle C \rangle_{\text{quad}} &= 2Dr \mathcal{L}_{s \rightarrow t}^{-1} \left[+\frac{1}{rs^2} + \frac{1}{r^2(\mathbf{r} + \mathbf{s})} - \frac{1}{r^2s} \right] \\ &= \frac{2Dt(\mathbf{R} + e^{-\mathbf{R}} - 1)}{\mathbf{R}}. \end{aligned} \quad (\text{E.11})$$

References

- [1] M. R. Evans and S. N. Majumdar, *Diffusion with stochastic resetting*, Phys. Rev. Lett. **106**, 160601 (2011), doi:10.1103/PhysRevLett.106.160601.
- [2] M. R. Evans, S. N. Majumdar and G. Schehr, *Stochastic resetting and applications*, J. Phys. A: Math.Theor. **53**, 193001 (2020), doi:10.1088/1751-8121/ab7cfe.
- [3] M. Montero, A. Masó-Puigdellosas, J. Villarroel, *Continuous-time random walks with reset events: historical background and new perspectives*, Eur. Phys. J. B **90**, 176 (2017), <https://doi.org/10.1140/epjb/e2017-80348-4>
- [4] S. Gupta and A. M. Jayannavar, *Stochastic resetting: A (very) brief review*, Frontiers in Physics **10**, 789097 (2022), doi:10.3389/fphy.2022.789097
- [5] O. Bénichou, C. Loverdo, M. Moreau and R. Voituriez, *Intermittent search strategies*, Reviews of Modern Physics, **83**, 81 (2011) doi:10.1103/RevModPhys.83.81
- [6] S. Reuveni, *Optimal Stochastic Restart Renders Fluctuations in First Passage Times Universal*, Phys. Rev. Lett. **116**, 170601 (2016), doi:10.1103/PhysRevLett.116.170601.
- [7] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series and Products*, New York:Academic (1980), doi:10.1016/C2010-0-64839-5.
- [8] S. Redner, *A guide to First-Passage Processes* Cambridge University Press, Cambridge (2001), doi:10.1017/CBO9780511606014.
- [9] S. N. Majumdar, S. Sabhapandit and G. Schehr, *Dynamical transition in the temporal relaxation of stochastic processes under resetting*, Phys. Rev. E **91**, 052131 (2015), doi:10.1103/PhysRevE.91.052131.
- [10] M. R. Evans and S. N. Majumdar, *Run and tumble particle under resetting: a renewal approach*, J. Phys. A: Math. Theor. **51**, 475003 (2018), doi:10.1088/1751-8121/aae74e.
- [11] M. R. Evans and S. N. Majumdar, *Diffusion with resetting in arbitrary spatial dimension*, J. Phys. A: Math. Theor. **47**, 285001 (2014), doi:10.1088/1751-8113/47/28/285001.

- [12] P. Singh and A. Pal, *First-passage Brownian functionals with stochastic resetting*, *J. Phys. A: Math. Theor.* **55**, 234001 (2022), doi:10.1088/1751-8121/ac677c.
- [13] M. R. Evans, S.N. Majumdar, G. Schehr, *An exactly solvable predator prey model with resetting*, *J. Phys. A: Math. Theor.* **55**, 274005 (2022) doi:10.1088/1751-8121/ac7269
- [14] J. M. Meylahn, S. Sabhapandit, and H. Touchette, *Large deviations for Markov processes with resetting*, *Phys. Rev. E* **92**, 062148 (2015), doi:PhysRevE.92.062148.
- [15] F. Den Hollander, S. N. Majumdar, J. M. Meylahn, H. Touchette, *Properties of additive functionals of Brownian motion with resetting*, *J. Phys. A: Math. Theor.* **52**, 175001 (2019), doi:10.1088/1751-8121/ab0efd
- [16] C. Monthus, *Large deviations for Markov processes with stochastic resetting: analysis via the empirical density and flows or via excursions between resets*, *J. Stat. Mech.* **2021**, 033201 (2021), doi:10.1088/1742-5468/abdeaf.
- [17] J. C. Sunil, R. A. Blythe, M. R. Evans and S. N. Majumdar, *The cost of stochastic resetting*, *J. Phys. A: Math. Theor.* **56**, 395001 (2023), doi:10.1088/1751-8121/acf3bb
- [18] J. C. Sunil, R. A. Blythe, M. R. Evans and S. N. Majumdar, *Minimizing the Profligacy of Searches with Reset*, *Phys. Rev. E.* **110**, 054122 (2024), doi:10.1103/PhysRevE.110.054122.
- [19] M. R. Evans and S. N. Majumdar, *Effects of refractory period on stochastic resetting*, *J. Phys. A: Math. Theor.* **52**, 01LT01 (2019), doi:10.1088/1751-8121/aaf080
- [20] A. Pal, L. Kuśmierz, S. Reuveni, *Search with home returns provides advantage under high uncertainty*, *Phys. Rev. Res.* **2**, 043174 (2020), doi:10.1103/PhysRevResearch.2.043174
- [21] F. Mori, K. S. Olsen, S. Krishnamurthy, *Entropy production of resetting processes*, *Phys. Rev. Res.* **5**, 023103 (2023), doi:10.1103/PhysRevResearch.5.023103
- [22] S. Eule and J. J. Metzger, *Non-equilibrium steady states of stochastic processes with intermittent resetting*, *New J. Phys.* **18**, 033006 (2016), doi:10.1088/1367-2630/18/3/033006.
- [23] A. Pal, A. Kundu and M. R. Evans, *Diffusion under time-dependent resetting*, *J. Phys. A: Math. Theor.* **49**, 225001 (2016), doi:10.1088/1751-8113/49/22/225001.
- [24] A. Pal and S. Reuveni, *First Passage under Restart*, *Phys. Rev. Lett.* **118**, 030603 (2017), doi:10.1103/PhysRevLett.118.030603.
- [25] A. Chechkin and I. M. Sokolov, *Random Search with Resetting: A Unified Renewal Approach*, *Phys. Rev. Lett.* **121**, 050601 (2018), doi:10.1103/PhysRevLett.121.050601.
- [26] M. R. Evans and S. N. Majumdar, *Diffusion with optimal resetting*, *J. Phys. A: Math. Theor.* **44**, 435001 (2011), doi:10.1088/1751-8113/44/43/435001.
- [27] M. R. Evans and S. Ray 2024, *Stochastic resetting prevails over sharp restart for broad target distribution*, arXiv preprint arXiv:2410.01941 (2024), doi:10.48550/arXiv.2410.01941.