

Generalized symmetry enriched criticality in (3+1)d

Benjamin Moy

*Department of Physics and Anthony J. Leggett Institute for Condensed Matter Theory,
University of Illinois Urbana-Champaign, Urbana, Illinois 61801, USA*

July 30, 2025

Abstract

We construct two classes of continuous phase transitions in 3+1 dimensions between phases that break distinct generalized global symmetries. Our analysis focuses on $SU(N)/\mathbb{Z}_N$ gauge theory coupled to N_f flavors of Majorana fermions in the adjoint representation. For N even and sufficiently large odd N_f , upon imposing time-reversal symmetry and an $SO(N_f)$ flavor symmetry, the massless theory realizes a quantum critical point between two gapped phases: one in which a \mathbb{Z}_N one-form symmetry is completely broken and another where it is broken to \mathbb{Z}_2 , leading to $\mathbb{Z}_{N/2}$ topological order. We provide an explicit lattice model that exhibits this transition. The critical point has an enhanced symmetry, which includes non-invertible analogues of time-reversal symmetry. Enforcing a non-invertible time-reversal symmetry and the $SO(N_f)$ flavor symmetry, for N and N_f both odd, we demonstrate that this critical point can appear between a topologically ordered phase and a phase that spontaneously breaks the non-invertible time-reversal symmetry, furnishing an analogue of deconfined quantum criticality for generalized symmetries.

Contents

1	Introduction	3
2	Topological phases in pure Yang-Mills	7
2.1	$SU(N)$ gauge theory	7
2.2	$SU(N)/\mathbb{Z}_N$ gauge theory	11
3	$SU(N)$ adjoint QCD: SPT transition	15
3.1	Global symmetries	16
3.2	Massive phases: SPTs	18
3.3	Phase transition	19
4	$SU(N)/\mathbb{Z}_N$ adjoint QCD: SET transition	20
4.1	Global symmetries	21
4.2	\mathbb{Z}_N topological order	22
4.3	$\mathbb{Z}_{N/2}$ topological order	24
4.4	String tension critical exponent	26
4.5	Topological transition on the lattice	28
5	SET to non-invertible SSB transition	33
5.1	Non-invertible time-reversal	34
5.2	SET phase	35
5.3	Non-invertible SSB	37
5.4	Critical exponents	39
6	Discussion	40
A	Twisted BF theory	42
B	Charge conjugation and Majorana condition	45
C	Thermal response for fermions	46
D	Other gauge groups	48

1 Introduction

The Landau theory of phases and phase transitions is a cornerstone of twentieth-century physics. In this framework, phases are distinguished by the global symmetries they preserve or spontaneously break, as detected by the expectation value of a local order parameter. Continuous transitions are associated with universal critical exponents that depend on the symmetries and number of dimensions of the system but not on its microscopic details.

Much of modern condensed matter physics is dedicated to phases and transitions that are not captured by Landau theory. Topologically ordered phases do not have any local order parameter but are characterized by long-range entanglement and excitations with fractional statistics [1, 2]. There also exist symmetry-protected topological phases (SPTs), which do not have topological order or symmetry breaking but nonetheless cannot be deformed into a trivial product state as long as a given symmetry is preserved [3–8]. These states are characterized by their response to background gauge fields or by the ’t Hooft anomalies of their boundary states. Similarly, with a global symmetry, topological orders may be distinct as symmetry-enriched topological phases (SETs) [9, 10]. They may have a different response to background gauge fields that probe the symmetry or have distinct patterns of symmetry fractionalization—the excitations can carry different fractional quantum numbers of the symmetry.

While substantial progress has been made on topological phases themselves, a general theory of phase transitions between them is lacking. An important first step in developing such a theory is to identify examples of these continuous topological transitions. Some examples of continuous transitions between phases with distinct topological orders are well known in (2+1)d. For example, gauging the \mathbb{Z}_2 global symmetry of the (2+1)d Ising transition leads to the confinement-deconfinement transition of \mathbb{Z}_2 lattice gauge theory [1, 11]. However, in (3+1)d continuous transitions between phases with different topological orders remain elusive since most of these transitions turn out to be first order [12–15]. Although topological transitions may still arise from gauging the discrete symmetry of an ordinary symmetry-breaking transition [16, 17], examples beyond these are rare. Therefore, it is of great importance to identify models with continuous transitions between phases with distinct topological orders in (3+1)d.

A possible source of insight on this problem comes from generalized global symmetries [18–26]. Two kinds of generalized symmetries will be important for us. The first is higher-form symmetry. The symmetry operators for an ordinary global symmetry are topological, unitary (or antiunitary), are supported on codimension one manifolds, and act on

local operators. In contrast, the operators for a q -form global symmetry are topological but are defined on codimension $q+1$ manifolds and act on operators supported on q -dimensional manifolds. Another generalization, non-invertible symmetry, has symmetry operators that are topological but have no inverse and are thus not unitary. An ordinary global symmetry is then said to be an invertible zero-form symmetry.

Generalized symmetries can be useful since this perspective allows us to draw analogies with conventional spontaneous symmetry breaking (SSB).¹ For example, Abelian topological orders may be viewed as spontaneously breaking an emergent discrete invertible one-form global symmetry.² In $(2+1)d$, the braiding of two Abelian anyons results in a phase factor, which can be interpreted as the action of a one-form symmetry associated with the worldline of one anyon acting on the other. Similarly, non-Abelian topological orders arise from the breaking of emergent non-invertible one-form symmetries.

Topological phase transitions can thus be viewed as part of a broader effort to understand transitions between phases that break different generalized global symmetries. This perspective has led to the development of an analogue of Ginzburg-Landau theory for invertible one-form symmetries known as mean string field theory (MSFT) [27]. While MSFT accurately captures phases and topological defects, it is difficult to use this framework to analyze phase transitions reliably, primarily because string observables have many more degrees of freedom than local operators.

In this work, we present two families of continuous phase transitions between phases that spontaneously break different generalized global symmetries. Our analysis focuses on $PSU(N) = SU(N)/\mathbb{Z}_N$ gauge theory coupled to N_f odd flavors of Majorana fermions in the adjoint representation, a theory we refer to as $PSU(N)$ adjoint QCD. Imposing an $SO(N_f)$ flavor symmetry and time-reversal symmetry constrains the allowed deformations to a single relevant operator: a common mass term for all fermions. Tuning the mass m can then induce a transition between distinct gapped phases. Similar constructions in $SU(N)$ gauge theory, which partially inspired this work, were found to host unconventional SPT transitions [28] and symmetry-breaking transitions [29, 30].

¹Of course there are also important differences. In realistic condensed matter systems, generalized symmetries are often either emergent or appear only in fine-tuned limits.

²To define SSB for a zero-form symmetry, we perturb the system with a local symmetry-breaking field $h(x)$ and examine whether the expectation value of the order parameter is nonzero if we first take the thermodynamic limit and then $h(x) \rightarrow 0$. This definition does not generalize to one-form symmetries since the observables are nonlocal. Instead, the appropriate criterion involves the expectation value of pairs of oppositely oriented loop operators wrapping nontrivial cycles; nonvanishing correlation at large separation indicates spontaneous symmetry breaking.

For sufficiently large N_f , when the fermions are massless, the theory will be either an interacting conformal field theory (CFT) [31, 32] or an infrared (IR) free theory, with the gauge coupling becoming small at low energies. The IR free theories are typically of limited interest to high energy theorists since their IR dynamics are simple by construction, and the gauge coupling becomes large in the ultraviolet (UV), rendering the field theory ill-defined at high enough energy scales. However, this issue is unimportant from the perspective of a low energy physicist, who regards the $PSU(N)$ adjoint QCD theory as an effective field theory valid near a phase transition, just as ϕ^4 theory describes the (3+1)d Ising transition. Moreover, in the IR free case, since the gauge coupling becomes small in the IR, we can be sure that there is a direct continuous transition at $m = 0$ and can easily compute critical exponents reliably. For this reason, we focus on the IR free theories in this work, but we note that the topological transitions we observe may also occur for a theory with a smaller value of N_f with an interacting CFT at the transition.

For $m > 0$, at energies far below the fermion mass, we can integrate out the fermions to obtain pure $PSU(N)$ gauge theory with vanishing theta angle. This phase has a \mathbb{Z}_N magnetic one-form symmetry, which becomes spontaneously broken at low energies, leading to \mathbb{Z}_N topological order (i.e., \mathbb{Z}_N toric code topological order). The point-like anyons are magnetic monopoles of the $PSU(N)$ gauge theory, but since we view the $PSU(N)$ gauge theory as an emergent theory of a condensed matter system, we emphasize that these quasiparticles need not be magnetic monopoles of the electromagnetic field.

For N even and N_f odd, the $m < 0$ phase flows to $PSU(N)$ gauge theory with³ $\theta = \pi N N_f$. As we will show, the theta term induces partial breaking of the \mathbb{Z}_N one-form symmetry to \mathbb{Z}_2 , resulting in $\mathbb{Z}_{N/2}$ topological order enriched by the $SO(N_f)$ flavor symmetry and time-reversal symmetry. For $N \geq 4$, this transition is between SETs with different topological orders. The $N = 2$ case, corresponding to gauge group $PSU(2) \cong SO(3)$, is a transition between a phase with \mathbb{Z}_2 topological order and an SPT state, providing a novel example of a continuous confinement-deconfinement transition. To our knowledge, this class of models is the first in (3+1)d with an exact one-form symmetry that have an unambiguously continuous transition between phases with different one-form symmetry breaking.

When the fermions are massless, the symmetry of $PSU(N)$ adjoint QCD is enhanced by a family of non-invertible time-reversal symmetries [33, 34], which are associated with topological operators T_n that act on the $PSU(N)$ theta angle as $\theta \rightarrow -\theta + 2\pi n$, where n is an integer mod N . For $n \in N\mathbb{Z}$, this transformation reduces to the usual invertible time-reversal symmetry. This observation raises a natural question: What phases arise if

³The normalization of the $PSU(N)$ theta term is such that θ has periodicity $2\pi N$.

we perturb the critical point by operators that preserve both the flavor symmetry and a non-invertible time-reversal symmetry \mathbb{T}_n ?

Once again, the symmetries permit only a single relevant local operator that can tune a continuous transition between two gapped phases. An important non-perturbative constraint indicating a potentially rich phase diagram is that, for N odd and $n > 0$ such that $\gcd(N, n) > 1$, the theory exhibits an anomaly: no single ground state can preserve both the \mathbb{Z}_N magnetic one-form symmetry and the non-invertible time-reversal symmetry. Indeed, when these anomaly conditions are met and N_f is odd, we show that the relevant deformation controls a transition between a phase with topological order and a phase that spontaneously breaks the non-invertible time-reversal symmetry. In the special case $n = 0$, the time-reversal symmetry is invertible, and the transition will be between a phase with \mathbb{Z}_N topological order and a phase that breaks time-reversal symmetry spontaneously.

To illustrate this kind of transition in a class of models for which the time-reversal symmetry is non-invertible, we take $N = k^2$ and $n = 2k$ with odd $k > 1$ for concreteness. At low energies, where the fermions may be integrated out, one phase becomes pure $PSU(N)$ Yang-Mills at $\theta = \pi n$, and the other phase has $\theta = \pi(n + NN_f)$. The $\theta = 2\pi k$ phase preserves the non-invertible time-reversal symmetry but spontaneously breaks the \mathbb{Z}_{k^2} magnetic one-form symmetry to \mathbb{Z}_k , leading to \mathbb{Z}_k topological order. In contrast, the $\theta = \pi(2k + k^2 N_f)$ phase retains the full one-form symmetry but spontaneously breaks the non-invertible time-reversal symmetry, resulting in non-invertible domain walls that interpolate between distinct one-form SPT states. In both phases, the $SO(N_f)$ flavor symmetry remains unbroken. This transition can be viewed as an analogue of deconfined quantum criticality [35–37] but for generalized symmetries. Recent works have identified unconventional critical points in (1+1)d involving phases that spontaneously break non-invertible symmetries [38–40]. Our construction introduces a novel example in (3+1)d, expanding the landscape of unconventional critical phenomena.

We proceed as follows. To keep this work self-contained, we begin in Section 2 by reviewing the topological phases that occur in $SU(N)$ and $SU(N)/\mathbb{Z}_N$ gauge theories, and we discuss an SPT transition in $SU(N)$ adjoint QCD in Section 3. We then combine these ingredients in Section 4 to find a continuous topological transition in $SU(N)/\mathbb{Z}_N$ adjoint QCD, which in general is a transition between SETs that have different topological orders. In Section 5, we introduce the critical point between a topologically ordered phase and a phase that spontaneously breaks a non-invertible zero-form symmetry. We conclude in Section 6 with a discussion of our results and possible future directions. Additional details and technical background are included in the appendices.

2 Topological phases in pure Yang-Mills

Before we discuss non-Abelian gauge theories coupled to matter in the adjoint representation, we review the topological phases that occur in pure $SU(N)$ and $PSU(N) = SU(N)/\mathbb{Z}_N$ gauge theory since these phases will be realized at low energies in adjoint QCD when the fermions are massive. As we will explain below, pure $SU(N)$ gauge theory with a theta term is a \mathbb{Z}_N one-form SPT at $\theta \in 2\pi\mathbb{Z}$. Correspondingly, $PSU(N)$ gauge theory at $\theta \in 2\pi\mathbb{Z}$ is characterized at low energies by a particular TQFT, which can describe a topologically ordered phase depending on θ .

2.1 $SU(N)$ gauge theory

We start by reviewing the physics of pure $SU(N)$ gauge theory in (3+1)d with a theta term. The action is

$$S_{\text{YM}}[\theta] = -\frac{1}{g^2} \int \text{Tr}(f \wedge \star f) + \frac{\theta}{8\pi^2} \int \text{Tr}(f \wedge f), \quad (2.1)$$

where $f = da - i a \wedge a$ is the field strength for the dynamical $SU(N)$ gauge field a_μ and the trace is taken in the fundamental representation. A Wilson loop represents the worldline of a probe electric quark-antiquark pair, and can be defined for any representation \mathcal{R} of $SU(N)$ as

$$W_{\mathcal{R}}(\gamma) = \text{Tr}_{\mathcal{R}} \mathcal{P} \exp \left(i \oint_{\gamma} a \right), \quad (2.2)$$

where \mathcal{P} denotes path ordering and γ is a closed loop in spacetime. For the purposes of topological physics, the electric loop operators are organized into classes labeled the center of the gauge group. Thus, for $SU(N)$, which has a center of \mathbb{Z}_N , these classes are labeled by an integer $q_e \bmod N$ [41, 42]. For example, the fundamental representation has $q_e = 1$, but more generally, q_e is the number of boxes (mod N) for the Young tableau diagram of the representation \mathcal{R} . The integer q_e is defined mod N because of screening; for instance, a Wilson loop in the adjoint representation can be opened and end on the field strength.

We can also introduce 't Hooft operators [43],

$$T_{q_m}(\gamma, \Omega), \quad (2.3)$$

which represent worldlines of magnetic probe particles characterized by an integer charge $q_m \bmod N$ [41, 42, 44]. While the Wilson loop is a genuine loop operator, a generic 't Hooft operator is actually supported on a loop γ that must be attached to an open surface Ω such that $\gamma = \partial\Omega$. Physically, this surface Ω is the worldsheet of a Dirac string that is detectable

by Wilson loops. Upon canonical quantizing the theory, at equal times the Wilson loop in the fundamental representation and the 't Hooft operator obey [43]

$$W_F(\gamma) T_{q_m}(\gamma', \Omega) = e^{2\pi i q_m \varphi(\gamma, \gamma')/N} T_{q_m}(\gamma', \Omega) W_F(\gamma), \quad (2.4)$$

where γ and $\gamma' = \partial\Omega$ are loops in space and $\varphi(\gamma, \gamma')$ is their linking number. The Dirac string is thus detectable unless $q_m = 0 \bmod N$. Indeed, for a gauge theory with gauge group G , the 't Hooft operators that are genuine loop operators (i.e., do not require a choice of surface Ω) are classified by the first homotopy group $\pi_1(G)$, which is trivial for $G = SU(N)$.

Combining Wilson loops and 't Hooft operators allows us to form a more general class of dyonic loop operators,

$$D_{(q_e, q_m)}(\gamma, \Omega) = W_{\mathcal{R}}(\gamma) T_{q_m}(\gamma, \Omega), \quad (2.5)$$

which carry both electric and magnetic charge. We denote their charges collectively as (q_e, q_m) . These operators are genuine loop operators in $SU(N)$ gauge theory only if $q_m = 0 \bmod N$.

The pure $SU(N)$ gauge theory has a global \mathbb{Z}_N electric one-form symmetry, also known as the “center symmetry”, that acts on Wilson loops. The symmetry operator is a surface operator $U_{q_m}(\Sigma)$, where Σ is a closed surface and q_m is an integer mod N . Physically, $U_{q_m}(\Sigma)$ can be thought of as the insertion of the worldsheet of a background magnetic flux tube. This operator may be constructed by taking the limit of the 't Hooft operator $T_{q_m}(\gamma, \Sigma)$ as $\gamma = \partial\Sigma$ is shrunk to a point so that Σ is a closed surface. This surface operator acts on Wilson loops as

$$\langle U_{q_m}(\Sigma) W_{\mathcal{R}}(\gamma) \dots \rangle = e^{2\pi i q_e q_m \Phi(\gamma, \Sigma)/N} \langle W_{\mathcal{R}}(\gamma) \dots \rangle, \quad (2.6)$$

where $\Phi(\gamma, \Sigma)$ is the linking number in spacetime of the loop γ and the closed surface Σ , and the \dots denote insertions of other operators.

Pure $SU(N)$ gauge theory is believed to be gapped and non-degenerate at zero temperature for any $\theta \neq (2k+1)\pi$ with $k \in \mathbb{Z}$. Moreover, electric charges are confined, as signaled by the area law for a Wilson loop in the fundamental representation [45],

$$\langle W_F(\gamma) \rangle \sim e^{-\sigma \text{Area}(\gamma)}, \quad (2.7)$$

where $\text{Area}(\gamma)$ is the area of the minimal surface that bounds the loop γ and σ is a constant called the string tension. The area law signals that the \mathbb{Z}_N electric one-form symmetry is unbroken.

While the 't Hooft operator is not a genuine loop operator, it can still be used as a probe to characterize the phase of the gauge theory. At $\theta = 0$, where conventional confinement is expected, the basic 't Hooft operator (i.e., with $q_m = 1$) has a perimeter law [43],

$$\langle T_1(\gamma, \Omega) \rangle \sim e^{-\rho \text{Length}(\gamma)}, \quad (2.8)$$

where $\text{Length}(\gamma)$ is the length of the loop γ and ρ is a constant. This perimeter law is consistent with the notion of confinement arising from the condensation of magnetic monopoles [43, 46–48]. Specifically, because monopoles with charge $q_m = N$ are screened, we view this phase as arising from the condensation of charge N monopoles.

The parameter θ is 2π periodic (on a spin manifold), so the spectrum of the theory at $\theta = 2\pi k$, where $k \in \mathbb{Z}$, is the same as at $\theta = 0$. However, the θ parameter gives an electric polarization charge to magnetic monopoles—a phenomenon known as the Witten effect [49]. After a change in theta by $\Delta\theta = 2\pi k$, a dyon that originally had charges (q_e, q_m) becomes

$$(q_e, q_m) \rightarrow (q_e + k q_m, q_m). \quad (2.9)$$

As argued by 't Hooft, the Witten effect can lead to phases with condensed dyons known as oblique confinement [50, 51], which, as we discuss below, can lead to rich topological physics [18, 52–60]. Since the 't Hooft operators have a perimeter law at $\theta = 0$, a dyon with charges (q_e, q_m) at $\theta = 0$ has a perimeter law at $\theta = 2\pi k$ if

$$(q_e + k q_m, q_m) = (0, q_m) \mod N \quad (2.10)$$

so that these loops are purely magnetic once the Witten effect is taken into account. At $\theta = 2\pi k$, we then have

$$\langle D_{(-k q_m, q_m)}(\gamma, \Omega) \rangle \sim e^{-\tilde{\rho} \text{Length}(\gamma)}, \quad (2.11)$$

where we use the convention of labeling the dyon charges prior to the Witten effect (i.e., we label by the charges of dyons at $\theta = 0$). The dyon condensed at $\theta = 2\pi k$ likewise has charges $(-Nk, N)$.

Another important way to characterize an oblique confining phase is by its response upon coupling to a background field that probes the \mathbb{Z}_N one-form symmetry. Following Ref. [61], we first embed the original $SU(N)$ gauge field a_μ into a $U(N)$ gauge field α_μ , which has a field strength $(f_\alpha)_{\mu\nu}$. We then use a two-form Lagrange multiplier $\beta_{\mu\nu}$ to constrain $\text{Tr}(f_\alpha)$ to be trivial so that the theory is still $SU(N)$ gauge theory. Introducing a background $U(1)$ two-form gauge field $B_{\mu\nu}$ and a background $U(1)$ one-form gauge field C_μ such that

$dC = NB$, we then couple these background fields so that the partition function is invariant under the gauge transformations,

$$B \rightarrow B + d\lambda, \quad C \rightarrow C + N\lambda + d\xi, \quad \alpha \rightarrow \alpha - \lambda \mathbb{I}_N, \quad (2.12)$$

where ξ is a 2π periodic scalar field and λ_μ is a $U(1)$ one-form gauge field. Upon coupling to these probes, the action is now

$$\begin{aligned} S_{\text{YM}}[\theta, B] = & -\frac{1}{g^2} \int \text{Tr}[(f_\alpha + B \mathbb{I}_N) \wedge \star(f_\alpha + B \mathbb{I}_N)] + \frac{\theta}{8\pi^2} \int \text{Tr}[(f_\alpha + B \mathbb{I}_N) \wedge (f_\alpha + B \mathbb{I}_N)] \\ & - \frac{\theta}{8\pi^2} \int [\text{Tr}(f_\alpha) + NB] \wedge [\text{Tr}(f_\alpha) + NB] + \frac{1}{2\pi} \int \beta \wedge [\text{Tr}(f_\alpha) + NB] \\ & + \frac{1}{2\pi} \int \tilde{\beta} \wedge (-dC + NB), \end{aligned} \quad (2.13)$$

where $\tilde{\beta}_{\mu\nu}$ is another two-form Lagrange multiplier that constrains $dC = NB$. Without the background fields present, the θ angle is 2π periodic in the pure $SU(N)$ gauge theory because of the quantization of the theta term. If we take $\theta \rightarrow \theta + 2\pi k$, where $k \in \mathbb{Z}$, then the action changes (modulo an integer multiple of 2π) by [61]

$$S_{\text{SPT}}[B] = \frac{N(N-1)k}{4\pi} \int B \wedge B, \quad (2.14)$$

Consequently, assuming that $SU(N)$ gauge theory at $\theta = 0$ is a trivial confining phase, then the oblique confining state at $\theta = 2\pi k$ is an SPT protected by the \mathbb{Z}_N electric one-form symmetry [18, 62] and characterized by the response in Eq. (2.14). This SPT is nontrivial unless $k \in N\mathbb{Z}$ on a spin manifold or $k \in 2N\mathbb{Z}$ on a generic manifold.

The physical meaning of the response, Eq. (2.14), is related to dyon condensation. Physically, the background field $B_{\mu\nu}$ can be constructed from a configuration of the \mathbb{Z}_N one-form symmetry operators $U_{q_m}(\Sigma)$ that end on loops. At $\theta = 0$, as discussed previously near Eq. (2.6), these operators are precisely the 't Hooft loops. At $\theta = 2\pi k$, the response, Eq. (2.14), indicates that the loops carry a charge $q_e = (N-1)k q_m = -k q_m \bmod N$ under the one-form symmetry. Thus, the loop at which the one-form symmetry operator $U_{q_m}(\Sigma)$ ends has charges $(-k q_m, q_m)$. As discussed near Eq. (2.11), dyons with these charges are precisely those that have a perimeter law at $\theta = 2\pi k$. Indeed, dyons with charges not of this form are confined and energetically suppressed, so the symmetry operators can only end on loops that have a perimeter law. We recall that the dyonic loops that have a perimeter law must be attached to surfaces (unless $k \in N\mathbb{Z}$, in which case the SPT is trivial), so these objects are analogous to string order parameters in (1+1)d zero-form SPTs [63].

Because there is a change in response to background fields as θ is tuned continuously from $\theta = 2\pi k$ to $\theta = 2\pi(k+1)$, there must be at least one phase transition as θ is varied. Based on evidence from lattice simulations, large N [64], 't Hooft anomalies [61], and deformations of supersymmetric theories [65], a single first order transition with spontaneously broken time-reversal symmetry [66] is believed to occur at $\theta = 2\pi k + \pi$, where the \mathbb{Z}_N one-form SPT states become degenerate. Although this picture is not fully settled, it holds rigorously for large enough N [64].

To summarize, pure $SU(N)$ Yang-Mills theory has a \mathbb{Z}_N electric one-form symmetry and realizes SPT phases protected by this symmetry at $\theta \in 2\pi\mathbb{Z}$. These SPT phases are realized by the mechanism of dyon condensation and oblique confinement.

2.2 $SU(N)/\mathbb{Z}_N$ gauge theory

We next review the continuum description of $PSU(N) = SU(N)/\mathbb{Z}_N$ Yang-Mills theory [18, 42, 67, 68], which includes $PSU(2) \cong SO(3)$ as a special case. The $PSU(N)$ gauge theory can be constructed by gauging the \mathbb{Z}_N electric one-form symmetry of $SU(N)$ gauge theory. Hence, we promote the background fields in Eq. (2.13) to dynamical fields. The explicit form of the action is⁴

$$S_{PSU(N)}[\theta, B] = S_{YM}[\theta, b] + \frac{N}{2\pi} \int b \wedge B, \quad (2.15)$$

where $S_{YM}[\theta, b]$ is the same as Eq. (2.13) but with $C_\mu \rightarrow c_\mu$ and $B_{\mu\nu} \rightarrow b_{\mu\nu}$ promoted to dynamical $U(1)$ one-form and two-form gauge fields respectively.⁵ Naturally, the fields α_μ , c_μ , and $b_{\mu\nu}$ have gauge transformations descending from Eq. (2.12). We have also introduced a new \mathbb{Z}_N two-form background gauge field $B_{\mu\nu}$, which probes a \mathbb{Z}_N magnetic one-form symmetry.

We constructed $PSU(N)$ gauge theory by gauging the full electric one-form symmetry of $SU(N)$ gauge theory, so unsurprisingly, the center of $PSU(N)$ is trivial, and $PSU(N)$ gauge theory accordingly has no electric one-form symmetry. Because of the coupling to the dynamical two-form gauge field $b_{\mu\nu}$, a generic Wilson loop is no longer gauge invariant.

⁴We could have added a discrete theta term [42] (proportional to the integral of $b \wedge b$) to the action, but this term can be absorbed into the definition of θ .

⁵The two-form gauge field $b_{\mu\nu}$ is roughly $b = (2\pi/N) w_2^{PSU(N)}$, where $w_2^{PSU(N)}$ is the second Stiefel-Whitney class for $PSU(N)$, which characterizes the obstruction to lift the $PSU(N)$ gauge bundle to an $SU(N)$ bundle.

Rather, it is supported on a loop $\gamma = \partial\Omega$ attached to a surface Ω as

$$\mathcal{W}_{\mathcal{R}}(\gamma, \Omega) = \left[\text{Tr}_{\mathcal{R}} \mathcal{P} \exp \left(i \oint_{\gamma} \alpha \right) \right] \exp \left(i q_e \int_{\Omega} b \right), \quad (2.16)$$

where q_e is the one-form gauge charge for the representation \mathcal{R} . The attached surface is trivial only if $q_e \in N\mathbb{Z}$, which is true for the adjoint representation, but a Wilson loop in the adjoint representation is screened by the gluons.

The possible magnetic charges are classified by $\pi_1(PSU(N)) \cong \mathbb{Z}_N$, so there should be N distinct 't Hooft loops that are genuine loop operators. These 't Hooft loops are expressed most easily if we integrate out c_{μ} , which implements the constraint that $\tilde{\beta} = d\tilde{a}$, where \tilde{a}_{μ} is a $U(1)$ one-form gauge field. The 't Hooft loops are

$$\mathcal{T}(\gamma)^{q_m} = \exp \left(i q_m \oint_{\gamma} \tilde{a} \right), \quad (2.17)$$

where q_m is an integer mod N . The 't Hooft loop is now always a genuine loop operator. Indeed, by definition, gauging the electric one-form symmetry of $SU(N)$ gauge theory trivializes the symmetry operator $U_{q_m}(\Sigma)$. Because 't Hooft loops are attached to open surface versions of $U_{q_m}(\Sigma)$, gauging the electric one-form symmetry renders the attached surface trivial, turning the 't Hooft loops into genuine loop operators.

As with $SU(N)$ gauge theory, dyon operators can be formed from products of the Wilson and 't Hooft loops as

$$\mathcal{D}_{(q_e, q_m)}(\gamma, \Omega) = \mathcal{W}_{\mathcal{R}}(\gamma, \Omega) \mathcal{T}(\gamma)^{q_m}. \quad (2.18)$$

We again label the dyonic operators by their charges (q_e, q_m) at $\theta = 0$, prior to the Witten effect. These operators are genuine loop operators in $PSU(N)$ gauge theory only if $q_e = 0$ mod N .

A magnetic \mathbb{Z}_N one-form global symmetry, probed by the \mathbb{Z}_N two-form background field $B_{\mu\nu}$, acts on the 't Hooft loops. The operator, supported on a closed surface Σ , that acts with this symmetry is

$$\mathcal{U}(\Sigma) = \exp \left(-i \oint_{\Sigma} b \right). \quad (2.19)$$

This operator is a closed surface version of the Wilson loop, Eq. (2.16), so it can be viewed as a worldsheet of a string of electric flux⁶. This magnetic one-form symmetry is the dual symmetry that arises after gauging the electric one-form symmetry.

⁶The origin of the minus sign in the exponent of Eq. (2.19) can be traced to the ordering of operators in Eq. (2.4).

Next, we characterize the phases realized in $PSU(N)$ gauge theory. Gauging the \mathbb{Z}_N electric one-form symmetry does not modify whether operators have a perimeter law or area law. Therefore, at $\theta = 0$ the basic 't Hooft loop has a perimeter law, signaling deconfinement of monopoles, and the charge N monopoles are condensed. But in $PSU(N)$ gauge theory, since the 't Hooft loops are genuine loop operators, their deconfinement signals that the \mathbb{Z}_N magnetic one-form symmetry is spontaneously broken at low energies, resulting in topological order. Indeed, assuming that $SU(N)$ gauge theory at $\theta = 0$ is a trivial confining phase, then gauging its \mathbb{Z}_N electric one-form symmetry results in a phase with a TQFT of

$$S_{\text{BF}}[B] = \frac{N}{2\pi} \int b \wedge (d\tilde{a} + B), \quad (2.20)$$

which is BF theory at level N . This relation between the phases of $SU(N)$ gauge theory and $PSU(N)$ gauge theory has an analogue in (3+1)d \mathbb{Z}_N lattice gauge theory, which has a \mathbb{Z}_N electric one-form symmetry. Under gauging this \mathbb{Z}_N one-form symmetry and modifying the gauge coupling in a particular way, the partition function for \mathbb{Z}_N lattice gauge theory remains the same, but the confining phase, which is topologically trivial, is exchanged with the topologically ordered deconfined phase.

Following the discussion from Section 2.1, shifting $\theta \rightarrow \theta + 2\pi k$, where $k \in \mathbb{Z}$, changes the action by Eq. (2.14) but now with $B_{\mu\nu} \rightarrow b_{\mu\nu}$ dynamical. Consequently, the periodicity of θ in $PSU(N)$ gauge theory is $2\pi N$. At $\theta = 2\pi k$, the condensed dyons have charges $(-Nk, N)$, and the dyons obeying a perimeter law have charges of the form

$$(q_e, q_m) = (-k q_m, q_m), \quad (2.21)$$

which is consistent with the $2\pi N$ periodicity of θ since q_e and q_m are integers defined mod N . However, not all of these dyons in Eq. (2.21) correspond to genuine loop operators. The one-form symmetry can act nontrivially only on operators supported on noncontractible loops, but loops attached to a surface are always contractible.

The subset of dyons in Eq. (2.21) associated with genuine loop operators must have electric charges satisfying $q_e = -k q_m = 0 \pmod{N}$. Such operators must thus have magnetic charges such that $q_m \in \frac{N}{L}\mathbb{Z}$, where $L = \text{gcd}(N, k)$. The dyonic loop operators that have a perimeter law and are genuine loop operators therefore have charges

$$(q_e, q_m) = \left(-\frac{Nk}{L}q, \frac{N}{L}q \right) = \left(0, \frac{N}{L}q \right) \pmod{N}, \quad (2.22)$$

where q is an integer mod $L = \text{gcd}(N, k)$. The minimal magnetic charge of these deconfined dyons is N/L , so the \mathbb{Z}_N magnetic one-form symmetry is spontaneously broken to $\mathbb{Z}_{N/L}$ in

this phase. This phase is characterized by a topological order that is the same as in the deconfined phase of $\mathbb{Z}_N/\mathbb{Z}_{N/L} \cong \mathbb{Z}_L$ gauge theory. Indeed, the braiding phase between a deconfined magnetic charge $\mathcal{T}(\gamma)^{Nq/L}$ and an electric flux tube $\mathcal{U}(\Sigma)^{q'}$ is

$$\langle \mathcal{T}(\gamma)^{Nq/L} \mathcal{U}(\Sigma)^{q'} \rangle = \exp \left(\frac{2\pi i}{L} q q' \Phi(\gamma, \Sigma) \right), \quad (2.23)$$

where $\Phi(\gamma, \Sigma)$ is the linking number of γ and Σ .

Naturally, the effective TQFT that describes the phase at $\theta = 2\pi k$ is the gauged version of the SPT action, Eq. (2.14). The specific action for the TQFT is

$$S_{\text{TQFT}}[p, \tilde{a}, b, B] = \frac{N}{2\pi} \int b \wedge (d\tilde{a} + B) + \frac{Np}{4\pi} \int b \wedge b, \quad (2.24)$$

where $p = (N-1)k$. As reviewed in Appendix A, this TQFT indeed describes a topological order realized by spontaneously breaking a \mathbb{Z}_N one-form global symmetry to $\mathbb{Z}_{N/L}$ at low energies, where $L = \gcd(N, p) = \gcd(N, k)$. Phases of this type are also realized [57, 69] in a lattice model introduced by Cardy and Rabinovici [51, 70].

The unbroken $\mathbb{Z}_{N/L}$ one-form symmetry also has a nontrivial response to a background field $B_{\mu\nu}$, given by

$$S_{\text{resp}}[B] = \frac{Nr}{4\pi L} \int B \wedge B, \quad (2.25)$$

where r is an integer mod N/L such that $rp = r(N-1)k = -L \bmod N$. The interpretation of this response is similar to that of Eq. (2.14). A surface operator that acts with the unbroken $\mathbb{Z}_{N/L}$ magnetic one-form symmetry is $\mathcal{U}(\Sigma)^L$. The configurations of the background field $B_{\mu\nu}$ are constructed by allowing these surface operators to end on loops. Because $\mathcal{U}(\Sigma)^L$ represents an electric flux tube, it must terminate on a loop with electric charge $q_e = -L$. This loop must have a perimeter law, so its charges must be of the form in Eq. (2.21) so that

$$q_e = -k q_m = -L \bmod N. \quad (2.26)$$

A solution to this equation for the magnetic charge is then $q_m = r \bmod N$ so that the loop on which $\mathcal{U}(\Sigma)^L$ ends has charges $(-L, r)$. Likewise, the symmetry operator $[\mathcal{U}(\Sigma)^L]^{Mk/L}$, where M is an integer mod N/L , must end on a loop with charges

$$(q_e, q_m) = (-k M, M). \quad (2.27)$$

These dyons are those that have a perimeter law but are not genuine line operators. Thus, although these operators do not contribute to the topological order, they can have physical consequences that manifest in the response, Eq. (2.25).

Similar to $SU(N)$ gauge theory, as θ is varied from $\theta = 2\pi k$ to $\theta = 2\pi(k+1)$, there must be a phase transition. It is believed that a first order transition occurs at $\theta = 2\pi k + \pi$, where the ground states of the TQFT, Eq. (2.24), with $p = (N-1)k$ are degenerate with the ground states of the same TQFT but with $p = (N-1)(k+1)$. This statement is rigorous for large N [64]. If $\gcd(N, k) = \gcd(N, k+1) = 1$, then there are two degenerate \mathbb{Z}_N one-form SPTs at the first order transition. Otherwise, there will be topologically ordered ground states degenerate with an SPT state.

3 $SU(N)$ adjoint QCD: SPT transition

Now that we have established the gapped phases of pure $SU(N)$ and $PSU(N)$ gauge theories, we are equipped to couple to matter and examine the transitions between these topological phases. We begin with $SU(N)$ gauge theory coupled to N_f flavors of Majorana fermions in the adjoint representation, which we refer to as $SU(N)$ adjoint QCD. We focus on the case in which N is even and N_f is odd. Because the adjoint representation is real, coupling to the $SU(N)$ gauge field is consistent with imposing the Majorana condition. The relation of this theory to SPT transitions was previously studied in Ref. [28], but in that case, the authors explicitly broke the electric one-form symmetry and found transitions between distinct zero-form SPTs. In this work, because we are ultimately interested in $SU(N)/\mathbb{Z}_N$ gauge theory coupled to fermions, we will leave the electric one-form symmetry unbroken in anticipation of Section 4 where we will gauge this symmetry.

Since the fermions are in the adjoint representation of $SU(N)$, it is convenient to express the fermions as matrices, given by⁷ $\psi_J(x) = \psi_J^\alpha(x) T^\alpha$, where J is the flavor index, α is the gauge index, and T^α are the generators of the $\mathfrak{su}(N)$ Lie algebra in the fundamental representation. The action is

$$S_{\text{AQCD}} = \int d^4x \sum_{J=1}^{N_f} \text{Tr} [\psi_J^T(x) \mathcal{C} (i \not{D}_a - m) \psi_J(x)] - \frac{1}{g^2} \int \text{Tr} (f \wedge \star f), \quad (3.1)$$

where we define $\not{D}_a \psi_J = \gamma^\mu (\partial_\mu \psi_J - i [a_\mu, \psi_J])$, $\psi_J^T(x)$ denotes the transpose of $\psi_J(x)$ for spinor indices only, γ^μ are Dirac matrices, and m is the fermion mass. The charge conjugation matrix \mathcal{C} is a unitary matrix acting on spinor indices that obeys $\mathcal{C} \gamma^\mu \mathcal{C}^{-1} = -(\gamma^\mu)^T$ and $\mathcal{C}^T = -\mathcal{C}$. The $\psi_J(x)$ are Majorana fermions obeying the constraint, $\bar{\psi}_J(x) = \psi_J^T(x) \mathcal{C}$. A brief review of Majorana fermions may be found in Appendix B.

⁷We suppress spinor indices for simplicity.

3.1 Global symmetries

We proceed by describing the global symmetries of the theory, Eq. (3.1). Since the fermionic matter is coupled to the gauge field in the adjoint representation, the \mathbb{Z}_N electric one-form symmetry of the pure $SU(N)$ gauge theory is retained. The zero-form symmetry that will be important for us is

$$G = SO(N_f) \times \mathbb{Z}_2^T, \quad (3.2)$$

where \mathbb{Z}_2^T is time-reversal symmetry and $SO(N_f)$ is the global flavor symmetry⁸ of the fermions at a generic mass m , acting as

$$\psi_J(x) \rightarrow \mathcal{O}_{JK} \psi_K(x), \quad (3.3)$$

for some matrix $\mathcal{O} \in SO(N_f)$. Time-reversal symmetry is associated with an antiunitary operator T and acts on the fields as

$$\psi_J^\alpha(\mathbf{x}, t) \rightarrow \mathcal{C} \gamma^5 \psi_J^\alpha(\mathbf{x}, -t), \quad a_0^\alpha(\mathbf{x}, t) \rightarrow -a_0^\alpha(\mathbf{x}, -t), \quad a_j^\alpha(\mathbf{x}, t) \rightarrow a_j^\alpha(\mathbf{x}, -t). \quad (3.4)$$

Hence, electric charge is odd under time-reversal while magnetic charge is even. From the properties of \mathcal{C} mentioned above, we can also deduce that $\mathsf{T}^2 = (-1)^F$, where $(-1)^F$ is fermion parity.⁹ Since fermion parity acts as a global symmetry, adjoint QCD describes a fermionic theory in the sense that there exist local gauge invariant fermionic operators, such as $\text{Tr}[(\psi_J^T \mathcal{C} \psi_J) \psi_K]$. An important observation is that every gauge invariant bosonic local operator that is a Kramers singlet, whereas every gauge invariant fermionic local operator transforms under time-reversal as a Kramers doublet. We refer to this observation as a spin-Kramers relation.

Parity and charge conjugation symmetries are not essential for our purposes, but we mention them for completeness. Parity is associated with a unitary operator P that acts on the fields as

$$\psi_J^\alpha(\mathbf{x}, t) \rightarrow i\gamma^0 \psi_J^\alpha(-\mathbf{x}, t), \quad a_0^\alpha(\mathbf{x}, t) \rightarrow a_0^\alpha(-\mathbf{x}, t), \quad a_j^\alpha(\mathbf{x}, t) \rightarrow -a_j^\alpha(-\mathbf{x}, t). \quad (3.5)$$

⁸For $m \neq 0$, the full flavor symmetry is $O(N_f)$. For odd N_f , this group factorizes to $O(N_f) = SO(N_f) \times \mathbb{Z}_2^F$ (it is a semidirect product for even N_f), but the \mathbb{Z}_2^F factor is equivalent to fermion parity. As we discuss below, this symmetry is already included in time-reversal.

⁹A formal consequence is that the Wick rotated theory (in Euclidean spacetime) may be placed on a non-orientable manifold that admits a $\text{Pin}^+(4)$ structure [71]. A manifold X of this type must have a trivial second Stiefel-Whitney class, $w_2(X) = 0$.

Acting with parity twice then also gives $P^2 = (-1)^F$. For $N > 2$, there is a global charge conjugation symmetry, which is associated with a unitary operator C and acts as

$$\psi_J(\mathbf{x}, t) = \psi_J^\alpha(\mathbf{x}, t) T^\alpha \rightarrow -\psi_J^\alpha(\mathbf{x}, t) (T^\alpha)^*, \quad a_\mu(\mathbf{x}, t) = a_\mu^\alpha(\mathbf{x}, t) T^\alpha \rightarrow -a_\mu^\alpha(\mathbf{x}, t) (T^\alpha)^*, \quad (3.6)$$

but for $N = 2$ this transformation is equivalent to a gauge transformation. These discrete symmetries are not entirely independent because of the CPT theorem.

At the massless point, $m = 0$, the zero-form flavor symmetry is enhanced to

$$G_{SU(N)} = \frac{SU(N_f) \times \mathbb{Z}_{2NN_f}}{\mathbb{Z}_{N_f}}. \quad (3.7)$$

The \mathbb{Z}_{2NN_f} factor is a remnant of the classical $U(1)$ axial symmetry,

$$\psi_J(x) \rightarrow e^{i\vartheta\gamma^5} \psi_J(x). \quad (3.8)$$

By the Adler-Bell-Jackiw anomaly [72, 73], this transformation is equivalent to a shift of the $SU(N)$ theta angle by

$$\theta \rightarrow \theta + 2NN_f\vartheta. \quad (3.9)$$

Since the $SU(N)$ theta angle has periodicity 2π , the action is invariant mod 2π under the transformation, Eq. (3.8), only if $NN_f\vartheta \in \pi\mathbb{Z}$. The axial global symmetry is thus anomalously broken to \mathbb{Z}_{2NN_f} quantum mechanically.

This \mathbb{Z}_{2NN_f} symmetry has a mixed 't Hooft anomalies. Suppose we couple Eq. (3.1) to a background \mathbb{Z}_N two-form gauge field $B_{\mu\nu}$, a background gauge field A_μ for the $SO(N_f)$ flavor symmetry,¹⁰ and a background metric $g_{\mu\nu}$ (to keep track of thermal response [74–77]). Transforming $\psi_J(x)$ by the discrete axial transformation,

$$\psi_J(x) \rightarrow e^{i\frac{2\pi k}{2NN_f}\gamma^5} \psi_J(x), \quad (3.10)$$

for some integer k , is equivalent to changing the action by

$$\Delta S = \frac{N(N-1)k}{4\pi} \int B \wedge B + \frac{\frac{2\pi k}{NN_f}(N^2-1)}{8\pi^2} \int \text{Tr}(F \wedge F) + \frac{\frac{2\pi k}{N}(N^2-1)}{384\pi^2} \int \text{Tr}(R \wedge R), \quad (3.11)$$

where $F_{\mu\nu}$ is the two-form field strength of the $SO(N_f)$ background field A_μ and R is the curvature two-form.¹¹ Eq. (3.11) signals a mixed 't Hooft anomaly for the \mathbb{Z}_{2NN_f} axial

¹⁰We couple only to a background field for the $SO(N_f)$ subgroup rather than $SU(N_f)$ because this subgroup is the symmetry that remains when we introduce masses for the fermions.

¹¹See Appendix C for a more detailed definition of R and a review of how to couple fermions to gravity in (3+1)d.



Figure 1: Schematic depiction of the phase diagram of $SU(N)$ adjoint QCD, Eq. (3.1), for N even and N_f odd flavors of Majorana fermions, as a function of the mass m . For $m > 0$, the theory flows to $SU(N)$ gauge theory at $\theta = 0$, which is adiabatically connected to a trivial product state. For $m < 0$, the state is a nontrivial fermionic SPT protected by the $\mathbb{Z}_N^{(1)}$ electric one-form symmetry and $G = SO(N_f) \times \mathbb{Z}_2^T$, which consists of a flavor symmetry and time-reversal.

symmetry with the \mathbb{Z}_N one-form symmetry, the flavor symmetry, and gravity. The first term in Eq. (3.11) arises because Eq. (3.10) is equivalent to a change in the $SU(N)$ theta angle by $2\pi k$ (cf. Eq. (3.9)). As discussed in Section 2.1, upon coupling to a background two-form gauge field $B_{\mu\nu}$, a shift in the θ angle of $SU(N)$ gauge theory by $2\pi k$ is equivalent to stacking with an SPT, Eq. (2.14), which matches the first term of Eq. (3.11).

3.2 Massive phases: SPTs

Next, we discuss the phases that occur when the fermions are massive. At low energies, well below the mass scale $|m|$, the fermions may be integrated out. For $m > 0$, the physics is governed by pure $SU(N)$ gauge theory with $\theta = 0$, which is gapped, confining, and topologically trivial. For $m < 0$, integrating out the fermions gives pure $SU(N)$ gauge theory with $\theta = \pi N N_f$, which is an integer multiple of 2π for N even. As discussed in Section 2.1, $SU(N)$ gauge theory is also gapped and topologically trivial at this value of θ , but it can differ from the $\theta = 0$ state as an SPT.

To understand precisely what SPT this state is, we again couple to a background metric $g_{\mu\nu}$, a background field A_μ for the $SO(N_f)$ flavor symmetry, and a background field $B_{\mu\nu}$ for the \mathbb{Z}_N electric one-form symmetry. The theory with negative mass $m = -|m|$ is equivalent to the theory with a positive mass $m = |m|$ acted upon by an axial transformation, Eq. (3.10), with $k = N N_f / 2$. Thus, if we choose a regularization such that the positive mass phase is a trivial SPT, then the negative mass phase is a nontrivial SPT with classical action,

$$S_{\text{SPT}} = \frac{N^2(N-1)N_f}{8\pi} \int B \wedge B + \frac{\pi(N^2-1)}{8\pi^2} \int \text{Tr}(F \wedge F) + \frac{\pi N_f(N^2-1)}{384\pi^2} \int \text{Tr}(R \wedge R). \quad (3.12)$$

The negative mass and positive mass phases thus differ as fermionic SPTs protected by the \mathbb{Z}_N electric one-form symmetry, the $SO(N_f)$ flavor symmetry, and time-reversal symmetry.

If we introduce a (2+1)d boundary to the SPT state in Eq. (3.12), several different

boundary states are possible. The bulk anomaly from the second and third terms of Eq. (3.12) can be matched at the boundary by $N_f(N^2 - 1)$ free massless Majorana fermions. The anomaly for the one-form symmetry can be matched by $SU(N)$ Chern-Simons theory with level $NN_f/2$, though other boundary states are also possible [61, 68]. A simple way to check that this topological order matches the anomaly is to observe that the bulk theory is $SU(N)$ gauge theory at $\theta = \pi NN_f$, which can be coupled to the two-form gauge field in a manifestly gauge invariant way as in Section 2.1. Applying Stokes' theorem to the $SU(N)$ theta term yields $SU(N)$ Chern-Simons theory at the boundary with level $NN_f/2$.

3.3 Phase transition

Having established the SPT phases that arise for $m \neq 0$, we address the transition between these phases, which is related to the IR fate of the massless point, $m = 0$. The low energy dynamics crucially depends on the number of flavors N_f . If the non-Abelian gauge field is coupled to $N_f(\mathcal{R})$ flavors of Majorana fermions in representation \mathcal{R} is, then under the scale change $x_\mu \rightarrow e^\ell x_\mu$, the one-loop beta function¹² for the Yang-Mills coupling g is [78, 79]

$$\beta(g) = \frac{dg}{d\ell} = (c_{\text{YM}} - c_{\text{M}}) g^3, \quad c_{\text{M}} = \frac{2}{3(4\pi)^2} \sum_{\mathcal{R}} I_{\mathcal{R}} N_f(\mathcal{R}), \quad c_{\text{YM}} = \frac{11}{3(4\pi)^2} I_{\text{adj}}, \quad (3.13)$$

where the group theoretic factor $I_{\mathcal{R}}$ is given by $\text{Tr}(T_{\mathcal{R}}^\alpha T_{\mathcal{R}}^\beta) = I_{\mathcal{R}} \delta^{\alpha\beta}$ for generators $T_{\mathcal{R}}^\alpha$ of the gauge group in representation \mathcal{R} . For $SU(N)$, we have $I_f = 1/2$ for the fundamental representation and $I_{\text{adj}} = N$ for the adjoint representation.

For N_f flavors of Majorana fermions in the adjoint representation, the beta function in Eq. (3.13) indicates that the theory, Eq. (3.1), is infrared free for $N_f \geq 6$. As discussed in the introduction, since the gauge coupling becomes small in the IR, we can be sure that there is a direct continuous transition at $m = 0$ from the trivial SPT to the nontrivial SPT (cf. Eq. (3.12)), as depicted in Figure 1. The critical exponents can also be easily determined reliably in this case. Because we are interested in N_f odd, the smallest value for which the transition is unambiguously continuous is $N_f = 7$.

The cases $2 \leq N_f \leq 5$ are less well understood¹³, but we briefly summarize what is known about them here for completeness. The standard lore is (see Ref. [82], for example) that

¹²Our sign convention is such that $\beta(g) > 0$ signals a flow to strong coupling at low energies.

¹³CFTs in 4d have a universal quantity a that is similar to the central charge of a 2d CFT [80]. For the IR free theories, we can compute that $a = (62 + N_f/2)(N^2 - 1)$, where we use units in which $a = 1$ for a single real free massless scalar field. For the asymptotically free theories, which have $N_f \leq 5$, the UV value of a is also $a_{\text{UV}} = (62 + N_f/2)(N^2 - 1)$. Since a monotonically decreases along RG flows [81], a useful constraint is that the IR value for $N_f \leq 5$ must satisfy $a_{\text{IR}} < a_{\text{UV}}$.

higher values of N_f are interacting conformal field theories of the Banks-Zaks type [31, 32] while smaller values of N_f ultimately confine and spontaneously break $G_{SU(N)} \rightarrow O(N_f)$, resulting in N copies of an $SU(N_f)/SO(N_f)$ sigma model. But this picture is not firmly established, so other infrared behaviors are possible [83, 84] as long as they are consistent with anomaly constraints [85]. Current evidence suggests that $N_f = 4$ and $N_f = 5$ are conformal [86–93]. Thus, the $N_f = 5$ theory may also have a continuous SPT transition associated with an interacting CFT.¹⁴

The $N_f = 1$ theory is better understood since it has supersymmetry. This theory confines and breaks $G_{SU(N)} = \mathbb{Z}_{2N} \rightarrow \mathbb{Z}_2^F$ spontaneously [94, 95], resulting in N degenerate ground states. Each of these N vacua is associated with the condensation of a different dyon [96–98] and a \mathbb{Z}_N one-form SPT of the form in Eq. (2.14) [18]. In this case the transition is first order since two of the N ground states are associated with the trivial state at $m > 0$ and the nontrivial SPT at $m < 0$.

Hence, the global flavor symmetry $G = SO(N_f) \times \mathbb{Z}_2^T$ plays an important role. At the $m = 0$ point, the mass term in Eq. (3.1) is the only relevant perturbation that respects the symmetry G . For large enough N_f , if this symmetry is preserved, then there is a continuous transition directly between the two SPT states as in Figure 1. If the symmetry G is not mandated, then other perturbations can drive the $m = 0$ theory to various intermediate phases or to a first-order transition.

4 $SU(N)/\mathbb{Z}_N$ adjoint QCD: SET transition

We now turn to our first main result—a topological transition between different (3+1)d SETs that have distinct patterns of one-form symmetry breaking (i.e., different topological orders). The theory in which this transition occurs is $SU(N)/\mathbb{Z}_N = PSU(N)$ gauge theory coupled to N_f flavors of Majorana fermions in the adjoint representation, where we again

¹⁴There does not necessarily have to be a direct transition between the two SPT phases in this case. It is possible that there could be an intermediate “quantum phase” that is not obvious from the Lagrangian description, or criticality could possibly extend to small nonzero m .

take N even and N_f odd. The action for $PSU(N)$ adjoint QCD theory is

$$\begin{aligned}
S_0 = & \int d^4x \sum_{J=1}^{N_f} \text{Tr} [\psi_J^T(x) \mathcal{C}(i\not{D}_{\alpha+c\mathbb{I}_N} - m) \psi_J(x)] - \frac{1}{g^2} \int \text{Tr}[(f_\alpha + b\mathbb{I}_N) \wedge \star(f_\alpha + b\mathbb{I}_N)] \\
& + \frac{1}{2\pi} \int \beta \wedge [\text{Tr}(f_\alpha) + \text{dc}] + \frac{1}{2\pi} \int \tilde{\beta} \wedge (-\text{dc} + Nb) + \frac{N}{2\pi} \int b \wedge B,
\end{aligned} \tag{4.1}$$

where we have gauged the \mathbb{Z}_N one-form symmetry of Eq. (3.1) using the same methods as in Section 2.2. Namely, α_μ is a dynamical $U(N)$ gauge field with field strength $(f_\alpha)_{\mu\nu}$, c_μ is a dynamical $U(1)$ one-form gauge field, and $b_{\mu\nu}$, $\beta_{\mu\nu}$, and $\tilde{\beta}_{\mu\nu}$ are all dynamical $U(1)$ two-form gauge fields. We have also coupled to a background \mathbb{Z}_N two-form gauge field $B_{\mu\nu}$, which probes a \mathbb{Z}_N magnetic one-form symmetry that acts on 't Hooft lines.

Because gauging the discrete one-form symmetry does not change the renormalization group flow of the gauge coupling g , the low energy physics of the theory at $m = 0$ can be deduced from our knowledge of $SU(N)$ adjoint QCD. In particular, for $m = 0$ the theory is IR free for $N_f \geq 7$ flavors, and $N_f = 5$ is believed to be an interacting CFT. Hence, the transition we discuss below will be continuous or first order for the same values of N_f as in the $SU(N)$ theory. However, as we discuss below, the gapped phases for $m \neq 0$ will be rather different from the $SU(N)$ case.

4.1 Global symmetries

The zero-form global symmetry of the theory at generic m is still $G = SO(N_f) \times \mathbb{Z}_2^T$. The $SO(N_f)$ flavor symmetry acts on the fermions $\psi_J(x)$ in the same way as in Eq. (3.3). Under time-reversal symmetry, the fermion fields and the $U(N)$ gauge field transform analogously to Eq. (3.4),

$$\psi_J(\mathbf{x}, t) \rightarrow \mathcal{C}\gamma^5 \psi_J(\mathbf{x}, -t), \quad \alpha_0(\mathbf{x}, t) \rightarrow -\alpha_0(\mathbf{x}, -t), \quad \alpha_j(\mathbf{x}, t) \rightarrow \alpha_j(\mathbf{x}, -t). \tag{4.2}$$

Again, we have $\mathbb{T}^2 = (-1)^F$. The transformations of $b_{\mu\nu}$ and c_μ can be inferred from the local constraint, $-\text{Tr}(f_\alpha) = Nb = \text{dc}$. The Lagrange multiplier $\beta_{\mu\nu}$ changes as

$$\beta_{0j}(\mathbf{x}, t) \rightarrow \beta_{0j}(\mathbf{x}, -t), \quad \beta_{jk}(\mathbf{x}, t) \rightarrow -\beta_{jk}(\mathbf{x}, -t), \tag{4.3}$$

and $\tilde{\beta}_{\mu\nu}$ changes in the same way since $\beta_{\mu\nu} = \tilde{\beta}_{\mu\nu}$ locally. As observed previously, magnetic charge is even under time-reversal while electric charge is odd. Again, there is a spin-Kramers relation—all local gauge invariant operators are either bosonic Kramers singlets or fermionic Kramers doublets.

Under parity P , the fields transform as

$$\begin{aligned}\psi_J(\mathbf{x}, t) &\rightarrow i\gamma^0 \psi_J(-\mathbf{x}, t), & \alpha_0(\mathbf{x}, t) &\rightarrow \alpha_0(-\mathbf{x}, t), & \alpha_j(\mathbf{x}, t) &\rightarrow -\alpha_j(-\mathbf{x}, t), \\ \beta_{0j}(\mathbf{x}, t) &\rightarrow \beta_{0j}(-\mathbf{x}, t), & \beta_{jk}(\mathbf{x}, t) &\rightarrow -\beta_{jk}(-\mathbf{x}, t),\end{aligned}\tag{4.4}$$

so magnetic charge is odd under parity while electric charge is even. Charge conjugation C (for $N > 2$) acts on the fields as

$$\psi_J(\mathbf{x}, t) = -[\psi_J(\mathbf{x}, t)]^*, \quad \alpha_\mu(\mathbf{x}, t) \rightarrow -[\alpha_\mu(\mathbf{x}, t)]^*, \quad \beta_{\mu\nu}(\mathbf{x}, t) \rightarrow -\beta_{\mu\nu}(\mathbf{x}, t), \tag{4.5}$$

where, as in Eq. (3.6), the complex conjugation acts only on complex numbers and not the Grassmann fields. We see that both electric and magnetic charges are odd under C .

At $m = 0$, the flavor symmetry is enhanced to

$$G_{PSU(N)} = \frac{SU(N_f) \times \mathbb{Z}_{2N_f}}{\mathbb{Z}_{N_f}}. \tag{4.6}$$

While $SU(N)$ adjoint QCD has a \mathbb{Z}_{2N_f} axial symmetry, this symmetry is reduced to \mathbb{Z}_{2N_f} in the $PSU(N)$ theory because of the mixed anomaly, Eq. (2.15), or equivalently, because the theta angle of $PSU(N)$ gauge theory has periodicity $2\pi N$ rather than 2π . However, upon gauging the \mathbb{Z}_N electric one-form symmetry of $SU(N)$ adjoint QCD, the $\mathbb{Z}_N \subset \mathbb{Z}_{2N_f}$ subgroup of the axial symmetry can be viewed as a non-invertible symmetry in the $PSU(N)$ theory [33, 99, 100].

As long as the subset $G = SO(N_f) \times \mathbb{Z}_2^T$ of the symmetry at $m = 0$ is preserved, then the only relevant deformation of the $m = 0$ point that respects the symmetry G is the mass in Eq. (4.1). We are now ready to examine the gapped phases that result from introducing a nonzero fermion mass m , which can be deduced from gauging the \mathbb{Z}_N one-form symmetries of the phases in Section 3.2.

4.2 \mathbb{Z}_N topological order

For positive m , after integrating out the fermions, the gapped phase at low energies is pure $PSU(N)$ Yang-Mills theory at $\theta = 0$. As discussed in Section 2.2, the 't Hooft loop $\mathcal{T}(\gamma)$ has a perimeter law in this phase. Because all magnetic charges are deconfined, the full \mathbb{Z}_N magnetic one-form symmetry is spontaneously broken in this phase. The low energy physics is described by BF theory at level N , Eq. (2.20).

Since charge N monopoles are screened, this phase may be viewed as a condensate of charge N monopoles, analogous to a superconductor. The analogue of an Abrikosov vortex is

an electric flux tube, which is associated with the Wilson surface operator $\mathcal{U}(\Sigma)$ for $b_{\mu\nu}$. The magnetic monopoles experience a dual Aharonov-Bohm effect when they are adiabatically transported around an electric flux tube, as captured by the correlation function,

$$\langle \mathcal{T}(\gamma) \mathcal{U}(\Sigma) \rangle = \exp \left(\frac{2\pi i}{N} \Phi(\gamma, \Sigma) \right), \quad (4.7)$$

where $\Phi(\gamma, \Sigma)$ is the linking number of the loop γ and surface Σ in spacetime.

Because $PSU(N)$ adjoint QCD also has the zero-form symmetry $G = SO(N_f) \times \mathbb{Z}_2^T$, this \mathbb{Z}_N topological order is enriched by G . Thus, there are several possible symmetry fractionalization classes [9] for this phase, which are determined by specifying how the $PSU(N)$ magnetic monopoles transform under the zero-form symmetry G .¹⁵ The monopoles do not have to transform linearly under G since they are not formed by local operators. For example, under the $SO(N_f)$ symmetry, the monopoles can transform either as tensors or spinors.

The fractionalization of time-reversal symmetry is more subtle. For a (3+1)d time-reversal invariant bosonic SET, we would specify independently whether the point-line anyons are bosons or fermions and whether they transform under time-reversal as Kramers singlets or doublets. However, we recall that adjoint QCD has a spin-Kramers relation—local operators are either bosonic Kramers singlets or fermionic Kramers doublets. Thus, attaching a fermionic Kramers doublet to a point-like anyon should not be viewed as changing the symmetry fractionalization class. Rather, the monopoles can either be consistent with the spin-Kramers relation or inconsistent with it. Hence, if a monopole is a bosonic Kramers singlet or fermionic Kramers doublet, the time-reversal symmetry is not fractionalized. If a monopole is a bosonic Kramers doublet or a fermionic Kramers singlet, then its time-reversal symmetry is fractionalized.

Finally, for a general (3+1)d SET, the loop excitations can also fractionalize. However, for the fermionic SET here, the electric flux loops are not fractionalized because in $PSU(N)$ gauge theory the surface operators $\mathcal{U}(\Sigma)$ can be opened and end on Wilson loops that do not carry any fractional quantum numbers under G .

¹⁵Formally, a change in symmetry fractionalization class can be induced by modifying the background two-form gauge field $B_{\mu\nu}$ for the \mathbb{Z}_N one-form symmetry by terms depending on G background fields. For example, taking $B \rightarrow B + \pi (w_1)^2$, where w_1 is the first Stiefel-Whitney class of the spacetime manifold, turns a bosonic Kramers singlet monopole into a bosonic Kramers doublet. Similarly, the transformation $B \rightarrow B + \pi w_2^{SO(N_f)}$, where $w_2^{SO(N_f)}$ is the second Stiefel-Whitney class for $SO(N_f)$, transmutes monopoles that transform as $SO(N_f)$ tensors into $SO(N_f)$ spinors.

4.3 $\mathbb{Z}_{N/2}$ topological order

The other phase, which occurs for negative m , can be deduced from Eq. (3.12) by promoting the \mathbb{Z}_N two-form gauge field in Eq. (3.12) to a dynamical field. Thus, the low energy physics of this phase is governed by a TQFT with effective action,

$$S_{\text{SET}} = \frac{N(N-1)NN_f/2}{4\pi} \int b \wedge b + \frac{N}{2\pi} \int b \wedge (d\tilde{a} + B) + \frac{\pi(N^2-1)}{8\pi^2} \int \text{Tr}(F \wedge F) + \frac{\pi N_f(N^2-1)}{384\pi^2} \int \text{Tr}(R \wedge R), \quad (4.8)$$

where $b_{\mu\nu}$ is a dynamical $U(1)$ two-form gauge field, \tilde{a}_μ is a Lagrange multiplier that constrains $b_{\mu\nu}$ to be a \mathbb{Z}_N gauge field, $B_{\mu\nu}$ is a background \mathbb{Z}_N two-form gauge field, F is the field strength of the background gauge field probing the $SO(N_f)$ flavor symmetry, and R is the curvature two-form. The topological order for the two-form gauge theory TQFT is worked out in Appendix A. The \mathbb{Z}_N magnetic one-form symmetry is spontaneously broken to \mathbb{Z}_2 , resulting in $\mathbb{Z}_N/\mathbb{Z}_2 \cong \mathbb{Z}_{N/2}$ topological order, and the unbroken \mathbb{Z}_2 magnetic one-form symmetry has nontrivial SPT order. For $N = 2$, there is no topological order, so in that case, this phase is a \mathbb{Z}_2 one-form SPT state.

Let us understand this topological order from a physical perspective. If m is negative, then integrating out the fermions at low energy results in pure $PSU(N)$ gauge theory at $\theta = \pi NN_f$. Because the theta term is odd under time-reversal symmetry and θ has periodicity $2\pi N$ in $PSU(N)$ gauge theory, the theory is indeed time-reversal invariant at $\theta = \pi NN_f \sim \pi N$, where we have used that N_f is an odd integer.

For N even, $\theta = \pi N$ is an integer multiple of 2π , and according to Eq. (2.21), the charges of dyons with a perimeter law are of the form

$$\left(-\frac{N}{2}q_m, q_m \right) \quad (4.9)$$

for some integer $q_m \bmod N$. As a consistency check, we note that time-reversal symmetry acts on dyon charges as

$$\mathbb{Z}_2^T : (q_e, q_m) \rightarrow (-q_e, q_m). \quad (4.10)$$

Hence, each dyon with a perimeter law, Eq. (4.9), is essentially mapped to itself under time-reversal symmetry. A minor caveat is that the $(-N/2, 1)$ dyon is actually mapped to $(N/2, 1)$, but these dyons simply differ by a neutral fermion, so they should be identified with each other since the topological order is fermionic.




Figure 2: Schematic depiction of the phase diagram of $PSU(N) = SU(N)/\mathbb{Z}_N$ adjoint QCD, Eq. (4.1), for N even and N_f odd flavors of Majorana fermions as a function of the mass m . The $m > 0$ phase breaks the full $\mathbb{Z}_N^{(1)}$ magnetic one-form symmetry at low energies, giving rise to \mathbb{Z}_N topological order described by BF theory at level N , Eq. (2.20). For $m < 0$, the low energy physics is described by Eq. (4.8). The \mathbb{Z}_N magnetic one-form symmetry, denoted $\mathbb{Z}_N^{(1)}$ is spontaneously broken to $\mathbb{Z}_2^{(1)}$, resulting in $\mathbb{Z}_{N/2}$ topological order. Additionally, the $m < 0$ phase is stacked with a nontrivial fermionic SPT for the unbroken $\mathbb{Z}_2^{(1)}$ and $G = SO(N_f) \times \mathbb{Z}_2^T$ symmetries. The G symmetry may be fractionalized on the point-like anyon excitations in both phases.

The subset of the dyons in Eq. (4.9) that are associated with genuine loop operators are the purely magnetic charges,

$$(q_e, q_m) = (0, 2q) \mod N, \quad (4.11)$$

where $q \in \{0, 1, \dots, \frac{N}{2} - 1\}$. The \mathbb{Z}_N magnetic one-form symmetry is therefore spontaneously broken to \mathbb{Z}_2 at low energies, resulting in a $\mathbb{Z}_N/\mathbb{Z}_2 = \mathbb{Z}_{N/2}$ topological order. Just as in Section 4.2, this topological order has several symmetry fractionalization classes, which are specified by how the monopoles transform under G . Of course, the symmetry fractionalization class for this phase will be correlated with the fractionalization class for the $m > 0$ phase.

If we couple to a background field $B_{\mu\nu}$ that probes the unbroken \mathbb{Z}_2 one-form symmetry, according to Eq. (2.25), we obtain the response,

$$S_{\text{resp}}[B] = \frac{2}{4\pi} \int B \wedge B, \quad (4.12)$$

resulting from the dyon $(-N/2, 1)$ that has a perimeter law but is not a genuine loop operator. Indeed, the symmetry operator for the unbroken \mathbb{Z}_2 magnetic one-form symmetry is $\mathcal{U}(\Sigma)^{N/2}$, and if we open this surface, it must end on a loop with electric charge $-N/2$. The response, Eq. (4.12), indicates that the loop must also have magnetic charge 1.

Now suppose we place the bulk phase on a manifold X with a boundary ∂X . The second and third terms of Eq. (4.8) are the same as in Eq. (3.12), so the topological order in X is stacked with a fermionic SPT protected by the zero-form symmetry $G = SO(N_f) \times \mathbb{Z}_2^T$. Thus, as in Section 3.2, a possible boundary state on ∂X that saturates the anomaly for this SPT sector of G is $N_f(N^2 - 1)$ free massless Majorana fermions.

The boundary states associated with the first term of Eq. (4.8) that preserve the \mathbb{Z}_N magnetic one-form symmetry have been studied previously [68]. The minimal topological order on ∂X must include the deconfined bulk particles (cf. Eq. (4.11)), a semion, and an anti-semion. The semion can be formed by fusing the anti-semion and a transparent fermion from the bulk. Physically, the anti-semion is the $(-N/2, 1)$ dyon, associated with operator $\mathcal{D}_{(-N/2,1)}(\gamma, \Omega)$ in Eq. (2.18), which has a perimeter law but is not a genuine loop operator. If the bulk phase, Eq. (4.8), is placed on a closed manifold, then because the dyon operator $\mathcal{D}_{(-N/2,1)}(\gamma, \Omega)$ is supported on a loop γ attached to an open surface Ω , the loop γ is always contractible. However, if Eq. (4.8) is placed on an open manifold X with a boundary ∂X , then γ can be noncontractible on ∂X with Ω extending into the bulk. Since $\mathcal{D}_{(-N/2,1)}(\gamma, \Omega)$ has a perimeter law, it will be deconfined on ∂X . Fusing the $(-N/2, 1)$ anti-semion with itself generates the dyons in Eq. (4.11) that contribute to the bulk topological order. Because these dyons braid trivially with all other anyons on ∂X , the topological order along ∂X is non-modular and cannot exist in a purely (2+1)d theory, just as in boundary topological orders of Walker-Wang models [101, 102].

To summarize, the negative m phase has $\mathbb{Z}_{N/2}$ topological order that results from spontaneously breaking the \mathbb{Z}_N magnetic one-form symmetry to \mathbb{Z}_2 . This topological order is enriched by the $G = SO(N_f) \times \mathbb{Z}_2^T$ zero-form symmetry and the unbroken \mathbb{Z}_2 magnetic one-form symmetry. The topological order is stacked with a nontrivial SPT for the symmetry G and the \mathbb{Z}_2 one-form symmetry, and it may be further enriched by G if the $PSU(N)$ monopoles carry fractional quantum numbers under G . The transition between this phase and the positive m phase discussed in Section 4.2 is depicted schematically in Figure 2. While we have focused on $SU(N)/\mathbb{Z}_N$ gauge theory coupled to N_f Majorana fermions in the adjoint representation, similar constructions exist for other gauge groups, which we discuss in Appendix D.

4.4 String tension critical exponent

Since we have an example of a critical point between phases with different patterns of one-form symmetry breaking, it is natural to consider critical exponents for loop operators. In the $PSU(N)$ theory coupled to adjoint fermions, the basic 't Hooft line $\mathcal{T}(\gamma)$ has an area law in the negative mass phase, but this monopole becomes deconfined both at the critical point, $m = 0$, and in the positive mass phase. As m approaches zero from the negative mass phase, the string tension of $\mathcal{T}(\gamma)$ goes to zero as

$$\sigma \sim |m|^\mu \tag{4.13}$$

for some critical exponent μ , which can be reliably computed for $N_f > 5$ so that the $m = 0$ theory is infrared free [103, 104].

For $m < 0$, the low energy physics, well below the fermion mass $|m|$, is governed by pure $PSU(N)$ Yang-Mills at $\theta = \pi N N_f$. On grounds of dimensional analysis, the string tension of $\mathcal{T}(\gamma)$ scales as

$$\sigma \sim (\Lambda_{\text{YM}})^2, \quad (4.14)$$

where Λ_{YM} is the energy scale dynamically generated in the pure non-Abelian gauge theory. To compute the critical exponent μ , we must then relate Λ_{YM} to the mass scale $|m|$ of the fermions. For energies far above $|m|$, the beta function for g is well-approximated by Eq. (3.13). In contrast, at energies far below $|m|$, the fermions may be integrated out, and the beta function is governed by that of pure Yang-Mills. To estimate the string tension, we approximate the beta function as

$$\beta(g) = \frac{dg}{d\ell} = \begin{cases} (c_{\text{YM}} - c_{\text{M}}) g^3, & \Lambda > |m|, \\ c_{\text{YM}} g^3, & \Lambda < |m|, \end{cases} \quad (4.15)$$

where $\Lambda = \Lambda_{\text{UV}} e^{-\ell}$ is the energy associated with scale ℓ and Λ_{UV} is a high energy reference scale.

Integrating Eq. (4.15) from Λ_{UV} to the mass scale gives

$$g^2(|m|) = \frac{g^2(\Lambda_{\text{UV}})}{1 + 2(c_{\text{M}} - c_{\text{YM}}) g^2(\Lambda_{\text{UV}}) \ln(\Lambda_{\text{UV}}/|m|)}. \quad (4.16)$$

We also integrate Eq. (4.15) from $|m|$ to an energy scale Λ_0 below $|m|$, resulting in

$$g^2(\Lambda_0) = \frac{g^2(|m|)}{1 - 2c_{\text{YM}} g^2(|m|) \ln(|m|/\Lambda_0)}. \quad (4.17)$$

The scale Λ_{YM} is determined by setting $1/g^2(\Lambda_{\text{YM}}) = 0$ using Eq. (4.17). Solving for Λ_{YM} and using the expression in Eq. (4.16) for $g^2(|m|)$ gives

$$\Lambda_{\text{YM}} = |m| \exp\left(-\frac{1}{2c_{\text{YM}} g^2(|m|)}\right) = \Lambda_{\text{UV}} e^{-1/(2c_{\text{YM}} g^2(\Lambda_{\text{UV}}))} \left(\frac{|m|}{\Lambda_{\text{UV}}}\right)^{c_{\text{M}}/c_{\text{YM}}}. \quad (4.18)$$

Although this calculation is based on Eq. (4.15), this approximation becomes reliable as $|m|/\Lambda_{\text{UV}} \rightarrow 0$ if the $m = 0$ point is infrared free [103, 104]. In this case, the coupling g is small at energy scales near $|m|$.

Thus, the critical exponent μ for the string tension of the 't Hooft loop $\mathcal{T}(\gamma)$ is

$$\mu = \frac{2c_{\text{M}}}{c_{\text{YM}}} = \frac{4}{11} \sum_{\mathcal{R}} \frac{I_{\mathcal{R}}}{I_{\text{adj}}} N_f(\mathcal{R}) = \frac{4N_f}{11}, \quad (4.19)$$

where we specialized to N_f flavors of adjoint fermions in the last step. Note that we did not have to specialize to a particular gauge group, so this critical exponent is the same for the transitions in other gauge groups discussed in Appendix D. It is interesting to compare this exponent with the prediction $\mu = 1/2$ from mean string field theory [27]. For the values of N_f where we expect Eq. (4.19) to be valid (i.e., $N_f \geq 7$), we have $\mu \geq 28/11 \approx 2.55$. Thus, in these cases, the string tension vanishes more strongly as the critical point is approached than mean string field theory predicts.

4.5 Topological transition on the lattice

Here, we provide a lattice gauge theory model that realizes the topological phase transition using the Hamiltonian formalism [105]. See Refs. [106–108] for a review of lattice gauge theory.¹⁶ First, we review how to place $PSU(N) = SU(N)/\mathbb{Z}_N$ gauge theory on the lattice [67, 109, 110] by gauging the \mathbb{Z}_N electric one-form symmetry of $SU(N)$ lattice gauge theory.

We work on a cubic lattice. On each link $(\mathbf{r}; j)$, which is labeled by a lattice point \mathbf{r} and a direction j , we place an $SU(N)$ gauge variable $U_j(\mathbf{r}) = e^{iT^\alpha A_j^\alpha(\mathbf{r})}$, where T^α are the generators of the $\mathfrak{su}(N)$ Lie algebra in the fundamental representation. On each plaquette $(\mathbf{r}; j, k)$, we define the operator,

$$W_{jk}(\mathbf{r}) = U_j(\mathbf{r}) U_k(\mathbf{r} + \mathbf{e}_j) U_j^{-1}(\mathbf{r} + \mathbf{e}_k) U_k^{-1}(\mathbf{r}), \quad (4.20)$$

where \mathbf{e}_j is a unit vector in the j direction (and we set the lattice constant to unity). The Hamiltonian for $SU(N)$ lattice gauge theory is

$$H_{SU(N)} = \frac{g^2}{2} \sum_{\mathbf{r}; j} E_j^\alpha(\mathbf{r}) E_j^\alpha(\mathbf{r}) - \frac{1}{2g^2} \sum_{\mathbf{r}; j, k} \text{Tr} \left(W_{jk}(\mathbf{r}) + W_{jk}^\dagger(\mathbf{r}) \right), \quad (4.21)$$

where the self-adjoint operators $E_j^\alpha(\mathbf{r})$ that are defined on links obey

$$[A_j^\alpha(\mathbf{r}), E_k^\beta(\mathbf{r}')] = i \delta_{\mathbf{r}, \mathbf{r}'} \delta_{j, k} \delta^{\alpha, \beta}, \quad [E_j^\alpha(\mathbf{r}), E_k^\beta(\mathbf{r}')] = i f^{\alpha\beta\gamma} E_j^\gamma(\mathbf{r}) \delta_{\mathbf{r}, \mathbf{r}'} \delta_{j, k}, \quad (4.22)$$

which also implies

$$\begin{aligned} [E_j^\alpha(\mathbf{r}), U_k(\mathbf{r}')] &= T^\alpha U_j(\mathbf{r}) \delta_{\mathbf{r}, \mathbf{r}'} \delta_{j, k}, \\ e^{i\xi^\alpha E_j^\alpha(\mathbf{r})} U_k(\mathbf{r}') e^{-i\xi^\beta E_j^\beta(\mathbf{r})} &= \exp(i\xi^\alpha T^\alpha \delta_{\mathbf{r}, \mathbf{r}'} \delta_{j, k}) U_k(\mathbf{r}'), \end{aligned} \quad (4.23)$$

¹⁶We note, however, that the Gauss law for $SU(2)$ lattice gauge theory is incorrect in Ref. [108]. The author of this (otherwise excellent) textbook is aware of this issue.

for some parameters ξ^α . The generators of $SU(N)$ gauge transformations are

$$Q^\alpha(\mathbf{r}) = \sum_{j=1}^3 \left[E_j^\alpha(\mathbf{r}) - (U_j^{\text{adj}})^{\beta\alpha}(\mathbf{r} - \mathbf{e}_j) E_j^\beta(\mathbf{r} - \mathbf{e}_j) \right], \quad (4.24)$$

where $(U_j^{\text{adj}})^{\alpha\beta}(\mathbf{r}) = e^{i(T_{\text{adj}}^\gamma)^{\alpha\beta} A_j^\gamma(\mathbf{r})} = e^{f^{\alpha\beta\gamma} A_j^\gamma(\mathbf{r})}$ are matrix elements of the gauge variables in the adjoint representation. The gauge charges $Q^\alpha(\mathbf{r})$ commute with the Hamiltonian, $[Q^\alpha(\mathbf{r}), H_{SU(N)}] = 0$, and annihilate physical states so that $Q^\alpha(\mathbf{r}) |\text{phys}\rangle = 0$.

To obtain $PSU(N)$ lattice gauge theory, we now gauge the \mathbb{Z}_N electric one-form symmetry of $SU(N)$ lattice gauge theory. To couple to a \mathbb{Z}_N two-form gauge field, we introduce unitary operators $\sigma_{jk}(\mathbf{r})$ and $\tau_{jk}(\mathbf{r})$ on plaquettes that satisfy

$$[\sigma_{jk}(\mathbf{r})]^N = [\tau_{jk}(\mathbf{r})]^N = 1, \quad \sigma_{jk}(\mathbf{r}) \tau_{j'k'}(\mathbf{r}') = \exp\left(\frac{2\pi i}{N} \delta_{\mathbf{r}, \mathbf{r}'} \delta_{j, j'} \delta_{k, k'}\right) \tau_{j'k'}(\mathbf{r}') \sigma_{jk}(\mathbf{r}). \quad (4.25)$$

In addition, we have $\tau_{kj}(\mathbf{r}) = \tau_{jk}^\dagger(\mathbf{r})$ and $\sigma_{kj}(\mathbf{r}) = \sigma_{jk}^\dagger(\mathbf{r})$. The three-form analogue of a field strength, defined on cubes $(\mathbf{r}; 1, 2, 3)$, is

$$h_{123}(\mathbf{r}) = \sigma_{12}(\mathbf{r}) \sigma_{12}^\dagger(\mathbf{r} + \mathbf{e}_3) \sigma_{23}(\mathbf{r}) \sigma_{23}^\dagger(\mathbf{r} + \mathbf{e}_1) \sigma_{31}(\mathbf{r}) \sigma_{31}^\dagger(\mathbf{r} + \mathbf{e}_2). \quad (4.26)$$

The Hamiltonian for $PSU(N)$ lattice gauge theory is

$$\begin{aligned} H_{PSU(N)} = & \frac{g^2}{2} \sum_{\mathbf{r}; j} E_j^\alpha(\mathbf{r}) E_j^\alpha(\mathbf{r}) - \frac{1}{2g^2} \sum_{\mathbf{r}; j, k} \text{Tr} \left(W_{jk}(\mathbf{r}) \sigma_{jk}(\mathbf{r}) + W_{jk}^\dagger(\mathbf{r}) \sigma_{jk}^\dagger(\mathbf{r}) \right) \\ & - \frac{1}{2\tilde{g}^2} \sum_{\mathbf{r}} \left(h_{123}(\mathbf{r}) + h_{123}^\dagger(\mathbf{r}) \right) - \frac{1}{2} \sum_{\mathbf{r}; j, k} \left(\tau_{jk}(\mathbf{r}) + \tau_{jk}^\dagger(\mathbf{r}) \right). \end{aligned} \quad (4.27)$$

The \mathbb{Z}_N two-form gauge field has its own Gauss law. To define the operator for this gauge charge, we must identify the operator that generates the \mathbb{Z}_N center transformation of $SU(N)$ on each link. We can always take a generator of $SU(N)$ to be diagonal, which we take to be

$$T^{\alpha_0} = \sqrt{\frac{N}{2(N-1)}} \text{diag} \left(\frac{1}{N}, \frac{1}{N}, \dots, \frac{1}{N}, -1 + \frac{1}{N} \right) \quad (4.28)$$

for some α_0 . A \mathbb{Z}_N center transformation on $U_k(\mathbf{r})$ is then given by

$$e^{i\eta E_j^{\alpha_0}(\mathbf{r})} U_k(\mathbf{r}') e^{-i\eta E_j^{\alpha_0}(\mathbf{r})} = e^{2\pi i \delta_{j,k} \delta_{\mathbf{r}, \mathbf{r}'}/N} U_k(\mathbf{r}'), \quad (4.29)$$

where $\eta = 2\pi\sqrt{2(N-1)/N}$. For each link (\mathbf{r}, j) , the generator of gauge transformations for the two-form gauge field is

$$\mathcal{Q}_j(\mathbf{r}) = e^{i\eta E_j^{\alpha_0}(\mathbf{r})} \prod_{k \neq j} \tau_{jk}(\mathbf{r}) \tau_{jk}^\dagger(\mathbf{r} - \mathbf{e}_k). \quad (4.30)$$

This operator commutes with the Hamiltonian, $[\mathcal{Q}_j(\mathbf{r}), H_{PSU(N)}] = 0$, and physical states must be invariant under this operator, $\mathcal{Q}_j(\mathbf{r}) |\text{phys}\rangle = |\text{phys}\rangle$. This constraint also ensures that the 't Hooft loop, which must be attached to a surface in $SU(N)$ gauge theory, becomes a genuine loop operator in $PSU(N)$ gauge theory. Indeed, the 't Hooft loop in $SU(N)$ gauge theory, which defined on a loop $\tilde{\gamma}$ on the dual lattice attached to a surface $\tilde{\Sigma}$, is

$$\mathcal{T}(\tilde{\gamma}) = \prod_{(\mathbf{r}; j) \in \tilde{\Sigma}} e^{-i\eta E_j^{\alpha_0}(\mathbf{r})}, \quad (4.31)$$

where the product is over links intersecting $\tilde{\Sigma}$. However, the invariance of states under $\mathcal{Q}_j(\mathbf{r})$ in $PSU(N)$ gauge theory ensures that the choice of surface $\tilde{\Sigma}$ does not matter since this operator acts on physical states in the same way as

$$\mathcal{T}(\tilde{\gamma}) = \prod_{(\mathbf{r}; j, k) \in \tilde{\gamma}} \tau_{jk}(\mathbf{r}), \quad (4.32)$$

where the product is now over plaquettes intersecting the loop $\tilde{\gamma}$.

Before coupling the gauge fields to fermions, let us check that \mathbb{Z}_N topological order is produced in a certain limit of the $PSU(N)$ lattice gauge theory. We take the limit $\tilde{g}^2 \rightarrow 0$ so that the terms in the Hamiltonian with $\tau_{jk}(\mathbf{r})$ and its Hermitian conjugate may be ignored. Now the ground state $|\psi_0\rangle$ must satisfy,

$$h_{123}(\mathbf{r}) |\psi_0\rangle = h_{123}^\dagger(\mathbf{r}) |\psi_0\rangle = |\psi_0\rangle, \quad (4.33)$$

so the 't Hooft loops cannot end. In this limit, there is an exact \mathbb{Z}_N magnetic one-form global symmetry, as in the continuum $PSU(N)$ gauge theory. For finite \tilde{g}^2 (but not too large), we expect that this symmetry will be emergent. The operator that acts with the \mathbb{Z}_N one-form symmetry, defined on a noncontractible surface Σ ,

$$\mathcal{U}(\Sigma) = \prod_{(\mathbf{r}; j, k) \in \Sigma} \sigma_{jk}(\mathbf{r}), \quad (4.34)$$

now commutes with the Hamiltonian. Taking $g^2 \rightarrow \infty$ also, the ground state must satisfy,

$$E_j^\alpha(\mathbf{r}) |\psi_0\rangle = 0, \quad (4.35)$$

so that

$$\langle \psi_0 | \mathcal{T}(\tilde{\gamma}) | \psi_0 \rangle = 1 \quad (4.36)$$

for any contractible loop $\tilde{\gamma}$ on the dual lattice. For finite g , the 't Hooft loop will instead have a perimeter law. Hence, the \mathbb{Z}_N magnetic one-form symmetry will ultimately be spontaneously broken at low energies by deconfined monopoles, leading to \mathbb{Z}_N topological order.

Another way to observe the topological order is to note that for $g^2 \rightarrow \infty$ the 't Hooft loop $\mathcal{T}(\tilde{\gamma})$ commutes with the Hamiltonian. For noncontractible $\tilde{\gamma}$ this operator generates a \mathbb{Z}_N two-form global symmetry that acts on the surface operator $\mathcal{U}(\Sigma)$ for noncontractible Σ . For example, we take periodic boundary conditions in all directions. Let $\tilde{\gamma}$ be a noncontractible loop in the j direction and Σ be a plane perpendicular to the j direction. The operators $\mathcal{U}(\Sigma)$ and $\mathcal{T}(\tilde{\gamma})$ both commute with the Hamiltonian and obey

$$\mathcal{U}(\Sigma) \mathcal{T}(\tilde{\gamma}) = e^{2\pi i/N} \mathcal{U}(\Sigma) \mathcal{T}(\tilde{\gamma}), \quad (4.37)$$

which implies a ground state degeneracy of N^3 on the torus, thus confirming the \mathbb{Z}_N topological order.

Finally, we introduce fermions. On each site \mathbf{r} , we place a four-component Majorana spinor $\psi_{\mathbf{a}}(\mathbf{r})$, which obeys the anticommutation relations,

$$\{\psi_{\mathbf{a}}(\mathbf{r}), \psi_{\mathbf{b}}(\mathbf{r}')\} = (\gamma^0 \mathcal{C}^{-1})_{\mathbf{ab}} \delta_{\mathbf{r}, \mathbf{r}'}, \quad (4.38)$$

where \mathbf{a} and \mathbf{b} are spinor indices and \mathcal{C} is the charge conjugation matrix (see Appendix B). The Hamiltonian for a single free Majorana fermion $\psi(\mathbf{r})$ of mass m is

$$H_M = \sum_{\mathbf{r}} \frac{1}{2} \psi^T(\mathbf{r}) \mathcal{C} (D_W + m) \psi(\mathbf{r}), \quad (4.39)$$

where $\psi^T(\mathbf{r})$ is the transpose of the spinor $\psi(\mathbf{r})$ and D_W is the Wilson operator [45],

$$D_W = \frac{1}{2} \sum_{j=1}^3 [-i \gamma^j (\Delta_j^+ + \Delta_j^-) - (\Delta_j^+ - \Delta_j^-)], \quad (4.40)$$

where $\Delta_j^+ \psi(\mathbf{r}) = \psi(\mathbf{r} + \mathbf{e}_j) - \psi(\mathbf{r})$ and $\Delta_j^- \psi(\mathbf{r}) = \psi(\mathbf{r}) - \psi(\mathbf{r} - \mathbf{e}_j)$. This model describes a time-reversal invariant topological superconductor in class DIII, characterized by a topological invariant ν , which is classified by \mathbb{Z} for free fermions [111–113] but collapses to \mathbb{Z}_{16} when interactions are taken into account [71, 114–116]. The topological invariant for Eq. (4.39) as a function of m is

$$\nu = \begin{cases} 0, & m < -6, m > 0, \\ -1, & -6 < m < -4, -2 < m < 0, \\ 2, & -4 < m < -2. \end{cases} \quad (4.41)$$

In particular, notice that the $m > 0$ phase is a trivial superconductor, and there is a transition at $m = 0$, where the Majorana fermion becomes massless, to the phase at $-2 < m < 0$, which is a topological superconductor with a gravitational response of $\theta_g = \pi$ (see Appendix C).

Thus, in the continuum limit, the transition at $m = 0$ becomes the transition of a single Majorana fermion whose mass changes sign. If we take $N_f (N^2 - 1)$ copies of this Majorana fermion and couple them to a $PSU(N)$ gauge field in the adjoint representation, then for N even and a sufficiently large odd number of flavors N_f , there will be a continuous transition at $m = 0$ that corresponds to the topological transition discussed for the continuum $PSU(N)$ adjoint QCD theory.

To couple fermion fields $\psi(\mathbf{r})$ that transform in a representation \mathcal{R} of the gauge field, we define

$$\begin{aligned} D_j^+(\mathcal{R}) \psi(\mathbf{r}) &= U_j^{\mathcal{R}}(\mathbf{r}) \psi(\mathbf{r} + \mathbf{e}_j) - \psi(\mathbf{r}), \\ D_j^-(\mathcal{R}) \psi(\mathbf{r}) &= \psi(\mathbf{r}) - (U_j^{\mathcal{R}})^{-1}(\mathbf{r} - \mathbf{e}_j) \psi(\mathbf{r} - \mathbf{e}_j), \end{aligned} \quad (4.42)$$

where $U_j^{\mathcal{R}}(\mathbf{r}) = e^{i T_{\mathcal{R}}^{\alpha} A_j^{\alpha}(\mathbf{r})}$ are the gauge variables in representation \mathcal{R} . We also define

$$\mathcal{D}_W^{\mathcal{R}} = \frac{1}{2} \sum_{j=1}^3 [-i \gamma^j (D_j^+(\mathcal{R}) + D_j^-(\mathcal{R})) - (D_j^+(\mathcal{R}) - D_j^-(\mathcal{R}))]. \quad (4.43)$$

Taking \mathcal{R} to be the adjoint representation and introducing N_f flavors of Majorana fermions, indexed by J , the Hamiltonian is

$$H = \sum_{J=1}^{N_f} \sum_{\mathbf{r}} \frac{1}{2} \psi_J^T(\mathbf{r}) \mathcal{C} (\mathcal{D}_W^{\text{adj}} + m) \psi_J(\mathbf{r}) + H_{PSU(N)}, \quad (4.44)$$

where $H_{PSU(N)}$ is given in Eq. (4.27). Even with the coupling to fermions, the one-form gauge charge $\mathcal{Q}_j(\mathbf{r})$, defined in Eq. (4.30), remains the same. However, the generator of zero-form gauge transformations, Eq. (4.24), must be modified to

$$Q^{\alpha}(\mathbf{r}) = \sum_{j=1}^3 \left[E_j^{\alpha}(\mathbf{r}) - (U_j^{\text{adj}})^{\beta\alpha}(\mathbf{r} - \mathbf{e}_j) E_j^{\beta}(\mathbf{r} - \mathbf{e}_j) \right] - \frac{1}{2} \sum_{J=1}^{N_f} \psi_J^T(\mathbf{r}) \mathcal{C} \gamma^0 T_{\text{adj}}^{\alpha} \psi_J(\mathbf{r}). \quad (4.45)$$

This operator commutes with the Hamiltonian, Eq. (4.44). As discussed above, for N even and sufficiently large odd N_f , there will be a continuous transition in the lattice Hamiltonian, Eq. (4.44), at $m = 0$. Taking g^2 large and \tilde{g}^2 small, the phase for $m > 0$ will be the \mathbb{Z}_N topological order discussed in Section 4.2, and the $-2 < m < 0$ phase is the SET with $\mathbb{Z}_{N/2}$ topological order explained in Section 4.3.

As noted previously, for finite \tilde{g}^2 the \mathbb{Z}_N magnetic one-form symmetry that is present in the continuum theory is explicitly broken by dynamical magnetic monopoles. This symmetry is expected to be emergent for small (but still finite) \tilde{g}^2 . Because the magnetic monopoles in this lattice model, Eq. (4.44), transform trivially under the $SO(N_f)$ flavor symmetry and

time-reversal symmetry, the SET orders arising from the Hamiltonian in Eq. (4.44) will have the trivial symmetry fractionalization class—the deconfined anyons will be bosonic Kramers singlets and $SO(N_f)$ tensors.

5 SET to non-invertible SSB transition

As we have established, $PSU(N)$ adjoint QCD with a sufficiently large odd number of flavors N_f can appear at a critical point between two SETs with different topological orders, provided that both the $SO(N_f)$ flavor symmetry and time-reversal symmetry are preserved. Recent work has introduced non-invertible analogues of time-reversal symmetry [33, 34], which are also symmetries of massless $PSU(N)$ adjoint QCD. This observation then raises the question of what kinds of phases can emerge if we add perturbations that respect the $SO(N_f)$ flavor symmetry and the non-invertible time-reversal symmetry.

In this section, we will analyze the theory with action,

$$S_n = S_0 + \frac{\pi n}{8\pi^2} \int \{ \text{Tr}[(f_\alpha + b \mathbb{I}_N) \wedge (f_\alpha + b \mathbb{I}_N)] - [\text{Tr}(f_\alpha) + Nb] \wedge [\text{Tr}(f_\alpha) + Nb] \}, \quad (5.1)$$

where S_0 is the $PSU(N)$ adjoint QCD action defined in Eq. (4.1), which is invariant under the standard invertible time-reversal symmetry. The additional term is a $PSU(N)$ theta term with $\theta = \pi n$.

Below, we will review the notion of non-invertible time-reversal symmetry and show that Eq. (5.1) respects this symmetry. Imposing this non-invertible time-reversal symmetry and the $SO(N_f)$ flavor symmetry, for large enough odd N_f , we can again have a direct continuous transition tuned by a single parameter. For N and N_f both odd, as we demonstrate below, this transition can be between a topologically ordered phase, which breaks the magnetic one-form symmetry to a subgroup, and a phase that breaks the non-invertible time-reversal symmetry spontaneously. We will primarily focus on the case $N = k^2$ and $n = 2k$, where $k > 1$ is odd, to keep the discussion simple, though these choices for N and n are not the only ones that give transitions between a topologically order phase and a non-invertible SSB phase.

5.1 Non-invertible time-reversal

To establish the non-invertible symmetry, it is useful to define the following transformations on a theory with a \mathbb{Z}_N one-form global symmetry with action $S[B]$,

$$\begin{aligned}\mathbf{S} : \quad & S[B] \rightarrow S[b] + \frac{N}{2\pi} \int b \wedge B, \\ \mathbf{T} : \quad & S[B] \rightarrow S[B] + \frac{N}{4\pi} \int B \wedge B, \\ \mathbf{C} : \quad & S[B] \rightarrow S[-B],\end{aligned}\tag{5.2}$$

where $B_{\mu\nu}$ is a background \mathbb{Z}_N two-form gauge field for the \mathbb{Z}_N one-form global symmetry and $b_{\mu\nu}$ is a dynamical \mathbb{Z}_N two-form gauge field. These transformations obey

$$\mathbf{S}^2 = \mathbf{C}, \quad \mathbf{C}^2 = 1, \quad \mathbf{T}^N = 1.\tag{5.3}$$

To summarize, \mathbf{S} denotes gauging of the \mathbb{Z}_N one-form symmetry, \mathbf{T} stacks a \mathbb{Z}_N one-form SPT, and \mathbf{C} simply changes the sign of the background field.

A non-invertible time-reversal transformation can be defined by [33, 34]

$$\mathbf{K}_n = \mathbf{CST}^{(N-1)n}\mathbf{SK},\tag{5.4}$$

where \mathbf{K} is the standard invertible time-reversal transformation and n is an integer mod N . From the relations, Eq. (5.3), we observe that if $n \in N\mathbb{Z}$, then \mathbf{K}_n reduces to \mathbf{K} . We recall that the theta term is odd under \mathbf{K} , and we observe that $\mathbf{CST}^{(N-1)n}\mathbf{S}$ shifts $\theta \rightarrow \theta + 2\pi n$. Hence, the non-invertible time-reversal transformation maps

$$\mathbf{K}_n : \quad \theta \rightarrow -\theta + 2\pi n.\tag{5.5}$$

Since θ has periodicity $2\pi N$, the values of θ that are invariant under \mathbf{K}_n are $\theta = \pi n$ and $\theta = \pi(n + N)$. Because the theta term added to Eq. (5.1) has $\theta = \pi n$, \mathbf{K}_n leaves Eq. (5.1) invariant as promised.

For $n \notin N\mathbb{Z}$, \mathbf{K}_n cannot be a transformation associated with a unitary operator because it involves gauging \mathbf{S} . However, it may be associated with an operator $\mathbf{T}_n = \mathbf{D}_n\mathbf{T}$ that is topological. Here, \mathbf{T} is the standard invertible time-reversal operator (see Section 4.1) and \mathbf{D}_n is defined as [33, 34, 99, 100]

$$\mathbf{D}_n = \int \mathcal{D}a_1 \mathcal{D}a_2 \exp \left[i \oint_Y \left(\frac{N(N-1)n}{4\pi} a_1 \wedge da_1 + \frac{N}{2\pi} a_1 \wedge da_2 + \frac{N}{2\pi} a_2 \wedge b \right) \right],\tag{5.6}$$

where Y is a closed three-dimensional manifold in spacetime while $(a_1)_\mu$ and $(a_2)_\mu$ are dynamical $U(1)$ one-form gauge fields defined only on Y . Hence, \mathbf{T}_n is constructed by decorating

the invertible time-reversal operator T with a particular fractional quantum Hall state. The operator T_n obeys the fusion rules,

$$\begin{aligned}\mathsf{T}_n \times (\mathsf{T}_n)^\dagger &= \mathsf{D}_n \times \mathsf{D}_{-n} = (\mathcal{Z}_N)_{N(N-1)n} \mathcal{C}_0, \\ \mathsf{T}_n \times \mathsf{T}_n &= \mathsf{D}_n \times \mathsf{D}_{-n} \times \mathsf{T}^2 = (\mathcal{Z}_N)_{N(N-1)n} \mathcal{C}_0 (-1)^F,\end{aligned}\tag{5.7}$$

where $(-1)^F$ is fermion parity, $(\mathcal{Z}_N)_{N(N-1)n}$ is the partition function for \mathbb{Z}_N Chern-Simons theory at level $N(N-1)n$,

$$(\mathcal{Z}_N)_{(N-1)nN} = \int \mathcal{D}a_1 \mathcal{D}a_2 \exp \left[i \oint_Y \left(\frac{N}{2\pi} a_1 \wedge da_2 + \frac{N(N-1)n}{4\pi} a_1 \wedge da_1 \right) \right], \tag{5.8}$$

and \mathcal{C}_0 is a condensation defect [117–121],

$$\mathcal{C}_0 = \int \mathcal{D}a_1 \mathcal{D}a_2 \exp \left(i \oint_Y \frac{N}{2\pi} (a_1 \wedge da_2 + a_1 \wedge b) \right), \tag{5.9}$$

where the \mathbb{Z}_N one-form symmetry is gauged only along Y .

The operator T_n is topological in the theory, Eq. (5.1), for any value of the fermion mass m , so we refer to T_n as a non-invertible time-reversal symmetry. In the remainder of Section 5, we will examine the phases of Eq. (5.1) as a function of the T_n preserving mass m . Importantly, the non-invertible symmetry can have important non-perturbative constraints on the phase diagram, a notion explored for (3+1)d systems in a number of recent works [56, 122–128]. As we demonstrate in Appendix E, for N odd and n such that $\gcd(N, n) > 1$, the non-invertible time-reversal symmetry and \mathbb{Z}_N magnetic one-form symmetry have a mixed anomaly in the sense that no single ground state can preserve both these symmetries. The phase diagram we find, Figure 3, is consistent with this constraint.

5.2 SET phase

For $m > 0$, at energies far below the mass scale of the fermions, the physics is governed by pure $PSU(N)$ gauge theory with $\theta = \pi n$. Taking n even with $n = 2k$, the low energy physics, as established in Section 2.2, is described by the TQFT,

$$S_{\text{TQFT}}[(N-1)k, \tilde{a}, b, B] = \frac{N(N-1)k}{4\pi} \int b \wedge b + \frac{N}{2\pi} \int b \wedge (d\tilde{a} + B), \tag{5.10}$$

where \tilde{a}_μ is a dynamical $U(1)$ one-form gauge field, $b_{\mu\nu}$ is a dynamical $U(1)$ two-form gauge field, and $B_{\mu\nu}$ is a background \mathbb{Z}_N two-form gauge field. As discussed in Section 2.2 and Appendix A, the physics of this phase depends on $\gcd(N, k)$. If $\gcd(N, k) = 1$, then this

phase is an SPT protected by the \mathbb{Z}_N magnetic one-form symmetry and is also invariant under the non-invertible time-reversal symmetry. Otherwise, there is topological order. For $n = 0$, the time-reversal symmetry is invertible, and this phase has \mathbb{Z}_N topological order described by BF theory with level N .

To focus on a class of examples in which this phase has topological order enriched by a *non-invertible* time-reversal symmetry, we take $N = k^2$ with $k > 1$ odd, which we assume for the remainder of Section 5.2. Within the $m > 0$ phase, the $\mathbb{Z}_N = \mathbb{Z}_{k^2}$ magnetic one-form symmetry is spontaneously broken to \mathbb{Z}_k at low energies, resulting in $\mathbb{Z}_{k^2}/\mathbb{Z}_k \cong \mathbb{Z}_k$ topological order, and the unbroken \mathbb{Z}_k one-form symmetry has nontrivial SPT order. This topological order is also enriched by the unbroken $SO(N_f)$ symmetry.

To understand the \mathbb{Z}_k topological order more physically, we observe from Eq. (2.21) that the dyons in this phase with a perimeter law are of the form $(-k q_m, q_m)$ for an integer $q_m \bmod N$. The non-invertible time-reversal symmetry T_{2k} maps the charges of a dyon as

$$(q_e, q_m) \rightarrow (-q_e - 2k q_m, q_m). \quad (5.11)$$

Thus, each dyon with a perimeter law in this phase is mapped to itself (up to a neutral fermion) under T_{2k} .

The dyons with a perimeter law that are also associated with genuine loop operators are the purely magnetic charges,

$$(q_e, q_m) = (0, kq) \bmod k^2, \quad (5.12)$$

where $q \in \{0, 1, \dots, k-1\}$. Thus, there are k anyon particles associated with the 't Hooft loops $\mathcal{T}(\gamma)^{kq}$. These 't Hooft loops can also have nontrivial correlation functions with the electric flux tubes associated with the surface operators $\mathcal{U}(\Sigma)$,

$$\langle \mathcal{T}(\gamma)^k \mathcal{U}(\Sigma) \rangle = \exp \left(\frac{2\pi i}{k} \Phi(\gamma, \Sigma) \right), \quad (5.13)$$

where $\Phi(\gamma, \Sigma)$ is the linking number of the loop γ and surface Σ in spacetime.

Following the analysis of the TQFT, Eq. (5.10), in Appendix A, the response to a background field $B_{\mu\nu}$ for the unbroken \mathbb{Z}_k one-form symmetry is

$$S_{\text{resp}}[B] = \frac{k}{4\pi} \int B \wedge B. \quad (5.14)$$

To interpret this response, we note that the symmetry operator for the unbroken \mathbb{Z}_k one-form symmetry is $\mathcal{U}(\Sigma)^k$, which describes the worldsheet of an electric flux loop with flux $-k$.

When this operator ends on a loop, the loop must carry electric charge $-k$, and the response indicates that it also carries magnetic charge 1. Indeed, this operator will be nontrivial in the low energy limit only if the loop on which it ends has a perimeter law. Hence, the physical meaning of Eq. (5.14) is that the $(-k, 1)$ dyon, which is not genuine loop operator, has a perimeter law.

5.3 Non-invertible SSB

Next, we determine the phase that arises for $m < 0$. In this case, at energies well below the scale of the fermion mass $|m|$, the fermions may be integrated out to give an effective action of

$$S_{\text{eff}} = S_{PSU(N)}[\pi(n + NN_f), B] + \frac{\pi(N^2 - 1)}{8\pi^2} \int \text{Tr}(F \wedge F) + \frac{\pi N_f(N^2 - 1)}{384\pi^2} \int \text{Tr}(R \wedge R), \quad (5.15)$$

where $F_{\mu\nu}$ is the field strength of the background field A_μ for the $SO(N_f)$ flavor symmetry, R is the curvature two-form (see Appendix C), and $S_{PSU(N)}[\pi(n + NN_f), B]$ is the action for pure $PSU(N)$ gauge theory (cf. Eqs. (2.13) and (2.15)) with $\theta = \pi(n + NN_f)$, which is equivalent to $\theta = \pi(n + N)$ for odd N_f . For odd N and even n , then θ is an odd multiple of π . As we discuss below, this phase spontaneously breaks the non-invertible time-reversal symmetry T_n . We can then ignore the theta terms for the background fields $F_{\mu\nu}$ and R in Eq. (5.15). Since this phase does not have unbroken time-reversal symmetry (invertible or not), the theta terms for these background fields may be continuously tuned to zero without encountering a phase transition.

As we reviewed in Section 2.2, there is evidence that at $\theta = \pi\tilde{n}$ for odd integer \tilde{n} , the ground states at $\theta = \pi(\tilde{n} - 1)$ and $\theta = \pi(\tilde{n} + 1)$ are degenerate, which may be established more rigorously for large N [64]. This IR behavior may be interpreted as the spontaneous breaking of the non-invertible time-reversal symmetry associated with the operator T_n , which exchanges the ground state(s) at $\theta = \pi(\tilde{n} - 1)$ with the state(s) at $\theta = \pi(\tilde{n} + 1)$ for $\tilde{n} = n + N$. In the special case $n = 0$ (or equivalently, $\tilde{n} = N$), the time-reversal symmetry is invertible and spontaneously broken in this phase, at least for large enough N .

For the remainder of Section 5.3, let us take $N = k^2$ and $n = 2k$ for odd $k > 1$. The time-reversal symmetry operator T_{2k} is non-invertible, and we can still in principle access the large N limit by taking k large. We then have $\tilde{n} = 2k + N = 2k + k^2$. The TQFTs describing the states at $\theta = \pi(\tilde{n} \pm 1)$ are

$$S^\pm = \frac{N(N - 1)(2k + N \pm 1)}{8\pi} \int b \wedge b + \frac{N}{2\pi} \int b \wedge (d\tilde{a} + B), \quad (5.16)$$




Figure 3: Schematic phase diagram of $PSU(N) = SU(N)/\mathbb{Z}_N$ adjoint QCD with a theta term $\theta = \pi n$, Eq. (5.1), for N_f odd flavors of Majorana fermions as a function of the fermion mass m . Here, we take $N = k^2$ and $n = 2k$ with $k > 1$ odd. In the $m > 0$ phase, described by Eq. (5.10), the $\mathbb{Z}_N = \mathbb{Z}_{k^2}$ magnetic one-form symmetry is spontaneously broken to \mathbb{Z}_k , resulting in \mathbb{Z}_k topological order. The unbroken \mathbb{Z}_k one-form symmetry also has nontrivial SPT order. For $m < 0$, the \mathbb{Z}_N one-form symmetry is unbroken, but the non-invertible time-reversal symmetry associated with operator $\mathsf{T}_{2k} = \mathsf{D}_{2k}\mathsf{T}$ is spontaneously broken, resulting in two ground states that are distinct \mathbb{Z}_N one-form SPTs with responses given by Eq. (5.19). The domain walls in this phase obey the non-invertible fusion rules in Eq. (5.7). The $SO(N_f)$ flavor symmetry remains unbroken throughout the phase diagram.

where \tilde{a}_μ is a dynamical $U(1)$ one-form gauge field, $b_{\mu\nu}$ is a dynamical $U(1)$ two-form gauge field, and $B_{\mu\nu}$ is a background \mathbb{Z}_N gauge field. The ground states described by these TQFTs become degenerate at $\theta = \pi\tilde{n} = \pi(2k + N)$, and they are exchanged by the non-invertible time-reversal symmetry T_{2k} . Indeed, the dyons with a perimeter law at $\theta = \pi(2k + N \pm 1)$ are

$$(q_e, q_m) = \left(-\frac{N + 2k \pm 1}{2} q_m, q_m \right) \mod k^2. \quad (5.17)$$

According to Eq. (5.11), the non-invertible time-reversal symmetry T_{2k} maps these dyons as

$$\left(-\frac{N + 2k \pm 1}{2} q_m, q_m \right) \rightarrow \left(-\frac{N + 2k \mp 1}{2} q_m, q_m \right). \quad (5.18)$$

Thus, at $\theta = \pi(2k + N)$ the non-invertible time-reversal symmetry T_{2k} is spontaneously broken. The domain walls obey the non-invertible fusion rules of Eq. (5.7).

There is no topological order for either of the TQFTs¹⁷ in Eq. (5.16) because of the identity¹⁸ $\gcd(N, (\tilde{n} \pm 1)/2) = \gcd(k^2, (k^2 + 2k \pm 1)/2) = 1$, which holds for any integer k . Thus, the \mathbb{Z}_N magnetic one-form symmetry is unbroken, and the ground state degeneracy

¹⁷This statement does not always hold for generic odd N and k . It is also possible for one of the TQFTs to have topological order so that the \mathbb{Z}_N magnetic one-form symmetry is spontaneously broken to a subgroup. But since $k + (N - 1)/2$ and $k + (N + 1)/2$ are coprime, if S_\pm has topological order, then S_\mp is necessarily topologically trivial.

¹⁸This identity follows from

$$1 = k^2(2k + 4 \mp 1) + \left(\frac{k^2 + 2k \pm 1}{2} \right) (-2)(2k \mp 1).$$

in this phase is two on any manifold. Integrating out the dynamical fields in Eq. (5.16), we obtain responses for the background \mathbb{Z}_{k^2} two-form gauge field $B_{\mu\nu}$ of

$$S_{\text{SPT}}^{\pm}[B] = -\frac{k^2 2(2k \mp 1)}{4\pi} \int B \wedge B, \quad (5.19)$$

so the degenerate ground states are \mathbb{Z}_{k^2} one-form SPTs.

To summarize the conclusions of Section 5, we have found that for N odd, sufficiently large odd N_f , and even n , there can be a continuous transition between a topologically ordered phase and a phase that spontaneously breaks a non-invertible time-reversal symmetry. If we take $N = k^2$ and $n = 2k$ with $k > 1$ odd, the $m < 0$ phase spontaneously breaks the non-invertible time-reversal symmetry \mathbb{T}_{2k} while the \mathbb{Z}_{k^2} one-form symmetry is unbroken. The $m > 0$ phase preserves the non-invertible time-reversal symmetry but breaks the magnetic \mathbb{Z}_{k^2} one-form symmetry to \mathbb{Z}_k , resulting in \mathbb{Z}_k topological order, and the unbroken \mathbb{Z}_k one-form symmetry also has nontrivial SPT order. The $SO(N_f)$ flavor symmetry is unbroken in both phases. For large enough N_f , there is a continuous transition between these phases at $m = 0$ (see Figure 3). In the special case $n = 0$, the time-reversal symmetry is invertible, and the $m > 0$ phase has \mathbb{Z}_N topological order while the $m < 0$ phase spontaneously breaks the invertible time-reversal symmetry.

5.4 Critical exponents

Following methods of Section 4.4 and Refs. [29, 103, 104], the critical exponents for this transition may be easily computed if N_f is large enough that the $m = 0$ point is IR free. For the $m > 0$ phase, the 't Hooft loop has an area law, and following the same reasoning as in Section 4.4, we will find that the string tension σ goes as $\sigma \sim |m|^\mu$ with μ given in Eq. (4.19).

Turning our attention to critical exponents associated with the non-invertible time-reversal symmetry, we note that an operator odd under \mathbb{T}_n , which can serve as a local order parameter, is

$$\phi(x) = \sum_{J=1}^{N_f} i \text{Tr} [\psi_J^T(x) \mathcal{C} \gamma^5 \psi_J(x)]. \quad (5.20)$$

Unlike the \mathbb{Z}_N magnetic one-form global symmetry, the non-invertible zero-form symmetry has a local order parameter, and thus, the kinds of critical exponents we can study for this symmetry have closer analogues in conventional Landau theory. Because the critical point is IR free, the order parameter has scaling dimension $\Delta_\phi = 3$, so its two-point function scales

as [29]

$$\langle \phi(x) \phi(x') \rangle \sim \frac{1}{|x - x'|^6} \quad (5.21)$$

at the critical point.

Next, we examine the critical exponent for how the order parameter vanishes as the critical point is approached from the ordered phase. For $m < 0$, consider weakly perturbing the action S_n , Eq. (5.1), by a term that explicitly breaks the non-invertible time-reversal symmetry,

$$S_n \rightarrow S_n + \int d^4x \, h_\epsilon \phi(x), \quad (5.22)$$

where the coefficient h_ϵ is small, $|h_\epsilon| \ll |m|$. Here, h_ϵ is the analogue of an external symmetry-breaking field. At low energies, where the fermions may be integrated out, the effect of the perturbation in Eq. (5.22) is to modify the theta angle of the $m < 0$ phase to

$$\theta = \pi(n + N) \rightarrow \pi(n + N) + \frac{h_\epsilon}{|m|} N N_f. \quad (5.23)$$

Hence, for small $|m|$, we expect the expectation value of the order parameter to scale as [29]

$$\langle \phi(x) \rangle \sim \frac{1}{|m|} \langle \text{Tr} (\epsilon^{\mu\nu\lambda\sigma} f_{\mu\nu} f_{\lambda\sigma}) \rangle \sim \frac{(\Lambda_{\text{YM}})^4}{|m|} \sim |m|^{(8N_f/11)-1}, \quad (5.24)$$

where we used Eq. (4.18). This critical exponent is analogous to $\beta = (8N_f/11) - 1$ in Landau theory. For similar reasons, if we add the symmetry-breaking perturbation h_ϵ but with $m = 0$, we obtain

$$\langle \phi(x) \rangle \sim |h_\epsilon|^{(8N_f/11)-1}, \quad (5.25)$$

which is analogous to an exponent of $\delta = 11/(8N_f - 11)$ in Landau theory. These critical exponents characterize the universality class of the transition and highlight its analogy with conventional Landau transitions despite that both the phases and the transition are beyond Landau.

6 Discussion

In this work, we have introduced two families of exotic transitions between phases that break different generalized symmetries, both of which involve $PSU(N)$ gauge theory coupled to N_f odd flavors of Majorana fermions in the adjoint representation. Our first example is a critical point between SET phases that have distinct topological orders—a phase with \mathbb{Z}_N topological order and another with $\mathbb{Z}_{N/2}$ topological order for N even. While we have

focused on this transition for a $PSU(N)$ gauge group, a topological transition of this type can occur for adjoint QCD with other gauge groups, as discussed in Appendix D. To our knowledge, this is the first clear example in (3+1)d of a model with an exact one-form symmetry that has a continuous transition between phases with different one-form symmetry breaking.

The second kind of unconventional transition we have explored is a continuous transition between a topologically ordered phase, which breaks a discrete one-form symmetry, and a phase that spontaneously breaks a non-invertible time-reversal symmetry, providing an analogue of deconfined quantum criticality for generalized symmetries. This critical point represents a “beyond Landau” transition between phases that also lie beyond Landau. Taken together, our two examples of exotic transitions can serve as guides for discovering other topological critical points and developing a more general theory classifying these transitions.

There are several promising directions for future work. One is to analyze symmetry fractionalization of the non-invertible time-reversal symmetry in the SET phase discussed in Section 4.2. While fractionalization of non-invertible symmetries has been studied in some examples in (2+1)d [129], there is currently no general framework for fractionalization of non-invertible symmetries. The (3+1)d \mathbb{Z}_k topological order enriched by non-invertible time-reversal symmetry in Section 5.2 provides a concrete example where this analysis can be done.

Another possible avenue is to investigate the fate of the SET phase discussed in Section 4.3 when the magnetic one-form symmetry is explicitly broken. Concretely, this question can be explored by studying the lattice model of Section 4.5, where the degree to which the one-form symmetry is explicitly broken is determined by the two-form gauge coupling \tilde{g}^2 . For small enough \tilde{g}^2 , the phase discussed in Section 4.3 will have topological orders both in the bulk and on the boundary. At large \tilde{g}^2 , these topological orders are expected to disappear. A natural question is whether the transitions must occur simultaneously, or if there can be an analogue of the extraordinary transition for one-form symmetries. Similarly, it would also be interesting to study boundary criticality in the transition between the $\mathbb{Z}_{N/2}$ and \mathbb{Z}_N topologically ordered phases.

Finally, all the topological transitions discussed in this work involve magnetic one-form symmetries. It would be interesting to develop a dual description in which these magnetic one-form symmetries are mapped to electric one-form symmetries. Such a duality would not only offer a complementary perspective on the dynamics of these transitions, but would also constitute a rare example of a non-supersymmetric duality in (3+1)d.

Note: While completing this manuscript, we became aware of a related work [130] that also studies transitions of invertible one-form symmetries in theories with local quantum fields but without imposing a flavor symmetry on matter fields.

Acknowledgments

The author is especially grateful to Eduardo Fradkin for numerous insightful discussions and for comments on the manuscript. We also thank Fiona Burnell, Junyi Cao, Clay Córdova, Hart Goldman, and Raman Sohal for useful discussions, as well as John McGreevy and Matthew O’Brien for feedback on the draft. This work was supported in part by the National Science Foundation under grant DMR-2225920 at the University of Illinois.

A Twisted BF theory

In this appendix, we review the topological quantum field theory (TQFT) for twisted BF theory [18, 67], given by the action,

$$S_{\text{TQFT}}[p, \tilde{a}, b, B] = \frac{N}{2\pi} \int b \wedge (d\tilde{a} + B) + \frac{Np}{4\pi} \int b \wedge b + \frac{N}{2\pi} \int B \wedge d\tilde{a}, \quad (\text{A.1})$$

where \tilde{a}_μ and $\tilde{\alpha}_\mu$ are dynamical $U(1)$ one-form gauge fields, $b_{\mu\nu}$ is a dynamical $U(1)$ two-form gauge field, and $B_{\mu\nu}$ is a background $U(1)$ two-form gauge field. The gauge field $\tilde{\alpha}_\mu$ is a Lagrange multiplier that constrains $B_{\mu\nu}$ to be a \mathbb{Z}_N gauge field. The parameters N and p are integers, and p is defined mod N on a spin manifold (mod $2N$ on a generic manifold). We first remove the background field, setting $B_{\mu\nu} = 0$. Consider the gauge transformation,

$$b \rightarrow b + d\lambda, \quad \tilde{a} \rightarrow \tilde{a} - p\lambda + d\xi, \quad (\text{A.2})$$

where ξ is a 2π periodic scalar and λ_μ is a $U(1)$ one-form gauge field. If we work on a closed four-manifold, then the partition function is invariant under Eq. (A.2) if N and p are integers and Np is even. If the manifold is a spin manifold, as we always assume in the main text, then N and p can be arbitrary integers.

We can form gauge invariant operators of the form,

$$\mathcal{D}(\gamma, \Omega)^K = \exp \left(iK \oint_\gamma \tilde{a} + ipK \int_\Omega b \right), \quad (\text{A.3})$$

where γ is a loop and Ω is an open surface bounding γ . The local equation of motion for $b_{\mu\nu}$ is that

$$d\tilde{a} + pb = 0, \quad (\text{A.4})$$

which renders $\mathcal{D}(\gamma, \Omega)^K$ trivial unless it is a genuine loop operator. Unlike loops attached to surfaces, genuine loops can be noncontractible and are not necessarily trivial globally. The surface in $\mathcal{D}(\gamma, \Omega)^K$ is trivial only if $pK \in N\mathbb{Z}$. This condition is satisfied by $K = \frac{N}{L}q$ where $L = \gcd(N, p)$ and q is an integer mod L . Thus, there are L physically distinct genuine loop operators, given by

$$\mathcal{T}(\gamma)^{Nq/L} = \exp\left(i\frac{Nq}{L} \oint_{\gamma} \tilde{a}\right), \quad (\text{A.5})$$

where γ can now be a non-contractible loop.

We can also form an operator supported on a closed surface Σ , given by

$$\mathcal{U}(\Sigma)^{q'} = \exp\left(-iq' \oint_{\Sigma} b\right). \quad (\text{A.6})$$

Here, q' is equivalent to $q' + N$ since \tilde{a}_{μ} is a Lagrange multiplier that constrains $b_{\mu\nu}$ to be a \mathbb{Z}_N gauge field. However, q' is also equivalent to $q' + p$ because of Eq. (A.4). Hence, q' is defined mod L , and there are L distinct surface operators.

The genuine loop operators and surface operators have correlation functions given by

$$\langle \mathcal{T}(\gamma)^{Nq/L} \mathcal{U}(\Sigma)^{q'} \rangle = \exp\left(\frac{2\pi i}{L} q q' \Phi(\gamma, \Sigma)\right), \quad (\text{A.7})$$

where $\Phi(\gamma, \Sigma)$ is the linking number of γ and Σ . Thus, the topological order realized by the loop and surface operators is equivalent to a topological \mathbb{Z}_L gauge theory. A key distinction, however, is that $\mathcal{T}(\gamma)^{N/L}$ can represent the worldline of a fermionic particle if Np/L^2 is odd.

The TQFT has a \mathbb{Z}_N one-form symmetry that acts as

$$\tilde{a} \rightarrow \tilde{a} + \eta, \quad (\text{A.8})$$

where η is a flat connection, satisfying $d\eta = 0$ locally and $\oint \eta \in (2\pi/N)\mathbb{Z}$ globally. The surface operator that acts with this one-form symmetry is $\mathcal{U}(\Sigma)$, and the background field $B_{\mu\nu}$ probes this symmetry. The loop operators $\mathcal{T}(\gamma)^{N/L}$ transform nontrivially under this global symmetry, so the \mathbb{Z}_N one-form symmetry is spontaneously broken to $\mathbb{Z}_{N/L}$ at low energies.

Next, we reintroduce the background field $B_{\mu\nu}$, which will allow us to observe consequences of the unbroken $\mathbb{Z}_{N/L}$ one-form symmetry. The theory is invariant under gauge transformations,

$$b \rightarrow b + d\lambda, \quad \tilde{a} \rightarrow \tilde{a} - p\lambda + d\xi - \tilde{\lambda}, \quad B \rightarrow B + d\tilde{\lambda}, \quad \tilde{\alpha} \rightarrow \tilde{\alpha} - \lambda + d\tilde{\xi}, \quad (\text{A.9})$$

where ξ and $\tilde{\xi}$ are 2π periodic scalar fields while λ_μ and $\tilde{\lambda}_\mu$ are $U(1)$ one-form gauge fields. Integrating out \tilde{a}_μ and $\tilde{\alpha}_\mu$ respectively constrain $b_{\mu\nu}$ and $B_{\mu\nu}$ to be \mathbb{Z}_N gauge fields, so they are locally trivial, but globally they satisfy

$$\oint_{\Sigma} b = \frac{2\pi\ell}{N}, \quad \oint_{\Sigma} B \in \frac{2\pi\tilde{\ell}}{N}, \quad \ell, \tilde{\ell} \in \mathbb{Z}, \quad (\text{A.10})$$

for any closed surface Σ . The action that remains after integrating out the Lagrange multipliers is

$$S_{\text{eff}}[B] = \frac{N}{2\pi} \int b \wedge B + \frac{Np}{4\pi} \int b \wedge b. \quad (\text{A.11})$$

We next integrate out $b_{\mu\nu}$. Both $b_{\mu\nu}$ and $B_{\mu\nu}$ are locally trivial, so we only have the global equation of motion,

$$\frac{2\pi}{N} (\tilde{\ell} + p\ell) = \left(\oint_{\Sigma} B + p \oint_{\Sigma} b \right) \in 2\pi\mathbb{Z}, \quad (\text{A.12})$$

for any closed surface Σ . We thus find that

$$(\tilde{\ell} + p\ell) \in N\mathbb{Z}, \quad (\text{A.13})$$

which is consistent only if $\tilde{\ell} \in L\mathbb{Z}$, where $L = \text{gcd}(N, p)$. Hence, we must have

$$\oint_{\Sigma} B \in \frac{2\pi}{N/L} \mathbb{Z}, \quad (\text{A.14})$$

which implies that $B_{\mu\nu}$ must be probing the unbroken $\mathbb{Z}_{N/L}$ subgroup of the \mathbb{Z}_N one-form symmetry. This restriction of the background field to configurations consistent with the unbroken symmetry is analogous to the Meissner effect in a superconductor. To integrate out $b_{\mu\nu}$, we must solve Eq. (A.12) for ℓ . Writing $\tilde{\ell} = L\ell'$, where $\ell' \in \mathbb{Z}$, a solution is

$$\frac{N}{2\pi} \oint_{\Sigma} b = \ell = r\ell' = \frac{rN}{2\pi L} \oint_{\Sigma} B \mod \frac{N}{L}, \quad (\text{A.15})$$

where r is an integer such that $rp = -L \mod N$. (Such an r must always exist.) The response after integrating out all dynamical fields is then

$$S_{\text{resp}}[B] = \frac{\frac{N}{L}r}{4\pi} \int B \wedge B, \quad (\text{A.16})$$

signaling nontrivial SPT order for the unbroken $\mathbb{Z}_{N/L}$ one-form symmetry.

B Charge conjugation and Majorana condition

We briefly review Majorana fermions in this appendix. A more detailed recent discussion may be found in Ref. [131]. For the $\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1)$ metric, the Dirac equation admits real spinor solutions for purely imaginary Dirac matrices, which is a property of the Majorana basis,

$$\begin{aligned}\gamma_M^0 &= \begin{pmatrix} 0 & \sigma^2 \\ \sigma^2 & 0 \end{pmatrix}, & \gamma_M^1 &= \begin{pmatrix} i\sigma^3 & 0 \\ 0 & i\sigma^3 \end{pmatrix}, & \gamma_M^2 &= \begin{pmatrix} 0 & -\sigma^2 \\ \sigma^2 & 0 \end{pmatrix}, \\ \gamma_M^3 &= \begin{pmatrix} -i\sigma^1 & 0 \\ 0 & -i\sigma^1 \end{pmatrix}, & \gamma_M^5 &= \begin{pmatrix} \sigma^2 & 0 \\ 0 & -\sigma^2 \end{pmatrix}.\end{aligned}\tag{B.1}$$

The Dirac matrices γ^μ in a generic basis are related to the above γ_M^μ by a similarity transformation M as

$$M\gamma^\mu M^{-1} = \gamma_M^\mu.\tag{B.2}$$

Because the γ_M^μ are purely imaginary, there must be a matrix U_C such that

$$U_C \gamma^\mu U_C^{-1} = -(\gamma^\mu)^*,\tag{B.3}$$

where $U_C = (M^*)^{-1}M$.

In the Majorana basis, a Majorana fermion ψ_M obeys the constraint,

$$(\psi_M^\dagger)^T = \psi_M.\tag{B.4}$$

To determine the analogue of this condition in a generic basis, we express $\psi_M = M\psi$ to obtain the condition,

$$(\psi^\dagger)^T = U_C \psi.\tag{B.5}$$

In terms of the matrix $\mathcal{C} = (U_C)^T \gamma^0$, the Majorana condition on ψ is

$$\bar{\psi} = \psi^\dagger \gamma^0 = \psi^T (U_C)^T \gamma^0 = \psi^T \mathcal{C}.\tag{B.6}$$

This version of the Majorana condition is what generalizes most naturally to other spacetime metrics. For Dirac matrices in any basis, it can be shown that there exists a unitary matrix \mathcal{C} that satisfies

$$\mathcal{C} \gamma^\mu \mathcal{C}^{-1} = -(\gamma^\mu)^T, \quad \mathcal{C}^T = -\mathcal{C},\tag{B.7}$$

for both sign conventions of Lorentzian signature and for Euclidean signature.

As an example, we can take $\mathcal{C} = \gamma^0$ in the Majorana basis. Similarly, if we use the Weyl basis for Dirac matrices,

$$\gamma^\mu = \begin{pmatrix} 0 & \sigma^\mu \\ \bar{\sigma}^\mu & 0 \end{pmatrix}, \quad \gamma^5 = i\gamma^0\gamma^1\gamma^2\gamma^3 = \begin{pmatrix} -\mathbb{I}_2 & 0 \\ 0 & \mathbb{I}_2 \end{pmatrix}, \quad (\text{B.8})$$

where $\sigma^\mu = (\mathbb{I}_2, \sigma^j)$ and $\bar{\sigma}^\mu = (\mathbb{I}_2, -\sigma^j)$, then

$$\mathcal{C} = i\gamma^0\gamma^2 = \begin{pmatrix} -i\sigma^2 & 0 \\ 0 & i\sigma^2 \end{pmatrix} \quad (\text{B.9})$$

is a suitable charge conjugation matrix.

The charge conjugation matrix \mathcal{C} may be used to define the action for a Majorana fermion in an arbitrary basis for the Dirac matrices. The Lagrangian density for a single Majorana fermion of mass m is

$$\mathcal{L}_M = \frac{1}{2} \psi^T(x) \mathcal{C}(i\rlap{\not{\partial}} - m) \psi(x). \quad (\text{B.10})$$

The partition function is

$$Z_M = \int \mathcal{D}\psi \exp \left(i \int d^4x \mathcal{L}_M \right) = \text{Pf}[\mathcal{C}(i\rlap{\not{\partial}} - m)], \quad (\text{B.11})$$

where Pf denotes the Pfaffian.

C Thermal response for fermions

Here, we summarize our conventions for curved spacetime and review how to couple fermions to a background metric $g_{\mu\nu}$, which is useful for keeping track of thermal response [74–77]. The spacetime metric $g_{\mu\nu}$ can be expressed in terms of a set of local Lorentz frame fields e_μ^a as

$$g_{\mu\nu} = e_\mu^a e_\nu^b \eta_{ab}, \quad (\text{C.1})$$

where η_{ab} is the Minkowski metric. The Christoffel symbols are constructed from the metric as

$$\Gamma^\mu_{\nu\lambda} = \frac{1}{2} g^{\mu\sigma} (\partial_\nu g_{\lambda\sigma} + \partial_\lambda g_{\nu\sigma} - \partial_\sigma g_{\nu\lambda}). \quad (\text{C.2})$$

The spin connection is

$$\omega_\mu^{ab} = e_\nu^a (\partial_\mu e^{\nu b} + \Gamma^\nu_{\mu\lambda} e^{\lambda b}), \quad (\text{C.3})$$

which can be used to define a one-form,

$$\omega^{ab} = \omega_\mu^{ab} dx^\mu, \quad (\text{C.4})$$

and a corresponding curvature two-form,

$$R^a{}_b = d\omega^a{}_b + \omega^a{}_c \wedge \omega^c{}_b. \quad (\text{C.5})$$

Using the notation $\text{Tr}(R \wedge R) = R^a{}_b \wedge R^b{}_a$, the gravitational theta term is

$$S_g = \frac{\theta_g}{384\pi^2} \int \text{Tr}(R \wedge R). \quad (\text{C.6})$$

This term is quantized so that θ_g has periodicity 2π on a spin manifold and periodicity 32π on a generic four-manifold. This bulk response leads to a boundary gravitational Chern-Simons response with chiral central charge $c = \theta_g/4\pi$. The boundary thermal Hall conductivity is then

$$\kappa_{xy} = c \frac{\pi k_B^2 T}{6\hbar}, \quad (\text{C.7})$$

where T is the temperature.

Next, we review how to place fermionic fields in curved spacetime. The Dirac matrices in flat spacetime γ^a satisfy

$$\{\gamma_a, \gamma_b\} = 2\eta_{ab}. \quad (\text{C.8})$$

The generators of $\text{Spin}(1, 3)$, the double cover of the Lorentz group, are

$$\sigma_{ab} = \frac{i}{4} [\gamma_a, \gamma_b], \quad (\text{C.9})$$

and obey the algebra,

$$[\sigma_{ab}, \sigma_{cd}] = i (g_{ad} \sigma_{bc} + g_{bc} \sigma_{ad} - g_{ac} \sigma_{bd} - g_{bd} \sigma_{ac}). \quad (\text{C.10})$$

To couple a Majorana fermion $\psi(x)$ to a background metric $g_{\mu\nu}(x)$, we introduce the covariant derivative,

$$\nabla_\mu \psi = \left(\partial_\mu - \frac{i}{2} \omega_\mu{}^{ab} \sigma_{ab} \right) \psi. \quad (\text{C.11})$$

Additionally, we define

$$\not{\nabla} = e^\mu{}_a(x) \gamma^a \nabla_\mu. \quad (\text{C.12})$$

The action of the fermion coupled to the metric is

$$S = \int d^4x \sqrt{-g} \frac{1}{2} \psi^T(x) \mathcal{C} (i\not{\nabla} - m) \psi(x), \quad (\text{C.13})$$

where g is the determinant of $g_{\mu\nu}$ and m is the fermion mass.

D Other gauge groups

In Section 4, we discuss a transition between SETs with different topological orders for $PSU(N)$ adjoint QCD. An analogue of this topological transition can occur for adjoint QCD with other gauge groups. Consider a gauge field with a gauge group G_g coupled to N_f odd flavors of Majorana fermions in the adjoint representation. As in Section 4, we impose an $SO(N_f)$ flavor symmetry and time-reversal symmetry, and we take N_f large enough so that the transition is continuous. Below, k will always be a nonnegative integer, and we regularize the theory so that the phase with positive fermion mass m is the pure gauge theory with $\theta = 0$. The full magnetic one-form symmetry will be spontaneously broken in this phase.

For a gauge group of $G_g = \text{Sp}(4k+1)/\mathbb{Z}_2$, the magnetic one-form symmetry is \mathbb{Z}_2 . The $m > 0$ phase has \mathbb{Z}_2 topological order, and at low energies, the $m < 0$ phase has a theta angle for G_g of $\theta = \pi(4k+2)N_f$, leading to the effective action,

$$S_{\text{eff}} = \frac{2N_f(2k+1)(4k+1)}{4\pi} \int b \wedge b + \frac{(2k+1)(4k+1)\pi}{8\pi^2} \int \text{Tr}(F \wedge F) + \frac{N_f(2k+1)(4k+1)\pi}{384\pi^2} \int \text{Tr}(R \wedge R), \quad (\text{D.1})$$

where $b_{\mu\nu}$ is a dynamical \mathbb{Z}_2 two-form gauge field, $F_{\mu\nu}$ is a background field for the $SO(N_f)$ flavor symmetry, and R is the curvature two-form. Because $N_f(2k+1)(4k+1)$ is odd, there is no topological order in this phase. If we couple to a background field $B_{\mu\nu}$ for the \mathbb{Z}_2 magnetic one-form symmetry and integrate out $b_{\mu\nu}$, we find that this phase has nontrivial SPT order for the unbroken \mathbb{Z}_2 magnetic one-form symmetry. Given that $\text{Sp}(1) \cong SU(2)$, this analysis is consistent with our results from Section 4.

The gauge group $G_g = \text{Spin}(8k+6)/\mathbb{Z}_4$ has a \mathbb{Z}_4 magnetic one-form symmetry, so the $m > 0$ phase has \mathbb{Z}_4 topological order. The $m < 0$ phase has a theta angle for G_g of $\theta = \pi(8k+4)N_f$, giving the effective action,

$$S_{\text{eff}} = \frac{4(4k+3)(4k+2)N_f}{4\pi} \int b \wedge b + \frac{(4k+3)(8k+5)\pi}{8\pi^2} \int \text{Tr}(F \wedge F) + \frac{N_f(4k+3)(8k+5)\pi}{384\pi^2} \int \text{Tr}(R \wedge R), \quad (\text{D.2})$$

where $b_{\mu\nu}$ is a dynamical \mathbb{Z}_4 two-form gauge field. Since $\text{gcd}(4, (4k+3)(4k+2)N_f) = 2$, the \mathbb{Z}_4 one-form symmetry is spontaneously broken to \mathbb{Z}_2 , and the unbroken \mathbb{Z}_2 one-form symmetry has nontrivial SPT order. For $\text{Spin}(6) \cong SU(4)$, this conclusion is consistent with Section 4.

The gauge groups $\text{Spin}(8k)/(\mathbb{Z}_2 \times \mathbb{Z}_2)$ and $\text{Spin}(8k+4)/(\mathbb{Z}_2 \times \mathbb{Z}_2)$ have $\mathbb{Z}_2 \times \mathbb{Z}_2$ magnetic one-form symmetries. The $m > 0$ phase thus has $\mathbb{Z}_2 \times \mathbb{Z}_2$ topological order in both cases. For the gauge group $G_g = \text{Spin}(8k)/(\mathbb{Z}_2 \times \mathbb{Z}_2)$, the $m < 0$ phase has a G_g theta angle of $\theta = \pi(8k-2)N_f$, which leads to an effective action of

$$S_{\text{eff}} = \frac{2(4k-1)N_f}{2\pi} \int b_1 \wedge b_2 + \frac{4k(8k-1)\pi}{8\pi^2} \int \text{Tr}(F \wedge F) + \frac{N_f 4k(8k-1)\pi}{384\pi^2} \int \text{Tr}(R \wedge R), \quad (\text{D.3})$$

where $(b_1)_{\mu\nu}$ and $(b_2)_{\mu\nu}$ are dynamical \mathbb{Z}_2 two-form gauge fields. Since $(4k-1)N_f$ is odd, there is no topological order in this phase. If we couple to background fields, $(B_1)_{\mu\nu}$ and $(B_2)_{\mu\nu}$, for the two \mathbb{Z}_2 magnetic one-form symmetries and integrate out $(b_1)_{\mu\nu}$ and $(b_2)_{\mu\nu}$, we find that the \mathbb{Z}_2 magnetic one-form symmetries have a mixed SPT response,

$$S_{\text{mixed}}[B_1, B_2] = \frac{2}{2\pi} \int B_1 \wedge B_2, \quad (\text{D.4})$$

in addition to the zero-form SPT response in Eq. (D.3).

For $G_g = \text{Spin}(8k+4)/(\mathbb{Z}_2 \times \mathbb{Z}_2)$, the $m < 0$ phase has a G_g theta angle of $\theta = \pi(8k+2)N_f$ so that the effective action is

$$S_{\text{eff}} = \frac{2(4k+1)(2k+1)N_f}{4\pi} \int (b_1 \wedge b_1 + b_2 \wedge b_2) + \frac{(4k+2)(8k+3)\pi}{8\pi^2} \int \text{Tr}(F \wedge F) + \frac{N_f(4k+2)(8k+3)\pi}{384\pi^2} \int \text{Tr}(R \wedge R). \quad (\text{D.5})$$

Since $(4k+1)(2k+1)N_f$ is odd, this phase is also an SPT. However, it has a different response for the magnetic one-form symmetry, given by

$$S_{\text{unmixed}}[B_1, B_2] = \frac{2}{4\pi} \int (B_1 \wedge B_1 + B_2 \wedge B_2), \quad (\text{D.6})$$

which does not couple the two background fields.

E Anomaly of non-invertible time-reversal

Here, we identify the conditions under which the non-invertible time-reversal symmetry associated with the operator \mathbb{T}_n , defined in Section 5.1, has a mixed anomaly with the \mathbb{Z}_N one-form symmetry. Specifically, the anomaly implies that no trivially gapped phase can simultaneously preserve both symmetries.

We first consider the most general fermionic \mathbb{Z}_N one-form SPT in (3+1)d, given by the action

$$S_{\text{SPT}}[B] = \frac{Np}{4\pi} \int B \wedge B, \quad (\text{E.1})$$

where p is an integer mod N and $B_{\mu\nu}$ is a background \mathbb{Z}_N two-form gauge field. The action for this SPT transforms under the non-invertible time-reversal transformation \mathbf{K}_n to

$$S_{\mathbf{K}_n} = -\frac{Np}{4\pi} \int b \wedge b + \frac{N}{2\pi} \int b \wedge \beta + \frac{N(N-1)n}{4\pi} \int \beta \wedge \beta - \frac{N}{2\pi} \int \beta \wedge B, \quad (\text{E.2})$$

where $b_{\mu\nu}$ and $\beta_{\mu\nu}$ are dynamical \mathbb{Z}_N two-form gauge fields. To integrate out $b_{\mu\nu}$, we must have $\gcd(N, p) = 1$. Integrating out $b_{\mu\nu}$ then gives the action

$$S_{\text{eff}} = \frac{N(\ell + (N-1)n)}{4\pi} \int \beta \wedge \beta - \frac{N}{2\pi} \int \beta \wedge B, \quad (\text{E.3})$$

where ℓ is an integer such that $\ell p = 1 \pmod{N}$. Finally, integrating out $\beta_{\mu\nu}$, which requires $\gcd(N, \ell - n) = 1$, results in the SPT,

$$\tilde{S}_{\text{SPT}}[B] = -\frac{N\tilde{\ell}}{4\pi} \int B \wedge B, \quad (\text{E.4})$$

where $\tilde{\ell}$ is an integer such that $\tilde{\ell}(\ell - n) = 1 \pmod{N}$.

The original SPT state, Eq. (E.1), is then invariant under the non-invertible time-reversal transformation \mathbf{K}_n if $\tilde{\ell} = -p \pmod{N}$, which implies that

$$1 = \tilde{\ell}(\ell - n) = -p(\ell - n) = -1 + pn \pmod{N}. \quad (\text{E.5})$$

Hence, we must have

$$pn = 2 \pmod{N} \quad (\text{E.6})$$

for some p , indicating that $\gcd(N, n)$ is either 1 or 2. If N or n is odd, a solution for p exists only if $\gcd(N, n) = 1$. If N and n are both even, there exists a solution for p only if $\gcd(N/2, n/2) = 1$. Thus, there is an anomaly for $\gcd(N, n) > 1$ if N or n is odd and for $\gcd(N/2, n/2) > 1$ if N and n are both even. In these cases, no trivially gapped state can preserve both the \mathbb{Z}_N one-form symmetry and non-invertible time-reversal symmetry.

References

- [1] F. J. Wegner, *Duality in generalized Ising models and phase transitions without local order parameters*, Journal of Mathematical Physics **12** (1971), doi:[10.1063/1.1665530](https://doi.org/10.1063/1.1665530).
- [2] X.-G. Wen, *Topological orders in rigid states*, International Journal of Modern Physics B **04** (1990), doi:[10.1142/s0217979290000139](https://doi.org/10.1142/s0217979290000139).
- [3] X. Chen, Z.-C. Gu, Z.-X. Liu and X.-G. Wen, *Symmetry-protected topological orders in interacting bosonic systems*, Science **338**(6114), 1604 (2012), doi:[10.1126/science.1227224](https://doi.org/10.1126/science.1227224).

- [4] X. Chen, Z.-C. Gu, Z.-X. Liu and X.-G. Wen, *Symmetry protected topological orders and the group cohomology of their symmetry group*, Phys. Rev. B **87**, 155114 (2013), doi:[10.1103/PhysRevB.87.155114](https://doi.org/10.1103/PhysRevB.87.155114).
- [5] F. Pollmann, E. Berg, A. M. Turner and M. Oshikawa, *Symmetry protection of topological phases in one-dimensional quantum spin systems*, Physical Review B - Condensed Matter and Materials Physics **85** (2012), doi:[10.1103/PhysRevB.85.075125](https://doi.org/10.1103/PhysRevB.85.075125).
- [6] L. Fidkowski and A. Kitaev, *Topological phases of fermions in one dimension*, Physical Review B - Condensed Matter and Materials Physics **83** (2011), doi:[10.1103/PhysRevB.83.075103](https://doi.org/10.1103/PhysRevB.83.075103).
- [7] A. Vishwanath and T. Senthil, *Physics of three-dimensional bosonic topological insulators: Surface-deconfined criticality and quantized magnetoelectric effect*, Phys. Rev. X **3**, 011016 (2013), doi:[10.1103/PhysRevX.3.011016](https://doi.org/10.1103/PhysRevX.3.011016).
- [8] T. Senthil, *Symmetry-protected topological phases of quantum matter*, Annual Review of Condensed Matter Physics **6**(1), 299 (2015), doi:[10.1146/annurev-conmatphys-031214-014740](https://doi.org/10.1146/annurev-conmatphys-031214-014740).
- [9] M. Barkeshli, P. Bonderson, M. Cheng and Z. Wang, *Symmetry fractionalization, defects, and gauging of topological phases*, Phys. Rev. B **100**, 115147 (2019), doi:[10.1103/PhysRevB.100.115147](https://doi.org/10.1103/PhysRevB.100.115147).
- [10] X. Chen, *Symmetry fractionalization in two dimensional topological phases*, doi:[10.1016/j.revip.2017.02.002](https://doi.org/10.1016/j.revip.2017.02.002) (2017).
- [11] E. Fradkin and S. H. Shenker, *Phase diagrams of lattice gauge theories with higgs fields*, Physical Review D **19** (1979), doi:[10.1103/PhysRevD.19.3682](https://doi.org/10.1103/PhysRevD.19.3682).
- [12] M. Creutz, L. Jacobs and C. Rebbi, *Monte Carlo study of Abelian lattice gauge theories*, Physical Review D **20** (1979), doi:[10.1103/PhysRevD.20.1915](https://doi.org/10.1103/PhysRevD.20.1915).
- [13] M. Creutz, L. Jacobs and C. Rebbi, *Monte Carlo computations in lattice gauge theories*, doi:[10.1016/0370-1573\(83\)90016-9](https://doi.org/10.1016/0370-1573(83)90016-9) (1983).
- [14] M. Vettorazzo and P. de Forcrand, *Electromagnetic fluxes, monopoles, and the order of 4d compact $U(1)$ phase transition*, Nuclear Physics B **686** (2004), doi:[10.1016/j.nuclphysb.2004.02.038](https://doi.org/10.1016/j.nuclphysb.2004.02.038).
- [15] O. Akerlund and P. de Forcrand, *$U(1)$ lattice gauge theory with a topological action*, Journal of High Energy Physics **2015** (2015), doi:[10.1007/JHEP06\(2015\)183](https://doi.org/10.1007/JHEP06(2015)183).
- [16] T. Filk, M. Marcu and K. Fredenhagen, *Line of second-order phase transitions in the four-dimensional Z_2 gauge theory with matter fields*, Physics Letters B **169** (1986), doi:[10.1016/0370-2693\(86\)90381-3](https://doi.org/10.1016/0370-2693(86)90381-3).
- [17] Y. Blum, P. K. Coyle, S. Elitzur, E. Rabinovici, H. Rubinstein and S. Solomon, *In-*

- vestigation of the critical behavior of the critical point of the \mathbb{Z}_2 gauge lattice*, Nuclear Physics B **535** (1998), doi:[10.1016/S0550-3213\(98\)00636-1](https://doi.org/10.1016/S0550-3213(98)00636-1).
- [18] D. Gaiotto, A. Kapustin, N. Seiberg and B. Willett, *Generalized global symmetries*, J. High Energ. Phys. **2015**(2), 172 (2015), doi:[10.1007/JHEP02\(2015\)172](https://doi.org/10.1007/JHEP02(2015)172).
 - [19] J. McGreevy, *Generalized symmetries in condensed matter*, Annual Review of Condensed Matter Physics **14** (2023), doi:[10.1146/annurev-conmatphys-040721-021029](https://doi.org/10.1146/annurev-conmatphys-040721-021029).
 - [20] P. R. Gomes, *An introduction to higher-form symmetries*, SciPost Physics Lecture Notes (2023), doi:[10.21468/SciPostPhysLectNotes.74](https://doi.org/10.21468/SciPostPhysLectNotes.74).
 - [21] C. Córdova, T. T. Dumitrescu, K. Intriligator and S.-H. Shao, *Snow-mass white paper: Generalized symmetries in quantum field theory and beyond*, doi:[10.48550/arXiv.2205.09545](https://doi.org/10.48550/arXiv.2205.09545) (2022), <https://arxiv.org/abs/2205.09545>.
 - [22] T. D. Brennan and S. Hong, *Introduction to generalized global symmetries in QFT and particle physics*, doi:[10.48550/arXiv.2306.00912](https://doi.org/10.48550/arXiv.2306.00912) (2023), <https://arxiv.org/abs/2306.00912>.
 - [23] S. Schäfer-Nameki, *ICTP lectures on (non-)invertible generalized symmetries*, Physics Reports **1063**, 1 (2024), doi:<https://doi.org/10.1016/j.physrep.2024.01.007>, ICTP lectures on (non-)invertible generalized symmetries.
 - [24] L. Bhardwaj, L. E. Bottini, L. Fraser-Taliente, L. Gladden, D. S. Gould, A. Platschorre and H. Tillim, *Lectures on generalized symmetries*, doi:[10.1016/j.physrep.2023.11.002](https://doi.org/10.1016/j.physrep.2023.11.002) (2024).
 - [25] R. Luo, Q. R. Wang and Y. N. Wang, *Lecture notes on generalized symmetries and applications*, doi:[10.1016/j.physrep.2024.02.002](https://doi.org/10.1016/j.physrep.2024.02.002) (2024).
 - [26] S.-H. Shao, *What's done cannot be undone: TASI lectures on non-invertible symmetries*, doi:[10.48550/arXiv.2308.00747](https://doi.org/10.48550/arXiv.2308.00747) (2024), <https://arxiv.org/abs/2308.00747>.
 - [27] N. Iqbal and J. McGreevy, *Mean string field theory: Landau-Ginzburg theory for 1-form symmetries*, SciPost Physics **13**, 114 (2022), doi:[10.21468/SciPostPhys.13.5.114](https://doi.org/10.21468/SciPostPhys.13.5.114).
 - [28] Z. Bi and T. Senthil, *Adventure in topological phase transitions in 3 + 1-D: Non-Abelian deconfined quantum criticalities and a possible duality*, Phys. Rev. X **9** (2019), doi:[10.1103/PhysRevX.9.021034](https://doi.org/10.1103/PhysRevX.9.021034).
 - [29] Z. Bi, E. Lake and T. Senthil, *Landau ordering phase transitions beyond the Landau paradigm*, Physical Review Research **2** (2020), doi:[10.1103/PhysRevResearch.2.023031](https://doi.org/10.1103/PhysRevResearch.2.023031).
 - [30] J. Wang, Y. Z. You and Y. Zheng, *Gauge enhanced quantum criticality and time reversal deconfined domain wall: $SU(2)$ Yang-Mills dynamics with topological terms*, Physical Review Research **2** (2020), doi:[10.1103/PhysRevResearch.2.013189](https://doi.org/10.1103/PhysRevResearch.2.013189).
 - [31] W. E. Caswell, *Asymptotic behavior of non-abelian gauge theories to two-loop order*,

- Physical Review Letters **33** (1974), doi:[10.1103/PhysRevLett.33.244](https://doi.org/10.1103/PhysRevLett.33.244).
- [32] T. Banks and A. Zaks, *On the phase structure of vector-like gauge theories with massless fermions*, Nuclear Physics, Section B **196** (1982), doi:[10.1016/0550-3213\(82\)90035-9](https://doi.org/10.1016/0550-3213(82)90035-9).
 - [33] J. Kaidi, K. Ohmori and Y. Zheng, *Kramers-Wannier-like duality defects in (3+1)d gauge theories*, Physical Review Letters **128** (2022), doi:[10.1103/PhysRevLett.128.111601](https://doi.org/10.1103/PhysRevLett.128.111601).
 - [34] Y. Choi, H. T. Lam and S.-H. Shao, *Noninvertible time-reversal symmetry*, Physical Review Letters **130** (2023), doi:[10.1103/PhysRevLett.130.131602](https://doi.org/10.1103/PhysRevLett.130.131602).
 - [35] T. Senthil, A. Vishwanath, L. Balents, S. Sachdev and M. P. Fisher, *Deconfined quantum critical points*, Science **303** (2004), doi:[10.1126/science.1091806](https://doi.org/10.1126/science.1091806).
 - [36] T. Senthil, L. Balents, S. Sachdev, A. Vishwanath and M. P. Fisher, *Quantum criticality beyond the Landau-Ginzburg-Wilson paradigm*, Physical Review B - Condensed Matter and Materials Physics **70** (2004), doi:[10.1103/PhysRevB.70.144407](https://doi.org/10.1103/PhysRevB.70.144407).
 - [37] T. Senthil, *Deconfined Quantum Critical Points: A Review*, chap. Chapter 14, pp. 169–195, World Scientific, doi:[10.1142/9789811282386_0014](https://doi.org/10.1142/9789811282386_0014) (2024), https://www.worldscientific.com/doi/pdf/10.1142/9789811282386_0014.
 - [38] A. Chatterjee, Ömer M. Aksoy and X.-G. Wen, *Quantum phases and transitions in spin chains with non-invertible symmetries*, SciPost Physics **17**, 115 (2024), doi:[10.21468/SciPostPhys.17.4.115](https://doi.org/10.21468/SciPostPhys.17.4.115).
 - [39] L. Li, R.-Z. Huang and W. Cao, *Noninvertible symmetry-enriched quantum critical point*, doi:[10.48550/arXiv.2411.19034](https://doi.org/10.48550/arXiv.2411.19034) (2024), <https://arxiv.org/abs/2411.19034>.
 - [40] Y.-H. Chen and T. Grover, *Analogues of deconfined quantum criticality for non-invertible symmetry breaking in 1d*, doi:[10.48550/arXiv.2506.01131](https://doi.org/10.48550/arXiv.2506.01131) (2025), <https://arxiv.org/abs/2506.01131>.
 - [41] A. Kapustin, *Wilson-'t Hooft operators in four-dimensional gauge theories and S-duality*, Physical Review D - Particles, Fields, Gravitation and Cosmology **74** (2006), doi:[10.1103/PhysRevD.74.025005](https://doi.org/10.1103/PhysRevD.74.025005).
 - [42] O. Aharony, N. Seiberg and Y. Tachikawa, *Reading between the lines of four-dimensional gauge theories*, Journal of High Energy Physics **2013** (2013), doi:[10.1007/JHEP08\(2013\)115](https://doi.org/10.1007/JHEP08(2013)115).
 - [43] G. 't Hooft, *On the phase transition towards permanent quark confinement*, Nuclear Physics, Section B **138** (1978), doi:[10.1016/0550-3213\(78\)90153-0](https://doi.org/10.1016/0550-3213(78)90153-0).
 - [44] P. Goddard, J. Nuyts and D. Olive, *Gauge theories and magnetic charge*, Nuclear Physics, Section B **125** (1977), doi:[10.1016/0550-3213\(77\)90221-8](https://doi.org/10.1016/0550-3213(77)90221-8).

- [45] K. G. Wilson, *Confinement of quarks*, Physical Review D **10** (1974), doi:[10.1103/PhysRevD.10.2445](https://doi.org/10.1103/PhysRevD.10.2445).
- [46] Y. Nambu, *Strings, monopoles, and gauge fields*, Physical Review D **10** (1974), doi:[10.1103/PhysRevD.10.4262](https://doi.org/10.1103/PhysRevD.10.4262).
- [47] A. M. Polyakov, *Compact gauge fields and the infrared catastrophe*, Physics Letters B **59** (1975), doi:[10.1016/0370-2693\(75\)90162-8](https://doi.org/10.1016/0370-2693(75)90162-8).
- [48] S. Mandelstam, *II. Vortices and quark confinement in non-Abelian gauge theories*, Physics Reports **23** (1976), doi:[10.1016/0370-1573\(76\)90043-0](https://doi.org/10.1016/0370-1573(76)90043-0).
- [49] E. Witten, *Dyons of charge $e\theta/2\pi$* , Physics Letters B **86**, 283 (1979), doi:[10.1016/0370-2693\(79\)90838-4](https://doi.org/10.1016/0370-2693(79)90838-4).
- [50] G. 't Hooft, *Topology of the gauge condition and new confinement phases in non-abelian gauge theories*, Nuclear Physics B **190**(3), 455 (1981), doi:[10.1016/0550-3213\(81\)90442-9](https://doi.org/10.1016/0550-3213(81)90442-9).
- [51] J. L. Cardy and E. Rabinovici, *Phase structure of \mathbb{Z}_p models with a θ parameter*, Nuclear Physics B **205**(1), [FS5] 1 (1982), doi:[10.1016/0550-3213\(82\)90463-1](https://doi.org/10.1016/0550-3213(82)90463-1).
- [52] S. Gukov and A. Kapustin, *Topological quantum field theory, nonlocal operators, and gapped phases of gauge theories*, doi:[10.48550/arXiv.1307.4793](https://doi.org/10.48550/arXiv.1307.4793) (2013), <https://arxiv.org/abs/1307.4793>.
- [53] A. Kapustin and R. Thorngren, *Topological field theory on a lattice, discrete theta-angles and confinement*, Advances in Theoretical and Mathematical Physics **18** (2014), doi:[10.4310/ATMP.2014.v18.n5.a4](https://doi.org/10.4310/ATMP.2014.v18.n5.a4).
- [54] M. Honda and Y. Tanizaki, *Topological aspects of 4D Abelian lattice gauge theories with the θ parameter*, J. High Energ. Phys. **2020**, 154 (2020), doi:[10.1007/JHEP12\(2020\)154](https://doi.org/10.1007/JHEP12(2020)154).
- [55] P.-S. Hsin, W. Ji and C. M. Jian, *Exotic invertible phases with higher-group symmetries*, SciPost Physics **12** (2022), doi:[10.21468/SCIPOSTPHYS.12.2.052](https://doi.org/10.21468/SCIPOSTPHYS.12.2.052).
- [56] Y. Hayashi and Y. Tanizaki, *Non-invertible self-duality defects of Cardy-Rabinovici model and mixed gravitational anomaly*, J. High Energ. Phys. **2022**, 36 (2022), doi:[10.1007/JHEP08\(2022\)036](https://doi.org/10.1007/JHEP08(2022)036).
- [57] B. Moy, H. Goldman, R. Sohal and E. Fradkin, *Theory of oblique topological insulators*, SciPost Physics **14** (2023), doi:[10.21468/SciPostPhys.14.2.023](https://doi.org/10.21468/SciPostPhys.14.2.023).
- [58] S. D. Pace and X. G. Wen, *Emergent higher-symmetry protected topological orders in the confined phase of $U(1)$ gauge theory*, Physical Review B **107** (2023), doi:[10.1103/PhysRevB.107.075112](https://doi.org/10.1103/PhysRevB.107.075112).
- [59] B. Moy and E. Fradkin, *Intertwined order of generalized global symmetries*, SciPost

- Physics **18**, 157 (2025), doi:[10.21468/SciPostPhys.18.5.157](https://doi.org/10.21468/SciPostPhys.18.5.157).
- [60] N. Katayama and Y. Tanizaki, *2d Cardy-Rabinovici model with the modified Villain lattice: Exact dualities and symmetries*, doi:[10.48550/arXiv.2505.19412](https://doi.org/10.48550/arXiv.2505.19412) (2025), <https://arxiv.org/abs/2505.19412>.
 - [61] D. Gaiotto, A. Kapustin, Z. Komargodski and N. Seiberg, *Theta, time reversal and temperature*, J. High Energ. Phys. **2017**, 091 (2017), doi:[10.1007/JHEP05\(2017\)091](https://doi.org/10.1007/JHEP05(2017)091).
 - [62] A. Kapustin and R. Thorngren, *Higher Symmetry and Gapped Phases of Gauge Theories*, pp. 177–202, Springer International Publishing, Cham, ISBN 978-3-319-59939-7, doi:[10.1007/978-3-319-59939-7_5](https://doi.org/10.1007/978-3-319-59939-7_5) (2017).
 - [63] F. Pollmann and A. M. Turner, *Detection of symmetry-protected topological phases in one dimension*, Physical Review B - Condensed Matter and Materials Physics **86** (2012), doi:[10.1103/PhysRevB.86.125441](https://doi.org/10.1103/PhysRevB.86.125441).
 - [64] E. Witten, *Theta dependence in the large N limit of four-dimensional gauge theories*, Physical Review Letters **81** (1998), doi:[10.1103/PhysRevLett.81.2862](https://doi.org/10.1103/PhysRevLett.81.2862).
 - [65] K. Konishi, *Confinement, soft supersymmetry breaking and θ parameter dependence in the Seiberg-Witten model*, Physics Letters, Section B: Nuclear, Elementary Particle and High-Energy Physics **392** (1997), doi:[10.1016/S0370-2693\(96\)01527-4](https://doi.org/10.1016/S0370-2693(96)01527-4).
 - [66] R. Dashen, *Some features of chiral symmetry breaking*, Physical Review D **3** (1971), doi:[10.1103/PhysRevD.3.1879](https://doi.org/10.1103/PhysRevD.3.1879).
 - [67] A. Kapustin and N. Seiberg, *Coupling a QFT to a TQFT and duality*, J. High Energ. Phys. **2014**, 001 (2014), doi:[10.1007/JHEP04\(2014\)001](https://doi.org/10.1007/JHEP04(2014)001).
 - [68] P.-S. Hsin, H. T. Lam and N. Seiberg, *Comments on one-form global symmetries and their gauging in 3d and 4d*, SciPost Physics **6**, 039 (2019), doi:[10.21468/SciPostPhys.6.3.039](https://doi.org/10.21468/SciPostPhys.6.3.039).
 - [69] C. W. Von Keyserlingk and F. J. Burnell, *Walker-Wang models and axion electrodynamics*, Phys. Rev. B **91**, 045134/1 (2015), doi:[10.1103/PhysRevB.91.045134](https://doi.org/10.1103/PhysRevB.91.045134).
 - [70] J. L. Cardy, *Duality and the θ parameter in Abelian lattice models*, Nuclear Physics B **205**, [FS5] 17 (1982), doi:[10.1016/0550-3213\(82\)90464-3](https://doi.org/10.1016/0550-3213(82)90464-3).
 - [71] A. Kapustin, R. Thorngren, A. Turzillo and Z. Wang, *Fermionic symmetry protected topological phases and cobordisms*, Journal of High Energy Physics **2015** (2015), doi:[10.1007/JHEP12\(2015\)052](https://doi.org/10.1007/JHEP12(2015)052).
 - [72] S. L. Adler, *Axial-vector vertex in spinor electrodynamics*, Physical Review **177** (1969), doi:[10.1103/PhysRev.177.2426](https://doi.org/10.1103/PhysRev.177.2426).
 - [73] J. S. Bell and R. Jackiw, *A PCAC puzzle: $\pi^0 \rightarrow \gamma\gamma$ in the σ -model*, Il Nuovo Cimento A **60** (1969), doi:[10.1007/bf02823296](https://doi.org/10.1007/bf02823296).

- [74] J. M. Luttinger, *Theory of thermal transport coefficients*, Physical Review **135** (1964), doi:[10.1103/PhysRev.135.A1505](https://doi.org/10.1103/PhysRev.135.A1505).
- [75] G. E. Volovik, *The gravitational-topological Chern-Simons term in a film of superfluid $^3\text{He-A}$* (1990).
- [76] N. Read and D. Green, *Paired states of fermions in two dimensions with breaking of parity and time-reversal symmetries and the fractional quantum Hall effect*, Physical Review B - Condensed Matter and Materials Physics **61** (2000), doi:[10.1103/PhysRevB.61.10267](https://doi.org/10.1103/PhysRevB.61.10267).
- [77] S. Ryu, J. E. Moore and A. W. Ludwig, *Electromagnetic and gravitational responses and anomalies in topological insulators and superconductors*, Physical Review B - Condensed Matter and Materials Physics **85** (2012), doi:[10.1103/PhysRevB.85.045104](https://doi.org/10.1103/PhysRevB.85.045104).
- [78] D. J. Gross and F. Wilczek, *Asymptotically free gauge theories. I*, Physical Review D **8** (1973), doi:[10.1103/PhysRevD.8.3633](https://doi.org/10.1103/PhysRevD.8.3633).
- [79] H. D. Politzer, *Reliable perturbative results for strong interactions?*, Physical Review Letters **30** (1973), doi:[10.1103/PhysRevLett.30.1346](https://doi.org/10.1103/PhysRevLett.30.1346).
- [80] J. L. Cardy, *Is there a c-theorem in four dimensions?*, Physics Letters B **215** (1988), doi:[10.1016/0370-2693\(88\)90054-8](https://doi.org/10.1016/0370-2693(88)90054-8).
- [81] Z. Komargodski and A. Schwimmer, *On renormalization group flows in four dimensions*, Journal of High Energy Physics **2011** (2011), doi:[10.1007/JHEP12\(2011\)099](https://doi.org/10.1007/JHEP12(2011)099).
- [82] M. Shifman, *Remarks on adjoint QCD with k flavors, $k \geq 2$* , Modern Physics Letters A **28** (2013), doi:[10.1142/S0217732313501794](https://doi.org/10.1142/S0217732313501794).
- [83] A. Athenodorou, E. Bennett, G. Bergner and B. Lucini, *Infrared regime of $SU(2)$ with one adjoint Dirac flavor*, Physical Review D - Particles, Fields, Gravitation and Cosmology **91** (2015), doi:[10.1103/PhysRevD.91.114508](https://doi.org/10.1103/PhysRevD.91.114508).
- [84] C. Córdova and T. T. Dumitrescu, *Candidate phases for $SU(2)$ adjoint QCD_4 with two flavors from $\mathcal{N} = 2$ supersymmetric Yang-Mills theory*, SciPost Physics **16**, 139 (2024), doi:[10.21468/SciPostPhys.16.5.139](https://doi.org/10.21468/SciPostPhys.16.5.139).
- [85] C. Córdova and K. Ohmori, *Anomaly obstructions to symmetry preserving gapped phases*, doi:[10.48550/arXiv.1910.04962](https://doi.org/10.48550/arXiv.1910.04962) (2020), <https://arxiv.org/abs/1910.04962>.
- [86] D. D. Dietrich and F. Sannino, *Conformal window of $SU(N)$ gauge theories with fermions in higher dimensional representations*, Physical Review D - Particles, Fields, Gravitation and Cosmology **75** (2007), doi:[10.1103/PhysRevD.75.085018](https://doi.org/10.1103/PhysRevD.75.085018).
- [87] T. A. Ryttov and R. Shrock, *Higher-loop corrections to the infrared evolution of a gauge theory with fermions*, Physical Review D - Particles, Fields, Gravitation and Cosmology **83** (2011), doi:[10.1103/PhysRevD.83.056011](https://doi.org/10.1103/PhysRevD.83.056011).

- [88] T. Degrand, Y. Shamir and B. Svetitsky, *Near the sill of the conformal window: Gauge theories with fermions in two-index representations*, Physical Review D - Particles, Fields, Gravitation and Cosmology **88** (2013), doi:[10.1103/PhysRevD.88.054505](https://doi.org/10.1103/PhysRevD.88.054505).
- [89] R. Shrock, *Higher-loop structural properties of the β function in asymptotically free vectorial gauge theories*, Physical Review D - Particles, Fields, Gravitation and Cosmology **87** (2013), doi:[10.1103/PhysRevD.87.105005](https://doi.org/10.1103/PhysRevD.87.105005).
- [90] J. Rantaharju, T. Rantalaiho, K. Rummukainen and K. Tuominen, *Running coupling in $SU(2)$ gauge theory with two adjoint fermions*, Physical Review D **93** (2016), doi:[10.1103/PhysRevD.93.094509](https://doi.org/10.1103/PhysRevD.93.094509).
- [91] G. Bergner, P. Giudice, G. Münster, I. Montvay and S. Piemonte, *Spectrum and mass anomalous dimension of $SU(2)$ adjoint QCD with two Dirac flavors*, Physical Review D **96** (2017), doi:[10.1103/PhysRevD.96.034504](https://doi.org/10.1103/PhysRevD.96.034504).
- [92] G. Bergner and S. Piemonte, *Running coupling from gluon and ghost propagators in the Landau gauge: Yang-Mills theories with adjoint fermions*, Physical Review D **97** (2018), doi:[10.1103/PhysRevD.97.074510](https://doi.org/10.1103/PhysRevD.97.074510).
- [93] G. Bergner, G. Münster and S. Piemonte, *Exploring gauge theories with adjoint matter on the lattice*, Universe **8** (2022), doi:[10.3390/universe8120617](https://doi.org/10.3390/universe8120617).
- [94] E. Witten, *Constraints on supersymmetry breaking*, Nuclear Physics, Section B **202** (1982), doi:[10.1016/0550-3213\(82\)90071-2](https://doi.org/10.1016/0550-3213(82)90071-2).
- [95] K. Intriligator and N. Seiberg, *Lectures on supersymmetric gauge theories and electric-magnetic duality*, Nuclear Physics B - Proceedings Supplements **45** (1996), doi:[10.1016/0920-5632\(95\)00626-5](https://doi.org/10.1016/0920-5632(95)00626-5).
- [96] N. Seiberg and E. Witten, *Electric-magnetic duality, monopole condensation, and confinement in $N=2$ supersymmetric Yang-Mills theory*, Nuclear Physics, Section B **426** (1994), doi:[10.1016/0550-3213\(94\)90124-4](https://doi.org/10.1016/0550-3213(94)90124-4).
- [97] P. C. Argyres and A. E. Faraggi, *Vacuum structure and spectrum of $N=2$ supersymmetric $SU(n)$ gauge theory*, Physical Review Letters **74**, 3931 (1995), doi:[10.1103/PhysRevLett.74.3931](https://doi.org/10.1103/PhysRevLett.74.3931).
- [98] A. Klemm, W. Lerche, S. Yankielowicz and S. Theisen, *Simple singularities and $N = 2$ supersymmetric Yang-Mills theory*, Physics Letters B **344**, 169 (1995), doi:[10.1016/0370-2693\(94\)01516-F](https://doi.org/10.1016/0370-2693(94)01516-F).
- [99] Y. Choi, H. T. Lam and S.-H. Shao, *Noninvertible global symmetries in the standard model*, Physical Review Letters **129** (2022), doi:[10.1103/PhysRevLett.129.161601](https://doi.org/10.1103/PhysRevLett.129.161601).
- [100] C. Córdova and K. Ohmori, *Noninvertible chiral symmetry and exponential hierarchies*, Physical Review X **13** (2023), doi:[10.1103/PhysRevX.13.011034](https://doi.org/10.1103/PhysRevX.13.011034).

- [101] K. Walker and Z. Wang, *(3 + 1)-TQFTs and topological insulators*, Front. Phys. **7**, 150 (2012), doi:[10.1007/s11467-011-0194-z](https://doi.org/10.1007/s11467-011-0194-z).
- [102] C. W. V. Keyserlingk, F. J. Burnell and S. H. Simon, *Three-dimensional topological lattice models with surface anyons*, Phys. Rev. B **87**, 045107/1 (2013), doi:[10.1103/PhysRevB.87.045107](https://doi.org/10.1103/PhysRevB.87.045107).
- [103] J. Rantaharju, T. Rindlisbacher, K. Rummukainen, A. Salami and K. Tuominen, *Spectrum of $SU(2)$ gauge theory at large number of flavors*, Physical Review D **104** (2021), doi:[10.1103/PhysRevD.104.114504](https://doi.org/10.1103/PhysRevD.104.114504).
- [104] T. Rindlisbacher, K. Rummukainen, A. Salami and K. Tuominen, *Nonperturbative decoupling of massive fermions*, Physical Review Letters **129** (2022), doi:[10.1103/PhysRevLett.129.131601](https://doi.org/10.1103/PhysRevLett.129.131601).
- [105] J. Kogut and L. Susskind, *Hamiltonian formulation of Wilson's lattice gauge theories*, Physical Review D **11** (1975), doi:[10.1103/PhysRevD.11.395](https://doi.org/10.1103/PhysRevD.11.395).
- [106] J. B. Kogut, *An introduction to lattice gauge theory and spin systems*, Rev. Mod. Phys. **51**, 659 (1979), doi:[10.1103/RevModPhys.51.659](https://doi.org/10.1103/RevModPhys.51.659).
- [107] R. Savit, *Duality in field theory and statistical systems*, Rev. Mod. Phys. **52**, 453 (1980), doi:[10.1103/RevModPhys.52.453](https://doi.org/10.1103/RevModPhys.52.453).
- [108] E. Fradkin, *Quantum Field Theory: An Integrated Approach*, Princeton University Press, Princeton, N. J., ISBN 9780691149080 (2021).
- [109] I. G. Halliday and A. Schwimmer, *The phase structure of $SU(N)/Z(N)$ lattice gauge theories*, Physics Letters B **101** (1981), doi:[10.1016/0370-2693\(81\)90055-1](https://doi.org/10.1016/0370-2693(81)90055-1).
- [110] I. G. Halliday and A. Schwimmer, *$Z(2)$ monopoles in lattice gauge theories*, Physics Letters B **102** (1981), doi:[10.1016/0370-2693\(81\)90630-4](https://doi.org/10.1016/0370-2693(81)90630-4).
- [111] A. P. Schnyder, S. Ryu, A. Furusaki and A. W. W. Ludwig, *Classification of topological insulators and superconductors in three spatial dimensions*, Phys. Rev. B **78**, 195125 (2008), doi:[10.1103/PhysRevB.78.195125](https://doi.org/10.1103/PhysRevB.78.195125).
- [112] A. Kitaev, *Periodic table for topological insulators and superconductors*, In M. Feiguin, ed., *Advances in Theoretical Physics: Landau Memorial Conference*, vol. 1134, p. 22. American Institute of Physics, AIP Conference Proceedings, College Park, Maryland, doi:[10.1063/1.3149495](https://doi.org/10.1063/1.3149495) (2009).
- [113] S. Ryu, A. P. Schnyder, A. Furusaki and A. W. W. Ludwig, *Topological insulators and superconductors: tenfold way and dimensional hierarchy*, New Journal of Physics **12**, 065010 (2010), doi:[10.1088/1367-2630/12/6/065010](https://doi.org/10.1088/1367-2630/12/6/065010).
- [114] L. Fidkowski, X. Chen and A. Vishwanath, *Non-Abelian topological order on the surface of a 3d topological superconductor from an exactly solved model*, Phys. Rev. X

- 3** (2014), doi:[10.1103/PhysRevX.3.041016](https://doi.org/10.1103/PhysRevX.3.041016).
- [115] C. Wang and T. Senthil, *Interacting fermionic topological insulators/superconductors in three dimensions*, Physical Review B - Condensed Matter and Materials Physics **89** (2014), doi:[10.1103/PhysRevB.89.195124](https://doi.org/10.1103/PhysRevB.89.195124).
 - [116] M. A. Metlitski, L. Fidkowski, X. Chen and A. Vishwanath, *Interaction effects on 3D topological superconductors: surface topological order from vortex condensation, the 16 fold way and fermionic Kramers doublets*, doi:[10.48550/arXiv.1406.3032](https://doi.org/10.48550/arXiv.1406.3032) (2014), <https://arxiv.org/abs/1406.3032>.
 - [117] L. Kong, *Anyon condensation and tensor categories*, Nuclear Physics B **886** (2014), doi:[10.1016/j.nuclphysb.2014.07.003](https://doi.org/10.1016/j.nuclphysb.2014.07.003).
 - [118] L. Kong and X.-G. Wen, *Braided fusion categories, gravitational anomalies, and the mathematical framework for topological orders in any dimensions*, doi:[10.48550/arXiv.1405.5858](https://doi.org/10.48550/arXiv.1405.5858) (2014), <https://arxiv.org/abs/1405.5858>.
 - [119] D. V. Else and C. Nayak, *Cheshire charge in (3+1)-dimensional topological phases*, Physical Review B **96** (2017), doi:[10.1103/PhysRevB.96.045136](https://doi.org/10.1103/PhysRevB.96.045136).
 - [120] D. Gaiotto and T. Johnson-Freyd, *Generalized symmetries phase transitions with local quantum fields*, doi:[10.48550/arXiv.1905.09566](https://doi.org/10.48550/arXiv.1905.09566) (2019), <https://arxiv.org/abs/1905.09566>.
 - [121] K. Roumpedakis, S. Seifnashri and S.-H. Shao, *Higher gauging and non-invertible condensation defects*, Communications in Mathematical Physics **401** (2023), doi:[10.1007/s00220-023-04706-9](https://doi.org/10.1007/s00220-023-04706-9).
 - [122] C. M. Chang, Y. H. Lin, S.-H. Shao, Y. Wang and X. Yin, *Topological defect lines and renormalization group flows in two dimensions*, Journal of High Energy Physics **2019** (2019), doi:[10.1007/JHEP01\(2019\)026](https://doi.org/10.1007/JHEP01(2019)026).
 - [123] Y. Choi, C. Córdova, P.-S. Hsin, H. T. Lam and S.-H. Shao, *Noninvertible duality defects in 3 + 1 dimensions*, Phys. Rev. D **105**, 125016 (2022), doi:[10.1103/PhysRevD.105.125016](https://doi.org/10.1103/PhysRevD.105.125016).
 - [124] Y. Choi, C. Córdova, P. S. Hsin, H. T. Lam and S. H. Shao, *Non-invertible condensation, duality, and triality defects in 3+1 dimensions*, Communications in Mathematical Physics **402** (2023), doi:[10.1007/s00220-023-04727-4](https://doi.org/10.1007/s00220-023-04727-4).
 - [125] A. Apte, C. Córdova and H. T. Lam, *Obstructions to gapped phases from noninvertible symmetries*, Physical Review B **108** (2023), doi:[10.1103/PhysRevB.108.045134](https://doi.org/10.1103/PhysRevB.108.045134).
 - [126] J. Kaidi, E. Nardoni, G. Zafrir and Y. Zheng, *Symmetry TFTs and anomalies of non-invertible symmetries*, Journal of High Energy Physics **2023** (2023), doi:[10.1007/JHEP10\(2023\)053](https://doi.org/10.1007/JHEP10(2023)053).

- [127] A. Antinucci, F. Benini, C. Copetti, G. Galati and G. Rizi, *Anomalies of non-invertible self-duality symmetries: fractionalization and gauging*, doi:[10.48550/arXiv.2308.11707](https://arxiv.org/abs/2308.11707) (2023), <https://arxiv.org/abs/2308.11707>.
- [128] C. Córdova, P.-S. Hsin and C. Zhang, *Anomalies of non-invertible symmetries in $(3+1)d$* , SciPost Physics **17**, 131 (2024), doi:[10.21468/SciPostPhys.17.5.131](https://arxiv.org/abs/2308.11707).
- [129] P.-S. Hsin, R. Kobayashi and C. Zhang, *Fractionalization of coset non-invertible symmetry and exotic Hall conductance*, SciPost Physics **17**, 095 (2024), doi:[10.21468/SciPostPhys.17.3.095](https://arxiv.org/abs/2308.11707).
- [130] P.-S. Hsin, *Generalized symmetries phase transitions with local quantum fields*, doi:[10.48550/arXiv.2506.07688](https://arxiv.org/abs/2506.07688) (2025), <https://arxiv.org/abs/2506.07688>.
- [131] M. Stone, *Gamma matrices, Majorana fermions, and discrete symmetries in Minkowski and Euclidean signature*, Journal of Physics A: Mathematical and Theoretical **55** (2022), doi:[10.1088/1751-8121/ac61b7](https://arxiv.org/abs/2506.07688).